

LOWEST WEIGHT MODULES OF $\mathrm{Sp}_4(\mathbb{R})$ AND NEARLY HOLOMORPHIC SIEGEL MODULAR FORMS

AMEYA PITALE, ABHISHEK SAHA, AND RALF SCHMIDT

ABSTRACT. We undertake a detailed study of the lowest weight modules for the Hermitian symmetric pair (G, K) , where $G = \mathrm{Sp}_4(\mathbb{R})$ and K is its maximal compact subgroup. In particular, we determine K -types and composition series, and write down explicit differential operators that navigate all the highest weight vectors of such a module starting from the unique lowest-weight vector. By rewriting these operators in classical language, we show that the automorphic forms on G that correspond to the highest weight vectors are exactly those that arise from *nearly holomorphic* vector-valued Siegel modular forms of degree 2.

Further, by explicating the algebraic structure of the relevant space of n -finite automorphic forms, we are able to prove a *structure theorem* for the space of nearly holomorphic vector-valued Siegel modular forms of (arbitrary) weight $\det^\ell \mathrm{sym}^m$ with respect to an arbitrary congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$. We show that the *cuspidal* part of this space is the *direct sum* of subspaces obtained by applying explicit differential operators to *holomorphic* vector-valued cusp forms of weight $\det^{\ell'} \mathrm{sym}^{m'}$ with (ℓ', m') varying over a certain set. The structure theorem for the space of *all modular forms* is similar, except that we may now have an additional component coming from certain nearly holomorphic forms of weight $\det^3 \mathrm{sym}^{m'}$ that cannot be obtained from holomorphic forms.

As an application of our structure theorem, we prove several arithmetic results concerning nearly holomorphic modular forms that improve previously known results in that direction.

1. INTRODUCTION

1.1. Motivation. In a series of influential works [35, 36, 38, 39], Shimura defined the notion of a *nearly holomorphic function* on a Kähler manifold $\mathbb{K}$ and proved various properties of such functions. Roughly speaking, a nearly holomorphic function on such a manifold is a polynomial of some functions $r_1, \dots, r_m$ on $\mathbb{K}$ (determined by the Kähler structure), over the ring of all holomorphic functions. For example, if $\mathbb{K} = \mathbb{H}_n$, the symmetric space for the group $\mathrm{Sp}_{2n}(\mathbb{R})$, then r_i are the entries of $\mathrm{Im}(Z)^{-1}$. When there is a notion of holomorphic modular forms on $\mathbb{K}$, one can define nearly holomorphic (scalar or vector-valued) modular forms by replacing holomorphy by near-holomorphy in the definition of modular forms.

The prototype of a nearly holomorphic modular form in the simplest case when $\mathbb{K}$ equals the complex upper-half plane $\mathbb{H}$ is provided by the function

$$f(z) := \left(\sum_{(c,d) \neq (0,0)} (cz + d)^{-k} |cz + d|^{-2s} \right)_{s=0}. \quad (1)$$

Here k is a positive even integer. The function f transforms like a modular form of weight k with respect to $\mathrm{SL}_2(\mathbb{Z})$. If $k > 2$, the function is holomorphic, but the case $k = 2$ involves a non-holomorphic term of the form $\frac{c}{y}$, where c is a constant.

A.S. is partially supported by EPSRC grant EP/L025515/1. A.P. and R.S. are supported by NSF grant DMS-1100541.

More generally, special values of Eisenstein series¹, and their restrictions to lower-dimensional manifolds, provide natural examples of nearly holomorphic modular forms. On the other hand, such restrictions of Eisenstein series appear in the theory of L -functions via their presence in integrals of Rankin-Selberg type. Thus, the arithmetic theory of nearly holomorphic forms is closely related to the arithmetic theory of L -functions. The theory was developed by Shimura in substantial detail and was exploited by him and other authors to prove algebraicity and Galois-equivariance of critical values of various L -functions. We refer the reader to the papers [2, 6, 3, 32, 37, 39] for some examples. The theory of nearly holomorphic modular forms and the differential operators related to them has also been very fruitful in the study of p -adic measures related to modular L -functions [5, 10, 28] and in the derivation of various arithmetic identities [11, 24].

From now on, we restrict ourselves to the symplectic case, and we assume further that the base field is $\mathbb{Q}$. The relevant manifold $\mathbb{K}$ is then the degree n Siegel upper half space $\mathbb{H}_n$ consisting of symmetric n by n matrices $Z = X + iY$ with $Y > 0$. For each non-negative integer p , we let $N^p(\mathbb{H}_n)$ denote the space of all polynomials of degree $\leq p$ in the entries of Y^{-1} with holomorphic functions on $\mathbb{H}_n$ as coefficients. The space $N(\mathbb{H}_n) = \bigcup_{p \geq 0} N^p(\mathbb{H}_n)$ is the space of nearly holomorphic functions on $\mathbb{H}_n$. Note that $N^0(\mathbb{H}_n)$ is the space of holomorphic functions on $\mathbb{H}_n$.

Given any congruence subgroup Γ of $\mathrm{Sp}_{2n}(\mathbb{Q})$ and any irreducible finite-dimensional rational representation (η, V) of $\mathrm{GL}_n(\mathbb{C})$, we let $N_\eta^p(\Gamma)$ denote the space of functions $F : \mathbb{H}_n \rightarrow V$ such that

- (1) $F \in N^p(\mathbb{H}_n)$,
- (2) $F(\gamma Z) = \eta(CZ + D)(F(Z))$ for all $\gamma = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \Gamma$.
- (3) F satisfies the “no poles at cusps” condition. This condition states that the Fourier expansion of F at any cusp is supported on the positive semi-definite matrices.²

The set $N_\eta^p(\Gamma)$ (which is clearly a complex vector-space) is known as the space of nearly holomorphic vector-valued modular forms of weight η and nearly holomorphic degree p for Γ . In the special case $(\eta, V) = (\det^k, \mathbb{C})$, we denote the space $N_\eta^p(\Gamma)$ by $N_k^p(\Gamma)$. We let $N_\eta^p(\Gamma)^\circ \subset N_\eta^p(\Gamma)$ denote the subspace of cusp forms (the cusp forms can be defined in the usual way via a vanishing condition at all cusps for degenerate Fourier coefficients). We also denote $M_\eta(\Gamma) = N_\eta^0(\Gamma)$, $S_\eta(\Gamma) = N_\eta^0(\Gamma)^\circ$, $N_\eta(\Gamma) = \bigcup_{p \geq 0} N_\eta^p(\Gamma)$ and $N_\eta(\Gamma)^\circ = \bigcup_{p \geq 0} N_\eta^p(\Gamma)^\circ$.

In the case $n = 1$, Shimura proved [36, Thm. 5.2] a complete *structure theorem* that describes the set $N_k^p(\Gamma)$ precisely for every weight k and every congruence subgroup Γ of $\mathrm{SL}_2(\mathbb{Z})$. For simplicity, write $N_k(\Gamma) = \bigcup_{p \geq 0} N_k^p(\Gamma)$. Let R denote the classical *weight-raising operator* on $\bigcup_k N_k(\Gamma)$ that acts on elements of $N_k(\Gamma)$ via the formula $\frac{k}{y} + 2i\frac{\partial}{\partial z}$. It can be easily checked that R takes $N_k^p(\Gamma)$ to $N_{k+2}^{p+1}(\Gamma)$. Then a slightly simplified version of the structure theorem of Shimura says that $N_0(\Gamma) = \mathbb{C}$, and for $k > 0$,

$$N_k(\Gamma) = R^{\frac{k-2}{2}}(\mathbb{C}E_2) \oplus \bigoplus_{\ell \geq 1} R^{\frac{k-\ell}{2}}(M_\ell(\Gamma)), \quad N_k(\Gamma)^\circ = \bigoplus_{\ell \geq 1} R^{\frac{k-\ell}{2}}(S_\ell(\Gamma)), \quad (2)$$

where we understand $R^v = 0$ if $v \notin \mathbb{Z}_{\geq 0}$, and where E_2 denotes the weight 2 nearly holomorphic Eisenstein series obtained by putting $k = 2$ in (1). For the refined structure theorem taking into

¹The typical situation is as follows. Let $E(z, s)$ be an appropriately normalized Eisenstein series on some Hermitian symmetric space that converges absolutely for $\mathrm{Re}(s) > s_0$ and transforms like a modular form in the variable z . Suppose that $E(z, k)$ is holomorphic for some $k \in \mathbb{Z}$. Then $E(z, s')$ is typically a nearly holomorphic modular form for all s' such that $s_0 < s' \leq k$, $s' \in \mathbb{Z}$; see [36, Thm. 4.2].

²If $n > 1$, the “no poles at cusps” condition follows automatically from the previous two conditions by the Koecher principle.

account the nearly holomorphic degree, we refer the reader to [29], where we reprove Shimura's results using representation-theoretic methods.

Shimura used his structure theorem to prove that the cuspidal holomorphic projection map from $N_k(\Gamma)$ to $S_k(\Gamma)$ has nice $\text{Aut}(\mathbb{C})$ -equivariance properties, and he even extended these results to the half-integral case [37, Prop. 9.4]. As an application, Shimura obtained many arithmetic results for ratios of Petersson norms and critical values of L -functions.

In the case $n > 1$, Shimura showed [39, Prop. 14.2] that if the lowest weight of η is “large enough” compared to the nearly holomorphic degree, then the space $N_\eta^p(\Gamma)$ is *spanned* by the functions obtained by letting differential operators act on various spaces $M_{\eta'}(\Gamma)$. Using this, he was able to construct an analogue of the projection map under some additional assumptions. But the arithmetic results thus obtained are weaker than those for $n = 1$.

There is another aspect in which the state of our understanding of nearly holomorphic modular forms is unsatisfactory, namely that the precise meaning of these objects in the modern language of automorphic forms on reductive groups, à la Langlands, has not been worked out. Most work done so far for nearly holomorphic forms has been in the classical language. There has been some work in interpreting these forms from the point of view of vector bundles and sheaf theory, see [17, 18, 27, 42]. There has also been some work on interpreting the differential operators involved in the language of Lie algebra elements, but this has been carried out explicitly only in the case $n = 1$ [11, 15]. A detailed investigation from the point of view of automorphic representations has so far been lacking in the case $n > 1$.

The objective of this paper is to address the issues discussed above in the case $n = 2$, i.e., when Γ is a congruence subgroup of $\text{Sp}_4(\mathbb{Q})$. The relevant η 's in this case are the representations $\det^\ell \text{sym}^m$ for integers ℓ, m with $m \geq 0$, and it is natural to use $N_{\ell,m}(\Gamma)$ to denote the corresponding space of nearly holomorphic forms. We achieve the following goals.

- We prove a structure theorem for $N_{\ell,m}(\Gamma)$ that is (almost) as complete and explicit as the $n = 1$ case. As a consequence, we are able to prove arithmetic results for this space (as well as for certain associated “isotypic projection” maps, and ratios of Petersson inner products) that improve previously known results in this direction.
- We make a detailed study of the spaces $N_{\ell,m}(\Gamma)$ in the language of $(\mathfrak{g}, K)$ -modules and automorphic forms for the group $\text{Sp}_4(\mathbb{R})$. We analyze the K -types, weight vectors and composition series, write down *completely explicit* operators from the classical as well as Lie-theoretic points of view, explain exactly how nearly holomorphic forms arise in the Langlands framework, and describe the automorphic representations attached to them.

In the rest of this introduction we explain these results and ideas in more detail.

1.2. The structure theorem in degree 2. Let Γ be a congruence subgroup of $\text{Sp}_4(\mathbb{Q})$. In order to prove a structure theorem for $N_{\ell,m}(\Gamma)$, it is necessary to have suitable differential operators that generalize the weight-raising operator considered above. In fact, it turns out that one needs four operators, which we term X_+ , U , E_+ and D_+ .

Each of these four operators acts on the set $\bigcup_{\ell,m} N_{\ell,m}(\Gamma)$. They take the subspace $N_{\ell,m}^p(\Gamma)$ to the subspace $N_{\ell_1,m_1}^{p_1}(\Gamma)$, where the integers ℓ_1, m_1, p_1 are given by the following table.

operator	ℓ_1	m_1	p_1
X_+	ℓ	$m+2$	$p+1$
U	$\ell+2$	$m-2$	$p+1$
E_+	$\ell+1$	m	$p+1$
D_+	$\ell+2$	m	$p+2$

(3)

Note that, in the above list, E_+ is the only operator that changes the parity of ℓ . For the explicit formulas for the above differential operators, see (85) – (92) of this paper. We note that the operator D_+ was originally studied by Maass in his book [25] in the case of scalar-valued forms. The operator X_+ (for both scalar and vector-valued forms) was already defined in [4], where it was called $\delta_{\ell+m}$. Also, the operator U was considered by Satoh [34] (who called it D) in the very special case $m = 2$. To the best of our knowledge, explicit formulas for the operators (except in the cases mentioned above) had not been written out before this work.

More generally, if $\mathcal{X}_+$ denotes the free monoid consisting of finite strings of the above four operators, then each element $X \in \mathcal{X}_+$ takes $N_{\ell,m}^p(\Gamma)$ to $N_{\ell_1,m_1}^{p_1}(\Gamma)$ for some integers ℓ_1, m_1, p_1 (uniquely determined by ℓ, m, p and X) that can be easily calculated using the above table. In particular, the non-negative integer $v = p_1 - p$ depends only on X ; we call it the degree of X . For example, the operator $D_+^r U^s \in \mathcal{X}_+$ takes the space $N_{\ell,2s}^p(\Gamma)$ to $N_{\ell+2s+2r,0}^{2r+s+p}(\Gamma)$ and has degree $2r + s$.

Let $X, \ell, m, \ell_1, m_1, v$ be as above. We show that X has the following properties.

- (1) (Lemma 4.1) For all $\gamma \in \mathrm{GSp}_4(\mathbb{R})^+$, we have

$$(XF)|_{\ell_1,m_1}\gamma = X(F|_{\ell,m}\gamma).$$

- (2) (Lemma 4.15) X takes $N_{\ell,m}(\Gamma)^\circ$ to $N_{\ell_1,m_1}(\Gamma)^\circ$ and takes the orthogonal complement of $N_{\ell,m}(\Gamma)^\circ$ to the orthogonal complement of $N_{\ell_1,m_1}(\Gamma)^\circ$.

- (3) (Proposition 4.17) There exists a constant $c_{\ell,m,X}$ (depending only on ℓ, m, X) such that for all F, G in $S_{\ell,m}(\Gamma)$,

$$\langle XF, XG \rangle = c_{\ell,m,X} \langle F, G \rangle.$$

- (4) (Proposition 5.6) For all $\sigma \in \mathrm{Aut}(\mathbb{C})$, we have

$$\sigma((2\pi)^{-v} XF) = (2\pi)^{-v} X(\sigma F).$$

We now state a coarse version of our structure theorem for cusp forms.

Theorem 1.1 (Structure theorem for cusp forms, coarse version). *Let ℓ, m be integers with $m \geq 0$. For each pair of integers ℓ', m' , there is a (possibly empty³) finite subset $\mathcal{X}_{\ell',m'}^{\ell,m}$ of $\mathcal{X}_+$ such that the following hold.*

- (1) *Each element $X \in \mathcal{X}_{\ell',m'}^{\ell,m}$ acts injectively on $M_{\ell',m'}(\Gamma)$ and takes this space to $N_{\ell,m}(\Gamma)$.*
(2) *We have an orthogonal direct sum decomposition*

$$N_{\ell,m}(\Gamma)^\circ = \bigoplus_{\ell'=1}^{\ell} \bigoplus_{m'=0}^{\ell+m-\ell'} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(S_{\ell',m'}(\Gamma)).$$

³Indeed, $\mathcal{X}_{\ell',m'}^{\ell,m}$ is empty unless $m' \geq 0, 0 \leq \ell' \leq \ell, 0 \leq \ell' + m' \leq \ell + m$, and some additional parity conditions are satisfied. Moreover, $\mathcal{X}_{\ell',m'}^{\ell,m}$ is always the singleton set consisting of the identity map whenever ℓ, m are non-negative integers.

For the refined version of this result, see Theorem 4.8, which contains an exact description of the sets $\mathcal{X}_{\ell',m'}^{\ell,m}$. We also formulate a version of this theorem for scalar valued cusp forms (Corollary 4.10), as well as deduce a result for forms of a fixed nearly holomorphic degree (Corollary 4.11).

Next, we turn to a structure theorem for the whole space, including the non-cusp forms. This situation turns out to be more complicated. Indeed, we need to now also include certain non-holomorphic objects among our building blocks. This is to be expected from the $n = 1$ situation, where the nearly holomorphic Eisenstein series E_2 appears in the direct sum decomposition (2).

For each $m \geq 0$, we define a certain subspace $M_{3,m}^*(\Gamma)$ of $N_{3,m}^1(\Gamma)$ consisting of forms that are annihilated by two differential operators that we call L and E_- (see Section 3.4 for their explicit formulas). From the definition, it is immediate that $M_{3,m}^*(\Gamma)$ contains $M_{3,m}(\Gamma)$. However, it may potentially contain more objects. These extra elements in $M_{3,m}^*(\Gamma)$ cannot exist if $M_{1,m}(\Gamma) = \{0\}$ (which is the case, for instance, when $\Gamma = \mathrm{Sp}_4(\mathbb{Z})$); moreover, if they exist, they cannot be cuspidal, must lie inside $N_{3,m}^1(\Gamma)$, and cannot be obtained by applying our differential operators to holomorphic modular forms of any weight. Furthermore, we can prove that the space $M_{3,m}^*(\Gamma)$ is $\mathrm{Aut}(\mathbb{C})$ -invariant.

Now, we may state our general structure theorem as follows.

Theorem 1.2 (Structure theorem for all modular forms, coarse version). *Let ℓ, m be integers with $\ell > 0$ and $m \geq 0$. For each pair of integers ℓ', m' , let $\mathcal{X}_{\ell',m'}^{\ell,m}$ be as in Theorem 1.1. Then we have a direct sum decomposition*

$$N_{\ell,m}(\Gamma) = \bigoplus_{\substack{\ell'=1 \\ \ell' \neq 3}}^{\ell} \bigoplus_{m'=0}^{\ell+m-\ell'} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(M_{\ell',m'}(\Gamma)) \oplus \bigoplus_{m'=0}^{\ell+m-3} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(M_{3,m'}^*(\Gamma)).$$

This decomposition is orthogonal in the sense that forms lying in different constituents, and such that at least one of them is cuspidal, are orthogonal with respect to the Petersson inner product.

For a refined version of this result, see Theorem 4.33. We note that the restriction to $\ell > 0$ is not serious, since the only nearly holomorphic modular forms with $\ell \leq 0$ are the constant functions.

1.3. Lowest weight modules and $\mathfrak{n}$ -finite automorphic forms. We now describe the representation-theoretic results that form the foundation for Theorems 1.1 and 1.2. We hope that they are of independent interest, as they explain nearly holomorphic forms from the point of view of representation theory.

Let $\mathfrak{g}$ be the Lie algebra of $\mathrm{Sp}_4(\mathbb{R})$, and let $\mathfrak{g}_{\mathbb{C}}$ be its complexification. We fix a basis of the root system of $\mathfrak{g}_{\mathbb{C}}$, and let $\mathfrak{n}$ be the space spanned by the non-compact negative roots. It is well known that vector-valued holomorphic modular forms F correspond to (scalar-valued) automorphic forms⁴ Φ on $\mathrm{Sp}_4(\mathbb{R})$ that are annihilated by $\mathfrak{n}$. The $(\mathfrak{g}, K)$ -module $\langle \Phi \rangle$ generated by such a Φ is a lowest weight module, and Φ is a lowest weight vector in this module. In fact, it will follow from our results that $\langle \Phi \rangle$ is always an *irreducible* module (see Proposition 4.28).

We define a vector v in any representation of $\mathfrak{g}_{\mathbb{C}}$ to be $\mathfrak{n}$ -finite, if the space $\mathcal{U}(\mathfrak{n})v$ is finite-dimensional; here $\mathcal{U}(\mathfrak{n})$ is the universal enveloping algebra of $\mathfrak{n}$, which in our case is simply a polynomial ring in three variables. Applying this concept to the space of automorphic forms on $\mathrm{Sp}_4(\mathbb{R})$, we arrive at the notion of $\mathfrak{n}$ -finite automorphic form, which is central to this work. Let $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ be the space of $\mathfrak{n}$ -finite automorphic forms on $\mathrm{Sp}_4(\mathbb{R})$ with respect to a fixed

⁴Here, and elsewhere in this paper, we use the term “automorphic form” in the sense of Borel-Jacquet [9, 1.3]; in particular, our automorphic forms are always *scalar-valued* functions on $\mathrm{Sp}_4(\mathbb{R})$. For the precise correspondence between (nearly holomorphic) vector-valued modular forms for Γ , and automorphic forms on $\mathrm{Sp}_4(\mathbb{R})$ with respect to Γ , see Lemma 3.2 and Proposition 4.5 of this paper.

congruence subgroup Γ . Finiteness results from the classical theory imply that $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ is an *admissible* $(\mathfrak{g}, K)$ -module.

Clearly, the lowest weight module $\langle \Phi \rangle$ considered above is contained in $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$. We will prove the following:

- Every automorphic form in $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ gives rise to a vector-valued nearly holomorphic modular form on $\mathbb{H}_2$. See Lemma 3.2, Proposition 4.27 and the discussion following it.
- Conversely, the automorphic form corresponding to a vector-valued nearly holomorphic modular form on $\mathbb{H}_2$ lies in $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$. See Proposition 4.5.

In other words, the $\mathbf{n}$ -finite automorphic forms correspond *precisely* to nearly holomorphic modular forms. The lowest weight vectors in irreducible submodules of $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ correspond precisely to holomorphic modular forms.

The structure theorems 1.1 and 1.2 are reflections of the fact that, in a lowest weight module appearing in $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$, we can navigate from the lowest weight vector to any given K -type using certain elements X_+ , U , E_+ and D_+ in $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ that correspond to the differential operators given in table (3). See Proposition 2.15 for the precise statement.

To prove statements like Proposition 2.15, we need rather precise information about K -types and multiplicities occurring in lowest weight modules. Such information is in principle available in the literature, but it requires some effort to obtain it from general theorems. It turns out that *category* $\mathcal{O}$ provides a framework well-suited for our purposes. More precisely, we will work in a parabolic version called category $\mathcal{O}^{\mathbf{p}}$, whose objects consist precisely of the finitely generated $(\mathfrak{g}, K)$ -modules in which all vectors are $\mathbf{n}$ -finite. This category thus contains all the lowest weight modules relevant for the study of $\mathbf{n}$ -finite automorphic forms.

Basic building blocks in category $\mathcal{O}^{\mathbf{p}}$ are the parabolic Verma modules $N(\lambda)$ and their unique irreducible quotients $L(\lambda)$; here, λ is an integral weight.⁵ We determine which of the $N(\lambda)$ are irreducible (Proposition 2.5), composition series in each reducible case (Proposition 2.6), and which of the $L(\lambda)$ are square-integrable, tempered, or unitarizable (Proposition 2.2). This is slightly more information than needed for our applications to automorphic forms, but we found it useful to collect all this information in one place.

By general principles, the admissible $(\mathfrak{g}, K)$ -module $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ decomposes into a direct sum of indecomposable objects in category $\mathcal{O}^{\mathbf{p}}$. The subspace of cusp forms $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}^{\circ} \subseteq \mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ decomposes in fact into a direct sum of irreducibles $L(\lambda)$, due to the presence of an inner product. The multiplicities with which each $L(\lambda)$ occurs is given by the dimension of certain spaces of holomorphic modular forms. We can thus determine the complete algebraic structure of the space $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}^{\circ}$ in terms of these dimensions. See Proposition 4.6 for the precise statement, which may be viewed as a precursor to Theorem 1.1.

One cannot expect that the entire space $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ also decomposes into a direct sum of irreducibles. This is already not the case in the degree 1 situation, where the modular form E_2 generates an indecomposable but not irreducible module. Sections 4.5 and 4.6 are devoted to showing that only a very limited class of indecomposable but not irreducible modules can possibly occur in $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$. These modules account for the presence of the spaces $M_{3,m'}^*(\Gamma)$ in Theorem 1.2. The algebraic structure of the entire space $\mathcal{A}(\Gamma)_{\mathbf{n}\text{-fin}}$ in terms of dimensions of spaces of modular forms is given in Proposition 4.27. As in the cuspidal case, this proposition is a precursor to the structure theorem.

1.4. Applications of the structure theorem. The significance of the structure theorem is twofold. On the one hand, it builds up the space of nearly holomorphic forms from holomorphic forms using differential operators. As the differential operators have nice arithmetic properties, this essentially reduces all arithmetic questions about nearly holomorphic forms to the case of holomorphic forms. Since there is considerable algebraic geometry known for

⁵The automorphic forms corresponding to elements in $M_{\ell,m}(\Gamma)$ generate the lowest weight module $L(\ell+m, \ell)$. We note that in previous papers, we have used the notation $\mathcal{E}(\ell+m, \ell)$ instead of $L(\ell+m, \ell)$.

the latter, powerful results can be obtained. For example, in Section 5.2, we show that the “isotypic projection” map from $N_{\ell,m}(\Gamma)$ to $\sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(M_{\ell',m'}(\Gamma))$ (this is commonly called the “holomorphic projection” map when $\ell = \ell'$, $m = m'$) obtained from our structure theorem is $\text{Aut}(\mathbb{C})$ -equivariant (see Propositions 5.10 and 5.17). This is a considerable generalization of results of Shimura. We also prove a result on the arithmeticity of ratios of Petersson inner products (Proposition 5.18) that will be of importance in our subsequent work.

On the other hand, sometimes one prefers to deal with modular forms of scalar weight, rather than vector-valued objects. The structure theorem gives an explicit and canonical way to start with an element of $M_{\ell,m}(\Gamma)$ (with m even and $\ell \geq 2$) and produce a non-zero element of $N_{\ell+m}^{m/2}(\Gamma)$ lying in the same representation. (This does not work if $\ell = 1$.) On a related note, this will also allow one to write down a *scalar-valued* nearly holomorphic lift in cases where previously only holomorphic vector-valued lifts have been considered (e.g., the Yoshida lift of two classical cusp forms f and g both of weight bigger than 2).

Both these points of view will be combined in a forthcoming work where we will prove results in the spirit of Deligne’s conjecture for the standard L -function attached to a holomorphic vector valued cusp form with respect to an arbitrary congruence subgroup of $\text{Sp}_4(\mathbb{Q})$. Such results have so far been proved (in the vector-valued case) only for forms of full level. The main new ingredient of this forthcoming work will be to consider an integral representation consisting only of scalar-valued nearly holomorphic vectors. The results of this paper will be key to doing that.

There are many other potential applications of this work, some of which we plan to pursue elsewhere. For example, one can use our structure theorems to produce exact formulas for the dimensions of spaces of nearly holomorphic modular forms; to the best of our knowledge, no such formulas are currently known in degree 2. One could try to see if our explicit formulas could be used to deal with problems related to congruences or the construction of p -adic measures for vector-valued Siegel modular forms, similar to what was done in the scalar-valued case in [10]. One could apply our results to the study of nearly overconvergent modular forms for congruence subgroups of $\text{Sp}_4(\mathbb{Z})$, following the general framework of [42]. One could also explore applications of our work to arithmetic and combinatorial identities, à la [11].

Finally, it would be interesting to generalize the results of this paper to the case $n > 2$ and possibly to other groups. We hope to come back to this problem in the future.

Acknowledgements. We would like to thank Siegfried Böcherer, Marcela Hanzer, Michael Harris, and Jonathan Kujawa for helpful discussions. We would also like to thank Shuji Hori-naga, who had kindly informed us about a mistake in an earlier version of this paper.

Notation.

- (1) The symbols $\mathbb{Z}$, $\mathbb{Z}_{\geq 0}$, $\mathbb{Q}$, $\mathbb{R}$, $\mathbb{C}$, $\mathbb{Z}_p$ and $\mathbb{Q}_p$ have the usual meanings. The symbol $\mathbb{A}$ denotes the ring of adeles of $\mathbb{Q}$, and $\mathbb{A}^\times$ denotes its group of ideles. We let $\mathfrak{f}$ denote the set of finite places, and $\mathbb{A}_{\mathfrak{f}}$ the subring of $\mathbb{A}$ with trivial archimedean component.
- (2) For any commutative ring R and positive integer n , let $M_n(R)$ denote the ring of $n \times n$ matrices with entries in R , and let $M_n^{\text{sym}}(R)$ denote the subset of symmetric matrices. We let $\text{GL}_n(R)$ denote the group of invertible elements in $M_n(R)$, and we use $R^\times$ to denote $\text{GL}_1(R)$. If $A \in M_n(R)$, we let tA denote its transpose.
- (3) Define $J_n \in M_n(\mathbb{Z})$ by $J_n = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$, where I_n is the n by n identity matrix. Let GSp_4 and Sp_4 be the algebraic groups whose $\mathbb{Q}$ -points are given by

$$\text{GSp}_4(\mathbb{Q}) = \{g \in \text{GL}_4(\mathbb{Q}) \mid {}^t g J_2 g = \mu_2(g) J_2, \mu_2(g) \in \mathbb{Q}^\times\}, \quad (4)$$

$$\text{Sp}_4(\mathbb{Q}) = \{g \in \text{GSp}_4(\mathbb{Q}) \mid \mu_2(g) = 1\}. \quad (5)$$

Let $\text{GSp}_4(\mathbb{R})^+ \subset \text{GSp}_4(\mathbb{R})$ consist of the matrices with $\mu_2(g) > 0$.

- (4) For $\tau = x + iy$, we let

$$\frac{\partial}{\partial \tau} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{\tau}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right)$$

denote the usual Wirtinger derivatives.

- (5) The Siegel upper half space of degree n is defined by

$$\mathbb{H}_n = \{Z \in M_n(\mathbb{C}) \mid Z = {}^t Z, i(\bar{Z} - Z) \text{ is positive definite}\}.$$

For $g = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathrm{GSp}_4(\mathbb{R})^+$, $Z \in \mathbb{H}_2$, define $J(g, Z) = CZ + D$. We let I denote the element $\begin{bmatrix} & 1 \\ i & \end{bmatrix}$ of $\mathbb{H}_2$.

- (6) We let $\mathfrak{g} = \mathfrak{sp}_4(\mathbb{R})$ be the Lie algebra of $\mathrm{Sp}_4(\mathbb{R})$ and $\mathfrak{g}_{\mathbb{C}} = \mathfrak{sp}_4(\mathbb{C})$ the complexified Lie algebra. We let $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ denote the universal enveloping algebra and let $\mathcal{Z}$ be its center. We use the following basis for $\mathfrak{g}_{\mathbb{C}}$.

$$\begin{aligned} Z &= -i \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & Z' &= -i \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}, \\ N_+ &= \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & -i \\ -1 & 0 & -i & 0 \\ 0 & i & 0 & 1 \\ i & 0 & -1 & 0 \end{bmatrix}, & N_- &= \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & i \\ -1 & 0 & i & 0 \\ 0 & -i & 0 & 1 \\ -i & 0 & -1 & 0 \end{bmatrix}, \\ X_+ &= \frac{1}{2} \begin{bmatrix} 1 & 0 & i & 0 \\ 0 & 0 & 0 & 0 \\ i & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & X_- &= \frac{1}{2} \begin{bmatrix} 1 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 \\ -i & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \\ P_{1+} &= \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & i \\ 1 & 0 & i & 0 \\ 0 & i & 0 & -1 \\ i & 0 & -1 & 0 \end{bmatrix}, & P_{1-} &= \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & -i \\ 1 & 0 & -i & 0 \\ 0 & -i & 0 & -1 \\ -i & 0 & -1 & 0 \end{bmatrix}, \\ P_{0+} &= \frac{1}{2} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & i \\ 0 & 0 & 0 & 0 \\ 0 & i & 0 & -1 \end{bmatrix}, & P_{0-} &= \frac{1}{2} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -i \\ 0 & 0 & 0 & 0 \\ 0 & -i & 0 & -1 \end{bmatrix}. \end{aligned}$$

- (7) For all smooth functions $f : \mathrm{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$, $X \in \mathfrak{g}$, define $(Xf)(g) = \frac{d}{dt}|_0 f(\exp(tX))$. This action is extended $\mathbb{C}$ -linearly to $\mathfrak{g}_{\mathbb{C}}$. Further, it is extended to all elements $X \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ in the usual manner.

2. LOWEST WEIGHT REPRESENTATIONS

In this section we study the lowest weight representations of the Hermitian symmetric pair (G, K) , where $G = \mathrm{Sp}_4(\mathbb{R})$ and K is its maximal compact subgroup. We will determine composition series and K -types for each parabolic Verma module. Of course, lowest weight representations have been extensively studied in the literature, in the more general context of semisimple Lie groups. Much of our exposition will consist in making the general theorems explicit in our low-rank case.

2.1. Set-up and basic facts. The subgroup K of $\mathrm{Sp}_4(\mathbb{R})$ consisting of all elements of the form $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$ is a maximal compact subgroup. It is isomorphic to $U(2)$ via the map $\begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mapsto A + iB$.

Let $\mathfrak{g} = \mathfrak{sp}_4(\mathbb{R})$ be the Lie algebra of $\mathrm{Sp}_4(\mathbb{R})$, which we think of as a 10-dimensional space of 4×4 matrices. Let $\mathfrak{k}$ be the Lie algebra of K ; it is a four-dimensional subspace of $\mathfrak{g}$. Let $\mathfrak{g}_{\mathbb{C}}$ (resp. $\mathfrak{k}_{\mathbb{C}}$) be the complexification of $\mathfrak{g}$ (resp. $\mathfrak{k}$). A Cartan subalgebra $\mathfrak{h}_{\mathbb{C}}$ of $\mathfrak{k}_{\mathbb{C}}$ (and of $\mathfrak{g}_{\mathbb{C}}$) is spanned by the two elements Z and Z' . If λ is in the dual space $\mathfrak{h}_{\mathbb{C}}^*$, we identify λ with the element $(\lambda(Z), \lambda(Z'))$ of $\mathbb{C}^2$. The root system of $\mathfrak{g}_{\mathbb{C}}$ is $\Phi = \{(\pm 2, 0), (0, \pm 2), (\pm 1, \pm 1), (\pm 1, \mp 1)\}$. These vectors lie in the subspace $E := \mathbb{R}^2$ of $\mathbb{C}^2$, which we think of as a Euclidean plane. The analytically integral elements of $\mathfrak{h}_{\mathbb{C}}^*$ are those that identify with points of $\mathbb{Z}^2$. These are exactly the points of the weight lattice Λ . The following diagram indicates the weight lattice, as well

as the roots and the elements of the Lie algebra spanning the corresponding root spaces.

Here, $(1, -1)$ and $(-1, 1)$ are the compact roots, with the corresponding root spaces being spanned by N_+ and N_- . We declare the set

$$\Phi^+ = \{(-2, 0), (-1, -1), (0, -2), (1, -1)\}$$

to be a positive system of roots. We define an ordering on Λ by

$$\mu \preceq \lambda \iff \lambda \in \mu + \Upsilon, \quad (7)$$

where Υ is the set of all $\mathbb{Z}_{\geq 0}$ -linear combinations of elements of Φ^+ . Hence, under this ordering, $(0, -2)$ is maximal among the non-compact positive roots.

Let $\mathcal{Z}$ be the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$. A particular element in $\mathcal{Z}$ is the Casimir element

$$\begin{aligned} \Omega_2 = & \frac{1}{2}Z^2 + \frac{1}{2}Z'^2 - \frac{1}{2}(N_+N_- + N_-N_+) + X_+X_- + X_-X_+ \\ & + \frac{1}{2}(P_{1+}P_{1-} + P_{1-}P_{1+}) + P_{0+}P_{0-} + P_{0-}P_{0+}. \end{aligned} \quad (8)$$

Using the commutation relations, an alternative form is

$$\Omega_2 = \frac{1}{2}Z^2 + \frac{1}{2}Z'^2 - Z - 2Z' - N_-N_+ + 2X_+X_- + P_{1+}P_{1-} + 2P_{0+}P_{0-}. \quad (9)$$

The characters of $\mathcal{Z}$ are indexed by elements of $\mathfrak{h}_{\mathbb{C}}^*$ modulo Weyl group action; see Sects. 1.7–1.10 of [20]. Let χ_{λ} be the character of $\mathcal{Z}$ corresponding to $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$. We normalize this correspondence such that χ_{ϱ} is the trivial character (i.e., the central character of the trivial representation of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$); here, $\varrho = (-1, -2)$ is half the sum of the positive roots. Note that Humphrey's χ_{λ} is our $\chi_{\lambda+\varrho}$.

If $\mathfrak{k}_{\mathbb{C}}$ acts on a space V , and $v \in V$ satisfies $Zv = kv$ and $Z'v = \ell v$ for $k, \ell \in \mathbb{C}$, then we say that v has *weight* (k, ℓ) . If the weight lies in E , we indicate it as a point in this Euclidean plane. Let V be a finite-dimensional $\mathfrak{k}_{\mathbb{C}}$ -module. Then this representation of $\mathfrak{k}_{\mathbb{C}}$ can be integrated to a representation of K if and only if all occurring weights are analytically integral. The isomorphism classes of irreducible such $\mathfrak{k}_{\mathbb{C}}$ -modules, or the corresponding irreducible representations of K , are called *K-types*.

Let V be a K -type. A non-zero vector $v \in V$ is called a *highest weight vector* if $N_+v = 0$. Such a vector v is unique up to scalars. Let (k, ℓ) be its weight. Then the weights occurring in V are $(k-j, \ell+j)$ for $j = 0, 1, \dots, k-\ell$. In particular, the dimension of V is $k-\ell+1$. If we associate with each K -type its highest weight, then we obtain a bijection between K -types and analytically integral elements (k, ℓ) with $k \geq \ell$.

Definition 2.1. We let Λ^+ denote the subset of Λ consisting of pairs of integers (k, ℓ) with $k \geq \ell$. If $\lambda \in \Lambda^+$, we denote by ρ_{λ} the corresponding K -type.

Let $\mathfrak{p}_\pm = \langle X_\pm, P_{1\pm}, P_{0\pm} \rangle$. Then $\mathfrak{p}_+$ and $\mathfrak{p}_-$ are commutative subalgebras of $\mathfrak{g}_\mathbb{C}$, and we have $[\mathfrak{k}_\mathbb{C}, \mathfrak{p}_\pm] \subset \mathfrak{p}_\pm$. Let ρ_λ be a K -type. Let $F(\lambda)$ be any model for ρ_λ . We consider $F(\lambda)$ a module for $\mathfrak{k}_\mathbb{C} + \mathfrak{p}_-$ by letting $\mathfrak{p}_-$ act trivially. Let

$$N(\lambda) := \mathcal{U}(\mathfrak{g}_\mathbb{C}) \otimes_{\mathfrak{k}_\mathbb{C} + \mathfrak{p}_-} F(\lambda). \quad (10)$$

Then $N(\lambda)$ is a $\mathfrak{g}_\mathbb{C}$ -module in the obvious way. It also is a $(\mathfrak{g}, K)$ -module, with K -action given by $g.(X \otimes v) = \text{Ad}(g)(X) \otimes \rho_\lambda(g)v$ for $g \in K$, $X \in \mathcal{U}(\mathfrak{g}_\mathbb{C})$ and $v \in F(\lambda)$. The modules $N(\lambda)$ are often called *highest weight modules* in the literature. However, when we think of the K -type ρ_λ as the weight of a modular form, it will be more natural to think of the $N(\lambda)$ as *lowest weight modules*.

As vector spaces, we have

$$N(\lambda) = \mathcal{U}(\mathfrak{p}_+) \otimes_\mathbb{C} F(\lambda). \quad (11)$$

Since $\mathcal{U}(\mathfrak{p}_+)$ is simply a polynomial algebra in X_+, P_{1+}, P_{0+} , it follows that $N(\lambda)$ is spanned by the vectors

$$X_+^\alpha P_{1+}^\beta P_{0+}^\gamma N_-^\delta w_0, \quad \alpha, \beta, \gamma, \delta \geq 0, \delta \leq k - \ell, \quad (12)$$

where $\lambda = (k, \ell)$, and these vectors are linearly independent. Here, w_0 is a highest weight vector in $F(\lambda)$ (identified with the element $1 \otimes w_0$ in the tensor product (10)). Alternatively, $N(\lambda)$ is spanned by the vectors

$$N_-^\delta X_+^\alpha P_{1+}^\beta P_{0+}^\gamma w_0, \quad \alpha, \beta, \gamma, \delta \geq 0, \quad (13)$$

but these are not linearly independent.

It will be convenient to work in a parabolic version of category $\mathcal{O}$; see Sect. 9 of [20]. Let $\mathfrak{n} = \langle X_-, P_{1-}, P_{0-} \rangle$; this is the same as $\mathfrak{p}_-$, but we will use the symbol $\mathfrak{n}$ henceforth. Let M be a $\mathfrak{g}_\mathbb{C}$ -module. We say M lies in category $\mathcal{O}^p$ if it satisfies the following conditions:

- ($\mathcal{O}^p$ 1) M is a finitely generated $\mathcal{U}(\mathfrak{g}_\mathbb{C})$ -module.
- ($\mathcal{O}^p$ 2) M is the direct sum of K -types.
- ($\mathcal{O}^p$ 3) M is locally $\mathfrak{n}$ -finite, meaning: For each $v \in M$ the subspace $\mathcal{U}(\mathfrak{n})v$ is finite-dimensional.

Recall that, by definition, all the weights occurring in a K -type are analytically integral. It follows that all the weights occurring in any module in category $\mathcal{O}^p$ are integral.

Evidently, the modules $N(\lambda)$ defined in (10) satisfy these conditions. In fact, they are nothing but the *parabolic Verma modules* defined in Sect. 9.4 of [20]. From the theory developed there, we have the following basic properties of the modules $N(\lambda)$.

- (1) Each weight of $N(\lambda)$ occurs with finite multiplicity. These multiplicities can be determined from (12).
- (2) $N(\lambda)$ contains the K -type ρ_λ with multiplicity one.
- (3) The module $N(\lambda)$ has the following universal property: Let M be a $(\mathfrak{g}, K)$ -module which contains a vector v such that:
 - $M = \mathcal{U}(\mathfrak{g}_\mathbb{C})v$;
 - v has weight λ ;
 - v is annihilated by $\langle X_-, P_{1-}, P_{0-}, N_+ \rangle$.

Then there is a surjection $N(\lambda) \rightarrow M$ mapping a highest weight vector in $N(\lambda)$ to v .

- (4) $N(\lambda)$ admits a unique irreducible submodule, and a unique irreducible quotient $L(\lambda)$. In particular, $N(\lambda)$ is indecomposable.
- (5) $N(\lambda)$ has finite length. Each factor in a composition series is of the form $L(\mu)$ for some $\mu \preceq \lambda$.
- (6) $N(\lambda)$ admits a central character, given by $\chi_{\lambda+\varrho}$. Here, as before, $\varrho = (-1, -2)$ is half the sum of the positive roots.
- (7) $L(\lambda)$ is finite-dimensional if and only if $\lambda = (k, \ell)$ with $0 \geq k \geq \ell$.

The modules M in $\mathcal{O}^p$ enjoy properties analogous to those in category $\mathcal{O}$. In particular:

- M has finite length, and admits a filtration

$$0 = V_0 \subset V_1 \subset \dots \subset V_n \subset M, \quad (14)$$

with $V_i/V_{i-1} \cong L(\lambda)$ for some $\lambda \in \Lambda^+$.

- M can be written as a finite direct sum of indecomposable modules.
- If M is an indecomposable module, then there exists a character χ of $\mathcal{Z}$ such that $M = M(\chi)$. Here,

$$M(\chi) = \{v \in M \mid (z - \chi(z))^n v = 0 \text{ for some } n \text{ depending on } z\}. \quad (15)$$

The following result, which is deeper, follows from standard classification theorems. Its last part will imply that cusp forms must have positive weight (see Proposition 4.6).

Proposition 2.2. *Let $\lambda = (k, \ell) \in \Lambda^+$.*

- (1) $L(\lambda)$ is square-integrable if and only if $\ell \geq 3$.
- (2) $L(\lambda)$ is tempered if and only if $\ell \geq 2$.
- (3) $L(\lambda)$ is unitarizable if and only if $\ell \geq 1$ or $(k, \ell) = (0, 0)$.

Proof. (1) follows from the classification of discrete series representations; see Theorem 12.21 of [23]. (The $L(\lambda)$ with $\ell \geq 3$ are precisely the holomorphic discrete series representations.)

(2) follows from the classification of tempered representations; see Theorem 8.5.3 of [23]. (The $L(\lambda)$ with $\ell = 2$ are precisely the limits of holomorphic discrete series representations.)

For a more explicit description of these classifications in the case of $\mathrm{Sp}_4(\mathbb{R})$, see [26].

(3) follows from the classification of unitary highest weight modules; see [22], [12] or [13]. We omit the details. $\square$

Lemma 2.3. *The only irreducible, locally $\mathfrak{n}$ -finite $(\mathfrak{g}, K)$ -modules are the $L(\lambda)$ for $\lambda \in \Lambda^+$.*

Proof. Let R be a locally $\mathfrak{n}$ -finite $(\mathfrak{g}, K)$ -module. Then R lies in category $\mathcal{O}^{\mathfrak{p}}$. By (14), R has a finite composition series with the quotients being $L(\lambda)$'s. So if R is irreducible, it must be an $L(\lambda)$. $\square$

Lemma 2.4. *Let $\lambda = (k, \ell) \in \Lambda^+$. The Casimir operator Ω_2 defined in (8) acts on $N(\lambda)$, and hence on $L(\lambda)$, by the scalar $\frac{1}{2}(k(k-2) + \ell(\ell-4))$.*


Proof. Since Ω_2 lies in the center of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$, it is enough to prove that $\Omega_2 w_0 = \frac{1}{2}(k(k-2) + \ell(\ell-4))w_0$, where w_0 is a vector of weight (k, ℓ) . This follows from (9). $\square$

2.2. Reducibilities and K -types. In this section we will determine composition series for each of the modules $N(\lambda)$, and determine the K -types of each $N(\lambda)$ and $L(\lambda)$.

Proposition 2.5. *Let $\lambda = (k, \ell) \in \Lambda^+$. Then $N(\lambda)$ is irreducible if and only if one of the following conditions is satisfied:*

- (1) $\ell \geq 2$.
- (2) $k = 1$.
- (3) $k + \ell = 3$.

Hence $N(\lambda)$ is irreducible if and only if λ corresponds to one of the blackened points in the following diagram:



(16)

Proof. Most cases can be handled by Theorem 9.12 in [20]. The condition (*) in this theorem translates into $\ell \geq 2$. Hence, by part a) of the theorem, $N(\lambda)$ is irreducible if $\ell \geq 2$, and by part b) of the theorem, $N(\lambda)$ is reducible if $\ell \leq 1$ and $\lambda + \varrho$ is regular (does not lie on a wall).

Hence consider λ with $\ell \leq 1$ and $\lambda + \varrho$ singular. Then either $\lambda = (1, \ell)$ with $\ell \leq 1$ or $\lambda = (x + 1, -x + 2)$ with $x \geq 1$. In the second case it is clear that no $L(\lambda')$ with $\lambda' \neq \lambda$ has the same central character as $N(\lambda)$; thus $N(\lambda)$ is irreducible. In the case that $\lambda = (1, \ell)$ with $\ell \leq 1$ we may use Theorem 9.13 in [20] (Jantzen's simplicity criterion) to see that $N(\lambda)$ is irreducible, as follows. The set of roots Ψ_λ^+ defined on page 198 of [20] is easily calculated to be $\{(-1, -1), (0, -2)\}$. Hence, by Corollary 9.13 a) of [20], $N(\lambda)$ is irreducible if and only if

$$\theta(s_{(-1, -1)} \cdot \lambda) + \theta(s_{(0, -2)} \cdot \lambda) = 0. \quad (17)$$

Here, the dot action of the Weyl group is defined in Sect. 1.8 of [20], and

$$\theta(\mu) = \text{ch}(M(\mu)) - \text{ch}(M(s_{(1, -1)} \cdot \mu)) \quad (18)$$

for any $\mu \in \Lambda^+$. It is straightforward to calculate that the characters involved on the left hand side of (17) cancel each other out, so that (17) is satisfied. $\square$

We see from (16) that the $\lambda = (k, \ell)$, $k \geq \ell$, for which $N(\lambda)$ is reducible fall into one of three regions:

- Region A: $k \leq 0$; these are the dominant integral weights.
- Region B: $k \geq 2$ and $k + \ell \leq 2$.
- Region C: $\ell \leq 1$ and $k + \ell \geq 4$.

In addition, we will consider

- Region D: $\ell \geq 3$.

Note that the disjoint union of Regions A – D comprises precisely the regular integral weights with $k \geq \ell$.

The *dot action* of an element w of the Weyl group W on $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ is defined by $w \cdot \lambda = w(\lambda + \varrho) - \varrho$, where on the right side we have the usual action of W via reflections, and where $\varrho = (-1, -2)$ is half the sum of the positive roots. Let $s_1 \in W$ be the reflection corresponding to the short simple root, and let $s_2 \in W$ be the reflection corresponding to the long simple root. Explicitly, $s_1(x, y) = (y, x)$ and $s_2(x, y) = (-x, y)$. Under the dot action, we have

$$s_2 \cdot A = B, \quad s_2 s_1 \cdot A = C, \quad s_2 s_1 s_2 \cdot A = D, \quad (19)$$

where we wrote “A” for “Region A”, etc. Consequently, $s_2 s_1 s_2 \cdot B = C$ and $s_1 s_2 s_1 \cdot C = D$.

Proposition 2.6. *Let $\lambda = (k, \ell) \in \Lambda^+$.*

- (1) *Assume that λ is in Region A. Then there is an exact sequence*

$$0 \longrightarrow L(s_2 \cdot \lambda) \longrightarrow N(\lambda) \longrightarrow L(\lambda) \longrightarrow 0.$$

The weight $s_2 \cdot \lambda = (-k + 2, \ell)$ is in Region B.

- (2) *Assume that λ is in Region B. Then there is an exact sequence*

$$0 \longrightarrow L(s_2 s_1 s_2 \cdot \lambda) \longrightarrow N(\lambda) \longrightarrow L(\lambda) \longrightarrow 0.$$

The weight $s_2 s_1 s_2 \cdot \lambda = (-\ell + 3, -k + 3)$ is in Region C.

- (3) *Assume that λ is in Region C. Then there is an exact sequence*

$$0 \longrightarrow L(s_1 s_2 s_1 \cdot \lambda) \longrightarrow N(\lambda) \longrightarrow L(\lambda) \longrightarrow 0,$$

The weight $s_1 s_2 s_1 \cdot \lambda = (k, -\ell + 4)$ is in Region D.

Proof. In this proof we will make use of the fact that a composition series for any $N(\lambda)$ is multiplicity free, i.e., each $L(\mu)$ can occur at most once as a subquotient in such a series. This fact is generally true for Hermitian symmetric pairs $(\mathfrak{g}, \mathfrak{p})$ and other pairs for which $\mathfrak{p}$ is maximal parabolic; see [7] or [8].

We first prove (3). Thus, assume that λ is in Region C. By general properties, each factor in a composition series of $N(\lambda)$ is of the form $L(\mu)$ for some $\mu \preceq \lambda$. Also, $N(\lambda)$ and $L(\mu)$ have the same central character, which is equivalent to λ and μ being in the same W -orbit under the dot action. The only μ satisfying these properties, other than λ itself, is $s_1 s_2 s_1 \cdot \lambda = (k, -\ell + 4)$. Since $N(\lambda)$ is reducible by Proposition 2.5, the module $L(s_1 s_2 s_1 \cdot \lambda)$ occurs at least once in a composition series for $N(\lambda)$. By multiplicity one, $L(s_1 s_2 s_1 \cdot \lambda)$ occurs exactly once. The assertion follows.

To prove (1) and (2), assume that λ is in Region A. By Theorem 9.16 of [20], there is an exact sequence

$$\begin{aligned} 0 \longrightarrow N(s_2 s_1 s_2 \cdot \lambda) \longrightarrow N(s_2 s_1 \cdot \lambda) \\ \longrightarrow N(s_2 \cdot \lambda) \longrightarrow N(\lambda) \longrightarrow L(\lambda) \longrightarrow 0. \end{aligned} \quad (20)$$

Note that $N(s_2 s_1 s_2 \cdot \lambda) = L(s_2 s_1 s_2 \cdot \lambda)$ by Proposition 2.5. By the already proven part (3), we get an exact sequence

$$0 \longrightarrow L(s_2 s_1 \cdot \lambda) \longrightarrow N(s_2 \cdot \lambda) \longrightarrow N(\lambda) \longrightarrow L(\lambda) \longrightarrow 0. \quad (21)$$

It follows that $N(\lambda)$ and $N(s_2 \cdot \lambda)$ have the same length. By central character considerations and multiplicity one, the length of $N(s_2 \cdot \lambda)$ can be at most 3. Hence the common length of $N(\lambda)$ and $N(s_2 \cdot \lambda)$ is 2 or 3, and our proof, of both (1) and (2), will be complete if we can show this length is 2.

By Proposition 9.14 of [20], the socle of $N(\lambda)$ is simple. It follows that the length of $N(\lambda)$ coincides with its Loewy length. We are thus reduced to showing that the Loewy length of $N(\lambda)$ is 2.

For this we will employ Theorem 4.3 of [21]. In the notation of this paper we have ${}^S W_\lambda = \{1, s_2, s_2 s_1, s_2 s_1 s_2\}$. From (20) and (21) we conclude that the set ${}^S X_\lambda$ determining the *socular weights* contains at least $s_2 s_1$ and $s_2 s_1 s_2$. It follows from (21) that $N(\lambda)$ does not contain $L(s_2 s_1 \cdot \lambda)$ in its composition series. Hence $(s_2 s_1)^\vee = s_2$, where $w^\vee$ for $w \in {}^S X_\lambda$ is defined on p. 734 of [21]. A computation of the elements $\overline{w}$, defined on p. 743 of [21] for $w \in {}^S W_\lambda$, yields

$$\overline{1} = 1, \quad \overline{s_2} = s_2, \quad \overline{s_2 s_1} = s_1, \quad \overline{s_2 s_1 s_2} = s_2.$$

Thus the number t appearing in Theorem 4.3 of [21], defined as the maximal length of any $\overline{w}$, is 1. The hypothesis of this theorem is satisfied (set $x = s_2 s_1$). The theorem implies that the Loewy length of any $N(w \cdot \lambda)$ for $w \in {}^S W_\lambda$ is at most 2. In particular, the Loewy length of $N(\lambda)$ is 2, concluding our proof. $\square$

We will next determine the K -types in $N(\lambda)$. Let V be any admissible $(\mathfrak{g}, K)$ -module. For a weight $\lambda \in \Lambda$, let V_λ be the corresponding weight space. We denote by

$$m_\lambda(V) = \dim V_\lambda \quad (22)$$

the multiplicity of the weight λ in V . Let

$$\text{mult}_\lambda(V) = \text{the multiplicity of the } K\text{-type } \rho_\lambda \text{ in } V. \quad (23)$$

It follows from the weight structure of the K -types that

$$\text{mult}_\lambda(V) = m_\lambda(V) - m_{\lambda+(1,-1)}(V). \quad (24)$$

Let $Q(\lambda)$ be the number of ways to write $\lambda \in \Lambda$ as a $\mathbb{Z}_{\geq 0}$ linear combination of $(2, 0)$, $(1, 1)$ and $(0, 2)$. It is easy to see that, for $\lambda = (x, y)$ with integers x, y ,

$$Q(x, y) = \begin{cases} \left\lfloor \frac{\min(x, y) + 2}{2} \right\rfloor & \text{if } x, y \geq 0 \text{ and } x \equiv y \pmod{2}, \\ 0 & \text{otherwise.} \end{cases} \quad (25)$$

Lemma 2.7. *Let $\lambda = (k, \ell) \in \Lambda^+$. Let x, y be integers with $x \geq y$. Then*

$$\text{mult}_{(x,y)}(N(\lambda)) = 0 \quad \text{if } x < k, \text{ or } y < \ell, \text{ or } x - y \not\equiv k - \ell \pmod{2}.$$

If $x \geq k$ and $y \geq \ell$ and $x - y \equiv k - \ell \pmod{2}$, then

$$\text{mult}_{(x,y)}(N(\lambda)) = \begin{cases} \left\lfloor \frac{\min(x-k, y-\ell) + 2}{2} \right\rfloor & \text{if } y \leq k, \\ \left\lfloor \frac{\min(x-k, y-\ell)}{2} \right\rfloor - \left\lfloor \frac{y-k-1}{2} \right\rfloor & \text{if } y > k. \end{cases}$$

Proof. It follows from (12) that

$$m_{(x,y)}(N(\lambda)) = \sum_{n=0}^{k-\ell} Q(x-k+n, y-\ell-n). \quad (26)$$

The result now follows from (24) and some simplification. $\square$

Proposition 2.6 combined with Lemma 2.7 allows us to calculate the multiplicities of the K -types of any $L(\lambda)$. Note also that if λ is not in Region A, B or C, then $L(\lambda) = N(\lambda)$ by Proposition 2.5, so that Lemma 2.7 can be used directly to calculate the multiplicities. We record a few special cases in the following result; the details of the elementary proofs are omitted.

Proposition 2.8. *Let $x \geq y$ be integers.*

- (1) *Assume that $\lambda = (\ell, \ell)$ with an integer $\ell \geq 1$. Then*

$$\text{mult}_{(x,y)}(L(\lambda)) = 0 \quad \text{if } x < \ell, \text{ or } y < \ell, \text{ or } x \not\equiv y \pmod{2}.$$

If $x \geq \ell$ and $y \geq \ell$ and $x \equiv y \pmod{2}$, then

$$\text{mult}_{(x,y)}(L(\lambda)) = \begin{cases} 1 & \text{if } y \equiv \ell \pmod{2}, \\ 0 & \text{if } y \not\equiv \ell \pmod{2}. \end{cases}$$

- (2) *Assume that $\lambda = (k, 1)$ with an integer $k \geq 2$. Then*

$$\text{mult}_{(x,y)}(L(\lambda)) = 0 \quad \text{if } x < k, \text{ or } y < 1, \text{ or } x - y \not\equiv k - 1 \pmod{2}.$$

If $x \geq k$ and $y \geq 1$ and $x - y \equiv k - 1 \pmod{2}$, then


$$\text{mult}_{(x,y)}(L(\lambda)) = \begin{cases} 1 & \text{if } y \leq x - k + 1, \\ 0 & \text{if } y > x - k + 1. \end{cases}$$

- (3) *Assume that $\lambda = (k, \ell)$ is in Region C. Then $\text{mult}_{(x,y)}(L(\lambda)) = 0$ if $y \geq x - k - \ell + 4$. Hence, all the K -types of $L(\lambda)$ are strictly below the diagonal line running through the point $(k, -\ell + 4)$.*

- (4) *Assume that $\lambda = (2, 0)$. Then*

$$\text{mult}_{(x,y)}(L(\lambda)) = \begin{cases} 1 & \text{if } x \geq 2, y \geq 0 \text{ and } x \equiv y \equiv 0 \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

Finally, we consider the location of the boundary K -types in the modules $N(\lambda)$ for any $\lambda = (k, \ell)$ with $k \geq \ell$. By Lemma 2.7, all the boundary K -types occur with multiplicity one. There are no K -types $\rho_{(x,y)}$ for $x < k$ or $y < \ell$. For $x = k$ or $y = \ell$ the K -types occur in steps of 2. The top boundary is provided by the line $y = x$ if $k \equiv \ell \pmod{2}$, or the line $y = x - 1$ if $k \not\equiv \ell \pmod{2}$. In the first case, the K -types on this line occur in steps of 2, in the second case in steps of 1. The following diagrams illustrate these two cases.



2.3. Navigating the highest weight vectors. Let V be a $(\mathfrak{g}, K)$ -module. In this section we will investigate a collection of elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ that preserve the property of being a highest weight vector in some K -type. In other words, these elements X will have the property that $N_+Xv = 0$ if $N_+v = 0$. Evidently, elements that commute with N_+ , like X_+ and P_{0-} , have this property.

TABLE 1. Some elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ that take highest weight vectors to highest weight vectors. The last column shows the resulting weight after applying an operator to a vector of weight $(\ell + m, \ell)$.

name	definition	new weight
P_{0-}		$(\ell + m, \ell - 2)$
L	$m(m-1)X_- - (m-1)P_{1-}N_- + P_{0-}N_-^2$	$(\ell + m - 2, \ell)$
U	$m(m-1)P_{0+} + (m-1)P_{1+}N_- + X_+N_-^2$	$(\ell + m, \ell + 2)$
X_+		$(\ell + m + 2, \ell)$
E_+	$(m+2)P_{1+} + 2N_-X_+$	$(\ell + m + 1, \ell + 1)$
E_-	$(m+2)P_{1-} - 2N_-P_{0-}$	$(\ell + m - 1, \ell - 1)$
D_+	$P_{1+}^2 - 4X_+P_{0+}$	$(\ell + m + 2, \ell + 2)$
D_-	$P_{1-}^2 - 4X_-P_{0-}$	$(\ell + m - 2, \ell - 2)$

More specifically, we consider a vector $v \in V$ of weight $(\ell + m, \ell)$ for $m \geq 0$. The new elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ that we introduce are called U , L , E_+ , E_- , D_+ , and D_- . Their definitions appear in Table 1. These operators take v to another vector in V of the weight indicated in the “new weight” column. Note that the operators U , L , E_+ and E_- depend on m . However, for brevity, our notation will not reflect this dependence. The formulas for the operators U and L are given only for $m \geq 2$; we adopt the convention that $U = L = 0$ if $m < 2$.

Remark 2.9. As was pointed out by the referee, the element $Z - Z'$ of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ acts on a vector of weight $(\ell + m, \ell)$ by multiplication by m ; therefore the operators in Table 1 can be defined as single elements of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ without needing to involve the integer m . This leads to the alternate definitions below, which are equivalent to the ones given in Table 1:

$$\begin{aligned} L &= X_-(Z - Z')(Z - Z' - 1) - P_{1-}N_-(Z - Z' - 1) + P_{0-}N_-^2, \\ U &= P_{0+}(Z - Z')(Z - Z' - 1) + P_{1+}N_-(Z - Z' - 1) + X_+N_-^2, \\ E_+ &= P_{1+}(Z - Z' + 2) + 2N_-X_+, \\ E_- &= P_{1-}(Z - Z' + 2) - 2N_-P_{0-}. \end{aligned}$$

Lemma 2.10. *Let ℓ be an integer, and m a non-negative integer. Let v be a vector of weight $(\ell + m, \ell)$ in some $(\mathfrak{g}, K)$ -module V . Let $X \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ be one of the elements in Table 1. Then $N_+Xv = 0$ if $N_+v = 0$. The weight of Xv is indicated in the last column of Table 1. For the U and L operators we assume $m \geq 2$.*

Proof. In order to show that $N_+Xv = 0$ if $N_+v = 0$, it suffices to show that $N_+X \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})N_+$. This is easily verified using the commutation relations. The weight of Xv as indicated in the last column of Table 1 is also immediate from the commutation relations. $\square$

As we already mentioned, $[N_+, X_+] = [N_+, P_{0-}] = 0$. The two-step diagonal operators $D_{\pm}$ have in fact the property that $[N_+, D_{\pm}] = [N_-, D_{\pm}] = 0$. The other operators in Table 1 do not universally commute with N_+ . Using the commutation relations, one may further verify that

$$X_+E_+ = E_+X_+, \quad (28)$$

$$UE_+ = E_+U, \quad (29)$$

$$D_+E_+ = E_+D_+, \quad (30)$$

$$UD_+ = D_+U, \quad (31)$$

$$X_+U - UX_+ = (m + 1)D_+. \quad (32)$$

We remind the reader that the particular element in $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ that each operator appearing in the above equations corresponds to, depends on the weight of the vector that the operator will act on. For instance, consider both sides of (28) acting on a vector of weight $(\ell + m, \ell)$. Then the E_+ on the left side is given by the formula in Table 1 while the E_+ on the right side is obtained by the substitution $m \mapsto m + 2$ in the same formula.

Now consider a weight $\lambda = (\ell + m, \ell)$ with $\ell \in \mathbb{Z}$ and $m \geq 0$. By Lemma 2.7, if a K -type $\rho_{(x,y)}$ occurs in $N(\lambda)$, then $x \geq \ell + m$ and $y \geq \ell$. We may therefore hope to generate all highest weight vectors in the K -types of $N(\lambda)$ by applying appropriate powers of the operators X_+ , D_+ , U and E_+ to the lowest weight vector w_0 of $N(\lambda)$. We will see below that this is indeed the case.

As a first step in this direction, consider the K -types $\rho_{(x,y)}$ with $x = \ell + m$; these are the ones that are straight above the minimal weight. By Lemma 2.7, these are exactly the K -types $(\ell + m, \ell + 2i)$, $i \in \{0, 1, \dots, \lfloor \frac{m}{2} \rfloor\}$, and each of these occurs with multiplicity 1 in $N(\lambda)$. Let w_0 be a lowest weight vector (i.e., a highest weight vector in the minimal K -type of $N(\lambda)$); thus, w_0 has weight $(\ell + m, \ell)$, and $N_+w_0 = 0$. For $i \in \{0, 1, \dots, \lfloor \frac{m}{2} \rfloor\}$, let⁶

$$w_i = U^i w_0. \quad (33)$$

Then w_i has weight $(\ell + m, \ell + 2i)$, and $N_+w_i = 0$. If $w_i \neq 0$, then it is a highest weight vector in the K -type $\rho_{(\ell+m, \ell+2i)}$ of $N(\lambda)$.

⁶Once again, we remind the reader that the operator U^i in (33) is really shorthand for $U_{m+2-2i} \dots U_{m-2} U_m$; i.e., the integer m appearing in the definition of U changes at each step.

Lemma 2.11. *With the above notations,*

$$P_{0-}w_{i+1} = -(i+1)(\ell+i-1)(m-2i)(m-2i-1)w_i.$$

for $i \in \{0, 1, \dots, \lfloor \frac{m}{2} \rfloor - 1\}$. In particular, if $\ell \geq 2$, then $w_i \neq 0$ for all $i \in \{0, 1, \dots, \lfloor \frac{m}{2} \rfloor\}$.

Lemma 2.12. *Suppose $\lambda = (\ell+m, \ell)$ with $m \geq 0$, m even, and $\ell \geq 1$. If $\ell = 1$, assume further that $m = 0$. Let w_0 be a non-zero vector of weight $(\ell+m, \ell)$ in $N(\lambda)$ such that $N_+w_0 = 0$. Then, for all $\beta \geq 0$, $P_{0-}^{m/2}D_-^\beta D_+^\beta U^{m/2}w_0$ is a non-zero multiple of w_0 .*

The above two lemmas can be proved by routine calculation.

Before stating the next result, it will be convenient to introduce the concept of N_- -layers. Let $\lambda = (\ell+m, \ell) \in \Lambda$ with $\ell \in \mathbb{Z}$ and $m \geq 0$. Given a non-negative integer δ , the δ -th N_- -layer of $N(\lambda)$, denoted by $N(\lambda)^\delta$, is defined as the subspace spanned by all vectors of the form

$$X_+^\alpha P_{1+}^\beta P_{0+}^\gamma N_-^\delta w_0, \quad \alpha, \beta, \gamma \geq 0. \quad (34)$$

Here, as before, w_0 is a fixed non-zero vector of weight λ . Note that $N(\lambda)^\delta = 0$ for $\delta > m$. By (12), we have $N(\lambda) = N(\lambda)^0 \oplus \dots \oplus N(\lambda)^m$. We also introduce the notation $N(\lambda)^{\leq \delta} = N(\lambda)^0 \oplus \dots \oplus N(\lambda)^\delta$. Observe that, since N_- normalizes $\mathfrak{p}_+ = \langle P_{0+}, P_{1+}, X_+ \rangle$, in any expression involving these four operators we may always move the N_- 's to the right. In fact,

$$N_-Yw_0 \in YN_-w_0 + N(\lambda)^\delta \quad \text{for } Y \in \mathcal{U}(\mathfrak{p}_+)N_-^\delta. \quad (35)$$

It follows that the operator N_- maps $N(\lambda)^\delta$ to $N(\lambda)^\delta \oplus N(\lambda)^{\delta+1}$. In particular, N_- induces an endomorphism of the top layer $N(\lambda)^m$.

Lemma 2.13. *Let $\lambda = (\ell+m, \ell) \in \Lambda^+$.*

- (1) *Let $f \in \mathbb{C}[X, Y, Z]$ be a non-zero polynomial. Then the element $f(X_+, P_{1+}, P_{0+})$ of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ acts injectively on $N(\lambda)$, and it preserves N_- -layers.*
- (2) *The restriction of E_+ to $N(\lambda)^{\leq (m-1)}$ is injective.*

Proof. (1) is immediate from (11). (2) follows easily from (35) and the defining formula $E_+ = (m+2)P_{1+} + 2N_-X_+$. $\square$

Lemma 2.14. *Let $\lambda = (\ell+m, \ell) \in \Lambda$ with $\ell \geq 2$ and $m \geq 0$. Let the vectors $w_i \in N(\lambda)$ be defined as in (33). Then the vectors*

$$X_+^\alpha D_+^\beta w_i, \quad \alpha, \beta \geq 0, i \in \left\{0, 1, \dots, \left\lfloor \frac{m}{2} \right\rfloor\right\}, \quad (36)$$

are linearly independent.

Proof. First note that the w_i are non-zero by Lemma 2.11. By Lemma 2.13 (1), all the vectors (36) are non-zero. We see from the defining formula for the U operator in Table 1 that $X_+^\alpha D_+^\beta w_i$ lies in $N(\lambda)^{\leq 2i}$, but not in $N(\lambda)^{\leq (2i-1)}$. It follows that any linear combination between the vectors (36) can only involve a single i . But for fixed i the vectors (36) have distinct weights as α and β vary. Our assertion follows. $\square$

Recall from Lemma 2.7 that if a K -type $\rho_{(x,y)}$ occurs in $N(\lambda)$, where $\lambda = (\ell+m, \ell)$, then $x-y \equiv m \pmod{2}$. We say that such a K -type is of *parity 0* if $x \equiv \ell+m \pmod{2}$ and $y \equiv \ell \pmod{2}$. Otherwise, if $x \not\equiv \ell+m \pmod{2}$ and $y \not\equiv \ell \pmod{2}$, we say the K -type is of *parity 1*. We apply the same terminology to the highest weight vectors of such K -types. Clearly, the operators X_+ , P_{0-} , U , L and $D_\pm$ preserve the parity, while $E_\pm$ change the parity. Let $N(\lambda)_{\text{par}(0)}$ (resp. $N(\lambda)_{\text{par}(1)}$) be the subspace of $N(\lambda)$ spanned by highest weight vectors of parity 0 (resp. parity 1). We now state the main result of this section.

Proposition 2.15. *Let $\lambda = (\ell+m, \ell) \in \Lambda^+$ with $\ell \geq 2$ and $m \geq 0$.*

- (1) *$N(\lambda)_{\text{par}(0)}$ is precisely the space spanned by the vectors (36).*
- (2) *If m is odd, then the map $E_+ : N(\lambda)_{\text{par}(0)} \rightarrow N(\lambda)_{\text{par}(1)}$ is an isomorphism.*

- (3) If m is even, then the map $E_+ : N(\lambda)_{\text{par}(0)} \rightarrow N(\lambda)_{\text{par}(1)}$ is surjective, and its kernel is spanned by the vectors (36) with $i = m/2$.

Proof. (1) Clearly, the highest weight vectors (36) all have parity 0. By easy combinatorics we can determine the number of vectors (36) of a fixed weight (x, y) . Comparing with the formula from Lemma 2.7, we see that this number coincides with $\text{mult}_{(x,y)}(N(\lambda))$. This proves (1) in view of the linear independence of the vectors (36).

(2) If m is odd, then the vectors (36) are all contained in $N(\lambda)^{\leq(m-1)}$. Hence $E_+ : N(\lambda)_{\text{par}(0)} \rightarrow N(\lambda)_{\text{par}(1)}$ is injective by part (1) and Lemma 2.13 (2). To prove surjectivity, it is enough to show that $\text{mult}_{(x,y)}(N(\lambda)) = \text{mult}_{(x-1,y-1)}(N(\lambda))$ for all (x, y) of parity 1. This follows from the formula in Lemma 2.7.

(3) Assume that m is even. The vector $w_{m/2}$ has weight $(\ell + m, \ell + m)$. By Lemma 2.7, the K -type $\rho_{(\ell+m+1, \ell+m+1)}$ is not contained in $N(\lambda)$; see also (27). Hence $E_+ w_{m/2} = 0$. By (28) - (30), E_+ annihilates all vectors (36) with $i = m/2$. The vectors (36) with $i < m/2$ are all contained in $N(\lambda)^{\leq(m-1)}$. Therefore, the assertion about the kernel of E_+ follows from part (1) and Lemma 2.13 (2).

To prove the surjectivity assertion, first note that, by Lemma 2.7,

$$\text{mult}_{(x,y)}(N(\lambda)) = \begin{cases} \text{mult}_{(x-1,y-1)}(N(\lambda)) & \text{if } y \leq \ell + m, \\ \text{mult}_{(x-1,y-1)}(N(\lambda)) - 1 & \text{if } y > \ell + m, \end{cases}$$

for all K -types $\rho_{(x,y)}$ of parity 1. The K -type $\rho_{(x-1,y-1)}$ of parity 0 receives a contribution from a vector (36) with $i = m/2$ if and only if $y > \ell + m$. The surjectivity therefore follows by what we already proved about the kernel of E_+ . $\square$

Remark 2.16. Let $N(\lambda)_{\text{hw}} = N(\lambda)_{\text{par}(0)} \oplus N(\lambda)_{\text{par}(1)}$ be the subspace of highest weight vectors, and let $\mathcal{J}$ be the subalgebra of $\mathcal{U}(\mathfrak{p}_+)$ generated by X_+ and D_+ . Then Proposition 2.14 implies that $N(\lambda)_{\text{hw}}$ is a free $\mathcal{J}$ -module of rank $m + 1$, the dimension of the minimal K -type. For holomorphic discrete series representations, i.e., for $\ell \geq 3$, this also follows from the main result of [19].

The case of lowest weight $(1 + m, 1)$. In Proposition 2.15 we assumed $\ell \geq 2$ since otherwise some of the vectors w_i might be zero; see Lemma 2.11. However, for later applications we also require the following analogous result for the $L(\lambda)$ with $\lambda = (1 + m, 1)$.

Proposition 2.17. *Let $\lambda = (1 + m, 1)$ with $m \geq 0$. Let w_0 be a non-zero vector of weight $(1 + m, 1)$ in $L(\lambda)$.*

- (1) $L(\lambda)_{\text{par}(0)}$ is precisely the space spanned by the vectors

$$X_+^\alpha D_+^\beta w_0, \quad \alpha, \beta \geq 0. \quad (37)$$

- (2) If $m \geq 1$, then the map $E_+ : L(\lambda)_{\text{par}(0)} \rightarrow L(\lambda)_{\text{par}(1)}$ is an isomorphism. If $m = 0$, then $L(\lambda)_{\text{par}(1)} = 0$.

Proof. The proof is similar to the previous case, and is omitted. $\square$

3. DIFFERENTIAL OPERATORS

3.1. Functions on the group and functions on $\mathbb{H}_2$. Recall that $K \cong U(2)$ via $\begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mapsto A + iB$. On the Lie algebra level, this map induces an isomorphism $\mathfrak{k} \cong \mathfrak{u}(2)$ given by the same formula. Extending this map $\mathbb{C}$ -linearly, we get an isomorphism $\mathfrak{k}_{\mathbb{C}} \cong \mathfrak{gl}_2(\mathbb{C})$. Under this isomorphism,

$$Z \mapsto \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad Z' \mapsto \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad N_+ \mapsto \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad N_- \mapsto \begin{bmatrix} 0 & 0 \\ -1 & 0 \end{bmatrix}. \quad (38)$$

Let ℓ be an integer, and m a non-negative integer. Let $W_m \simeq \text{sym}^m(\mathbb{C}^2)$ be the space of all complex homogeneous polynomials of total degree m in the two variables S and T . For any $g \in \text{GL}_2(\mathbb{C})$, and $P(S, T) \in W_m$, define $\eta_{\ell, m}(g)P(S, T) = \det(g)^\ell P((S, T)g)$. Then $(\eta_{\ell, m}, W_m)$ gives a concrete realization of the irreducible representation $\det^\ell \text{sym}^m$ of $\text{GL}_2(\mathbb{C})$. We will denote the derived representation of $\mathfrak{gl}_2(\mathbb{C})$ by the same symbol $\eta_{\ell, m}$. Easy calculations show that, under the identification (38),

$$\eta_{\ell, m}(Z)S^{m-j}T^j = (\ell + m - j)S^{m-j}T^j, \quad (39)$$

$$\eta_{\ell, m}(Z')S^{m-j}T^j = (\ell + j)S^{m-j}T^j, \quad (40)$$

$$\eta_{\ell, m}(N_+)S^{m-j}T^j = jS^{m-j+1}T^{j-1}, \quad (41)$$

$$\eta_{\ell, m}(N_-)S^{m-j}T^j = -(m - j)S^{m-j-1}T^{j+1}. \quad (42)$$

In particular, $\eta_{\ell, m}(N_+)S^m = 0$ and $\eta_{\ell, m}(N_-)T^m = 0$. Since the vector S^m is a highest weight vector of weight $(\ell + m, \ell)$, we see that

$$\text{The restriction of } \eta_{\ell, m} \text{ to } U(2) \text{ is } \rho_{(\ell+m, \ell)}. \quad (43)$$

For a smooth function Φ on $\text{Sp}_4(\mathbb{R})$ of weight $(\ell + m, \ell)$ satisfying $N_+\Phi = 0$, we define a function $\vec{\Phi}$ taking values in the polynomial ring $\mathbb{C}[S, T]$ by

$$\vec{\Phi}(g) = \sum_{j=0}^m \frac{(-1)^j}{j!} (N_-^j \Phi)(g) S^{m-j} T^j, \quad g \in \text{Sp}_4(\mathbb{R}). \quad (44)$$

Evidently, $\vec{\Phi}$ takes values in the space $W_m \subset \mathbb{C}[S, T]$ of the representation $\eta_{\ell, m}$. Hence, an expression like $\eta_{\ell, m}(h)(\vec{\Phi}(g))$ makes sense, for any $h \in \text{GL}_2(\mathbb{C})$.

In the following lemma, for clarity of notation, we let ι be the transposition map on 2×2 complex matrices. We may interpret ι as an anti-involution of $\text{GL}_2(\mathbb{C})$. The derived map, also given by transposition and also denoted by ι , is an anti-involution of $\mathfrak{gl}_2(\mathbb{C})$. It extends to an anti-involution of the algebra $\mathcal{U}(\mathfrak{gl}_2(\mathbb{C}))$. When we write $\iota(h)$ for $h \in K$, we mean ι applied to the element of $U(2)$ corresponding to $h \in K$ via the map $\begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mapsto A + iB$.

Lemma 3.1. *Let ℓ be any integer, and m a non-negative integer. Let Φ be a K -finite function on $\text{Sp}_4(\mathbb{R})$ of weight $(\ell + m, \ell)$ satisfying $N_+\Phi = 0$ (right translation action). Let $\vec{\Phi}$ be the polynomial-valued function defined in (44). Then*

$$\vec{\Phi}(gh) = \eta_{\ell, m}(\iota(h))(\vec{\Phi}(g)), \quad \text{for } h \in K \quad (45)$$

and $g \in \text{Sp}_4(\mathbb{R})$. On the Lie algebra level,

$$(X\vec{\Phi})(g) = \eta_{\ell, m}(\iota(X))(\vec{\Phi}(g)) \quad (46)$$

for $X \in \mathcal{U}(\mathfrak{k}_{\mathbb{C}})$ and $g \in \text{Sp}_4(\mathbb{R})$. More generally,

$$(YX\vec{\Phi})(g) = \eta_{\ell, m}(\iota(X))((Y\vec{\Phi})(g)) \quad (47)$$

for $X \in \mathcal{U}(\mathfrak{k}_{\mathbb{C}})$, $Y \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ and $g \in \text{Sp}_4(\mathbb{R})$.

Proof. Fixing $g \in \text{Sp}_4(\mathbb{R})$, we first claim that (46) holds for $X \in \mathfrak{k}_{\mathbb{C}}$. In fact, this assertion is easily verified using the formulas (39) – (42). For $X = N_+$ the identity

$$N_+ N_-^j = N_-^j N_+ + j N_-^{j-1} (Z' - Z) + j(j-1) N_-^{j-1}$$

is helpful.

Replacing g by $g \exp(tY)$ and taking $\frac{d}{dt}|_0$ on both sides, one proves that (46) also holds for elements of degree 2 in $\mathcal{U}(\mathfrak{k}_{\mathbb{C}})$. Continuing in this manner, we see that (46) holds for any element $X \in \mathcal{U}(\mathfrak{k}_{\mathbb{C}})$. Now using that $\exp((d\eta)(X)) = \eta(\exp(X))$ for any representation η and $X \in \mathfrak{k}$, one can derive the identity (45).

To prove (47), replace g by $g \exp(tY)$ in (46) for some $Y \in \mathfrak{g}$. Taking $\frac{d}{dt}|_0$ on both sides, we see that (47) holds for $Y \in \mathfrak{g}$, and then also for $Y \in \mathfrak{g}_{\mathbb{C}}$. Continuing in this manner, we conclude that (47) holds for $Y \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ of any degree. $\square$

Evidently, the function Φ in Lemma 3.1 can be recovered as the S^m -component of $\vec{\Phi}$. It is easy to see that the map $\Phi \mapsto \vec{\Phi}$ establishes an isomorphism between the space of K -finite functions of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$, and the space of smooth functions $\vec{\Phi} : \mathrm{Sp}_4(\mathbb{R}) \rightarrow W_m$ satisfying (45).

For later use, we make the following observation. Recall from Sect. 2.1 that we have $\mathfrak{n} = \langle X_-, P_{1-}, P_{0-} \rangle$, and that this commutative Lie algebra is normalized by $\mathfrak{k}_{\mathbb{C}}$. For a smooth function Φ of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$, we then have

$$\mathfrak{n}\Phi = 0 \quad \Longleftrightarrow \quad \mathfrak{n}\vec{\Phi} = 0. \quad (48)$$

(on both sides we mean the right translation action of $\mathfrak{n}$ on smooth functions on the group). This follows from the definition (44), and the fact that N_- normalizes $\mathfrak{n}$.

Descending to the Siegel upper half space. From the vector-valued function $\vec{\Phi}$ we can construct a vector-valued function on $\mathbb{H}_2$, as follows. For $g \in \mathrm{Sp}_4(\mathbb{R})$ and $Z \in \mathbb{H}_2$, let

$$J(g, Z) = CZ + D, \quad g = \begin{bmatrix} A & B \\ C & D \end{bmatrix}. \quad (49)$$

Then $J(g_1 g_2, Z) = J(g_1, g_2 Z) J(g_2, Z)$. Since $\iota(h) = \bar{h}^{-1}$ for $h \in U(2)$, the transformation property (45) can be rewritten as $\vec{\Phi}(gh) = \eta_{\ell, m}(J(h, I))^{-1} \vec{\Phi}(g)$ for $h \in K$; recall that $I = \begin{bmatrix} & 1 \\ 1 & \end{bmatrix}$. It follows that the W_m -valued function $g \mapsto \eta_{\ell, m}(J(g, I)) \vec{\Phi}(g)$ is right K -invariant. Hence, this function descends to a function F on $\mathbb{H}_2 \cong \mathrm{Sp}_4(\mathbb{R})/K$. Explicitly, we define F by

$$F(Z) = \eta_{\ell, m}(J(g, I)) \vec{\Phi}(g), \quad (50)$$

where g is any element of $\mathrm{Sp}_4(\mathbb{R})$ satisfying $gI = Z$. Conversely, if F is a smooth W_m -valued function on $\mathbb{H}_2$, then we can define a smooth function $\vec{\Phi}$ on $\mathrm{Sp}_4(\mathbb{R})$ by the formula $\vec{\Phi}(g) = \eta_{\ell, m}(J(g, I))^{-1} F(gI)$. Clearly, $\vec{\Phi}$ satisfies the transformation property (45). Combining the maps $\Phi \mapsto \vec{\Phi}$ and $\vec{\Phi} \mapsto F$, we obtain the following result.

Lemma 3.2. *Let ℓ be any integer, and m a non-negative integer. Let $\mathcal{V}_{\ell, m}$ be the space of smooth K -finite functions $\Phi : \mathrm{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$. Then $\mathcal{V}_{\ell, m}$ is isomorphic to the space of smooth functions $F : \mathbb{H}_2 \rightarrow W_m$. If $\Phi \in \mathcal{V}_{\ell, m}$, then the corresponding function F is given by (50), where $\vec{\Phi}$ is defined in (44).*

Given any function $F : \mathbb{H}_2 \rightarrow W_m$, we will write F in the form

$$F(Z) = \sum_{j=0}^m F_j(Z) S^{m-j} T^j,$$

and call the complex-valued functions F_j the *component functions* of F . The component F_0 is obtained from F by setting $(S, T) = (1, 0)$. The component F_1 is obtained by taking $\frac{\partial}{\partial T}$ and then setting $(S, T) = (1, 0)$. In general,

$$F_j(Z) = \frac{1}{j!} \frac{\partial^j}{\partial T^j} F(Z) \Big|_{(S, T) = (1, 0)}. \quad (51)$$

Next, we introduce coordinates on $\mathbb{H}_2$, as follows. Let us write an element $Z \in \mathbb{H}_2$ as

$$Z = \begin{bmatrix} \tau & z \\ z & \tau' \end{bmatrix}, \quad \tau = x + iy, \quad z = u + iv, \quad \tau' = x' + iy', \quad (52)$$

where x, y, u, v, x', y' are real numbers, $y, y' > 0$, and $yy' - v^2 > 0$. We set

$$b_Z = \begin{bmatrix} 1 & x & u \\ & 1 & u \\ & & 1 \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & v/y' & & \\ & 1 & & \\ & & 1 & \\ & & -v/y' & 1 \end{bmatrix} \begin{bmatrix} a & & & \\ & b & & \\ & & a^{-1} & \\ & & & b^{-1} \end{bmatrix} \quad (53)$$

with

$$a = \sqrt{y - \frac{v^2}{y'}} \quad \text{and} \quad b = \sqrt{y'}. \quad (54)$$

Then b_Z is an element of the Borel subgroup of $\mathrm{Sp}_4(\mathbb{R})$, and $b_Z I = Z$. Every element of $\mathrm{Sp}_4(\mathbb{R})$ can be written as $b_Z h$ for a uniquely determined $Z \in \mathbb{H}_2$ and a uniquely determined $h \in K$.

If $F, \Phi, \vec{\Phi}$ are as above, then the following relation is immediate from (50).

$$F(Z) = \eta_{l,m}(J(b_Z, I)) \vec{\Phi}(b_Z). \quad (55)$$

3.2. The action of the root vectors. Let $\Phi, \vec{\Phi}$ and F be as in Lemma 3.2. In this section we will calculate $(X\vec{\Phi})(b_Z)$, where X is any of the root vectors $X_{\pm}, P_{1\pm}, P_{0\pm}, N_{\pm}$, and where b_Z is the element defined in (53). The result will be expressed in terms of differential operators applied to the function F . As a consequence, we will prove that F is holomorphic if and only if $\mathfrak{n}\vec{\Phi} = 0$.

For $Z \in \mathbb{H}_2$, let $D_Z = J(b_Z, I)$. Then D_Z is simply the lower right 2×2 -block of b_Z , explicitly,

$$D_Z = \begin{bmatrix} 1 & \\ -v/y' & 1 \end{bmatrix} \begin{bmatrix} a^{-1} & \\ & b^{-1} \end{bmatrix}, \quad a = \sqrt{y - \frac{v^2}{y'}}, \quad b = \sqrt{y'}. \quad (56)$$

Proposition 3.3. *Let (η, W) be a finite-dimensional holomorphic representation of $\mathrm{GL}_2(\mathbb{C})$. Let F be a W -valued smooth function on $\mathbb{H}_2$, and let $\vec{\Phi}$ be the corresponding W -valued function on $\mathrm{Sp}_4(\mathbb{R})$, i.e.,*

$$\vec{\Phi}(g) = \eta(J(g, I))^{-1} F(gI).$$

Let b_Z be as in (53), and D_Z as in (56). Then the following formulas hold.

$$\eta(D_Z)(N_+ \vec{\Phi})(b_Z) = \eta(D_Z) \eta \left(\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \right) \eta(D_Z)^{-1} F(Z). \quad (57)$$

$$\eta(D_Z)(N_- \vec{\Phi})(b_Z) = -\eta(D_Z) \eta \left(\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \right) \eta(D_Z)^{-1} F(Z). \quad (58)$$

$$\begin{aligned} \eta(D_Z)(P_{0+} \vec{\Phi})(b_Z) &= \eta(D_Z) \eta \left(\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \eta(D_Z)^{-1} F(Z) \\ &\quad + \frac{2i}{y'} \left(v^2 \frac{\partial F}{\partial \tau} + vy' \frac{\partial F}{\partial z} + y'^2 \frac{\partial F}{\partial \tau'} \right) (Z). \end{aligned} \quad (59)$$

$$\eta(D_Z)(P_{0-} \vec{\Phi})(b_Z) = -\frac{2i}{y'} \left(v^2 \frac{\partial F}{\partial \bar{\tau}} + vy' \frac{\partial F}{\partial \bar{z}} + y'^2 \frac{\partial F}{\partial \bar{\tau}'} \right) (Z). \quad (60)$$

$$\begin{aligned} \eta(D_Z)(P_{1+} \vec{\Phi})(b_Z) &= \eta(D_Z) \eta \left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right) \eta(D_Z)^{-1} F(Z) \\ &\quad + \frac{2i}{y'} \sqrt{\Delta} \left(2v \frac{\partial F}{\partial \tau} + y' \frac{\partial F}{\partial z} \right) (Z). \end{aligned} \quad (61)$$

$$\eta(D_Z)(P_{1-} \vec{\Phi})(b_Z) = -\frac{2i}{y'} \sqrt{\Delta} \left(2v \frac{\partial F}{\partial \bar{\tau}} + y' \frac{\partial F}{\partial \bar{z}} \right) (Z). \quad (62)$$

$$\eta(D_Z)(X_+ \vec{\Phi})(b_Z) = \eta(D_Z) \eta \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \right) \eta(D_Z)^{-1} F(Z) + \frac{2i}{y'} \Delta \frac{\partial F}{\partial \tau} (Z). \quad (63)$$

$$\eta(D_Z)(X_- \vec{\Phi})(b_Z) = -\frac{2i}{y'} \Delta \frac{\partial F}{\partial \bar{\tau}}(Z). \quad (64)$$

Here, we used the abbreviation $\Delta = yy' - v^2$.

Proof. To prove these formulas, one has to first compute the action of a basis of root vectors in the uncomplexified Lie algebra. This is relatively straightforward using the definitions, though somewhat tedious. Once that is done, the action of the root vectors above lying in the complexified Lie algebra follows by linearity. We omit the details. $\square$

Corollary 3.4. *Let ℓ be any integer, and m a non-negative integer. Let $\Phi : \mathrm{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ be a K -finite function of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$. Let $F : \mathbb{H}_2 \rightarrow W_m$ be the function corresponding to Φ according to Lemma 3.2. Then F is holomorphic if and only if $\mathfrak{n}\Phi = 0$.*

Proof. It follows from (60), (62) and (64) that F is holomorphic if and only if $\mathfrak{n}\vec{\Phi} = 0$. Now use (48). $\square$

3.3. The differential operators in classical language. Let ℓ be any integer, and m a non-negative integer. Let Φ be a K -finite complex-valued function on $\mathrm{Sp}_4(\mathbb{R})$ of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$. Let $F : \mathbb{H}_2 \rightarrow W_m$ be the function corresponding to Φ according to Lemma 3.2. Let X be one of the operators defined in Table 1, and set $\Psi = X\Phi$. Then Ψ is a K -finite function satisfying $N_+ \Psi = 0$, of weight indicated in Table 1. Hence, according to Lemma 3.2, there exists a vector-valued function G corresponding to Ψ . In this section, we write down G in terms of F for all elements X defined in Table 1. The proofs rely on our formulas for the action of the root vectors (Proposition 3.3). The calculations involved, while long and detailed, are essentially routine, and therefore omitted.

Going down. We start with $X = P_{0-}$. Hence, let $\Psi = P_{0-}\Phi$. Then Ψ has weight $(\ell + m, \ell - 2)$ and satisfies $N_+ \Psi = 0$. Let $G : \mathbb{H}_2 \rightarrow W_{m+2}$ be the function corresponding to Ψ according to Lemma 3.2. The following diagram illustrates the situation.

$$\begin{array}{ccccc} \text{(weight } (\ell + m, \ell)) & \Phi & \longrightarrow & \vec{\Phi} & \longrightarrow F & \text{(values in } W_m) \\ & P_{0-} \downarrow & & & & \\ \text{(weight } (\ell + m, \ell - 2)) & \Psi & \longrightarrow & \vec{\Psi} & \longrightarrow G & \text{(values in } W_{m+2}) \end{array} \quad (65)$$

Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0, \dots, G_{m+2}$ be the component functions of G ; see (51). We define three differential operators on $\mathbb{H}_2$,

$$\bar{\partial}_0 = 2i \left(v^2 \frac{\partial}{\partial \bar{\tau}} + vy' \frac{\partial}{\partial \bar{z}} + y'^2 \frac{\partial}{\partial \bar{\tau}'} \right), \quad (66)$$

$$\bar{\partial}_1 = -2i \left(2vy \frac{\partial}{\partial \bar{\tau}} + (yy' + v^2) \frac{\partial}{\partial \bar{z}} + 2vy' \frac{\partial}{\partial \bar{\tau}'} \right), \quad (67)$$

$$\bar{\partial}_2 = 2i \left(y^2 \frac{\partial}{\partial \bar{\tau}} + vy \frac{\partial}{\partial \bar{z}} + v^2 \frac{\partial}{\partial \bar{\tau}'} \right). \quad (68)$$

The following result expresses the G_j in terms of the F_i .

Proposition 3.5. *With the above notations,*

$$G_j = -(\bar{\partial}_2 F_{j-2} + \bar{\partial}_1 F_{j-1} + \bar{\partial}_0 F_j) \quad (69)$$

for $j = 0, \dots, m+2$. (We understand $F_i = 0$ for $i < 0$ or $i > m$.)

Going left. Next we write down the effect of the operator L , whose defining formula is given in Table 1. In order for L to be defined, we assume $m \geq 2$. Let $\Psi = L\Phi$. Then Ψ has weight $(\ell + m - 2, \ell)$, and $N_+\Psi = 0$. Let $G : \mathbb{H}_2 \rightarrow W_{m-2}$ be the function corresponding to Ψ . Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0, \dots, G_{m-2}$ be the component functions of G ; see (51).

Proposition 3.6. *With the above notations,*

$$\begin{aligned} G_j = & -(m-j)(m-j-1)\bar{\partial}_2 F_j \\ & + (m-j-1)(j+1)\bar{\partial}_1 F_{j+1} \\ & - (j+2)(j+1)\bar{\partial}_0 F_{j+2}. \end{aligned} \quad (70)$$

for $j = 0, \dots, m-2$.

Going up. Next, we consider U , whose defining formula is given in Table 1. We will assume $m \geq 2$, so that U is well-defined. Let $\Psi = U\Phi$. Then Ψ has weight $(\ell + m, \ell + 2)$, and $N_+\Psi = 0$. Let $G : \mathbb{H}_2 \rightarrow W_{m-2}$ be the function corresponding to Ψ . Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0, \dots, G_{m-2}$ be the component functions of G ; see (51). The following result expresses the G_j in terms of the F_i .

Proposition 3.7. *With the above notations,*

$$\begin{aligned} G_j = & (m-j)(m-j-1)\left((\ell-1)\frac{y}{\Delta} + 2i\frac{\partial}{\partial\tau'}\right)F_j \\ & + (m-j-1)(j+1)\left((\ell-1)\frac{2v}{\Delta} - 2i\frac{\partial}{\partial z}\right)F_{j+1} \\ & + (j+2)(j+1)\left((\ell-1)\frac{y'}{\Delta} + 2i\frac{\partial}{\partial\tau}\right)F_{j+2} \end{aligned} \quad (71)$$

for $j = 0, \dots, m-2$.

Going right. Next we calculate the effect of the operator X_+ . Let $\Psi = X_+\Phi$. Then Ψ has weight $(\ell + m + 2, \ell)$, and $N_+\Psi = 0$. Let $G : \mathbb{H}_2 \rightarrow W_{m+2}$ be the function corresponding to Ψ . Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0, \dots, G_{m+2}$ be the component functions of G ; see (51).

Proposition 3.8. *With the above notations,*

$$\begin{aligned} G_j = & \left((\ell+m)\frac{y}{\Delta} + 2i\frac{\partial}{\partial\tau'}\right)F_{j-2} \\ & - \left((\ell+m)\frac{2v}{\Delta} - 2i\frac{\partial}{\partial z}\right)F_{j-1} \\ & + \left((\ell+m)\frac{y'}{\Delta} + 2i\frac{\partial}{\partial\tau}\right)F_j \end{aligned} \quad (72)$$

for $j = 0, \dots, m+2$.

Remark 3.9. The operator X_+ is the same as the operator $\delta_{\ell+m}$ occurring in [4].

The degree 1 diagonal operators. Let $\Psi^\pm = E_\pm\Phi$. Then $\Psi^\pm$ has weight $(\ell + m \pm 1, \ell \pm 1)$, and $N_+\Psi^\pm = 0$. Let $G^\pm : \mathbb{H}_2 \rightarrow W_m$ be the function corresponding to $\Psi^\pm$. Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0^\pm, \dots, G_m^\pm$ be the component functions of $G^\pm$; see (51). The following result expresses the $G_j^\pm$ in terms of the F_i .

Proposition 3.10. *With the above notations,*

$$\begin{aligned} G_j^+ = & (m-j+1)\left((2\ell+m-2)\frac{y}{\Delta} + 4i\frac{\partial}{\partial\tau'}\right)F_{j-1} \\ & + (m-2j)\left(-(2\ell+m-2)\frac{v}{\Delta} + 2i\frac{\partial}{\partial z}\right)F_j \end{aligned}$$

$$-(j+1)\left((2\ell+m-2)\frac{y'}{\Delta}+4i\frac{\partial}{\partial\tau}\right)F_{j+1}, \quad (73)$$

$$G_j^- = 2(m+1-j)\bar{\partial}_2 F_{j-1} + (m-2j)\bar{\partial}_1 F_j - 2(j+1)\bar{\partial}_0 F_{j+1}, \quad (74)$$

for $j = 0, \dots, m$. (We understand $F_i = 0$ for $i < 0$ or $i > m$.) The differential operators $\bar{\partial}_0, \bar{\partial}_1, \bar{\partial}_2$ are the ones defined in (66) – (68).

The degree 2 diagonal operators. Let $\Psi^\pm = D_\pm \Phi$. Then Ψ has weight $(\ell + m \pm 2, \ell \pm 2)$, and $N_+ \Psi^\pm = 0$. Let $G^\pm : \mathbb{H}_2 \rightarrow W_m$ be the function corresponding to $\Psi^\pm$. Let $F_0, \dots, F_m$ be the component functions of F , and let $G_0^\pm, \dots, G_m^\pm$ be the component functions of $G^\pm$; see (51). For scalar-valued or vector-valued functions on $\mathbb{H}_2$, we define the differential operator

$$\bar{\partial}_3 = 2i\left(y\frac{\partial}{\partial\bar{\tau}} + v\frac{\partial}{\partial\bar{z}} + y'\frac{\partial}{\partial\bar{\tau}'}\right). \quad (75)$$

The following result expresses the $G_j^\pm$ in terms of the F_i .

Proposition 3.11. *With the above notations,*

$$\begin{aligned} G_j^+ &= (m-j+1)(m-j+2)\frac{y^2}{\Delta^2}F_{j-2} \\ &+ \left(4i(m-j+1)\left(\frac{y}{\Delta}\frac{\partial}{\partial z} + \frac{2v}{\Delta}\frac{\partial}{\partial\tau'}\right) - 2(m-2j+1)(m-j+1)\frac{vy}{\Delta^2}\right)F_{j-1} \\ &+ \left[(4j^2 + 4l^2 - 4jm + m(m-3) + l(4m-2))\frac{v^2}{\Delta^2} \right. \\ &\quad + (j^2 - jm - (2l-1)(l+m))\frac{2yy'}{\Delta^2} + 16\frac{\partial^2}{\partial\tau\partial\tau'} - 4\frac{\partial^2}{\partial z^2} \\ &\quad \left. - 4i\left((2j+2l-1)\frac{y}{\Delta}\frac{\partial}{\partial\tau} + (m+2l-1)\frac{v}{\Delta}\frac{\partial}{\partial z} + (2m-2j+2l-1)\frac{y'}{\Delta}\frac{\partial}{\partial\tau'}\right)\right]F_j \\ &+ \left(4i(j+1)\left(\frac{2v}{\Delta}\frac{\partial}{\partial\tau} + \frac{y'}{\Delta}\frac{\partial}{\partial z}\right) + 2(m-2j-1)(j+1)\frac{vy'}{\Delta^2}\right)F_{j+1} \\ &+ (j+1)(j+2)\frac{y'^2}{\Delta^2}F_{j+2}, \end{aligned} \quad (76)$$

$$G_j^- = \left(4\Delta^2\left(4\frac{\partial}{\partial\bar{\tau}}\frac{\partial}{\partial\bar{\tau}'} - \frac{\partial^2}{\partial\bar{z}^2}\right) - 2\Delta\bar{\partial}_3\right)F_j, \quad (77)$$

for $j = 0, \dots, m$.

Remark 3.12. The formula for D_+ in the special case that $m = 0$ (the scalar valued case) is given by

$$D_+F = \left(-\frac{2l(2l-1)}{\Delta} - \frac{4i(2l-1)}{\Delta}\left(y\frac{\partial}{\partial\tau} + v\frac{\partial}{\partial z} + y'\frac{\partial}{\partial\tau'}\right) + 4\left(4\frac{\partial^2}{\partial\tau\partial\tau'} - \frac{\partial^2}{\partial z^2}\right)\right)F. \quad (78)$$

In this case, the operator D_+ was originally defined by Maass in his book [25].

3.4. Nearly holomorphic functions. Let p be a non-negative integer. We will write elements $Z \in \mathbb{H}_2$ as $Z = X + iY$ with real X and Y . By definition of $\mathbb{H}_2$, the real symmetric matrix Y is positive definite. We let $N^p(\mathbb{H}_2)$ denote the space of all polynomials of degree $\leq p$ in the entries of Y^{-1} with holomorphic functions on $\mathbb{H}_2$ as coefficients. The space

$$N(\mathbb{H}_2) = \bigcup_{p \geq 0} N^p(\mathbb{H}_2)$$

is the space of *nearly holomorphic functions* on $\mathbb{H}_2$. Evidently, $N(\mathbb{H}_2)$ is a ring, and we have $N^p(\mathbb{H}_2)N^q(\mathbb{H}_2) \subset N^{p+q}(\mathbb{H}_2)$. For convenience, we let $N^p(\mathbb{H}_2) = 0$ for negative p . If $f \in N(\mathbb{H}_2)$ lies in $N^p(\mathbb{H}_2)$ but not in $N^{p-1}(\mathbb{H}_2)$, we say that f has *nearly holomorphic degree* p . Note that $N^0(\mathbb{H}_2)$ is the space of holomorphic functions on $\mathbb{H}_2$.

As before, we will use the coordinates (52) on $\mathbb{H}_2$, and set $\Delta = yy' - v^2$. The entries of Y^{-1} are then y/Δ , v/Δ and y'/Δ . Since $\frac{y}{\Delta}\frac{y'}{\Delta} - \frac{v^2}{\Delta^2} = \frac{1}{\Delta}$, the function $\frac{1}{\Delta}$ is a nearly holomorphic function. For a typical nearly holomorphic monomial we will use the notation

$$[\alpha, \beta, \gamma] := \left(\frac{y}{\Delta}\right)^\alpha \left(\frac{v}{\Delta}\right)^\beta \left(\frac{y'}{\Delta}\right)^\gamma; \quad (79)$$

here, α, β, γ are non-negative integers.

We may ask how the various differential operators we defined in previous sections behave with respect to nearly holomorphic functions. It is easy to see that the basic partial derivatives

$$\frac{\partial}{\partial \tau}, \quad \frac{\partial}{\partial z}, \quad \frac{\partial}{\partial \tau'}, \quad \frac{\partial}{\partial \bar{\tau}}, \quad \frac{\partial}{\partial \bar{z}}, \quad \frac{\partial}{\partial \bar{\tau}'}$$

map $N^p(\mathbb{H}_2)$ to $N^{p+1}(\mathbb{H}_2)$. The following lemma gives the action of differential operators, including those defined in (66) – (68) and (75), on a nearly holomorphic monomial. In particular, the lemma shows that the operators $\bar{\partial}_0, \bar{\partial}_1, \bar{\partial}_2$ act as “nearly holomorphic derivatives”.

Lemma 3.13. *The following formulas hold for all non-negative integers α, β, γ .*

$$\bar{\partial}_0[\alpha, \beta, \gamma] = \alpha[\alpha - 1, \beta, \gamma], \quad (80)$$

$$\bar{\partial}_1[\alpha, \beta, \gamma] = \beta[\alpha, \beta - 1, \gamma], \quad (81)$$

$$\bar{\partial}_2[\alpha, \beta, \gamma] = \gamma[\alpha, \beta, \gamma - 1], \quad (82)$$

$$\bar{\partial}_3[\alpha, \beta, \gamma] = (\alpha + \beta + \gamma)[\alpha, \beta, \gamma], \quad (83)$$

$$D_-[\alpha, \beta, \gamma] = \beta(\beta - 1)[\alpha, \beta - 2, \gamma] - 4\alpha\gamma[\alpha - 1, \beta, \gamma - 1]. \quad (84)$$

Proof. Everything follows from direct calculations. $\square$

As a consequence, we note that the operators $\bar{\partial}_0, \bar{\partial}_1, \bar{\partial}_2$ commute on $N(\mathbb{H}_2)$ (they do not commute on all of $C^\infty(\mathbb{H}_2)$).

Lemma 3.14. *Assume that $F = \sum_{\alpha, \beta, \gamma \geq 0} [\alpha, \beta, \gamma] F_{\alpha, \beta, \gamma}$ is a nearly holomorphic function, where the $F_{\alpha, \beta, \gamma}$ are holomorphic. Then F is zero if and only if all $F_{\alpha, \beta, \gamma}$ are zero.*

Proof. This can be proved by induction on the nearly holomorphic degree, using the formulas (80) – (82). $\square$

Operators on vector-valued functions. Let ℓ be any integer, and m a non-negative integer. Let $C_{\ell, m}^\infty(\mathbb{H}_2)$ be the space of smooth functions $F : \mathbb{H}_2 \rightarrow W_m$. Note that this space does not actually depend on ℓ ; nevertheless, it will be useful to carry this subindex along (the significance of this subindex will be seen in the next chapter, when we will restrict to the subspace of $C_{\ell, m}^\infty(\mathbb{H}_2)$ consisting of forms F which transform via $\eta_{\ell, m}$ with respect to some congruence subgroup).

For each of the operators X appearing in Table 1 we will define a linear map $X : C_{\ell, m}^\infty(\mathbb{H}_2) \rightarrow C_{\ell_1, m_1}^\infty(\mathbb{H}_2)$, where $(\ell_1 + m_1, \ell_1)$ is the “new weight” given in Table 1. Some of the operators X will depend on ℓ , (or m , or both) but, as before, our notation will not reflect this dependence.

If $m < 2$, we set $U = L = 0$. In all other cases, the definitions will be in terms of the component functions $F_0, \dots, F_m$ of F given by $F(Z) = \sum_{j=0}^m F_j(Z) S^{m-j} T^j$, and are as follows.

$$(P_0 F)_j = \text{right hand side of (69)}, \quad (85)$$

$$(L F)_j = \text{right hand side of (70)}, \quad (86)$$

$$(U F)_j = \text{right hand side of (71)}, \quad (87)$$

$$(X_+ F)_j = \text{right hand side of (72)}, \quad (88)$$

$$(E_+ F)_j = \text{right hand side of (73)}, \quad (89)$$

$$(E_- F)_j = \text{right hand side of (74)}, \quad (90)$$

$$(D_+ F)_j = \text{right hand side of (76)}, \quad (91)$$

$$(D_- F)_j = \text{right hand side of (77)}. \quad (92)$$

These formulas hold for *all* $j \in \mathbb{Z}$, but the expressions on the right hand sides are automatically zero if $j < 0$ or $j > m_1$.

For a non-negative integer p , let $N_{\ell,m}^p(\mathbb{H}_2)$ be the subspace of $C_{\ell,m}^\infty(\mathbb{H}_2)$ consisting of those F for which all component functions F_j are in $N^p(\mathbb{H}_2)$. Hence, these are nearly holomorphic W_m -valued functions. The space $N_{\ell,m}^0(\mathbb{H}_2)$ consists of the holomorphic W_m -valued functions.

TABLE 2. Let X be one of the operators given in the first column. Let $F \in N_{\ell,m}^p(\mathbb{H}_2)$. Then $XF \in N_{\ell_1,m_1}^{p_1}(\mathbb{H}_2)$, with ℓ_1, m_1, p_1 given in the last three columns of the table. The second column indicates the direction from the old weight $(\ell + m, \ell)$ to the new weight $(\ell_1 + m_1, \ell_1)$, assuming F corresponds to the K -finite function $\Phi : \text{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ of weight $(\ell + m, \ell)$. If $m < 2$, then by definition, $U = L = 0$.

operator	direction	new ℓ	new m	new p
P_{0-}	$\downarrow$	$\ell - 2$	$m + 2$	$p - 1$
L	$\leftarrow$	ℓ	$m - 2$	$p - 1$
U	$\uparrow$	$\ell + 2$	$m - 2$	$p + 1$
X_+	$\rightarrow$	ℓ	$m + 2$	$p + 1$
E_+	$\nearrow$	$\ell + 1$	m	$p + 1$
E_-	$\swarrow$	$\ell - 1$	m	$p - 1$
D_+	$\nearrow$	$\ell + 2$	m	$p + 2$
D_-	$\swarrow$	$\ell - 2$	m	$p - 2$

Proposition 3.15. *Let ℓ be any integer, and m a non-negative integer. Let X be one of the operators in Table 1. Let $F \in C_{\ell,m}^\infty(\mathbb{H}_2)$.*

- (1) *Assume that F corresponds, via Lemma 3.2, to the K -finite function Φ on $\text{Sp}_4(\mathbb{R})$ of weight $(\ell + m, \ell)$ satisfying $N_+ \Phi = 0$. Then XF corresponds to $X\Phi$. In other words, the diagram*

$$\begin{array}{ccc} \mathcal{V}_{\ell,m} & \xrightarrow{\sim} & C_{\ell,m}^\infty(\mathbb{H}_2) \\ X \downarrow & & \downarrow X \\ \mathcal{V}_{\ell_1,m_1} & \xrightarrow{\sim} & C_{\ell_1,m_1}^\infty(\mathbb{H}_2) \end{array} \quad (93)$$

is commutative. Here, ℓ_1, m_1 are given in Table 2, and the horizontal isomorphisms are those from Lemma 3.2.

- (2) *If $F \in N_{\ell,m}^p(\mathbb{H}_2)$, then $XF \in N_{\ell_1,m_1}^{p_1}(\mathbb{H}_2)$, where ℓ_1, m_1, p_1 are given in the last three columns of Table 2.*

Proof. (1) simply summarizes the content of Propositions 3.5, 3.6, 3.7, 3.8, 3.10 and 3.11.

(2) follows from the formulas (85) – (92), together with Lemma 3.13. $\square$

We see from (2) of this result that if we walk in the direction of one of the roots in $\mathfrak{n}$, then the nearly holomorphic degree decreases, while if we walk in the direction of one of the roots in $\mathfrak{p}_+$, then the nearly holomorphic degree (potentially) increases. In the next section, we will use the following holomorphy criterion to prove that spaces of nearly holomorphic modular forms are finite-dimensional.

Lemma 3.16. *Let ℓ be any integer, and m a non-negative integer. Let $F \in C_{\ell,m}^\infty(\mathbb{H}_2)$. Let $p \in \{0, 1\}$.*

(1) *If $m = 0$, then the following are equivalent:*

- (a) $F \in N^p(\mathbb{H}_2)$.
- (b) $P_0 F \in N^{p-1}(\mathbb{H}_2)$.

In particular, F is holomorphic if and only if $P_0 F = 0$.

(2) *If $m = 1$, then the following are equivalent:*

- (a) $F \in N^p(\mathbb{H}_2)$.
- (b) $P_0 F, E_- F \in N^{p-1}(\mathbb{H}_2)$.

In particular, F is holomorphic if and only if $P_0 F = E_- F = 0$.

(3) *If $m \geq 2$, then the following are equivalent:*

- (a) $F \in N^p(\mathbb{H}_2)$.
- (b) $P_0 F, E_- F, L F \in N^{p-1}(\mathbb{H}_2)$.

In particular, F is holomorphic if and only if $P_0 F = E_- F = L F = 0$.

Proof. These properties can be verified from the explicit formulas for the operators written above. The details are omitted. $\square$

4. THE STRUCTURE THEOREMS

4.1. Modular forms. Recall that, for a positive integer N , the *principal congruence subgroup* $\Gamma(N)$ consists of all elements of $\mathrm{Sp}_4(\mathbb{Z})$ that are congruent to the identity matrix modulo N . A *congruence subgroup* of $\mathrm{Sp}_4(\mathbb{Q})$ is a subgroup that, for some N , contains $\Gamma(N)$ with finite index. The reason that we do not restrict ourselves to subgroups of $\mathrm{Sp}_4(\mathbb{Z})$ is that we would like to include groups like the paramodular group.

Let ℓ be an integer, and m a non-negative integer. Recall from Sect. 3.1 that $\eta_{\ell,m}$ denotes the $(m+1)$ -dimensional representation $\det^\ell \mathrm{sym}^m$ of $\mathrm{GL}_2(\mathbb{C})$. As before, let $C_{\ell,m}^\infty(\mathbb{H}_2)$ be the space of smooth W_m -valued functions on $\mathbb{H}_2$. We define a right action of $\mathrm{Sp}_4(\mathbb{R})$ on $C_{\ell,m}^\infty(\mathbb{H}_2)$ by

$$(F|_{\ell,m} g)(Z) = \eta_{\ell,m}(J(g, Z))^{-1} F(gZ) \quad \text{for } g \in \mathrm{Sp}_4(\mathbb{R}), Z \in \mathbb{H}_2. \quad (94)$$

In the following we fix a congruence subgroup Γ of $\mathrm{Sp}_4(\mathbb{Q})$. Let $C_{\ell,m}^\infty(\Gamma)$ be the space of smooth functions $F : \mathbb{H}_2 \rightarrow W_m$ satisfying

$$F|_{\ell,m} \gamma = F \quad \text{for all } \gamma \in \Gamma. \quad (95)$$

It is easy to see that $F \in C_{\ell,m}^\infty(\mathbb{H}_2)$ has this transformation property if and only if the function $\Phi \in \mathcal{V}_{\ell,m}$ corresponding to F via Lemma 3.2 satisfies $\Phi(\gamma g) = \Phi(g)$ for all $g \in \mathrm{Sp}_4(\mathbb{R})$ and $\gamma \in \Gamma$. Let $\mathcal{V}_{\ell,m}(\Gamma)$ be the subspace of $\mathcal{V}_{\ell,m}$ consisting of Φ with this transformation property. If X is one of the operators in Table 1, then it follows from Proposition 3.15 that there is a commutative diagram

$$\begin{array}{ccc} \mathcal{V}_{\ell,m}(\Gamma) & \xrightarrow{\sim} & C_{\ell,m}^\infty(\Gamma) \\ X \downarrow & & \downarrow X \\ \mathcal{V}_{\ell_1,m_1}(\Gamma) & \xrightarrow{\sim} & C_{\ell_1,m_1}^\infty(\Gamma) \end{array} \quad (96)$$

Here, ℓ_1, m_1 are the integers given in Table 2. (One could verify directly that if F satisfies (95), then XF satisfies $(XF)|_{\ell_1, m_1} \gamma = F$ for all $\gamma \in \Gamma$, but the use of the diagrams is much easier.)

More generally, one has the following basic commutation relation.

Lemma 4.1. *Let $\mathcal{X}$ be the free monoid consisting of all (finite) strings of the symbols in the left column of Table 1. Suppose that X is an element of $\mathcal{X}$ and let (ℓ_1, m_1) be the integers (uniquely determined by ℓ, m and X) such that X takes $C_{\ell, m}^\infty(\Gamma)$ to $C_{\ell_1, m_1}^\infty(\Gamma)$. Let $\gamma \in \mathrm{Sp}_4(\mathbb{R})$. Then, for all $F \in C_{\ell, m}^\infty(\mathbb{H}_2)$, we have $(XF)|_{\ell_1, m_1} \gamma = X(F|_{\ell, m} \gamma)$.*

Proof. Let Φ be the function corresponding to F via Lemma 3.2. Then it follows from Proposition 3.15 that $X\Phi$ corresponds to XF . On the other hand, the operation $|_{\ell_1, m_1} \gamma$ corresponds to left multiplication of the argument by γ . Define a function Φ_1 on $\mathrm{Sp}_4(\mathbb{R})$ by $\Phi_1(g) = \Phi(\gamma g)$. Now the proof follows from the obvious identity $(X\Phi)(\gamma g) = (X\Phi_1)(g)$. $\square$

Fourier expansions. Now let $F \in C_{\ell, m}^\infty(\Gamma) \cap N_{\ell, m}^p(\mathbb{H}_2)$. Hence, F is nearly holomorphic and satisfies (95). Let $F_0, \dots, F_m$ be the component functions of F , as defined in (51). Suppose F_j is written as $F_j = \sum_{\alpha, \beta, \gamma} [\alpha, \beta, \gamma] F_{j, \alpha, \beta, \gamma}$ with holomorphic functions $F_{j, \alpha, \beta, \gamma}$; see (79) for notation. Since F is invariant under the translations $\tau \mapsto \tau + N$, $z \mapsto z + N$ and $\tau' \mapsto \tau' + N$ for some positive integer N , the same is true for F_j and each $F_{j, \alpha, \beta, \gamma}$; observe here Lemma 3.14. Thus $F_{j, \alpha, \beta, \gamma}$ admits a Fourier expansion

$$F_{j, \alpha, \beta, \gamma}(Z) = \sum_Q a_{j, \alpha, \beta, \gamma}(Q) e^{2\pi i \mathrm{Tr}(QZ)}, \quad (97)$$

where Q runs over matrices $\begin{bmatrix} a & b/2 \\ b/2 & c \end{bmatrix}$ with $a, b, c \in \frac{1}{N}\mathbb{Z}$. It follows that F_j admits a Fourier expansion

$$F_j(Z) = \sum_Q a_j(Q) e^{2\pi i \mathrm{Tr}(QZ)}, \quad a_j(Q) := \sum_{\alpha, \beta, \gamma} a_{j, \alpha, \beta, \gamma}(Q) [\alpha, \beta, \gamma], \quad (98)$$

and that F admits a Fourier expansion

$$F(Z) = \sum_{Q \in M_2^{\mathrm{sym}}(\mathbb{Q})} a(Q) e^{2\pi i \mathrm{Tr}(QZ)}, \quad (99)$$

where

$$a(Q) = \sum_{j=0}^m \sum_{\alpha, \beta, \gamma} a_{j, \alpha, \beta, \gamma}(Q) [\alpha, \beta, \gamma] S^{m-j} T^j. \quad (100)$$

Thus, the Fourier coefficients of F are polynomial functions in the entries of Y^{-1} taking values in W_m . For fixed Q , the complex-valued functions $a_j(Q)$ in (98) are nothing but the component functions of $a(Q)$. If X is one of the operators defined in (85), (86), (90) or (92), and if F has Fourier expansion (99), then XF has Fourier expansion

$$(XF)(Z) = \sum_Q (Xa(Q)) e^{2\pi i \mathrm{Tr}(QZ)}. \quad (101)$$

This follows directly from the definitions and the fact that $e^{2\pi i \mathrm{Tr}(QZ)}$ is holomorphic for all matrices Q . If X is one of the operators defined in (87), (88), (89) or (91), then the Fourier expansion of XF is more complicated. However, it is easy to see that

$$(XF)(Z) = \sum_Q b(Q) e^{2\pi i \mathrm{Tr}(QZ)}, \quad \text{with } b(Q) = 0 \text{ if } a(Q) = 0. \quad (102)$$

Hence, none of the eight operators introduces any “new” Fourier coefficients.

Nearly holomorphic modular forms. Let ℓ be an integer, and m, p be non-negative integers. For a congruence subgroup Γ , let $N_{\ell, m}^p(\Gamma)$ be the space of all functions $F : \mathbb{H}_2 \rightarrow W_m$ with the following properties.

- (1) $F \in N_{\ell, m}^p(\mathbb{H}_2)$.
- (2) F satisfies the transformation property (95).
- (3) F satisfies the “no poles at cusps” condition. This means: For any $g \in \mathrm{Sp}_4(\mathbb{Q})$ the function $F|_{\ell, m} g$ admits a Fourier expansion of the form (99) such that $a(Q) = 0$ unless Q is positive semidefinite.

Let $N_{\ell, m}(\Gamma) = \bigcup_{p \geq 0} N_{\ell, m}^p(\Gamma)$. We refer to $N_{\ell, m}(\Gamma)$ as the space of *nearly holomorphic Siegel modular forms* of weight $\det^\ell \mathrm{sym}^m$ with respect to Γ . We sometimes write $M_{\ell, m}(\Gamma)$ for $N_{\ell, m}^0(\Gamma)$; this is the usual space of holomorphic vector-valued Siegel modular forms taking values in $\eta_{\ell, m}$.

An element $F \in N_{\ell, m}(\Gamma)$ is called a *cusp form* if it vanishes at all cusps. By definition, this means: For any $g \in \mathrm{Sp}_4(\mathbb{Q})$ the function $F|_{\ell, m} g$ admits a Fourier expansion of the form (99), for some N , such that $a(Q) = 0$ unless Q is positive definite. We write $N_{\ell, m}(\Gamma)^\circ$ for the subspace of cusp forms. Let $N_{\ell, m}^p(\Gamma)^\circ = N_{\ell, m}(\Gamma)^\circ \cap N_{\ell, m}^p(\Gamma)$. We sometimes write $S_{\ell, m}(\Gamma)$ for $N_{\ell, m}^0(\Gamma)^\circ$; this is the usual space of holomorphic vector-valued Siegel cusp forms taking values in $\eta_{\ell, m}$.

Lemma 4.2. *The spaces $N_{\ell, m}^p(\Gamma)$ and $N_{\ell, m}^p(\Gamma)^\circ$ are finite-dimensional.*

Proof. Obviously, we only need to prove this for $N_{\ell, m}^p(\Gamma)$. It is well known, and can be proved using Harish-Chandra’s general finiteness result stated as Theorem 1.7 in [9], that the statement is true for $p = 0$, i.e., for holomorphic modular forms. Assume that $p > 0$. If $m = 0$, then, by (1) of Lemma 3.16, the map $F \mapsto P_{0-} F$ gives rise to an exact sequence

$$0 \longrightarrow M_{\ell, m}(\Gamma) \longrightarrow N_{\ell, m}^p(\Gamma) \longrightarrow N_{\ell-2, m+2}^{p-1}(\Gamma).$$

If $m = 1$, then, by (2) of Lemma 3.16, the map $F \mapsto (P_{0-} F, E_- F)$ gives rise to an exact sequence

$$0 \longrightarrow M_{\ell, m}(\Gamma) \longrightarrow N_{\ell, m}^p(\Gamma) \longrightarrow N_{\ell-2, m+2}^{p-1}(\Gamma) \oplus N_{\ell-1, m}^{p-1}(\Gamma).$$

If $m \geq 2$, then, by (3) of Lemma 3.16, the map $F \mapsto (P_{0-} F, E_- F, L F)$ gives rise to an exact sequence

$$0 \longrightarrow M_{\ell, m}(\Gamma) \longrightarrow N_{\ell, m}^p(\Gamma) \longrightarrow N_{\ell-2, m+2}^{p-1}(\Gamma) \oplus N_{\ell-1, m}^{p-1}(\Gamma) \oplus N_{\ell, m-2}^{p-1}(\Gamma).$$

Hence our assertion follows by induction on p . $\square$

4.2. Automorphic forms. Let Γ be a congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$. We denote by $\mathcal{A}(\Gamma)$ the space of automorphic forms on $\mathrm{Sp}_4(\mathbb{R})$ with respect to Γ . Recall that an automorphic form is a smooth function on $\mathrm{Sp}_4(\mathbb{R})$ that is left Γ -invariant, $\mathcal{Z}$ -finite, K -finite and slowly increasing; here $\mathcal{Z}$ is the center of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$. Let $\mathcal{A}(\Gamma)^\circ$ be the subspace of cuspidal automorphic forms. We refer to [9] for precise definitions of these notions. The spaces $\mathcal{A}(\Gamma)$ and $\mathcal{A}(\Gamma)^\circ$ are $(\mathfrak{g}, K)$ -modules under right translation.

Let dg be any Haar measure on $\mathrm{Sp}_4(\mathbb{R})$. For Φ_1 and Φ_2 in $\mathcal{A}(\Gamma)$, we define the integral

$$\langle \Phi_1, \Phi_2 \rangle := \frac{1}{\mathrm{vol}(\Gamma \backslash \mathrm{Sp}_4(\mathbb{R}))} \int_{\Gamma \backslash \mathrm{Sp}_4(\mathbb{R})} \Phi_1(g) \overline{\Phi_2(g)} dg \quad (103)$$

whenever it is *absolutely convergent*. This happens, for example, whenever at least one of Φ_1 and Φ_2 lies in $\mathcal{A}(\Gamma)^\circ$. In particular, $\langle, \rangle$ defines an inner product on $\mathcal{A}(\Gamma)^\circ$ invariant under right translations by $\mathrm{Sp}_4(\mathbb{R})$. For an element $X \in \mathfrak{g}$, we have

$$\langle X \Phi_1, \Phi_2 \rangle + \langle \Phi_1, X \Phi_2 \rangle = 0.$$

By general principles (see [9] and the references therein) $\mathcal{A}(\Gamma)^\circ$ decomposes into an orthogonal direct sum of irreducible $(\mathfrak{g}, K)$ -modules, each occurring with finite multiplicity.

Let $\lambda = (k, \ell)$ be an element of the weight lattice Λ . We say that $\Phi \in \mathcal{A}(\Gamma)$ has weight λ if $Z\Phi = k\Phi$ and $Z'\Phi = \ell\Phi$ (right translation action). Let $\mathcal{A}_\lambda(\Gamma)$ be the subspace of $\mathcal{A}(\Gamma)$ consisting of elements of weight λ , and let $\mathcal{A}_\lambda(\Gamma)^\circ$ be similarly defined.

Let $\mathfrak{n} \subset \mathfrak{g}$ be the span of the root vectors X_- , P_{1-} and P_{0-} . Then $\mathcal{U}(\mathfrak{n})$ is the polynomial algebra in the three variables X_- , P_{1-} and P_{0-} . An automorphic form Φ is called $\mathfrak{n}$ -finite if the space $\mathcal{U}(\mathfrak{n})\Phi$ is finite-dimensional. We denote by $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ the space of $\mathfrak{n}$ -finite automorphic forms, and by $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ the subspace of cusp forms. The following properties are easy to verify:

- $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is a $(\mathfrak{g}, K)$ -submodule of $\mathcal{A}(\Gamma)$.
- $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is the direct sum of its K -types, i.e.: If $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ and $\Phi = \Phi_1 + \dots + \Phi_m$, where Φ_i lies in the ρ_i -isotypical component of $\mathcal{A}(\Gamma)$ for different K -types ρ_i , then $\Phi_i \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ for each i .

Analogous statements hold for cusp forms.

Lemma 4.3. *$\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is an admissible $(\mathfrak{g}, K)$ -module.*

Proof. Assume that a K -type ρ_λ occurs infinitely often in $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ for some $\lambda = (\ell + m, \ell)$. We may assume that λ is maximal in the order (7). Let W be the space of highest weight vectors in the ρ_λ -isotypical component; by assumption, W is infinite-dimensional. By our maximality assumption, the kernel W_1 of P_{0-} on W is infinite-dimensional; note that N_+ commutes with P_{0-} . Similarly, the kernel W_2 of P_{1-} on W_1 is infinite-dimensional. Finally, the kernel W_3 of X_- on W_2 is infinite-dimensional. The vectors in W_3 correspond to holomorphic modular forms in $M_{\ell, m}(\Gamma)$. Since this space is finite-dimensional, we obtain a contradiction. $\square$

Modular forms and automorphic forms. We are going to prove that nearly holomorphic modular forms generate $\mathfrak{n}$ -finite automorphic forms. The following lemma will be useful.

Lemma 4.4. *Let V be a $\mathfrak{g}_\mathbb{C}$ -module, and $v_0 \in V$ a vector with the following properties:*

- $V = \mathcal{U}(\mathfrak{g}_\mathbb{C})v_0$.
- v_0 has weight $(\ell + m, \ell)$ for some integer ℓ and non-negative integer m .
- $N_+v_0 = 0$.
- $N_-^r v_0 = 0$ for some $r > 0$.
- $P_{0-}^s v_0 = 0$ for some $s > 0$.
- $D_-^t v_0 = 0$ for some $t > 0$.

Then v_0 is $\mathfrak{n}$ -finite, and V is an admissible $(\mathfrak{g}, K)$ -module in which each weight space is finite-dimensional.

Proof. Let $X = X_-$, $Y = P_{1-}$ and $Z = P_{0-}$, so that $\mathcal{U}(\mathfrak{n})$ is the polynomial ring $\mathbb{C}[X, Y, Z]$. In this ring, let I be the ideal generated by $D_-^t = (Y^2 - 4XZ)^t$ and Z^s . By our hypothesis, every element of I annihilates v_0 .

In affine three-space, consider the vanishing set $N(I)$. Clearly, a point (x, y, z) in $N(I)$ must have $y = z = 0$. Since the polynomial Y vanishes on all of $N(I)$, we have $Y^n \in I$ for some positive integer n by Hilbert's Nullstellensatz.

By the PBW theorem, $\mathcal{U}(\mathfrak{g}_\mathbb{C})$ is spanned by monomials of the form

$$(\text{monomial in } X_-, N_-, P_{0+}, P_{1+}, X_+, Z, Z') \times P_{1-}^\alpha P_{0-}^\beta N_+^\gamma$$

with $\alpha, \beta, \gamma \geq 0$. Since P_{1-} , P_{0-} , N_+ are the only root vectors with a downwards component, and since $P_{1-}^n v_0 = P_{0-}^s v_0 = N_+ v_0 = 0$, it follows that V cannot have weights (k, k') below a certain line $k' = k'_0$ for some $k'_0 < \ell$.

Now consider the vectors $X_-^q v_0$ for positive integers q . Since $[N_-, X_-] = 0$, all these vectors are annihilated by N_-^r . If $X_-^q v_0$ would be non-zero for very large q , then it would generate a

$\mathfrak{k}_{\mathbb{C}}$ -module containing weights below the line $k' = k'_0$; this is impossible. Hence there exists a q such that $X_-^q v_0 = 0$.

Now, in $\mathbb{C}[X, Y, Z]$, consider the ideal J generated by X^q and $D_-^t = (Y^2 - 4XZ)^t$ and Z^s . Clearly, its vanishing set in affine three-space consists of only the point $(0, 0, 0)$. It follows that $\mathbb{C}[X, Y, Z]/J$ is finite-dimensional as a $\mathbb{C}$ -vector space (see, e.g., Corollary 4 in Sect. 1.7 of [14]). Since the annihilator of v_0 contains J , it follows that v_0 is $\mathfrak{n}$ -finite.

Since we know that $X_-^q v_0 = 0$, an argument analogous to the above shows that V cannot have any weights (k, k') to the left of a certain line $k = k_0$. Thus, V contains only finite-dimensional $\mathfrak{k}_{\mathbb{C}}$ -modules. It also follows that each weight space is finite-dimensional. In particular, V is admissible. $\square$

Proposition 4.5. *Let ℓ be an integer, and m and p be non-negative integers. Let Γ be a congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$. Let $F \in N_{\ell, m}^p(\Gamma)$ be non-zero. Let $\Phi : \mathrm{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ be the function corresponding to F via Lemma 3.2. Then $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$. Furthermore, $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ if and only if F is a cusp form.*

Proof. Evidently, Φ is smooth, left Γ -invariant, K -finite and has weight $(\ell + m, \ell)$. Next, we briefly sketch a proof that Φ is slowly increasing (for further details, the reader can look at [30] where a generalisation of this to higher degree is proved). Using the fact that the Fourier expansion of each component function F_j is supported on the positive semi-definite matrices, an argument similar to [25, §13] shows that the function F is bounded in a Siegel set; use the fact that the functions $[\alpha, \beta, \gamma]$ are bounded in a Siegel set. This shows that there is a constant c such that $|(F|_{\ell, m} \gamma)(Z)| \leq c$ for all γ in $\mathrm{Sp}_4(\mathbb{Z})$ and all Z in the Siegel set.

For any point $Z \in \mathbb{H}_2$ there exists $\gamma \in \mathrm{Sp}_4(\mathbb{Z})$ such that $Z' := \gamma Z$ lies in the Siegel set. One can prove that there exists a positive integer n , depending on ℓ and m , such that

$$\|\eta_{\ell, m}(J(\gamma, Z))^{-1}\| \leq (\mathrm{tr}(Y) + \mathrm{tr}(Y^{-1}))^n$$

for all $\gamma \in \mathrm{Sp}_4(\mathbb{Z})$ and all $Z = X + iY \in \mathbb{H}_2$. This shows that $|F(Z)| \leq c(\mathrm{tr}(Y) + \mathrm{tr}(Y^{-1}))^n$ for all $Z \in \mathbb{H}_2$. It then follows easily that Φ is slowly increasing.

Since, by Table 2, the operators D_- and P_{0-} lower the nearly holomorphic degree, we have $P_{0-}^s F = D_-^t F = 0$ for some $s, t > 0$. By the diagram (93), it follows that $P_{0-}^s \Phi = D_-^t \Phi = 0$. Hence, we can apply Lemma 4.4 and conclude that Φ is $\mathfrak{n}$ -finite, and generates an admissible $(\mathfrak{g}, K)$ -module. Since each weight space is finite-dimensional by Lemma 4.4, it follows that Φ is $\mathbb{Z}$ -finite. This proves $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$.

Finally, we sketch a proof that $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ if and only if F is a cusp form. Since we already know that $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$, it suffices to show that Φ is a cusp form if and only if F is a cusp form. First, start with a cusp form $F \in N_{\ell, m}(\Gamma)^\circ$. For any $\gamma \in \mathrm{Sp}_4(\mathbb{Q})$, define $F_\gamma := F|_{\ell, m} \gamma$. Then $F_\gamma \in N_{\ell, m}(\Gamma(N_\gamma))^\circ$ for some integer N_γ , and its Fourier expansion is supported on positive definite matrices. For any $x \in \mathbb{R}$, define $n_x = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & x \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ and put $N'(\mathbb{R}) := \{n_x : x \in \mathbb{R}\}$. Note that $N'(\mathbb{R})$ is contained in the unipotent radicals of all the (standard-form) maximal parabolic subgroups of $\mathrm{Sp}_4(\mathbb{R})$. In order to show that Φ is a cusp form, it suffices to show that for all $\gamma \in \mathrm{Sp}_4(\mathbb{Q})$, $g \in \mathrm{Sp}_4(\mathbb{R})$, we have $\int_{N_\gamma \mathbb{Z} \backslash \mathbb{R}} \Phi(\gamma n_x g) dx = 0$. This is implied by $\int_{N_\gamma \mathbb{Z} \backslash \mathbb{R}} F_\gamma(Z + \begin{bmatrix} 0 & 0 \\ 0 & x \end{bmatrix}) dx = 0$, which in turn follows from the fact that Fourier coefficients of F_γ vanish at all matrices of the form $\begin{bmatrix} a & b/2 \\ b/2 & 0 \end{bmatrix}$.

Conversely, suppose that F is *not* a cusp form. We will show that Φ is not a cusp form. We can find $\gamma \in \mathrm{Sp}_4(\mathbb{Q})$ such that $F_\gamma := F|_{\ell, m} \gamma$ has a non-vanishing degenerate Fourier coefficient. By choosing γ appropriately, we can assume that the Fourier coefficient of F_γ is non-vanishing at a matrix of the form $Q_t := \frac{1}{N_\gamma} \begin{bmatrix} 0 & 0 \\ 0 & t \end{bmatrix}$ with $t \in \mathbb{Z}$. Fix such a matrix Q_t . Write the Fourier

expansion as

$$F_\gamma(Z) = \sum_{Q \in (1/N_\gamma)M_2^{\text{sym}}(\mathbb{Z})} a(Q) e^{2\pi i \text{Tr}(QZ)}, \quad (104)$$

where

$$a(Q) = \sum_{j=0}^m \sum_{\alpha, \beta, \gamma} a_{j, \alpha, \beta, \gamma}(Q) \left(\frac{y}{yy' - v^2} \right)^\alpha \left(\frac{v}{yy' - v^2} \right)^\beta \left(\frac{y'}{yy' - v^2} \right)^\gamma S^{m-j} T^j. \quad (105)$$

Since $a(Q_t) \neq 0$, we can find positive real numbers r_2, r_3 such that the element $f(y)$ in $\mathbb{C}(y)[S, T]$ given by

$$f(y) = \sum_{j=0}^m \sum_{\alpha, \beta, \gamma} a_{j, \alpha, \beta, \gamma}(Q_t) \left(\frac{y}{yr_3 - r_2^2} \right)^\alpha \left(\frac{r_2}{yr_3 - r_2^2} \right)^\beta \left(\frac{r_3}{yr_3 - r_2^2} \right)^\gamma S^{m-j} T^j$$

is not identically zero. Fix such r_2, r_3 . For $y > r_2^2/r_3$, define the matrix

$$M_y := \frac{1}{\sqrt{y + r_3 + 2\sqrt{yr_3 - r_2^2}}} \begin{bmatrix} y + \sqrt{yr_3 - r_2^2} & r_2 \\ r_2 & r_3 + \sqrt{yr_3 - r_2^2} \end{bmatrix}$$

which is a square root of the matrix $\begin{bmatrix} y & r_2 \\ r_2 & r_3 \end{bmatrix}$. Then there is a real number α and some positive real c , both independent of y , such that

$$|\eta_{\ell, m}(M_y)f(y)| > cy^\alpha \quad (106)$$

holds for all large enough y . Above, $|\cdot|$ is the norm on $\mathbb{C}[S, T]$ defined by $|\sum_{i,j} a_{i,j} S^i T^j| := \max_{i,j} |a_{i,j}|$, though any other norm on $\mathbb{C}[S, T]$ will work equally well.

Now let $\Phi_\gamma(g) = \Phi(\gamma g)$ be the automorphic form attached to F_γ . Consider the integral

$$\begin{aligned} & \int_{M_2^{\text{sym}}(N_\gamma \mathbb{Z} \backslash \mathbb{R})} \vec{\Phi}_\gamma \left(\begin{bmatrix} I_2 & X \\ 0 & I_2 \end{bmatrix} \begin{bmatrix} M_y & 0 \\ 0 & M_y^{-1} \end{bmatrix} \right) e^{-2\pi i \text{Tr}(Q_t X)} dX \\ &= \eta_{\ell, m}(M_y) \int_{M_2^{\text{sym}}(N_\gamma \mathbb{Z} \backslash \mathbb{R})} F_\gamma(X + i \begin{bmatrix} y & r_2 \\ r_2 & r_3 \end{bmatrix}) e^{-2\pi i \text{Tr}(Q_t X)} dX \\ &= \text{vol}(M_2^{\text{sym}}(N_\gamma \mathbb{Z} \backslash \mathbb{R})) e^{-\frac{2\pi i r_3}{N_\gamma}} (\eta_{\ell, m}(M_y)f(y)). \end{aligned}$$

Now, using the triangle inequality on the above integral, together with (106), we conclude that for all sufficiently large y , we can find $X \in M_2^{\text{sym}}(\mathbb{R})$ such that the matrix

$$g = \begin{bmatrix} I_2 & X \\ 0 & I_2 \end{bmatrix} \begin{bmatrix} M_y & 0 \\ 0 & M_y^{-1} \end{bmatrix}$$

has the property $|\vec{\Phi}_\gamma(g)| > ce^{-\frac{2\pi i r_3}{N_\gamma}} y^\alpha$. Letting $y \rightarrow \infty$, we conclude that $|\vec{\Phi}_\gamma|$ is *not* a rapidly decaying function on $\text{Sp}_4(\mathbb{R})$. Hence, Φ_γ is not a cusp form and therefore neither is Φ . This completes the proof. $\square$

4.3. The structure theorem for cusp forms. In this section we prove the structure theorem for cusp forms. It is based on the following decomposition of the space $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ into irreducibles.

Proposition 4.6. *As $(\mathfrak{g}, K)$ -modules, we have*

$$\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ = \bigoplus_{\ell=1}^{\infty} \bigoplus_{m=0}^{\infty} n_{\ell, m} L(\ell + m, \ell), \quad n_{\ell, m} = \dim S_{\ell, m}(\Gamma).$$

The lowest weight vectors in the isotypical component $n_{\ell, m} L(\ell + m, \ell)$ correspond to elements of $S_{\ell, m}(\Gamma)$ via the isomorphism from Lemma 3.2.

Proof. Since $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ is a $(\mathfrak{g}, K)$ -submodule of $\mathcal{A}(\Gamma)^\circ$, it decomposes into an orthogonal direct sum of irreducible $(\mathfrak{g}, K)$ -modules, each occurring with finite multiplicity. Recall from Lemma 2.3 that the only irreducible, locally $\mathfrak{n}$ -finite $(\mathfrak{g}, K)$ -modules are the $L(\lambda)$ for $\lambda \in \Lambda$. Since $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ admits the inner product (103), each $L(\lambda)$ occurring in the decomposition of $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ is unitarizable. The trivial $(\mathfrak{g}, K)$ -module $L(0, 0)$ cannot occur, since constant functions are not cuspidal. Proposition 2.2 (3) therefore implies that only $L(\ell + m, \ell)$ with $\ell \geq 1$ can occur. The module $L(\ell + m, \ell)$ must occur with multiplicity $\dim S_{\ell, m}(\Gamma)$, since every lowest weight vector in its isotypical component gives rise to an element of $S_{\ell, m}(\Gamma)$, and conversely. $\square$

Remark 4.7. By (2) of Proposition 2.2, the modules $L(1 + m, 1)$ are non-tempered. Still, it is possible for these modules to occur in $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ for certain Γ . After all, cusp forms of weight 1 do exist; see [44]. Globally, the modules $L(1 + m, 1)$ occur in CAP representations with respect to the Borel or Klingen parabolic subgroup, which were considered in [40]. Therefore, these modules have to be excluded from any correct formulation of the Ramanujan conjecture.

Recall from Lemma 4.1 that $\mathcal{X}$ denotes the free monoid consisting of all strings of the symbols in the left column of Table 1. For integers ℓ, m, ℓ', m' , we define the following subsets of $\mathcal{X}$. If $\ell \geq \ell' \geq 2$, $m \geq 0$, $m' \geq 0$, then let

$$\begin{aligned} \mathcal{X}_{\ell', m'}^{\ell, m} = & \left\{ X_+^\alpha D_+^\beta U^\gamma \mid \alpha, \beta, \gamma \in \mathbb{Z}_{\geq 0}, \gamma \leq m'/2, \right. \\ & \left. \ell' + m' + 2\alpha + 2\beta = \ell + m, \ell' + 2\beta + 2\gamma = \ell \right\} \\ & \cup \left\{ E_+ X_+^\alpha D_+^\beta U^\gamma \mid \alpha, \beta, \gamma \in \mathbb{Z}_{\geq 0}, \gamma < m'/2, \right. \\ & \left. \ell' + m' + 2\alpha + 2\beta + 1 = \ell + m, \ell' + 2\beta + 2\gamma + 1 = \ell \right\}. \end{aligned} \quad (107)$$

If $\ell \geq \ell' = 1$, $m \geq 0$, $m' \geq 0$, then let

$$\mathcal{X}_{\ell', m'}^{\ell, m} = \begin{cases} \emptyset & \text{if } m' > m \text{ or } m \not\equiv m' \pmod{2}, \\ \left\{ X_+^{\frac{m-m'}{2}} D_+^{\frac{\ell-1}{2}} \right\} & \text{if } m' \leq m, m \equiv m' \pmod{2}, \text{ and } \ell \text{ is odd,} \\ \left\{ E_+ X_+^{\frac{m-m'}{2}} D_+^{\frac{\ell-2}{2}} \right\} & \text{if } m' \leq m, m \equiv m' \pmod{2}, \text{ and } \ell \text{ is even.} \end{cases} \quad (108)$$

In every other case we put $\mathcal{X}_{\ell', m'}^{\ell, m} = \emptyset$, except for $\mathcal{X}_{0,0}^{0,0}$ which we put equal to $\{1\}$.

With these notations we are now ready to prove one of our main results.

Theorem 4.8 (Structure theorem for cusp forms). *Let ℓ be an integer, and m a non-negative integer. Then we have an orthogonal direct sum decomposition*

$$N_{\ell, m}(\Gamma)^\circ = \bigoplus_{\ell'=1}^{\ell} \bigoplus_{m'=0}^{\ell+m-\ell'} \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X(S_{\ell', m'}(\Gamma)). \quad (109)$$

Proof. Let $F \in N_{\ell, m}(\Gamma)^\circ$. Let $\Phi : \text{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ be the function corresponding to F via Lemma 3.2. By Proposition 4.5, we have $\Phi \in \mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$. According to Proposition 4.6, we can write $\Phi = \sum_{j=1}^r \Phi_j$ with non-zero Φ_j of weight $(\ell + m, \ell)$ and lying in an irreducible submodule $L(\ell_j + m_j, \ell_j)$ of $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$. Since $N_+ \Phi = 0$, we have $N_+ \Phi_j = 0$ for all j . Considering the possible K -types of the $L(\lambda)$ given in Lemma 2.7, we see $\ell_j \leq \ell$ and $\ell_j + m_j \leq \ell + m$ for all j .

Let Ψ_j be a vector of weight $(\ell_j + m_j, \ell_j)$ in $L(\ell_j + m_j, \ell_j)$. By Propositions 2.15 and 2.17, we can navigate from Ψ_j to Φ_j using the operators U , X_+ , D_+ and E_+ . The functions Ψ_j correspond to elements of $S_{\ell_j, m_j}(\Gamma)$. The commutativity of the diagram (96) allows us to rewrite the relations in terms of functions on $\mathbb{H}_2$. This proves the theorem. $\square$

Corollary 4.9. *Let ℓ be an integer, and m a non-negative integer. Then*

$$N_{\ell,m}(\Gamma)^\circ = N_{\ell,m}^p(\Gamma)^\circ \quad \text{with } p = \ell - 1 + \left\lfloor \frac{m}{2} \right\rfloor.$$

Proof. Consider a typical term $X_+^\alpha D_+^\beta U^\gamma S_{\ell',m'}(\Gamma)$ appearing in the structure theorem. By Table 2, such a term can produce nearly holomorphic degrees no larger than $\alpha + 2\beta + \gamma$. By the conditions in the first set in (107),

$$\alpha + 2\beta + \gamma = \ell - \ell' + \frac{m - m'}{2} \leq \ell - 2 + \frac{m}{2}.$$

Similarly we can estimate the nearly holomorphic degree of all the terms in the structure theorem. The maximal number is $\ell - 1 + \frac{m}{2}$, proving our result. $\square$

Corollary 4.10 (Structure theorem for scalar-valued cusp forms). *Let ℓ be an integer. Then we have an orthogonal direct sum decomposition*

$$N_{\ell,0}(\Gamma)^\circ = \bigoplus_{\substack{\ell'=2 \\ \ell' \equiv \ell \pmod{2}}}^{\ell} \bigoplus_{\substack{\ell'-\ell'=0 \\ m' \equiv 0 \pmod{2}}}^{\ell-\ell'} D_+^{(\ell-\ell'-m')/2} U^{m'/2} S_{\ell',m'}(\Gamma) \oplus N_{\ell,0}(\Gamma)_1^\circ,$$

where

$$N_{\ell,0}(\Gamma)_1^\circ = \begin{cases} D_+^{(\ell-1)/2} S_{1,0}(\Gamma) & \text{if } \ell \text{ is odd,} \\ 0 & \text{if } \ell \text{ is even.} \end{cases}$$

Proof. The terms of the decomposition in Theorem 4.8 simplify for $m = 0$. Note that all the E^+ terms are zero by (3) of Proposition 2.15 and (2) of Proposition 2.17. $\square$

Corollary 4.11 (Structure theorem for scalar-valued cusp forms of bounded nearly holomorphic degree). *Let ℓ be an integer. Then, for each $p \geq 0$, we have an orthogonal direct sum decomposition*

$$N_{\ell,0}^p(\Gamma)^\circ = \bigoplus_{\substack{\ell'=\max(2,\ell-2p) \\ \ell' \equiv \ell \pmod{2}}}^{\ell} \bigoplus_{\substack{\ell'-\ell'=0 \\ m' \equiv 0 \pmod{2}}}^{\ell-\ell'} D_+^{\frac{\ell-\ell'-m'}{2}} U^{\frac{m'}{2}} S_{\ell',m'}(\Gamma) \oplus N_{\ell,0}^p(\Gamma)_1^\circ,$$

where

$$N_{\ell,0}^p(\Gamma)_1^\circ = \begin{cases} D_+^{(\ell-1)/2} S_{1,0}(\Gamma) & \text{if } \ell \text{ is odd and } p \geq \ell - 1, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. The fact that the right side is contained in the left side follows immediately from Table 2. Next, let $F \in N_{\ell,0}^p(\Gamma)^\circ$. By Corollary 4.10, we can write

$$F = \sum_{\ell',m'} D_+^{\frac{\ell-\ell'-m'}{2}} U^{\frac{m'}{2}} F_{\ell',m'} + D_+^{(\ell-1)/2} F_{1,0},$$

where $F_{\ell',m'} \in S_{\ell',m'}(\Gamma)$ and $F_{1,0} \in S_{1,0}(\Gamma)$ (with $F_{1,0} = 0$ if ℓ is even). To complete the proof, it suffices to show that each ℓ', m' above with $F_{\ell',m'} \neq 0$ satisfies $\ell - \ell' - \frac{m'}{2} \leq p$, and furthermore, that $F_{1,0} \neq 0$ implies $p \geq \ell - 1$.

We show that $F_{\ell',m'} \neq 0$ implies $\ell - \ell' - \frac{m'}{2} \leq p$; the proof for the other inequality is similar. Suppose that $\ell - \ell' - \frac{m'}{2} > p$. Then, using Table 2, we see that $P_{0-}^{m'/2} D_-^{(\ell-\ell'-m')/2} F = 0$. This implies that

$$P_{0-}^{m'/2} D_-^{(\ell-\ell'-m')/2} D_+^{(\ell-\ell'-m')/2} U^{m'/2} F_{\ell',m'} = 0.$$

But this contradicts Lemma 2.12. $\square$

4.4. Petersson inner products. Let ℓ be an integer, and m, p be non-negative integers. We let $\langle \cdot, \cdot \rangle_m$ be the unique $U(2)$ -invariant inner product on W_m such that $\langle S^m, S^m \rangle_m = 1$. Let Γ be a congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$. For $F, G \in N_{\ell, m}(\Gamma)$, we define the Petersson inner product $\langle F, G \rangle$ by

$$\langle F, G \rangle = \mathrm{vol}(\Gamma \backslash \mathbb{H}_2)^{-1} \int_{\Gamma \backslash \mathbb{H}_2} \langle \eta_{\ell, m}(\mathrm{Im}(Z)) (F(Z)), G(Z) \rangle_m dZ,$$

where dZ is any invariant measure on $\mathbb{H}_2$, *provided the integral converges absolutely*. We denote this absolute convergence condition by $\langle F, G \rangle < \infty$. If the integral does not converge absolutely, we write $\langle F, G \rangle = \infty$.

Remark 4.12. Let $F, G \in N_{\ell, m}(\Gamma)$ such that at least one of F and G lies in $N_{\ell, m}(\Gamma)^\circ$. Then $\langle F, G \rangle < \infty$.

Lemma 4.13. *Let ℓ be an integer, and m a non-negative integer. Let $F, G \in N_{\ell, m}^p(\Gamma)$ and let Φ_F, Φ_G be the functions on $\mathrm{Sp}_4(\mathbb{R})$ corresponding to F, G respectively via Lemma 3.2. Suppose that $\langle F, G \rangle < \infty$. Then $\langle \Phi_F, \Phi_G \rangle < \infty$ and*

$$\langle F, G \rangle = \langle \Phi_F, \Phi_G \rangle, \quad (110)$$

where $\langle \Phi_F, \Phi_G \rangle$ is defined by (103).

Proof. This follows from a standard computation as in [1, p. 195]. We omit the details. $\square$

We define the subspace $\mathcal{E}_{\ell, m}(\Gamma)$ to be the orthogonal complement of $N_{\ell, m}(\Gamma)^\circ$ in $N_{\ell, m}(\Gamma)$ with respect to the Petersson inner product.

Lemma 4.14. *Let ℓ, m be non-negative integers. Let $F \in \mathcal{E}_{\ell, m}(\Gamma)$, and let $\Phi \in \mathcal{A}(\Gamma)_{\mathrm{n-fin}}$ be the function corresponding to F via Lemma 3.2. Then Φ is orthogonal to $\mathcal{A}(\Gamma)_{\mathrm{n-fin}}^\circ$.*

Proof. Let $\Psi \in \mathcal{A}(\Gamma)_{\mathrm{n-fin}}^\circ$; we have to show that $\langle \Phi, \Psi \rangle = 0$. We may assume that Ψ generates an irreducible module $L(\lambda)$ for some λ . Since, under the K -action, Φ generates the K -type $\rho_{(\ell+m, \ell)}$, we may assume that Ψ does as well. Writing Ψ as a sum of weight vectors, we may even assume that Ψ has the same weight as Φ , namely $(\ell + m, \ell)$. But then Ψ corresponds to an element G of $N_{\ell, m}(\Gamma)^\circ$. By hypothesis $\langle F, G \rangle = 0$. Hence $\langle \Phi, \Psi \rangle = 0$ by Lemma 4.13. $\square$

Lemma 4.15. *Let $X, \mathcal{X}$ be as in Lemma 4.1. Then X takes $N_{\ell, m}(\Gamma)^\circ$ to $N_{\ell', m'}(\Gamma)^\circ$ and $\mathcal{E}_{\ell, m}(\Gamma)$ to $\mathcal{E}_{\ell', m'}(\Gamma)$.*

Proof. The fact that X takes $N_{\ell, m}(\Gamma)^\circ$ to $N_{\ell', m'}(\Gamma)^\circ$ is an immediate consequence of the fact that X does not introduce new Fourier coefficients (this is true for each operator in Table 1 by (102) and is therefore true for all elements of $\mathcal{X}$).

To prove that X takes $\mathcal{E}_{\ell, m}(\Gamma)$ to $\mathcal{E}_{\ell', m'}(\Gamma)$, let $F \in \mathcal{E}_{\ell, m}(\Gamma)$, and let $\Phi \in \mathcal{A}(\Gamma)_{\mathrm{n-fin}}$ be the corresponding automorphic form. By Lemma 4.14, Φ is orthogonal to $\mathcal{A}(\Gamma)_{\mathrm{n-fin}}^\circ$. Hence the entire $(\mathfrak{g}, K)$ -module $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi$ is orthogonal to $\mathcal{A}(\Gamma)_{\mathrm{n-fin}}^\circ$. Since XF corresponds to $X\Phi \in \mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi$, our assertion follows. $\square$

Lemma 4.16. *Let ℓ be a positive integer, and m a non-negative integer. Let $X, \mathcal{X}$ be as in Lemma 4.1. There exists a constant $c_{\ell, m, X}$ (depending only on ℓ, m, X) such that for all $F \in S_{\ell, m}(\Gamma)$ we have $\langle XF, XF \rangle = c_{\ell, m, X} \langle F, F \rangle$.*

Proof. Set $\lambda = (\ell + m, \ell)$, and consider the $(\mathfrak{g}, K)$ -module $L(\lambda)$. Let v_0 be a highest weight vector in the minimal K -type of $L(\lambda)$; of course, v_0 is unique up to multiples. Since $L(\lambda)$ is unitary by Proposition 2.2, we may endow it with a $\mathfrak{g}$ -invariant inner product $\langle \cdot, \cdot \rangle$. By irreducibility, this inner product is unique up to multiples. Put $c_{\ell, m, X} = \langle Xv_0, Xv_0 \rangle / \langle v_0, v_0 \rangle$. Note that $c_{\ell, m, X}$ does not depend on the choice of model for $L(\lambda)$, the choice of v_0 , or the normalization of inner products.

Now all we need to observe is that the automorphic form $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ corresponding to F generates a module isomorphic to $L(\lambda)$, that Φ is a lowest weight vector in this module, and Lemma 4.13. $\square$

Proposition 4.17. *Let ℓ be a positive integer, and m a non-negative integer. Let $X, \mathcal{X}$ be as in Lemma 4.1. Then, for all $F \in S_{\ell,m}(\Gamma)$ and $G \in M_{\ell,m}(\Gamma)$, we have $\langle XF, XG \rangle = c_{\ell,m,X} \langle F, G \rangle$, where the constant $c_{\ell,m,X}$ is the same as in Lemma 4.16.*

Proof. Because of Lemma 4.15, we may assume that F and G both belong to $S_{\ell,m}(\Gamma)$. Now the proposition follows by applying the previous lemma to $F + G$. $\square$

4.5. Initial decomposition in the general case. As before, we fix a congruence subgroup Γ , and consider the space $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ of $\mathfrak{n}$ -finite automorphic forms. In this and the next sections we investigate the algebraic structure of this $(\mathfrak{g}, K)$ -module. We know from Proposition 4.6 that the subspace of cusp forms is completely reducible. Since there is no inner product defined on all of $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$, this may no longer be true for the entire space. The following vanishing result for Siegel modular forms will imply some basic restrictions on the possible K -types occurring in $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$.

Lemma 4.18. *Let $\ell, m \in \mathbb{Z}$ with $m \geq 0$. Assume that $M_{\ell,m}(\Gamma) \neq 0$. Then $\ell \geq 1$ or $\ell = m = 0$. The space $M_{0,0}(\Gamma)$ consists only of the constant functions.*

Proof. The first statement follows from the vanishing theorem Satz 2 of [43]. The second statement says that the only holomorphic modular forms of weight 0 are the constant functions; this is well known. $\square$

Lemma 4.19. *The space $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ does not contain any weights (k, ℓ) with negative ℓ . It contains the weight $(0, 0)$ with multiplicity one; the corresponding weight space consists precisely of the constant functions.*

Proof. To prove the first statement, suppose that $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ contains a non-zero vector Φ of weight (k, ℓ) with $\ell < 0$. After applying P_{0-} , P_{1-} and X_- finitely many times to Φ , we may assume that Φ is annihilated by all these operators. By Corollary 3.4, Φ corresponds to a non-zero element F of $M_{\ell,k-\ell}(\Gamma)$. But such F do not exist by Lemma 4.18.

To prove the second statement, let Φ be a vector of weight $(0, 0)$. By the first statement, $P_{1-}\Phi = P_{0-}\Phi = N_+\Phi = 0$. Hence also $N_-\Phi = 0$. Since $[N_-, P_{1-}] = 2X_-$, then also $X_-\Phi = 0$. Therefore Φ corresponds to an element of $M_{0,0}(\Gamma)$. By Lemma 4.18, Φ must be constant. $\square$

Lemma 4.20. *The space $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ does not contain the weight $(2, 0)$.*

Proof. Suppose that $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is a non-zero vector of weight $(2, 0)$; we will obtain a contradiction. Since $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ does not contain any weights (k, ℓ) with negative ℓ , we have $E_-\Phi = P_{1-}\Phi = P_{0-}\Phi = N_+\Phi = 0$. By Lemma 4.18, Φ cannot be annihilated by all of $\mathfrak{p}_-$. Hence $X_-\Phi \neq 0$. Since the formula for the L -operator in Table 1 can be rewritten as

$$L = m(m+1)X_- - (m+1)N_-P_{1-} + N_-^2P_{0-},$$

it follows that $L\Phi \neq 0$. Since $L\Phi$ has weight $(0, 0)$, it is a constant function by Lemma 4.19. We normalize such that $L\Phi = -6$; the reason for this normalization will become clear momentarily.

Let $F : \mathbb{H}_2 \rightarrow W_2$ be the function corresponding to Φ . Let $F = F_0S^2 + F_1ST + F_2T^2$, where F_j are the component functions. By Proposition 3.15 (1), the relations $E_-F = P_{0-}F = 0$ and $LF = -6$ hold. Looking at the definitions (85), (86), (90) of these differential operators, and solving the resulting linear system, we conclude

$$F_j = [1, 0, 0] - 2[0, 1, 0] + [0, 0, 1] + H_j \quad (111)$$

for $j \in \{0, 1, 2\}$, where H_j is holomorphic. (See (79) for notation.) Now consider the function on $\mathbb{H}_2$ given by $G(Z) := F(Z) - 2F(2Z)$. Then $G(Z)$ is a modular form with respect to a

smaller congruence subgroup Γ' . It is easy to see that G is non-zero. In view of (111), the nearly holomorphic parts of $F(Z)$ and $2F(2Z)$ cancel each other out, so that G is holomorphic. Hence G is a non-zero element of $M_{0,2}(\Gamma')$. By Lemma 4.18, this is impossible. $\square$

For the next lemma, recall that $\mathcal{A}_\lambda(\Gamma)_{\text{n-fin}}$ denotes the subspace of vectors of weight $\lambda \in \Lambda$.

Lemma 4.21. *Let ℓ be an integer, and m a non-negative integer.*

- (1) $\mathcal{A}_{(\ell+m,\ell)}(\Gamma)_{\text{n-fin}} = 0$ if $\ell < 0$.
- (2) $\mathcal{A}_{(0,0)}(\Gamma)_{\text{n-fin}} = \mathbb{C}$.
- (3) $\mathcal{A}_{(m,0)}(\Gamma)_{\text{n-fin}} = 0$ for all $m > 0$.

Proof. (1) and (2) were already noted in Lemma 4.19.

(3) By part (1) and Lemma 4.18, the operator X_- induces injective maps

$$\mathcal{A}_{(m+2,0)}(\Gamma)_{\text{n-fin}} \longrightarrow \mathcal{A}_{(m,0)}(\Gamma)_{\text{n-fin}} \quad (112)$$

for each $m \geq 0$. Clearly, $\mathcal{A}_{(1,0)}(\Gamma)_{\text{n-fin}} = 0$ by Lemma 4.18, and $\mathcal{A}_{(2,0)}(\Gamma)_{\text{n-fin}} = 0$ by Lemma 4.20. Hence $\mathcal{A}_{(m,0)}(\Gamma)_{\text{n-fin}}$ is zero for all $m > 0$. $\square$

For a character χ of $\mathcal{Z}$ (the center of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$) let $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$ be the subspace of $\mathcal{A}(\Gamma)_{\text{n-fin}}$ consisting of vectors Φ with the property $(z - \chi(z))^n \Phi = 0$ for all $z \in \mathcal{Z}$ and some n depending on z .

Lemma 4.22. *We have*

$$\mathcal{A}(\Gamma)_{\text{n-fin}} = \bigoplus_{\chi} \mathcal{A}(\Gamma, \chi)_{\text{n-fin}}. \quad (113)$$

Each space $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$ has finite length as a $(\mathfrak{g}, K)$ -module.

Proof. For a weight $\mu \in \Lambda$, let $\mathcal{A}_{\succcurlyeq \mu}(\Gamma)_{\text{n-fin}}$ be the subspace of $\mathcal{A}(\Gamma)_{\text{n-fin}}$ spanned by all vectors of weight $\lambda \succcurlyeq \mu$; see (7) for the definition of the order. Since $\mathcal{A}(\Gamma)_{\text{n-fin}}$ is admissible by Lemma 4.3, and since there are no weights below a horizontal line by Lemma 4.19, the space $\mathcal{A}_{\succcurlyeq \mu}(\Gamma)_{\text{n-fin}}$ is finite-dimensional. Therefore, the $(\mathfrak{g}, K)$ -module $\mathcal{A}_{\langle \succcurlyeq \mu \rangle}(\Gamma)_{\text{n-fin}}$ generated by $\mathcal{A}_{\succcurlyeq \mu}(\Gamma)_{\text{n-fin}}$ lies in category $\mathcal{O}^p$. By general properties of this category, it admits a decomposition into χ -isotypical components, as defined in (15), each of which has finite length. If we move μ farther up and farther to the right, we will exhaust the whole space $\mathcal{A}(\Gamma)_{\text{n-fin}}$. The assertion follows. $\square$

By Lemma 4.22 (and Lemma 2.3), each $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$ has a finite length composition series whose irreducible quotients are of the form $L(\lambda)$ for some $\lambda \in \Lambda^+$. Since $L(\lambda)$ has central character $\chi_{\lambda+\varrho}$, only those λ with $\chi_{\lambda+\varrho} = \chi$ can occur in $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$. For a given χ , this allows for only finitely many λ . Lemma 4.21 puts restrictions on the possible $L(\lambda)$'s that can occur; for example, $L(k, \ell)$ with $\ell < 0$ can never occur in $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$. We will go through the list of χ 's for which there exists at least one $L(\lambda)$ that is permitted by Lemma 4.21; evidently, only such χ 's can occur in the decomposition (113):

- The trivial character, i.e., $\chi = \chi_{\varrho}$, where $\varrho = (-1, -2)$. The irreducible modules $L(\lambda)$ that can occur as subquotients of $\mathcal{A}(\Gamma, \chi_{\varrho})_{\text{n-fin}}$ are $L(0, 0)$ (the trivial representation), $L(3, 1)$ and $L(3, 3)$. (The module $L(2, 0)$ also has central character χ_{ϱ} , but is not permitted by (3) of Lemma 4.21). Following terminology in the literature, we call χ_{ϱ} the *principal character*.
- The characters $\chi_{\lambda+\varrho}$ for $\lambda = (k, 1)$ with $k \geq 4$. The irreducible modules that can occur as subquotients of $\mathcal{A}(\Gamma, \chi_{\lambda+\varrho})_{\text{n-fin}}$ are $L(k, 1)$ and $L(k, 3)$. Since the modules $L(k, 1)$ are non-tempered by Proposition 2.2, we will refer to these $\chi_{\lambda+\varrho}$ as *non-tempered characters*.
- The character $\chi_{\lambda+\varrho}$ for $\lambda = (1, 1)$. The irreducible modules that can occur as subquotients of $\mathcal{A}(\Gamma, \chi_{\lambda+\varrho})_{\text{n-fin}}$ are $L(1, 1)$ and $L(2, 2)$.

- The character $\chi_{\lambda+\varrho}$ for $\lambda = (2, 1)$. The only irreducible module that can occur as a subquotient of $\mathcal{A}(\Gamma, \chi_{\lambda+\varrho})_{\text{n-fin}}$ is $L(2, 1)$.
- The characters $\chi_{\lambda+\varrho}$ for $\lambda = (\ell + m, \ell)$ with $(\ell \geq 4, m \geq 0)$, or $(\ell = 2, m \geq 1)$. The only irreducible module that can occur as a subquotient of $\mathcal{A}(\Gamma, \chi_{\lambda+\varrho})_{\text{n-fin}}$ is $L(\lambda)$. We will refer to these $\chi_{\lambda+\varrho}$ as the *tempered characters*.

Our task in the following will be to determine the structure of each $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$ occurring in (113). We can quickly treat the case of tempered χ . Since $L(\lambda)$ admits no non-trivial self-extensions by Proposition 3.1 (d) of [20], the component $\mathcal{A}(\Gamma, \chi)_{\text{n-fin}}$ for tempered $\chi = \chi_{\lambda+\varrho}$ is a direct sum of copies of $L(\lambda)$. The lowest weight vector in such an $L(\lambda)$ corresponds to an element of $M_{\ell, m}(\Gamma)$, where $\lambda = (\ell + m, \ell)$. Thus,

$$\mathcal{A}(\Gamma, \chi)_{\text{n-fin}} = n_{\lambda} L(\lambda), \quad n_{\lambda} = \dim M_{\ell, m}(\Gamma), \quad (114)$$

for tempered $\chi = \chi_{\lambda+\varrho}$ with $\lambda = (\ell + m, \ell)$.

The third and fourth cases above can also be dealt with easily. We get

$$\mathcal{A}(\Gamma, \chi)_{\text{n-fin}} = n_{\lambda} L(\lambda), \quad n_{\lambda} = \dim M_{1, 1}(\Gamma), \quad (115)$$

for $\chi = \chi_{\lambda+\varrho}$ with $\lambda = (2, 1)$, and

$$\mathcal{A}(\Gamma, \chi)_{\text{n-fin}} = n_1 L(1, 1) \oplus n_2 L(2, 2), \quad n_k = \dim M_{k, 0}(\Gamma), \quad (116)$$

for $\chi = \chi_{\lambda+\varrho}$ with $\lambda = (1, 1)$.

As for the principal character, note that, by (2) of Lemma 4.21, the trivial module $L(0, 0)$ occurs exactly once in $\mathcal{A}(\Gamma)_{\text{n-fin}}$, and it occurs as a submodule. It is easy to see that $L(0, 0)$ does not admit any non-trivial extensions with $L(3, 1)$ or $L(3, 3)$. It follows that

$$\mathcal{A}(\Gamma, \chi_{\varrho})_{\text{n-fin}} \cong L(0, 0) \oplus V_3, \quad (117)$$

where the module V_3 has a composition series with the only subquotients being $L(3, 1)$ and $L(3, 3)$. This module V_3 can be treated together with the non-tempered characters, which we will take up in the next section.

4.6. The non-tempered characters. In this section we investigate the contribution to the space $\mathcal{A}(\Gamma)_{\text{n-fin}}$ coming from non-tempered central characters, as defined in the previous section. Recall that these are the $\chi_{\lambda+\varrho}$ for $\lambda = (k, 1)$ with $k \geq 4$. The only irreducible $L(\lambda)$ that can occur as subquotients of such modules are $\lambda = (k, 1)$ and $\lambda = (k, 3)$.

In the following we will require the dual module $N(\lambda)^{\vee}$ of the module $N(\lambda)$. The basic properties of the duality functor on category $\mathcal{O}^{\text{p}}$ are given in Sects. 3.2 and 3.3 of [20]. In particular, $N(\lambda)^{\vee}$ contains the same K -types as $N(\lambda)$, and admits $L(\lambda)$ as its unique irreducible submodule.

Lemma 4.23. *Let $k \geq 3$ be an integer and $\lambda = (k, 1) \in \Lambda$. Then*

$$\text{Ext}_{\mathcal{O}}(N(\lambda), N(\lambda)) = 0, \quad (118)$$

$$\text{Ext}_{\mathcal{O}}(L(\lambda), L(\lambda)) = 0, \quad (119)$$

$$\text{Ext}_{\mathcal{O}}(N(\lambda), L(\lambda)) = 0, \quad (120)$$

$$\text{Ext}_{\mathcal{O}}(L(\lambda), N(\lambda)) = 0, \quad (121)$$

$$\text{Ext}_{\mathcal{O}}(N(\lambda)^{\vee}, N(\lambda)^{\vee}) = 0, \quad (122)$$

$$\text{Ext}_{\mathcal{O}}(L(\lambda), N(\lambda)^{\vee}) = 0, \quad (123)$$

$$\text{Ext}_{\mathcal{O}}(N(\lambda)^{\vee}, L(\lambda)) = 0. \quad (124)$$

Proof. Equations (118) – (120) are general properties; see Proposition 3.1 of [20]. The claim (121) follows exactly as in the first part of the proof of Proposition 3.12 of [20]. Equations (122) – (124) follow from the previous ones and the properties of duality. $\square$

Lemma 4.24. *Let $k \geq 3$ be an integer. Let $\lambda = (k, 1) \in \Lambda$ and $\mu = (k, 3) \in \Lambda$. Let V be a module in category $\mathcal{O}^{\mathfrak{p}}$ with the following properties:*

- *The only possible irreducible subquotients of V are $L(\lambda)$ and $L(\mu)$.*
- *V does not contain $N(\lambda)$ as a submodule.*

Then

$$V = aL(\lambda) \oplus bL(\mu) \oplus cN(\lambda)^{\vee}. \quad (125)$$

with non-negative integers a, b, c .

Proof. For $i = 1$ or $i = 3$, denote by H_i the space of vectors $v \in V$ of weight (k, i) that are annihilated by N_+ . Then

$$V = \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_1 + \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_3. \quad (126)$$

Indeed, the quotient $V/(\mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_1 + \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_2)$ does not contain the K -types ρ_{λ} or ρ_{μ} . Since its only possible irreducible subquotients are $L(\lambda)$ or $L(\mu)$, it must be zero.

Note that P_{0-} induces a linear map $H_3 \rightarrow H_1$. Let H_3'' be the kernel of this map. Let H_3' be any vector space complement to H_3'' in H_3 . Let $H_1' = P_{0-}(H_3')$, and let H_1'' be any vector space complement to H_1' in H_1 . Thus $H_3 = H_3' \oplus H_3''$ and $H_1 = H_1' \oplus H_1''$, and P_{0-} induces an isomorphism $H_3' \rightarrow H_1'$. From (126) we see that

$$V = \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_1'' + \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_3'' + \mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_3'. \quad (127)$$

Our hypothesis that V does not contain $N(\lambda)$ as a submodule implies that every non-zero $v \in H_1$ generates a copy of $L(\lambda)$. From (119) we thus conclude that $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_1'' = aL(\lambda)$ (a direct sum) for some $a \geq 0$. Similarly, $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_3'' = bL(\mu)$ for some $b \geq 0$. Our hypothesis, together with the PBW theorem, also implies that every non-zero $v \in H_3'$ generates a copy of $N(\lambda)^{\vee}$. Using (122), we get $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})H_3' = cN(\lambda)^{\vee}$ for some $c \geq 0$. By (127) we now have

$$V = aL(\lambda) + bL(\mu) + cN(\lambda)^{\vee}. \quad (128)$$

Lemma 4.23 implies that $aL(\lambda) + cN(\lambda)^{\vee} = aL(\lambda) \oplus cN(\lambda)^{\vee}$. Since

$$\mathrm{Hom}(L(\mu), aL(\lambda) \oplus cN(\lambda)^{\vee}) = 0,$$

the intersection of $bL(\mu)$ with $aL(\lambda) \oplus cN(\lambda)^{\vee}$ is zero. It follows that the sums in (128) are direct. $\square$

Let $\lambda = (k, 1)$ and $\mu = (k, 3)$ for some $k \geq 3$. Let $\chi = \chi_{\lambda+\varrho}$. If $k \geq 4$, then let $V_k = \mathcal{A}(\Gamma, \chi)_{\mathrm{n-fin}}$; hence, V_k is the component appearing in the decomposition (113) corresponding to the non-tempered character χ . Let V_3 be the module appearing in (117); hence, V_3 is “almost” $\mathcal{A}(\Gamma, \chi_{\varrho})_{\mathrm{n-fin}}$, but without the trivial module.

Lemma 4.25. *V_k does not contain $N(\lambda)$ as a submodule.*

Proof. Suppose that V_k does contain $N(\lambda)$ as a submodule; we will obtain a contradiction. Let $\Phi \in N(\lambda)$ be an automorphic form of weight $(k, 1)$. Clearly, Φ generates $N(\lambda)$. Let $F \in M_{1, k-1}(\Gamma)$ be the holomorphic modular form corresponding to Φ . By the Folgerung to Satz 3 of [43], the modular form F is square-integrable. By Lemma 4.13 the function Φ is square-integrable on $\mathrm{Sp}_4(\mathbb{R})$. Since square-integrable automorphic forms constitute a $(\mathfrak{g}, K)$ -submodule of $\mathcal{A}(\Gamma)_{\mathrm{n-fin}}$, it follows that $N(\lambda)$ consists entirely of square-integrable forms, and hence admits an invariant inner product. In particular, we obtain the contradiction that $N(\lambda)$ is semisimple. $\square$

For any $k \geq 3$, the module V_k admits only $L(\lambda)$ and $L(\mu)$ as irreducible subquotients. Therefore, by Lemma 4.24 and Lemma 4.25,

$$V_k \cong aL(\lambda) \oplus bL(\mu) \oplus cN(\lambda)^{\vee} \quad (129)$$

with certain multiplicities a, b, c . These multiplicities may be related to dimensions of spaces of modular forms, as follows. Any vector of weight $(k, 1)$ in either $L(\lambda)$ or $N(\lambda)^{\vee}$ gives rise to an

element of $M_{1,k-1}(\Gamma)$. Conversely, a non-zero element $F \in M_{1,k-1}(\Gamma)$ (or rather the function Φ on $\mathrm{Sp}_4(\mathbb{R})$ corresponding to F) generates a copy of $L(\lambda)$ (which may lie inside an $N(\lambda)^\vee$). This explains the first of the following three equations,

$$a + c = \dim M_{1,k-1}(\Gamma), \quad (130)$$

$$b = \dim M_{3,k-3}(\Gamma), \quad (131)$$

$$b + c = \dim M_{3,k-3}^*(\Gamma). \quad (132)$$

For the second equation, observe that any vector of weight $(k, 3)$ in $L(\mu)$ gives rise to an element of $M_{3,k-3}(\Gamma)$. Conversely, a non-zero element $F \in M_{3,k-3}(\Gamma)$ generates a copy of $L(\mu)$.

The space appearing in (132) is defined by

$$M_{3,k-3}^*(\Gamma) = \{F \in N_{3,k-3}(\Gamma) \mid LF = E_-F = 0, P_{0-}F \text{ is holomorphic}\}. \quad (133)$$

By (3) of Lemma 3.16, an alternative definition is

$$M_{3,k-3}^*(\Gamma) = \{F \in N_{3,k-3}^1(\Gamma) \mid LF = E_-F = 0\}. \quad (134)$$

Evidently,

$$M_{3,k-3}(\Gamma) \subset M_{3,k-3}^*(\Gamma) \subset N_{3,k-3}^1(\Gamma). \quad (135)$$

We already noted that a vector of weight $(k, 3)$ in $L(\mu)$ gives rise to an element of $M_{3,k-3}(\Gamma)$, and hence to an element of $M_{3,k-3}^*(\Gamma)$. We claim that a vector Φ of weight $(k, 3)$ in $N(\lambda)^\vee$ also gives rise to an element of $M_{3,k-3}^*(\Gamma)$. Let F be the smooth function on $\mathbb{H}_2$ corresponding to Φ . Then (3) of Lemma 3.16 implies that F is nearly holomorphic of degree 1. Hence $F \in N_{3,k-3}^1(\Gamma)$. Clearly $F \in M_{3,k-3}^*(\Gamma)$, as claimed. Conversely, a non-zero $F \in M_{3,k-3}^*(\Gamma)$ generates either a copy of $L(\mu)$ or a copy of $N(\lambda)^\vee$. This proves (132).

Solving the linear system (130) – (132), we obtain the following result.

Lemma 4.26. *For $k \geq 3$, let V_k be defined as above. Then we have the direct sum decomposition*

$$V_k \cong a_k L(\lambda) \oplus b_k L(\mu) \oplus c_k N(\lambda)^\vee, \quad (136)$$

where

$$a_k = \dim M_{1,k-1}(\Gamma) + \dim M_{3,k-3}(\Gamma) - \dim M_{3,k-3}^*(\Gamma), \quad (137)$$

$$b_k = \dim M_{3,k-3}(\Gamma), \quad (138)$$

$$c_k = \dim M_{3,k-3}^*(\Gamma) - \dim M_{3,k-3}(\Gamma). \quad (139)$$

We note that the component $cN(\lambda)^\vee$ in (136) is not well-defined as a subspace of V_k ; while the multiplicities of indecomposable modules are well-defined in category $\mathcal{O}^p$, isotypical components are in general not. For example, if Φ has weight $(k, 3)$ and generates an $N(\lambda)^\vee$, and if Ψ has the same weight and generates an $L(\mu)$, then $\Phi + \Psi$ also generates an $N(\lambda)^\vee$. Hence, the vectors of weight $(k, 3)$ generating the $N(\lambda)^\vee$ are only determined up to “holomorphic” vectors of the same weight.

In classical language, this means that we do not know of a canonical way to define a complement of $M_{3,k-3}(\Gamma)$ inside $M_{3,k-3}^*(\Gamma)$. We prefer not to choose any such complement, but work with the full space $M_{3,k-3}^*(\Gamma)$ instead. The modular forms in this space generate the component $b_k L(\mu) \oplus c_k N(\lambda)^\vee$, which is well-defined as a subspace of V_k .

Consider the map P_{0-} from $M_{3,k-3}^*(\Gamma)$ to $M_{1,k-1}(\Gamma)$. Recall from [43] that modular forms in the space $M_{1,k-1}(\Gamma)$ are square-integrable. Hence, we may consider the orthogonal complement $M_{1,k-1}^{**}(\Gamma)$ of $P_{0-}(M_{3,k-3}^*(\Gamma))$ inside $M_{1,k-1}(\Gamma)$. The various spaces are then connected by the exact sequence

$$0 \longrightarrow M_{3,k-3}(\Gamma) \longrightarrow M_{3,k-3}^*(\Gamma) \xrightarrow{P_{0-}} M_{1,k-1}(\Gamma) \longrightarrow M_{1,k-1}^{**}(\Gamma) \longrightarrow 0, \quad (140)$$

in which the fourth map is orthogonal projection. The quantity a_k in (137) is equal to $\dim M_{1,k-1}^{**}(\Gamma)$. Let V_k^* be the subspace of V_k generated by the elements of $M_{3,k-3}^*(\Gamma)$, and let V_k^{**} be the subspace of V_k generated by the elements of $M_{1,k-1}^{**}(\Gamma)$. Then

$$V_k = V_k^* \oplus V_k^{**}. \quad (141)$$

The subspaces V_k^* and V_k^{**} are canonically defined, and decompose according to

$$V_k^* \cong b_k L(\mu) \oplus c_k N(\lambda)^\vee, \quad V_k^{**} \cong a_k L(\lambda) \quad (142)$$

as abstract modules.

4.7. The structure theorem for all modular forms. Recall that in Proposition 4.6 we obtained a decomposition of the space $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ into irreducible $(\mathfrak{g}, K)$ -modules. The analogous statement for all $\mathfrak{n}$ -finite modular forms is slightly more complicated.

Proposition 4.27. *As $(\mathfrak{g}, K)$ -modules, we have*

$$\begin{aligned} \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}} &= \bigoplus_{\substack{\ell=2 \\ \ell \neq 3}}^{\infty} \bigoplus_{m=0}^{\infty} n_{\ell,m} L(\ell+m, \ell) \\ &\quad \oplus \bigoplus_{k=3}^{\infty} V_k^* \oplus \bigoplus_{k=1}^{\infty} V_k^{**} \oplus L(0,0). \end{aligned} \quad (143)$$

where $n_{\ell,m} = \dim M_{\ell,m}(\Gamma)$, the spaces V_k^*, V_k^{**} for $k \geq 3$ are as in (142), and $V_k^{**} = n_{1,k-1} L(k, 1)$ for $k = 1, 2$.

Proof. This follows by combining Lemma 4.22 with the results of the previous subsection. $\square$

Proposition 4.28. *Let ℓ be a positive integer, and m a non-negative integer. Let $F \in M_{\ell,m}(\Gamma)$ and let $\Phi_F : \mathrm{Sp}_4(\mathbb{R}) \rightarrow \mathbb{C}$ be the function of weight $(\ell+m, \ell)$ corresponding to F by Lemma 3.2. Then the submodule $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_F$ of $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is irreducible and isomorphic to $L(\ell+m, \ell)$.*

Proof. By Property (3) of the modules $N(\lambda)$ in Section 2.1, we see that $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_F$ is isomorphic to a quotient of $N(\ell+m, \ell)$. If $N(\ell+m, \ell) = L(\ell+m, \ell)$ there is nothing to prove. Otherwise assume that $N(\ell+m, \ell) \neq L(\ell+m, \ell)$. It suffices to prove that $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_F$ is not isomorphic to $N(\ell+m, \ell)$. But this follows from Proposition 4.27, as the module $N(\ell+m, \ell)$, when reducible, does not appear as a submodule of $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$. $\square$

Recall that the cuspidal structure theorem, Theorem 4.8, was based on Proposition 4.6, which is the cuspidal analogue of Proposition 4.27, and Propositions 2.15 and 2.17, which say that every highest weight vector in an $L(k, \ell)$ can be generated from the highest weight vector of its minimal K -type by applying U , X_+ , D_+ and E_+ operators. We therefore require a result similar to Propositions 2.15 and 2.17 for the indecomposable modules $N(k, 1)^\vee$ appearing in (143). For these modules we define $N(k, 1)_{\mathrm{par}(0)}^\vee$ and $N(k, 1)_{\mathrm{par}(1)}^\vee$ just as we did in the paragraph before Proposition 2.15 (set $\lambda = (k, 1)$). Recall that $N(k, 1)^\vee$ sits in an exact sequence

$$0 \longrightarrow L(k, 1) \longrightarrow N(k, 1)^\vee \xrightarrow{\varphi} L(k, 3) \longrightarrow 0.$$

For the submodule $L(k, 1)$ we have the spaces $L(k, 1)_{\mathrm{par}(0)}$ and $L(k, 1)_{\mathrm{par}(1)}$ of even and odd highest weight vectors, and clearly

$$L(k, 1)_{\mathrm{par}(0)} \subset N(k, 1)_{\mathrm{par}(0)}^\vee \quad \text{and} \quad L(k, 1)_{\mathrm{par}(1)} \subset N(k, 1)_{\mathrm{par}(1)}^\vee.$$

The spaces $L(k, 1)_{\mathrm{par}(0)}$ and $L(k, 1)_{\mathrm{par}(1)}$ originate from w_0 , the essentially unique vector of weight $(k, 1)$, by applying X_+ , D_+ and E_+ operators. Let w_1 be the essentially unique vector

of weight $(k, 3)$, so that $\varphi(w_1)$ is the highest weight vector in the minimal K -type of $L(k, 3)$. Then, by Proposition 2.15,

$$L(k, 3)_{\text{par}(0)} = \bigoplus_{\substack{\alpha, \beta \geq 0 \\ 0 \leq \gamma \leq \frac{k-3}{2}}} \mathbb{C} X_+^\alpha D_+^\beta U^\gamma \varphi(w_1), \quad L(k, 3)_{\text{par}(1)} = \bigoplus_{\substack{\alpha, \beta \geq 0 \\ 0 \leq \gamma < \frac{k-3}{2}}} \mathbb{C} E_+ X_+^\alpha D_+^\beta U^\gamma \varphi(w_1).$$

Now let

$$\tilde{L}(k, 3)_{\text{par}(0)} = \bigoplus_{\substack{\alpha, \beta \geq 0 \\ 0 \leq \gamma \leq \frac{k-3}{2}}} \mathbb{C} X_+^\alpha D_+^\beta U^\gamma w_1 \quad (144)$$

and

$$\tilde{L}(k, 3)_{\text{par}(1)} = \bigoplus_{\substack{\alpha, \beta \geq 0 \\ 0 \leq \gamma < \frac{k-3}{2}}} \mathbb{C} E_+ X_+^\alpha D_+^\beta U^\gamma w_1. \quad (145)$$

It is clear that φ maps $\tilde{L}(k, 3)_{\text{par}(i)}$ isomorphically onto $L(k, 3)_{\text{par}(i)}$; in particular, the sums in (144) and (145) are really direct.

Lemma 4.29. *With the above notations, we have $N(k, 1)_{\text{par}(i)}^\vee = L(k, 1)_{\text{par}(i)} \oplus \tilde{L}(k, 3)_{\text{par}(i)}$ for $i = 0, 1$.*

Proof. It is clear that the sum is direct, since $L(k, 1)_{\text{par}(i)}$ lies in the kernel of φ , while the restriction of φ to $\tilde{L}(k, 3)_{\text{par}(i)}$ is an isomorphism. Let $v \in N(k, 1)_{\text{par}(i)}^\vee$. Then $\varphi(v) \in L(k, 3)_{\text{par}(i)}$. Let $\tilde{v} \in \tilde{L}(k, 3)_{\text{par}(i)}$ be such that $\varphi(\tilde{v}) = \varphi(v)$. Then $v - \tilde{v} \in L(k, 1)_{\text{par}(i)}$. The assertion follows. $\square$

Theorem 4.30 (Structure theorem for all modular forms). *Let ℓ be a positive integer, and m a non-negative integer. Let the sets $\mathcal{X}_{\ell', m'}^{\ell, m}$ be defined as in (107) and (108). Then we have a direct sum decomposition*

$$N_{\ell, m}(\Gamma) = \bigoplus_{\ell'=1}^{\ell} \bigoplus_{m'=0}^{\ell+m-\ell'} \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X(M_{\ell', m'}^*(\Gamma)), \quad (146)$$

where

$$M_{\ell', m'}^*(\Gamma) = \begin{cases} M_{\ell', m'}(\Gamma) & \text{if } \ell' \neq 3, \\ \text{as in (133)} & \text{if } \ell' = 3. \end{cases} \quad (147)$$

The decomposition (146) is orthogonal in the following sense: If

$$F_1 \in \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X(S_{\ell', m'}(\Gamma)), \quad F_2 \in \sum_{X \in \mathcal{X}_{\ell'', m''}^{\ell, m}} X(M_{\ell'', m''}^*(\Gamma)), \quad (148)$$

and if $(\ell', m') \neq (\ell'', m'')$, then $\langle F_1, F_2 \rangle = 0$.

Proof. The proof of (146) is similar to that of Theorem 4.8. Instead of Proposition 4.6 one uses Proposition 4.27. In addition to Propositions 2.15 and 2.17, one also uses Lemma 4.29. We omit the details.

To prove the orthogonality statement, write $F_2 = \sum X_i F'_i + \sum X_j F''_j$, where $F'_i \in S_{\ell'', m''}(\Gamma)$, and the $F''_j \in M_{\ell'', m''}^*(\Gamma)$ are orthogonal to $S_{\ell'', m''}(\Gamma)$. Clearly, if $F' = \sum X_i F'_i$, then $\langle F_1, F_2 \rangle = \langle F_1, F' \rangle$. We are thus reduced to cusp forms, for which the statement follows from the orthogonality of the decomposition in Theorem 4.8. $\square$

Remark 4.31. Not contained in Theorem 4.30 is the case $\ell = 0$. But recall from Lemma 4.21 (or Proposition 4.27) that $N_{0,0}(\Gamma) = \mathbb{C}$, while $N_{0,m}(\Gamma) = 0$ for $m > 0$.

Modular forms orthogonal to cusp forms. We will introduce some notation involving orthogonal complements of cusp forms. First, let $E_{\ell,m}(\Gamma)$ be the orthogonal complement of $S_{\ell,m}(\Gamma)$ inside $M_{\ell,m}(\Gamma)$, so that

$$M_{\ell,m}(\Gamma) = S_{\ell,m}(\Gamma) \oplus E_{\ell,m}(\Gamma). \quad (149)$$

Recall from (134) that $M_{3,m}^*(\Gamma) = \{F \in N_{3,m}^1(\Gamma) \mid LF = E_-F = 0\}$. We let $E_{3,m}^*(\Gamma)$ be the orthogonal complement of $S_{3,m}(\Gamma)$ in $M_{3,m}^*(\Gamma)$, so that

$$M_{3,m}^*(\Gamma) = S_{3,m}(\Gamma) \oplus E_{3,m}^*(\Gamma). \quad (150)$$

Recall that in Sect. 4.4 we defined $\mathcal{E}_{\ell,m}(\Gamma)$ to be the orthogonal complement of $N_{\ell,m}(\Gamma)^\circ$ in $N_{\ell,m}(\Gamma)$, so that

$$N_{\ell,m}(\Gamma) = N_{\ell,m}(\Gamma)^\circ \oplus \mathcal{E}_{\ell,m}(\Gamma). \quad (151)$$

Lemma 4.32. *Let ℓ be a positive integer, and m a non-negative integer. Then:*

- (1) $E_{3,m}^*(\Gamma) \subset \mathcal{E}_{3,m}(\Gamma)$.
- (2) $\mathcal{E}_{3,m}(\Gamma) \cap M_{3,m}^*(\Gamma) = E_{3,m}^*(\Gamma)$.
- (3) $E_{\ell,m}(\Gamma) \subset \mathcal{E}_{\ell,m}(\Gamma)$.
- (4) $\mathcal{E}_{\ell,m}(\Gamma) \cap M_{\ell,m}(\Gamma) = E_{\ell,m}(\Gamma)$.

Proof. (1) Let $F \in E_{3,m}^*(\Gamma)$ and $G \in N_{3,m}(\Gamma)^\circ$; we have to show that $\langle F, G \rangle = 0$. We work instead with the corresponding automorphic forms Φ_F, Φ_G , and will show that $\langle \Phi_F, \Phi_G \rangle = 0$. We may assume that Φ_G generates an irreducible module $L(\kappa)$. Recall from the definition of the space $M_{3,m}^*(\Gamma)$ that Φ_F generates either a module $L(\mu)$, where $\mu = (m+3, 3)$, or a module $N(\lambda)^\vee$, where $\lambda = (m+3, 1)$. Assume that $\langle \Phi_F, \Phi_G \rangle \neq 0$; we will obtain a contradiction. Since the modules $\langle \Phi_F \rangle$ and $\langle \Phi_G \rangle \cong L(\kappa)$ pair non-trivially, we get a non-zero $\mathfrak{g}_\mathbb{C}$ -map

$$L(\mu) \longrightarrow L(\kappa) \quad \text{or} \quad N(\lambda)^\vee \longrightarrow L(\kappa).$$

In either case we conclude $L(\kappa) \cong L(\mu)$, hence $\kappa = \mu$. It follows that G is holomorphic, therefore an element of $S_{3,m}(\Gamma)$. Since $F \in E_{3,m}^*(\Gamma)$, we have $\langle F, G \rangle = 0$, contradicting our assumption $\langle \Phi_F, \Phi_G \rangle \neq 0$.

(2) follows from (1), (3) is proved in a way analogous to (1), and (4) follows from (3). $\square$

Theorem 4.33 (Structure theorem for modular forms orthogonal to cusp forms). *Let ℓ be a positive integer, and m a non-negative integer. Let the sets $\mathcal{X}_{\ell',m'}^{\ell,m}$ be defined as in (107) and (108). Then we have a direct sum decomposition*

$$\mathcal{E}_{\ell,m}(\Gamma) = \bigoplus_{\ell'=1}^{\ell} \bigoplus_{m'=0}^{\ell+m-\ell'} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(E_{\ell',m'}^*(\Gamma)), \quad (152)$$

where

$$E_{\ell',m'}^*(\Gamma) = \begin{cases} E_{\ell',m'}(\Gamma) & \text{if } \ell' \neq 3, \\ \text{as in (150)} & \text{if } \ell' = 3. \end{cases} \quad (153)$$

Proof. By Lemma 4.32, $E_{\ell',m'}^*(\Gamma) \subset \mathcal{E}_{\ell',m'}(\Gamma)$ for all ℓ', m' . Lemma 4.15 therefore implies that the right hand side is contained in the left hand side. The reverse inclusion follows in a straightforward way from Theorem 4.30. $\square$

5. ADELIZATION AND ARITHMETICITY

5.1. Adelization and automorphic representations. Throughout this section, we let G denote the group GSp_4 . Let K_∞ denote the maximal compact subgroup of $\mathrm{Sp}_4(\mathbb{R})$, and for each prime p , put $K_p = G(\mathbb{Z}_p)$. Recall that an automorphic form on $G(\mathbb{A})$ is a smooth function on $G(\mathbb{A})$ that is left $G(\mathbb{Q})$ -invariant, $\mathcal{Z}$ -finite, K_∞ -finite and slowly increasing; here $\mathcal{Z}$ is the center of $\mathcal{U}(\mathfrak{g}'_\mathbb{C})$, where $\mathfrak{g}' \cong \mathbb{R} \oplus \mathfrak{g}$ is the Lie algebra of $\mathrm{GSp}_4(\mathbb{R})$. We let $\mathcal{A}(G)$ denote the space

of automorphic forms on $G(\mathbb{A})$ and $\mathcal{A}(G)^\circ$ denote the subspace of cusp forms on $G(\mathbb{A})$. All the automorphic forms we will consider in the following will be annihilated by the center $\mathbb{R}$ of $\mathfrak{g}'$.

For each prime p , and each positive integer N , define a compact open subgroup K_p^N of $G(\mathbb{Z}_p)$ by

$$K_p^N = \left\{ g \in G(\mathbb{Z}_p) \mid g \equiv \begin{bmatrix} I_2 & \\ & aI_2 \end{bmatrix} \pmod{N}, a \in \mathbb{Z}_p^\times \right\}. \quad (154)$$

Note that our choice of K_p^N satisfies the following properties:

- $K_p^N = G(\mathbb{Z}_p)$ for all primes p not dividing N ,
- The multiplier map $\mu_2 : K_p^N \mapsto \mathbb{Z}_p^\times$ is surjective for all primes p ,
- $\Gamma(N) = G(\mathbb{Q}) \cap G(\mathbb{R})^+ \prod_{p < \infty} K_p^N$.

As always, let ℓ, m denote integers with $m \geq 0$. Let Γ be a congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$ and F be an element of $C_{\ell,m}^\infty(\Gamma)$. Let N be any integer such that $\Gamma(N) \subset \Gamma$. By Lemma 3.2, we can attach to F a function Φ on $\mathrm{Sp}_4(\mathbb{R})$ that is left invariant by Γ . By strong approximation, we can write any element $g \in G(\mathbb{A})$ as

$$g = \lambda g_{\mathbb{Q}} g_\infty k_{\mathfrak{f}}, \quad g_{\mathbb{Q}} \in G(\mathbb{Q}), \quad g_\infty \in \mathrm{Sp}_4(\mathbb{R}), \quad k_{\mathfrak{f}} \in \prod_p K_p^N, \quad \lambda \in Z_G(\mathbb{R})^+,$$

We define the *adelization* Φ_F of F to be the function on $G(\mathbb{A})$ defined by $\Phi_F(g) = \Phi(g_\infty)$. This is well defined because of the way the groups K_p^N were chosen. Furthermore, it is independent of the choice of N . By construction, it is clear that $\Phi_F(hg) = \Phi_F(g)$ for all $h \in G(\mathbb{Q}), g \in G(\mathbb{A})$. It is also easy to see that the mapping $F \mapsto \Phi_F$ is linear. The following is immediate from Proposition 4.5.

Proposition 5.1. *Let Γ be a congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$ and F be an element of $N_{\ell,m}(\Gamma)$. Let Φ_F be the adelization of F . Then $\Phi_F \in \mathcal{A}(G)$. If $F \in N_{\ell,m}(\Gamma)^\circ$, then $\Phi_F \in \mathcal{A}(G)^\circ$.*

Let $F \in N_{\ell,m}(\Gamma)$ and $\Phi_F \in \mathcal{A}(G)$ be its adelization as defined above. Then Φ_F generates a representation π_F under the natural right regular action⁷ of $G(\mathbb{A})$. From the results of the previous sections it follows that any irreducible subquotient of π_F is an irreducible *automorphic* representation of $G(\mathbb{A})$; it is cuspidal whenever $F \in N_{\ell,m}(\Gamma)^\circ$.

Let $X, \mathcal{X}$ be as in Lemma 4.1. Let $F \in N_{\ell,m}(\Gamma)$ be such that Φ_F generates a factorizable representation $\pi = \otimes_v \pi_v$ of $G(\mathbb{A})$, and suppose that Φ_F corresponds to a factorizable vector $\phi = \otimes_v \phi_v$ inside π . Then, if $G := XF \in N_{\ell',m'}(\Gamma)$, then Φ_G is the vector inside π corresponding to $\otimes_{p < \infty} \phi_v \otimes (X\phi_\infty)$. In particular, if π is an irreducible automorphic representation, then Φ_G generates π . This is immediate from (96), the definition of the adelization map, and the fact that X does not alter the components of F at any of the finite places.

Proposition 5.2. *Let $F \in M_{\ell,m}^*(\Gamma)$ and π_F be the $(\mathfrak{g}, K_\infty) \times G(\mathbb{A}_{\mathfrak{f}})$ -module generated by Φ_F . Let $\pi = \otimes_v \pi_v$ be any irreducible subquotient of π_F .*

- (1) *If $\ell \neq 3$, then $\pi_\infty \simeq L(\ell + m, \ell)$.*
- (2) *If $\ell = 3$, then π_∞ is isomorphic to either $L(3 + m, 3)$ or $L(3 + m, 1)$.*

Proof. This can be derived in a straightforward way from our results in Sect. 4.6. $\square$

5.2. Arithmeticity for nearly holomorphic forms. Recall that any $F \in N_{\ell,m}(\Gamma) = \bigcup_{p \geq 0} N_{\ell,m}^p(\Gamma)$ has a Fourier expansion as follows (note the difference in normalization between (156) and (100); this is for arithmetic purposes):

$$F(Z) = \sum_{Q \in M_2^{\mathrm{sym}}(\mathbb{Q})} a(Q) e^{2\pi i \mathrm{Tr}(QZ)}, \quad (155)$$

⁷More precisely, one takes the right regular action of $G(\mathbb{A}_{\mathfrak{f}})$ together with the action of the Lie algebra at the infinite place.

where

$$a(Q) = \sum_{\alpha, \beta, \gamma} a_{\alpha, \beta, \gamma}(Q) \left(\frac{y}{2\pi\Delta} \right)^\alpha \left(\frac{v}{2\pi\Delta} \right)^\beta \left(\frac{y'}{2\pi\Delta} \right)^\gamma, \quad a_{\alpha, \beta, \gamma}(Q) \in W_m. \quad (156)$$

We note that $a_{\alpha, \beta, \gamma}(Q) = 0$ unless $Q \in \frac{1}{N} M_2^{\text{sym}}(\mathbb{Z})$ for some integer N . Given any $\sigma \in \text{Aut}(\mathbb{C})$, we define a function ${}^\sigma F$ via the action of σ on the elements $a_{\alpha, \beta, \gamma}(Q)$:

$${}^\sigma F(Z) = \sum_{Q \in M_2^{\text{sym}}(\mathbb{Q})} {}^\sigma a(Q) e^{2\pi i \text{Tr}(QZ)},$$

where

$${}^\sigma a(Q) := \sum_{j=0}^m \sum_{\alpha, \beta, \gamma} \sigma(a_{j, \alpha, \beta, \gamma}(Q)) \left(\frac{y}{2\pi\Delta} \right)^\alpha \left(\frac{v}{2\pi\Delta} \right)^\beta \left(\frac{y'}{2\pi\Delta} \right)^\gamma S^{m-j} T^j. \quad (157)$$

For any subfield L of $\mathbb{C}$, define $N_{\ell, m}(\Gamma; L)$ to be the subspace of $N_{\ell, m}(\Gamma)$ consisting of the forms that are fixed by $\text{Aut}(\mathbb{C}/L)$. Define $N_{\ell, m}(\Gamma; L)^\circ$, $N_{\ell, m}^p(\Gamma; L)$, $N_{\ell, m}^p(\Gamma; L)^\circ$, $M_{\ell, m}(\Gamma; L)$, $S_{\ell, m}(\Gamma; L)$ similarly.

We say that a congruence subgroup Γ of $\text{Sp}_4(\mathbb{Q})$ is “nice” if there exists a compact open subgroup K_0 of $G(\mathbb{A}_f)$ with the following properties.

- (1) $K_0 = \prod_{p < \infty} K_{0, p}$, where, for each prime p , $K_{0, p}$ is a compact open subgroup of $G(\mathbb{Q}_p)$ with $K_{0, p} = G(\mathbb{Z}_p)$ for almost all primes.
- (2) For all p , and all $x \in \mathbb{Z}_p^\times$, we have

$$\text{diag}(1, 1, x, x) K_{0, p} \text{diag}(1, 1, x^{-1}, x^{-1}) = K_{0, p}.$$

- (3) $K_0 \text{GSp}_4(\mathbb{R})^+ \cap \text{GSp}_4(\mathbb{Q}) = \Gamma$.

We note that all congruence subgroups that are naturally encountered in the theory, such as the principal, Siegel, Klingen, Borel or paramodular congruence subgroups, are nice in the above sense. The following result follows from [39, Theorem 14.13].

Theorem 5.3 (Shimura). *Let Γ be a nice congruence subgroup of $\text{Sp}_4(\mathbb{Q})$. Then for all $p \geq 0$ we have the equalities*

$$\begin{aligned} N_{\ell, m}^p(\Gamma) &= N_{\ell, m}^p(\Gamma; \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C}, \\ N_{\ell, m}^p(\Gamma)^\circ &= N_{\ell, m}^p(\Gamma; \mathbb{Q})^\circ \otimes_{\mathbb{Q}} \mathbb{C}. \end{aligned}$$

In particular, the action of $\text{Aut}(\mathbb{C})$ preserves the above spaces.

Remark 5.4. Theorem 14.13 of [39] had the added condition that $M_{k, 0}(\Gamma; \overline{\mathbb{Q}}) \neq \{0\}$ for some $0 < k \in \mathbb{Z}$. This is clearly true in our case. Indeed, we have $\Gamma \subset \gamma^{-1} \Gamma^{\text{para}}(N) \gamma$ for some squarefree integer N and some $\gamma \in G(\mathbb{Q})$; this is because every compact open subgroup of $G(\mathbb{Q}_p)$ is either contained in a conjugate of $G(\mathbb{Z}_p)$ or in a conjugate of the local paramodular group at p . Let $F_1 \in S_{10, 0}(\text{Sp}_4(\mathbb{Z}); \mathbb{Q})$ be the unique weight 10 cusp form of full level. Then $F = (\prod_{p|N} \theta_p) F_1$ belongs to $S_{10, 0}(\Gamma^{\text{para}}(N); \overline{\mathbb{Q}})$, where θ_p is as in [31]; the fact that the Fourier coefficients are algebraic follow from the q -expansion principle. Observe that θ_p is injective by Theorem 7.2 of [31], so that $F \neq 0$. Finally, $F|_{k, 0} \gamma$ is an element of $S_{10, 0}(\Gamma; \overline{\mathbb{Q}})$.

Let $\mathcal{X}_+$ be the free monoid consisting of all (finite) strings of the symbols X_+ , U , E_+ , and D_+ in the left column of Table 1. Clearly $\mathcal{X}_+$ is a submonoid of the monoid $\mathcal{X}$ defined earlier, and furthermore contains all the subsets $\mathcal{X}_{\ell', m'}^{\ell, m}$ introduced for the purpose of stating the structure theorems. Each element $X \in \mathcal{X}_+$ is an operator that for any particular ℓ, m, p , takes $N_{\ell, m}^p(\Gamma)$ to $N_{\ell_1, m_1}^{p_1}(\Gamma)$, where the integers ℓ_1, m_1, p_1 (that depend on ℓ, m, p and X) can be easily calculated using Table 2. In particular, the non-negative integer $v = p_1 - p$ depends only on X . Precisely, $v = 1$ for X_+ , U , and E_+ ; $v = 2$ for D_+ . For longer strings, v can be calculated by adding up the contributions from the individual operators.

Definition 5.5. For any $X \in \mathcal{X}_+$, we define the degree of X to be the integer v described above.

The following key proposition, when combined with our structure theorems, allows us to transfer arithmeticity results from holomorphic forms to nearly holomorphic forms.

Proposition 5.6. Let $X \in \mathcal{X}_+$ and let v be the degree of X . Then, for all $F \in N_{\ell,m}(\Gamma)$, and all $\sigma \in \text{Aut}(\mathbb{C})$, we have

$$\sigma((2\pi)^{-v}XF) = (2\pi)^{-v}X(\sigma F).$$

Proof. It suffices to prove this for each of the basic operators X_+ , U , E_+ , and D_+ . Using equations (85) – (92), we note that the action of the operators X_+ , U , and E_+ on the component functions of F are given by rational linear combinations from the following set S of operators on $C^\infty(\mathbb{H}_2)$,

$$S = \left\{ \frac{y}{\Delta}, \frac{y'}{\Delta}, \frac{v}{\Delta}, 2i\frac{\partial}{\partial z}, 2i\frac{\partial}{\partial \tau}, 2i\frac{\partial}{\partial \tau'} \right\}.$$

Furthermore, the action of the operator D_+ on the component functions of F is given by rational linear combinations of the objects formed by taking the composition of exactly *two* operators from the set S .

Therefore, to complete the proof, it suffices to show that for each element $Q \in M_2^{\text{sym}}(\mathbb{Q})$, each triple of non-negative integers α, β, γ , and each operator $s \in S$, there exist *rational numbers* $a_{\alpha', \beta', \gamma'}(Q)$ indexed by a finite set of triples of non-negative integers α', β', γ' , such that

$$\begin{aligned} (2\pi)^{-1}s \left(\left(\frac{y}{2\pi\Delta} \right)^\alpha \left(\frac{v}{2\pi\Delta} \right)^\beta \left(\frac{y'}{2\pi\Delta} \right)^\gamma e^{2\pi i \text{Tr}(QZ)} \right) \\ = \sum_{\alpha', \beta', \gamma'} a_{\alpha', \beta', \gamma'}(Q) \left(\frac{y}{2\pi\Delta} \right)^{\alpha'} \left(\frac{v}{2\pi\Delta} \right)^{\beta'} \left(\frac{y'}{2\pi\Delta} \right)^{\gamma'} e^{2\pi i \text{Tr}(QZ)}. \end{aligned}$$

This is an elementary calculation and can be easily verified for each element s of S . We omit the details. $\square$

Isotypic projections. By our structure theorem, the space $N_{\ell,m}(\Gamma)$ decomposes as a direct sum as follows:

$$N_{\ell,m}(\Gamma) = \bigoplus_{\substack{0 \leq \ell' \leq \ell \\ 0 \leq \ell' + m' \leq \ell + m \\ m' \geq 0}} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(M_{\ell',m'}^*(\Gamma)), \quad (158)$$

where we adopt the convention that $M_{\ell',m'}^*(\Gamma) := M_{\ell',m'}(\Gamma)$ whenever $\ell' \neq 3$. The identical decomposition holds for the cuspidal subspace.

Definition 5.7. For each quadruple of integers ℓ, m, ℓ', m' with m, m' non-negative, define an endomorphism $\mathfrak{p}_{\ell',m'}$ of $N_{\ell,m}(\Gamma)$ by the projection map

$$N_{\ell,m}(\Gamma) \longrightarrow \left(\sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(M_{\ell',m'}^*(\Gamma)) \right) \subset N_{\ell,m}(\Gamma)$$

given by the direct sum decomposition (158). In particular, if the set $\mathcal{X}_{\ell',m'}^{\ell,m}$ is empty, we have $\mathfrak{p}_{\ell',m'} = 0$.

Lemma 5.8. Suppose that $F \in N_{\ell,m}(\Gamma)$. Then the following hold.

- (1) Suppose that $F \in N_{\ell,m}(\Gamma)^\circ$, resp. $F \in \mathcal{E}_{\ell,m}(\Gamma)$. Then, $\mathfrak{p}_{\ell',m'}(F) \in N_{\ell,m}(\Gamma)^\circ$, resp. $\mathfrak{p}_{\ell',m'}(F) \in \mathcal{E}_{\ell,m}(\Gamma)$.

(2) We have

$$F = \sum_{\ell' \geq 0, m' \geq 0} \mathfrak{p}_{\ell', m'}(F).$$

The above sum is orthogonal in the sense that if $(\ell'_1, m'_1) \neq (\ell'_2, m'_2)$, and $\mathfrak{p}_{\ell'_1, m'_1}(F) \in N_{\ell, m}(\Gamma)^\circ$, then

$$\langle \mathfrak{p}_{\ell'_1, m'_1}(F), \mathfrak{p}_{\ell'_2, m'_2}(F) \rangle = 0.$$

(3) Suppose that $F \in N_{\ell, m}(\Gamma)$, and $G \in S_{\ell', m'}(\Gamma)$. Then, for all $X \in \mathcal{X}_{\ell', m'}^{\ell, m}$,

$$\langle F, XG \rangle = \langle \mathfrak{p}_{\ell', m'}(F), XG \rangle.$$

Proof. All the parts follow directly from the structure theorems and our definition of the projection map. We omit the details. $\square$

Lemma 5.9. Let Γ be a nice congruence subgroup of $\mathrm{Sp}_4(\mathbb{Q})$. Then we have the equality

$$M_{\ell', m'}^*(\Gamma) = M_{\ell', m'}^*(\Gamma; \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C}.$$

In particular, the action of $\mathrm{Aut}(\mathbb{C})$ preserves the above space.

Proof. We only need to consider the case $\ell' = 3$, since otherwise $M_{\ell', m'}^*(\Gamma) = M_{\ell', m'}(\Gamma)$ and this case has already been covered by Theorem 5.3. So, assume $\ell' = 3$. Let $F \in M_{3, m'}^*(\Gamma)$ and $\sigma \in \mathrm{Aut}(\mathbb{C})$. It suffices to show that ${}^\sigma F \in M_{3, m'}^*(\Gamma)$; see Lemma 3.19 of [33]. We already know from Theorem 5.3 that ${}^\sigma F \in N_{3, m'}^1(\Gamma)$. So, to complete the proof, we only need to show that $L({}^\sigma F) = E_-({}^\sigma F) = 0$. But this is an immediate consequence of Proposition 5.6. $\square$

We now state our main arithmeticity result concerning this projection map.

Proposition 5.10. For all quadruples (ℓ, m, ℓ', m') , all $\sigma \in \mathrm{Aut}(\mathbb{C})$, and all $F \in N_{\ell, m}(\Gamma)$, we have

$$\mathfrak{p}_{\ell', m'}({}^\sigma F) = {}^\sigma(\mathfrak{p}_{\ell', m'}(F)).$$

Proof. By shrinking Γ if necessary, we may assume Γ is nice. Using the structure theorem 4.30, write

$$F = \sum_{\ell', m'} \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X'(F_X), \quad \text{where } X' = (2\pi)^{-v(X)} X$$

and $F_X \in M_{\ell', m'}^*(\Gamma)$. Then, by Proposition 5.6,

$${}^\sigma F = \sum_{\ell', m'} \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} {}^\sigma(X'(F_X)) = \sum_{\ell', m'} \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X'({}^\sigma F_X).$$

By Theorem 5.3 and Lemma 5.9, the modular form ${}^\sigma F_X$ lies in $M_{\ell', m'}^*(\Gamma)$. Hence

$$\mathfrak{p}_{\ell', m'}({}^\sigma F) = \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} X'({}^\sigma F_X) = \sum_{X \in \mathcal{X}_{\ell', m'}^{\ell, m}} {}^\sigma X'(F_X) = {}^\sigma(\mathfrak{p}_{\ell', m'}(F)).$$

This completes the proof. $\square$

Remark 5.11. In the special case $\ell' = \ell$, $m' = m$, Shimura defined the map $\mathfrak{p}_{\ell, m} : N_{\ell, m}(\Gamma) \rightarrow M_{\ell, m}(\Gamma)$ and called it the holomorphic projection map. He was able to prove results of $\mathrm{Aut}(\mathbb{C})$ -equivariance in this special case under the additional assumption that either $F \in N_{\ell, m}(\Gamma)^\circ$ or $m = 0$; see [39, Prop. 15.3, Prop. 15.6].

Definition 5.12. Let $\mathfrak{q}$ denote the natural projection map from nearly holomorphic modular forms to nearly holomorphic cusp forms, i.e., $\mathfrak{q} : \oplus_{\ell, m} N_{\ell, m}(\Gamma) \rightarrow \oplus_{\ell, m} N_{\ell, m}(\Gamma)^\circ$ is obtained from the orthogonal direct sum decomposition

$$N_{\ell, m}(\Gamma) = N_{\ell, m}(\Gamma)^\circ \oplus \mathcal{E}_{\ell, m}(\Gamma).$$

Definition 5.13. Define $\mathfrak{p}_{\ell',m'}^\circ = \mathfrak{q} \circ \mathfrak{p}_{\ell',m'}$.

Thus,

$$\mathfrak{p}_{\ell',m'}^\circ : N_{\ell,m}(\Gamma) \rightarrow \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(S_{\ell',m'}(\Gamma)) \subset N_{\ell,m}(\Gamma)^\circ.$$

If $F \in N_{\ell,m}(\Gamma)^\circ$, then $\mathfrak{p}_{\ell',m'}^\circ(F) = \mathfrak{p}_{\ell',m'}(F)$. It is clear that for all $F \in N_{\ell,m}(\Gamma)$, $G \in S_{\ell',m'}(\Gamma)$, $X \in \mathcal{X}_{\ell',m'}^{\ell,m}$, we have

$$\langle F, XG \rangle = \langle \mathfrak{p}_{\ell',m'}(F), XG \rangle = \langle \mathfrak{p}_{\ell',m'}^\circ(F), XG \rangle.$$

Furthermore, if $F \in N_{\ell,m}(\Gamma)$ and we write, using the structure theorem,

$$F = \sum_{\ell',m'} \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(F_X),$$

then

$$\mathfrak{p}_{\ell',m'}^\circ(F) = \sum_{X \in \mathcal{X}_{\ell',m'}^{\ell,m}} X(\mathfrak{q}(F_X)).$$

Recall that $E_{\ell,m}(\Gamma)$ denotes the orthogonal complement of $S_{\ell,m}(\Gamma)$ in $M_{\ell,m}(\Gamma)$ and has the property that $E_{\ell,m}(\Gamma) = \mathcal{E}_{\ell,m}(\Gamma) \cap M_{\ell,m}(\Gamma)$; see Lemma 4.32.

Definition 5.14. Given a number field L , we say that $E_{\ell,m}(\Gamma)$ is L -rational if

$$E_{\ell,m}(\Gamma) = E_{\ell,m}(\Gamma; L) \otimes_L \mathbb{C}.$$

Remark 5.15. If $E_{\ell,m}(\Gamma)$ is L -rational, then for all $F \in M_{\ell,m}(\Gamma)$, $\sigma \in \text{Aut}(\mathbb{C}/L)$, we have ${}^\sigma(\mathfrak{q}(F)) = \mathfrak{q}({}^\sigma F)$.

Remark 5.16. The results of Harris (see [16]) imply that if $\ell > 4$ (so that we are in the absolutely convergent range, and so $E_{\ell,m}(\Gamma)$ is spanned by holomorphic Siegel and Klingen Eisenstein series) and Γ is nice, then $E_{\ell,m}(\Gamma)$ is L -rational for some number field L . It is unclear to us if we can always take $L = \mathbb{Q}$ in this case, though we suspect this to be the case.

Proposition 5.17. Suppose that $\ell' > 3$ and $E_{\ell',m'}(\Gamma)$ is L -rational. Then, for all $F \in N_{\ell,m}(\Gamma)$ and $\sigma \in \text{Aut}(\mathbb{C}/L)$,

$$\mathfrak{p}_{\ell',m'}^\circ({}^\sigma F) = {}^\sigma(\mathfrak{p}_{\ell',m'}^\circ(F)).$$

Proof. The proof is essentially identical to that of Proposition 5.10. $\square$

We end this section with an arithmeticity result for ratios of Petersson inner products.

Proposition 5.18. Let $F \in S_{\ell,m}(\Gamma)$ have the property that for all $G \in S_{\ell,m}(\Gamma)$ and all $\sigma \in \text{Aut}(\mathbb{C})$, we have

$$\sigma \left(\frac{\langle G, F \rangle}{\langle F, F \rangle} \right) = \frac{\langle {}^\sigma G, {}^\sigma F \rangle}{\langle {}^\sigma F, {}^\sigma F \rangle}.$$

Let ℓ_1, m_1 be integers such that $\mathcal{X}_{\ell_1, m_1}^{\ell_1, m_1}$ is a singleton set equal to $\{X\}$. Then for all $G \in N_{\ell_1, m_1}(\Gamma)^\circ$, and all $\sigma \in \text{Aut}(\mathbb{C})$, we have

$$\sigma \left(\frac{\langle G, XF \rangle}{\langle XF, XF \rangle} \right) = \frac{\langle {}^\sigma G, {}^\sigma XF \rangle}{\langle {}^\sigma XF, {}^\sigma XF \rangle}.$$

Remark 5.19. It is expected that whenever $\ell \geq 6$, all Hecke eigenforms F in $S_{\ell,m}(\Gamma)$ with coefficients in a CM field have the property required in the above proposition. This has been proved in many special cases, e.g., when $\Gamma = \text{Sp}_4(\mathbb{Z})$ (see [41]).

Proof. By (3) of Lemma 5.8,

$$\frac{\langle G, XF \rangle}{\langle XF, XF \rangle} = \frac{\langle \mathfrak{p}_{\ell,m}(G), XF \rangle}{\langle XF, XF \rangle}.$$

Now, $\mathfrak{p}_{\ell,m}(G) = XG'$ for some $G' \in S_{\ell,m}(\Gamma)$. By Proposition 4.17,

$$\sigma \left(\frac{\langle G, XF \rangle}{\langle XF, XF \rangle} \right) = \sigma \left(\frac{\langle X(G'), XF \rangle}{\langle XF, XF \rangle} \right) = \sigma \left(\frac{\langle G', F \rangle}{\langle F, F \rangle} \right).$$

Similarly, using Proposition 5.6,

$$\frac{\langle {}^\sigma G, {}^\sigma XF \rangle}{\langle {}^\sigma XF, {}^\sigma XF \rangle} = \frac{\langle {}^\sigma G', {}^\sigma F \rangle}{\langle {}^\sigma F, {}^\sigma F \rangle}.$$

Now the result follows from the property of F assumed in the statement of the proposition. $\square$

Remark 5.20. The condition that $\mathcal{X}_{\ell,m}^{\ell_1,m_1}$ is a singleton set is satisfied when $\ell_1 = \ell + m$ and $m_1 = 0$, provided m is even. In this case, we have $\mathcal{X}_{\ell,m}^{\ell_1,m_1} = \{U^{m/2}\}$. The application of the above proposition in this special case will be of crucial importance in our upcoming work.

Proposition 5.21. *Let F be as in Proposition 5.18. Assume further that $\ell > 3$ and $E_{\ell,m}(\Gamma)$ is L -rational for some number field L .*

Let ℓ_1, m_1 be integers such that $\mathcal{X}_{\ell,m}^{\ell_1,m_1}$ is a singleton set equal to $\{X\}$. Then for all $G \in N_{\ell_1,m_1}(\Gamma)$, and all $\sigma \in \text{Aut}(\mathbb{C}/L)$, we have

$$\sigma \left(\frac{\langle G, XF \rangle}{\langle XF, XF \rangle} \right) = \frac{\langle {}^\sigma G, {}^\sigma XF \rangle}{\langle {}^\sigma XF, {}^\sigma XF \rangle}.$$

Proof. The proof is identical to Proposition 5.18, except that we use $\mathfrak{p}_{\ell,m}^\circ$. $\square$

REFERENCES

- [1] Mahdi Asgari and Ralf Schmidt. Siegel modular forms and representations. *Manuscripta Math.*, 104(2):173–200, 2001.
- [2] Antonia Blüher. Near holomorphy, arithmeticity, and the theta correspondence. In *Automorphic forms, automorphic representations, and arithmetic (Fort Worth, TX, 1996)*, volume 66 of *Proc. Sympos. Pure Math.*, pages 9–26. Amer. Math. Soc., Providence, RI, 1999.
- [3] Siegfried Böcherer and Bernhard Heim. Critical values of L -functions on $\text{GSp}_2 \times \text{GL}_2$. *Math. Z.*, 254(3):485–503, 2006.
- [4] Siegfried Böcherer, Takakazu Satoh, and Tadashi Yamazaki. On the pullback of a differential operator and its application to vector valued Eisenstein series. *Comment. Math. Univ. St. Paul.*, 41(1):1–22, 1992.
- [5] Siegfried. Böcherer and Claus-Günther Schmidt. p -adic measures attached to Siegel modular forms. *Ann. Inst. Fourier (Grenoble)*, 50(5):1375–1443, 2000.
- [6] Siegfried Böcherer and Rainer Schulze-Pillot. On the central critical value of the triple product L -function. In *Number theory (Paris, 1993–1994)*, volume 235 of *London Math. Soc. Lecture Note Ser.*, pages 1–46. Cambridge Univ. Press, Cambridge, 1996.
- [7] Brian Boe and David Collingwood. A multiplicity one theorem for holomorphically induced representations. *Math. Z.*, 192(2):265–282, 1986.
- [8] Brian Boe and David Collingwood. Multiplicity free categories of highest weight representations. I, II. *Comm. Algebra*, 18(4):947–1032, 1033–1070, 1990.
- [9] Armand Borel and Hervé Jacquet. Automorphic forms and automorphic representations. In *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977)*, Part 1, Proc. Sympos. Pure Math., XXXIII, pages 189–207. Amer. Math. Soc., Providence, R.I., 1979. With a supplement “On the notion of an automorphic representation” by R. P. Langlands.
- [10] Michel Courtieu and Alexei Panchishkin. *Non-Archimedean L -functions and arithmetical Siegel modular forms*, volume 1471 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, second edition, 2004.
- [11] Amine El Gradechi. The Lie theory of certain modular form and arithmetic identities. *Ramanujan J.*, 31(3):397–433, 2013.
- [12] Thomas Enright, Roger Howe, and Nolan Wallach. A classification of unitary highest weight modules. In *Representation theory of reductive groups (Park City, Utah, 1982)*, volume 40 of *Progr. Math.*, pages 97–143. Birkhäuser Boston, Boston, MA, 1983.

- [13] Thomas Enright and Anthony Joseph. An intrinsic analysis of unitarizable highest weight modules. *Math. Ann.*, 288(4):571–594, 1990.
- [14] William Fulton. *Algebraic curves*. Advanced Book Classics. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989. An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original.
- [15] Michael Harris. A note on three lemmas of Shimura. *Duke Math. J.*, 46(4):871–879, 1979.
- [16] Michael Harris. Eisenstein series on Shimura varieties. *Ann. of Math. (2)*, 119(1):59–94, 1984.
- [17] Michael Harris. Arithmetic vector bundles and automorphic forms on Shimura varieties. I. *Invent. Math.*, 82(1):151–189, 1985.
- [18] Michael Harris. Arithmetic vector bundles and automorphic forms on Shimura varieties. II. *Compositio Math.*, 60(3):323–378, 1986.
- [19] Roger Howe and Hanspeter Kraft. Principal covariants, multiplicity-free actions, and the K -types of holomorphic discrete series. In *Geometry and representation theory of real and p -adic groups (Córdoba, 1995)*, volume 158 of *Progr. Math.*, pages 147–161. Birkhäuser Boston, Boston, MA, 1998.
- [20] James Humphreys. *Representations of semisimple Lie algebras in the BGG category $\mathcal{O}$* , volume 94 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [21] Ronald Irving. Projective modules in the category $\mathcal{O}_S$: Loewy series. *Trans. Amer. Math. Soc.*, 291(2):733–754, 1985.
- [22] Hans Plesner Jakobsen. Hermitian symmetric spaces and their unitary highest weight modules. *J. Funct. Anal.*, 52(3):385–412, 1983.
- [23] Anthony Knapp. *Representation theory of semisimple groups*, volume 36 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1986. An overview based on examples.
- [24] Dominic Lanphier. Combinatorics of Maass-Shimura operators. *J. Number Theory*, 128(8):2467–2487, 2008.
- [25] Hans Maass. *Siegel’s modular forms and Dirichlet series*. Lecture Notes in Mathematics, Vol. 216. Springer-Verlag, Berlin-New York, 1971. Dedicated to the last great representative of a passing epoch. Carl Ludwig Siegel on the occasion of his seventy-fifth birthday.
- [26] Goran Muić. Intertwining operators and composition series of generalized and degenerate principal series for $\mathrm{Sp}(4, \mathbb{R})$. *Glas. Mat. Ser. III*, 44(64)(2):349–399, 2009.
- [27] Mark Alan Nappari. *Holomorphic forms canonically attached to nearly holomorphic automorphic forms*. ProQuest LLC, Ann Arbor, MI, 1992. Thesis (Ph.D.)—Brandeis University.
- [28] Alexei Panchishkin. The Maass-Shimura differential operators and congruences between arithmetical Siegel modular forms. *Mosc. Math. J.*, 5(4):883–918, 973–974, 2005.
- [29] Ameya Pitale, Abhishek Saha, and Ralf Schmidt. Representations of $\mathrm{SL}_2(\mathbf{R})$ and nearly holomorphic modular forms. In *Proceedings of RIMS Workshop on Modular forms and automorphic representations*, RIMS Kôkyûroku, No.1973, pages 141–153. Res. Inst. Math. Sci. (RIMS), Kyoto, 2015.
- [30] Ameya Pitale, Abhishek Saha, and Ralf Schmidt. A note on the growth of nearly holomorphic vector-valued Siegel modular forms. In Jan Hendrik Bruinier and Winfried Kohnen, editors, *L -Functions and Automorphic Forms*, pages 185–193. Springer International Publishing, 2017.
- [31] Brooks Roberts and Ralf Schmidt. On modular forms for the paramodular groups. In *Automorphic forms and zeta functions*, pages 334–364. World Sci. Publ., Hackensack, NJ, 2006.
- [32] Abhishek Saha. Pullbacks of Eisenstein series from $\mathrm{GU}(3, 3)$ and critical L -values for $\mathrm{GSp}_4 \times \mathrm{GL}_2$. *Pacific J. Math.*, 246(2):435–486, 2010.
- [33] Abhishek Saha. On ratios of Petersson norms for Yoshida lifts. *Forum Math.*, 27(4):2361–2412, 2015.
- [34] Takakazu Satoh. On certain vector valued Siegel modular forms of degree two. *Math. Ann.*, 274(2):335–352, 1986.
- [35] Goro Shimura. On a class of nearly holomorphic automorphic forms. *Ann. of Math. (2)*, 123(2):347–406, 1986.
- [36] Goro Shimura. Nearly holomorphic functions on Hermitian symmetric spaces. *Math. Ann.*, 278(1-4):1–28, 1987.
- [37] Goro Shimura. On Hilbert modular forms of half-integral weight. *Duke Math. J.*, 55(4):765–838, 1987.
- [38] Goro Shimura. Differential operators, holomorphic projection, and singular forms. *Duke Math. J.*, 76(1):141–173, 1994.
- [39] Goro Shimura. *Arithmeticity in the theory of automorphic forms*, volume 82 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2000.
- [40] David Soudry. The CAP representations of $\mathrm{GSp}(4, \mathbf{A})$. *J. Reine Angew. Math.*, 383:87–108, 1988.
- [41] Yoshinori Takei. On algebraicity of vector valued Siegel modular forms. *Kodai Math. J.*, 15(3):445–457, 1992.
- [42] Eric Urban. Nearly overconvergent modular forms. *Preprint*, 2013.
- [43] Rainer Weissauer. Vektorwertige Siegelsche Modulformen kleinen Gewichtes. *J. Reine Angew. Math.*, 343:184–202, 1983.
- [44] Rainer Weissauer. Modular forms of genus 2 and weight 1. *Math. Z.*, 210(1):91–96, 1992.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019, USA

E-mail address: `apitale@math.ou.edu`

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY UNIVERSITY OF LONDON, LONDON E1 4NS, UK

E-mail address: `abhishek.saha@qmul.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019, USA

E-mail address: `rschmidt@math.ou.edu`