

# Leray-Hopf and Continuity Properties for All Weak Solutions for the 3D Navier-Stokes Equations

December 17, 2021

Nataliia V. Gorban<sup>1</sup>, Pavlo O. Kasyanov<sup>2</sup>, Olha V. Khomenko<sup>3</sup>, and Luisa Toscano<sup>4</sup>.

## Abstract

In this note we prove that each weak solution for the 3D Navier-Stokes system satisfies Leray-Hopf property. Moreover, each weak solution is rightly continuous in the standard phase space  $H$  endowed with the strong convergence topology.

## 1 Introduction and Main Result

Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with rather smooth boundary  $\Gamma = \partial\Omega$ , and  $[\tau, T]$  be a fixed time interval with  $-\infty < \tau < T < +\infty$ . We consider 3D Navier-Stokes system in  $\Omega \times [\tau, T]$

$$\begin{cases} \frac{\partial y}{\partial t} - \nu \Delta y + (y \cdot \nabla)y = -\nabla p + f, & \operatorname{div} y = 0, \\ y|_{\Gamma} = 0, & y|_{t=\tau} = y_{\tau}, \end{cases} \quad (1.1)$$

where  $y(x, t)$  means the unknown velocity,  $p(x, t)$  the unknown pressure,  $f(x, t)$  the given exterior force, and  $y_{\tau}(x)$  the given initial velocity with  $t \in [\tau, T]$ ,  $x \in \Omega$ ,  $\nu > 0$  means the viscosity constant.

Throughout this note we consider generalized setting of Problem (1.1). For this purpose define the usual function spaces

$$\mathcal{V} = \{u \in (C_0^\infty(\Omega))^3 : \operatorname{div} u = 0\}, \quad V_\sigma = \operatorname{cl}_{(H_0^\sigma(\Omega))^3} \mathcal{V}, \quad \sigma \geq 0,$$

where  $\operatorname{cl}_X$  denotes the closure in the space  $X$ . Set  $H := V_0$ ,  $V := V_1$ . It is well known that each  $V_\sigma$ ,  $\sigma > 0$ , is a separable Hilbert space and identifying  $H$  and its dual  $H^*$  we have  $V_\sigma \subset H \subset V_\sigma^*$  with dense and

<sup>1</sup>Institute for Applied System Analysis, National Technical University of Ukraine “Kyiv Polytechnic Institute”, Peremogy ave., 37, build, 35, 03056, Kyiv, Ukraine, nata\_gorban@i.ua

<sup>2</sup>Institute for Applied System Analysis, National Technical University of Ukraine “Kyiv Polytechnic Institute”, Peremogy ave., 37, build, 35, 03056, Kyiv, Ukraine, kasyanov@i.ua.

<sup>3</sup>Institute for Applied System Analysis, National Technical University of Ukraine “Kyiv Polytechnic Institute”, Peremogy ave., 37, build, 35, 03056, Kyiv, Ukraine, olgkhomenko@ukr.net

<sup>4</sup>University of Naples “Federico II”, Dep. Math. and Appl. R.Caccioppoli, via Claudio 21, 80125 Naples, Italy, luisatoscano@libero.it

compact embedding for each  $\sigma > 0$ . We denote by  $(\cdot, \cdot)$ ,  $\|\cdot\|$  and  $((\cdot, \cdot))$ ,  $\|\cdot\|_V$  the inner product and norm in  $H$  and  $V$ , respectively;  $\langle \cdot, \cdot \rangle$  will denote pairing between  $V$  and  $V^*$  that coincides on  $H \times V$  with the inner product  $(\cdot, \cdot)$ . Let  $H_w$  be the space  $H$  endowed with the weak topology. For  $u, v, w \in V$  we put

$$b(u, v, w) = \int_{\Omega} \sum_{i,j=1}^3 u_i \frac{\partial v_j}{\partial x_i} w_j dx.$$

It is known that  $b$  is a trilinear continuous form on  $V$  and  $b(u, v, v) = 0$ , if  $u, v \in V$ . Furthermore, there exists a positive constant  $C$  such that

$$|b(u, v, w)| \leq C \|u\|_V \|v\|_V \|w\|_V, \quad (1.2)$$

for each  $u, v, w \in V$ ; see, for example, Sohr [17, Lemma V.1.2.1] and references therein.

Let  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$  and  $y_{\tau} \in H$ . Recall that the function  $y \in L^2(\tau, T; V)$  with  $\frac{dy}{dt} \in L^1(\tau, T; V^*)$  is a *weak solution* of Problem (1.1) on  $[\tau, T]$ , if for all  $v \in V$

$$\frac{d}{dt} (y, v) + \nu ((y, v)) + b(y, y, v) = \langle f, v \rangle \quad (1.3)$$

in the sense of distributions, and

$$y(\tau) = y_{\tau}. \quad (1.4)$$

The weak solution  $y$  of Problem (1.1) on  $[\tau, T]$  is called a *Leray-Hopf* solution of Problem (1.1) on  $[\tau, T]$ , if  $y$  satisfies the energy inequality:

$$V_{\tau}(y(t)) \leq V_{\tau}(y(s)) \quad \text{for all } t \in [s, T], \text{ a.e. } s > \tau \text{ and } s = \tau, \quad (1.5)$$

where

$$V_{\tau}(y(\varsigma)) := \frac{1}{2} \|y(\varsigma)\|^2 + \nu \int_{\tau}^{\varsigma} \|y(\xi)\|_V^2 d\xi - \int_{\tau}^{\varsigma} \langle f(\xi), y(\xi) \rangle d\xi, \quad \varsigma \in [\tau, T]. \quad (1.6)$$

For each  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$  and  $y_{\tau} \in H$  there exists at least one Leray-Hopf solution of Problem (1.1); see, for example, Temam [18, Chapter III] and references therein. Moreover,  $y \in C([\tau, T], H_w)$  and  $\frac{dy}{dt} \in L^{\frac{4}{3}}(\tau, T; V^*) + L^1(\tau, T; H)$ . If  $f \in L^2(\tau, T; V^*)$ , then, additionally,  $\frac{dy}{dt} \in L^{\frac{4}{3}}(\tau, T; V^*)$ . In particular, the initial condition (1.4) makes sense.

The following Theorem 1.1 implies that each weak solution of the 3D Navier-Stokes system is Leray-Hopf one and it is rightly strongly continuous in  $H$  at all the points  $t \in [\tau, T]$ . This theorem is the main result of this note.

**Theorem 1.1.** *Let  $-\infty < \tau < T < +\infty$ ,  $y_{\tau} \in H$ ,  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$ , and  $y$  be a weak solution of Problem (1.1) on  $[\tau, T]$ . Then the following statements hold:*

- (a)  $y \in C([\tau, T], H_w)$  and the following energy inequality holds:

$$V_{\tau}(y(t)) \leq V_{\tau}(y(s)) \quad \text{for all } t, s \in [\tau, T], \quad t \geq s, \quad (1.7)$$

where  $V_{\tau}$  is defined in formula (1.6);

(b) for each  $t \in [\tau, T)$  the following convergence holds:

$$y(s) \rightarrow y(t) \text{ strongly in } H \text{ as } s \rightarrow t+;$$

(c) the function  $t \rightarrow \|y(t)\|^2$  is of bounded variation on  $[\tau, T]$ .

**Remark 1.2.** Since a real function of bounded variation has no more than countable set of discontinuity points, then statement (a) of Theorem 1.1, weak continuity in Hilbert space  $H$  of each weak solution of Problem (1.1) on  $[\tau, T]$ , yield that each weak solution of the 3D Navier-Stokes system has no more than countable set of discontinuity points in the phase space  $H$  endowed with the strong convergence topology. Theorem 1.1 partially clarifies the results provided in Ball [1]; Balibrea et al. [2]; Barbu et al. [3]; Cao and Titi [4]; Chepyzhov and Vishik [5]; Cheskidov and Shvydkoy [6]; Kapustyan et al. [9, 10]; Kloeden et al. [13]; Sohr [17] and references therein.

## 2 Topological Properties of Solutions for Auxiliary Control Problem

Let  $-\infty < \tau < T < +\infty$ . We consider the following space of parameters:

$$\mathbb{U}_{\tau, T} := (L^2(\tau, T; V)) \times (L^2(\tau, T; V^*) + L^1(\tau, T; H)) \times H.$$

Each triple  $(u, g, z_\tau) \in \mathbb{U}_{\tau, T}$  is called *admissible* for the following auxiliary control problem:

**Problem (C)** on  $[\tau, T]$  with  $(u, g, z_\tau) \in \mathbb{U}_{\tau, T}$ : find  $z \in L^2(\tau, T; V)$  with  $\frac{dz}{dt} \in L^1(\tau, T; V^*)$  such that  $z(\tau) = z_\tau$  and for all  $v \in V$

$$\frac{d}{dt}(z, v) + \nu((z, v)) + b(u, z, v) = \langle g, v \rangle \quad (2.1)$$

in the sense of distributions; cf. Kapustyan et al. [9, 10]; Kasyanov et al. [11, 12]; Melnik and Toscano [14]; Zgurovsky et al. [19, Chapter 6].

As usual, let  $A : V \rightarrow V^*$  be the linear operator associated with the bilinear form  $((u, v)) = \langle Au, v \rangle$ ,  $u, v \in V$ . For  $u, v \in V$  we denote by  $B(u, v)$  the element of  $V^*$  defined by  $\langle B(u, v), w \rangle = b(u, v, w)$ , for all  $w \in V$ . Then Problem (C) on  $[\tau, T]$  with  $(u, g, z_\tau) \in \mathbb{U}_{\tau, T}$  can be rewritten as: find  $z \in L^2(\tau, T; V)$  with  $\frac{dz}{dt} \in L^1(\tau, T; V^*)$  such that

$$\frac{dz}{dt} + \nu Az + B(u, z) = g, \text{ in } V^*, \text{ and } z(\tau) = z_\tau. \quad (2.2)$$

The following theorem establishes the uniqueness properties for solutions of Problem (C).

**Theorem 2.1.** Let  $-\infty < \tau < T < +\infty$  and  $u \in L^2(\tau, T; V)$ . Then Problem (C) on  $[\tau, T]$  with  $(u, \bar{0}, \bar{0}) \in \mathbb{U}_{\tau, T}$  has the unique solution  $z \equiv \bar{0}$ .

We recall, that  $\{w_1, w_2, \dots\} \subset \mathcal{V}$  is the *special basis*, if  $((w_j, v)) = \lambda_j(w_j, v)$  for each  $v \in V$  and  $j = 1, 2, \dots$ , where  $0 < \lambda_1 \leq \lambda_2 \leq \dots$  is the sequence of eigenvalues. Let  $P_m$  be the projection operator of  $H$  onto  $H_m := \text{span}\{w_1, \dots, w_m\}$ , that is  $P_m v = \sum_{i=1}^m (v, w_i) w_i$  for each  $v \in H$  and  $m = 1, 2, \dots$ . Of course we may consider  $P_m$  as a projection operator that acts from  $V_\sigma$  onto  $H_m$  for each  $\sigma > 0$  and, since  $P_m^* = P_m$ , we deduce that  $\|P_m\|_{\mathcal{L}(V_\sigma^*; V_\sigma^*)} \leq 1$ . Note that  $(w_j, v)_{V_\sigma} = \lambda_j^\sigma(w_j, v)$  for each  $v \in V_\sigma$  and  $j = 1, 2, \dots$ .

*Proof of Theorem 2.1.* Let  $-\infty < \tau < T < +\infty$ ,  $u \in L^2(\tau, T; V)$ , and  $z$  be a solution of Problem (C) on  $[\tau, T]$  with  $(u, \bar{0}, \bar{0}) \in \mathbb{U}_{\tau, T}$ . Prove that  $z \equiv \bar{0}$ .

Let us fix an arbitrary  $m = 1, 2, \dots$ . According to the definition of a solution for Problem (C) on  $[\tau, T]$  with  $(u, \bar{0}, \bar{0}) \in \mathbb{U}_{\tau, T}$ , the following equality holds:

$$\frac{1}{2} \frac{d}{dt} \|P_m z(t)\|^2 + \nu \|P_m z(t)\|_V^2 = b(u(t), P_m z(t), z(t)), \quad (2.3)$$

for a.e.  $t \in (\tau, T)$ . Since  $b(u(t), P_m z(t), P_m z(t)) = 0$  for a.e.  $t \in (\tau, T)$ , then inequality (1.2) yields that

$$b(u(t), P_m z(t), z(t)) \leq C \|u(t)\|_V \|P_m z(t)\|_V \|z(t) - P_m z(t)\|_V,$$

for a.e.  $t \in (\tau, T)$ . Therefore, equality (2.3) imply the following inequality

$$\frac{1}{2} \frac{d}{dt} \|P_m z(t)\|^2 + \|P_m z(t)\|_V (\nu \|P_m z(t)\|_V - C \|u(t)\|_V \|z(t) - P_m z(t)\|_V) \leq 0, \quad (2.4)$$

for a.e.  $t \in (\tau, T)$ .

Let us set  $\psi_m(t) := \|P_m z(t)\|_V (\nu \|P_m z(t)\|_V - C \|u(t)\|_V \|z(t) - P_m z(t)\|_V)$ , for each  $m = 1, 2, \dots$  and a.e.  $t \in (\tau, T)$ . The following statements hold:

- (i)  $\psi_m \in L^1(\tau, T)$  for each  $m = 1, 2, \dots$ ;
- (ii)  $\psi_m(t) \leq \psi_{m+1}(t)$  for each  $m = 1, 2, \dots$  and a.e.  $t \in (\tau, T)$ ;
- (iii)  $\psi_m(t) \rightarrow \nu \|z(t)\|_V^2$  as  $m \rightarrow \infty$ , for a.e.  $t \in (\tau, T)$ .

Indeed, statement (i) holds, because  $u, z \in L^2(\tau, T; V)$  and  $P_m z \in L^\infty(\tau, T; V)$  for each  $m = 1, 2, \dots$ . Statement (ii) holds, because  $\|P_m z(t)\|_V \leq \|P_{m+1} z(t)\|_V$  and  $-\|z(t) - P_m z(t)\|_V \leq -\|z(t) - P_{m+1} z(t)\|_V$  for each  $m = 1, 2, \dots$  and a.e.  $t \in (\tau, T)$ . Statement (iii) holds, because  $P_m z(t) \rightarrow z(t)$  strongly in  $V$  as  $m \rightarrow \infty$ , for a.e.  $t \in (\tau, T)$ .

Since  $\|z(\cdot)\|_V^2 \in L^1(\tau, T)$ , then statements (i)–(iii) and Lebesgue's monotone convergence theorem yield

$$\lim_{m \rightarrow \infty} \int_\tau^t \psi_m(s) ds = \int_\tau^t \lim_{m \rightarrow \infty} \psi_m(s) ds = \int_\tau^t \|z(s)\|_V^2 ds, \quad (2.5)$$

for each  $t \in [\tau, T]$ . Inequality (2.4) implies

$$\frac{1}{2} \|P_m z(t)\|^2 + \nu \int_\tau^t \psi_m(s) ds = \int_\tau^t \frac{1}{2} \frac{d}{dt} \|P_m z(t)\|^2 + \nu \int_\tau^t \psi_m(s) ds \leq 0, \quad (2.6)$$

for each  $m = 1, 2, \dots$  and  $t \in [\tau, T]$ . We note that the equality in (2.6) holds, because  $z(\tau) = \bar{0}$ .

Equality (2.5) and inequality (2.6) yield that

$$\frac{1}{2} \|z(t)\|^2 + \nu \int_\tau^t \|z(s)\|_V^2 ds \leq 0,$$

for a.e.  $t \in (\tau, T)$ , because  $P_m z(t) \rightarrow z(t)$  strongly in  $H$  for a.e.  $t \in (\tau, T)$ . Thus,  $z(t) = \bar{0}$  for a.e.  $t \in (\tau, T)$ . Since  $z \in C([\tau, T]; V^*)$ , then  $z \equiv \bar{0}$ , that is, Problem (C) on  $[\tau, T]$  with  $(u, \bar{0}, \bar{0}) \in \mathbb{U}_{\tau, T}$  has the unique solution  $z \equiv \bar{0}$ .  $\square$

The following theorem establishes sufficient conditions for the existence of an unique solution for Problem (C). This is the main result of this section.

**Theorem 2.2.** *Let  $-\infty < \tau < T < +\infty$ ,  $y_\tau \in H$ ,  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$ , and  $y$  be a weak solution of Problem (1.1) on  $[\tau, T]$ . Then  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$  and Problem (C) on  $[\tau, T]$  with  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$  has the unique solution  $z = y$ . Moreover,  $y$  satisfies inequality (1.5).*

Before the proof of Theorem 2.2 we remark that  $AC([\tau, T]; H_m)$ ,  $m = 1, 2, \dots$ , will denote the family of absolutely continuous functions acting from  $[\tau, T]$  into  $H_m$ ,  $m = 1, 2, \dots$

*Proof of Theorem 2.2.* Prove that  $z = y$  is the unique solution of Problem (C) on  $[\tau, T]$  with  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$ . Indeed,  $y$  is the solution of Problem (C) on  $[\tau, T]$  with  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$ , because  $y$  is a weak solution of Problem (1.1) on  $[\tau, T]$ . Uniqueness holds, because if  $z$  is a solution of Problem (C) on  $[\tau, T]$  with  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$ , then  $z - y \equiv \bar{0}$  is the unique solution of Problem (C) on  $[\tau, T]$  with  $(y, \bar{0}, \bar{0}) \in \mathbb{U}_{\tau, T}$  (see Theorem 2.1).

The rest of the proof establishes that  $y$  satisfies inequality (1.5). We note that  $y$  can be obtained via standard Galerkin arguments, that is, if  $y_m \in AC([\tau, T]; H_m)$  with  $\frac{d}{dt}y_m \in L^1(\tau, T; H_m)$ ,  $m = 1, 2, \dots$ , is the approximate solution such that

$$\frac{dy_m}{dt} + \nu Ay_m + P_m B(y, y_m) = P_m f, \text{ in } H_m, \quad y_m(\tau) = P_m y(\tau), \quad (2.7)$$

then the following statements hold:

(i)  $y_m$  satisfy the following energy equality:

$$\begin{aligned} \frac{1}{2} \|y_m(t_1)\|^2 + \nu \int_s^{t_1} \|y_m(\xi)\|_V^2 d\xi - \int_s^{t_1} \langle f(\xi), y_m(\xi) \rangle d\xi \\ = \frac{1}{2} \|y_m(t_2)\|^2 + \nu \int_s^{t_2} \|y_m(\xi)\|_V^2 d\xi - \int_s^{t_2} \langle f(\xi), y_m(\xi) \rangle d\xi, \end{aligned} \quad (2.8)$$

for each  $t_1, t_2 \in [\tau, T]$ , for each  $m = 1, 2, \dots$ ;

(ii) there exists a subsequence  $\{y_{m_k}\}_{k=1,2,\dots} \subseteq \{y_m\}_{m=1,2,\dots}$  such that the following convergence (as  $m \rightarrow \infty$ ) hold:

- (ii)<sub>1</sub>  $y_{m_k} \rightarrow y$  weakly in  $L^2(\tau, T; V)$ ;
- (ii)<sub>2</sub>  $y_{m_k} \rightarrow y$  weakly star in  $L^\infty(\tau, T; H)$ ;
- (ii)<sub>3</sub>  $P_{m_k} B(u, y_{m_k}) \rightarrow B(u, y)$  weakly in  $L^2(\tau, T; V_{\frac{3}{2}}^*)$ ;
- (ii)<sub>4</sub>  $P_{m_k} f \rightarrow f$  strongly in  $L^2(\tau, T; V^*) + L^1(\tau, T; H)$ ;
- (ii)<sub>5</sub>  $\frac{dy_{m_k}}{dt} \rightarrow \frac{dy}{dt}$  weakly in  $L^2(\tau, T; V_{\frac{3}{2}}^*) + L^1(\tau, T; H)$ .

Indeed, convergence (ii)<sub>1</sub> and (ii)<sub>2</sub> follow from (2.8) (see also Temam [18, Remark III.3.1, pp. 264, 282]) and Banach-Alaoglu theorem. Since there exists  $C_1 > 0$  such that  $|b(u, v, w)| \leq C \|u\|_V \|w\|_V \|v\|_V^{\frac{1}{2}} \|v\|_V^{\frac{1}{2}}$ , for each  $u, v, w \in V$  (see, for example, Sohr [17, Lemma V.1.2.1]), then (ii)<sub>1</sub>, (ii)<sub>2</sub> and Banach-Alaoglu

theorem imply (ii)<sub>3</sub>. Convergence (ii)<sub>4</sub> holds, because of the basic properties of the projection operators  $\{P_m\}_{m=1,2,\dots}$ . Convergence (ii)<sub>5</sub> directly follows from (ii)<sub>3</sub>, (ii)<sub>4</sub> and (2.7). We note that we may not pass to a subsequence in (ii)<sub>1</sub>–(ii)<sub>5</sub>, because  $z = y$  is the unique solution of Problem (C) on  $[\tau, T]$  with  $(y, f, y_\tau) \in \mathbb{U}_{\tau, T}$ .

Moreover, there exists a subsequence  $\{y_{k_j}\}_{j=1,2,\dots} \subseteq \{y_{m_k}\}_{k=1,2,\dots}$  such that

$$y_{k_j}(t) \rightarrow y(t) \text{ strongly in } H \text{ for a.e. } t \in (\tau, T) \text{ and } t = \tau, \quad j \rightarrow \infty. \quad (2.9)$$

Indeed, according to (2.7), (2.8) and (ii)<sub>3</sub>, the sequence  $\{y_{m_k} - F_{m_k}\}_{k=1,2,\dots}$ , where  $F_{m_k}(t) := \int_\tau^t P_{m_k} f(s) ds$ ,  $m = 1, 2, \dots$ ,  $t \in [\tau, T]$ , is bounded in a reflexive Banach space  $W_{\tau, T} := \{w \in L^2(\tau, T; V) : \frac{d}{dt} w \in L^2(\tau, T; V_{\frac{3}{2}}^*)\}$ . Compactness lemma yields that  $W_{\tau, T} \subset L^2(\tau, T; H)$  with compact embedding. Therefore, (ii)<sub>1</sub>–(ii)<sub>5</sub> imply that  $y_{m_k} \rightarrow y$  strongly in  $L^2(\tau, T; H)$  as  $m \rightarrow \infty$ . Thus, there exists a subsequence  $\{y_{k_j}\}_{j=1,2,\dots} \subseteq \{y_{m_k}\}_{k=1,2,\dots}$  such that (2.9) holds.

Due to convergence (ii)<sub>1</sub>–(ii)<sub>5</sub> and (2.9), if we pass to the limit in (2.8) as  $m_{k_j} \rightarrow \infty$ , then we obtain that  $y$  satisfies the inequality

$$\frac{1}{2} \|y(t)\|^2 + \nu \int_s^t \|y(\xi)\|_V^2 d\xi - \int_s^t \langle f(\xi), y(\xi) \rangle d\xi \leq \frac{1}{2} \|y(\tau)\|^2, \quad (2.10)$$

for a.e.  $t \in (s, T)$ , a.e.  $s \in (\tau, T)$  and  $s = \tau$ .

Since  $y \in L^\infty(\tau, T; H) \cap C([\tau, T]; V^*)$  and  $H \subset V^*$  with continuous embedding, then  $y \in C([\tau, T]; H_w)$ . Thus, equality (2.10) yields

$$\frac{1}{2} \|y(t)\|^2 + \nu \int_s^t \|y(\xi)\|_V^2 d\xi - \int_s^t \langle f(\xi), y(\xi) \rangle d\xi \leq \frac{1}{2} \|y(\tau)\|^2,$$

for each  $t \in [\tau, T]$ , a.e.  $s \in (\tau, T)$  and  $s = \tau$ . Therefore,  $y$  satisfies inequality (1.5).  $\square$

### 3 Proof Theorem 1.1

In this section we establish the proof of Theorem 1.1. Let  $\Pi_{t_1, t_2}$  be the restriction operator to the finite time subinterval  $[t_1, t_2] \subseteq [\tau, T]$ ; Chepyzhov and Vishik [5].

*Proof of Theorem 1.1.* Let  $-\infty < \tau < T < +\infty$ ,  $y_\tau \in H$ ,  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$ , and  $y$  be a weak solution of Problem (1.1) on  $[\tau, T]$ .

Let us prove statement (a). Fix an arbitrary  $s \in [\tau, T)$ . Since  $(\Pi_{s, T} y, \Pi_{s, T} f, y(s)) \in \mathbb{U}_{s, T}$ , then Theorem 2.2 yields that  $\Pi_{s, T} y \in L^\infty(s, T; H)$  and it satisfies the following inequality:

$$V_\tau(y(t)) \leq V_\tau(y(s)) \quad \text{for all } t \in [s, T],$$

where  $V_\tau$  is defined in formula (1.6). Since  $s \in [\tau, T)$  be an arbitrary, then statement (a) holds.

Let us prove statement (b). Statement (a) yields

$$\frac{1}{2} \|y(t)\|^2 + \nu \int_s^t \|y(\xi)\|_V^2 d\xi - \int_s^t \langle f(\xi), y(\xi) \rangle d\xi \leq \frac{1}{2} \|y(s)\|^2, \quad (3.1)$$

for each  $t \in [s, T]$ , for each  $s \in [\tau, T)$ . In particular,  $\limsup_{t \rightarrow s^+} \|y(t)\| \leq \|y(s)\|$  for all  $s \in [\tau, T)$ , and

$$y(t) \rightarrow y(s) \text{ strongly in } H \text{ as } t \rightarrow s^+ \text{ for each } s \in [\tau, T), \quad (3.2)$$

because  $y \in C([\tau, T]; H_w)$ .

Let us prove statement (c). Since  $y \in L^2(\tau, T; V) \cap L^\infty(\tau, T; H)$  and  $f \in L^2(\tau, T; V^*) + L^1(\tau, T; H)$ , then statements (a) and (b) imply that the mapping  $t \rightarrow \|y(t)\|^2$  is of bounded variation on  $[\tau, T]$ .  $\square$

## References

- [1] Ball, J.M.: Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equations. *Nonlinear Science*. 7, 475–502 (1997) Erratum, *ibid* 8:233,1998. Corrected version appears in ‘Mechanics: from Theory to Computation’. pp. 447–474. Springer Verlag, 2000
- [2] Balibrea F., Caraballo T., Kloeden P.E., Valero J., Recent developments in dynamical systems: three perspectives, *International Journal of Bifurcation and Chaos*, 2010, doi:10.1142/S0218127410027246
- [3] Barbu, V., Rodrigues, S.S., and Shirikyan, A.: Internal Exponential Stabilization to a Nonstationary Solution for 3D NavierStokes Equations. *SIAM J. Control Optim.* 2011, doi: 10.1137/100785739
- [4] Cao, Ch., Titi, E.S.: Global Regularity Criterion for the 3D NavierStokes Equations Involving One Entry of the Velocity Gradient Tensor. *Archive for Rational Mechanics and Analysis*, 2011, doi: 10.1007/s00205-011-0439-6
- [5] Chepyzhov, V.V., Vishik, M.I.: Trajectory and Global Attractors of Three-Dimensional Navier–Stokes Systems. *Mathematical Notes*, 2002. doi: 10.1023/A:1014190629738
- [6] Cheskidov, A., Shvydkoy, R.: A Unified Approach to Regularity Problems for the 3D Navier-Stokes and Euler Equations: the Use of Kolmogorovs Dissipation Range. *Journal of Mathematical Fluid Mechanics*, 2014, doi: 10.1007/s00021-014-0167-4
- [7] Gajewski H., Gröger K., Zacharias K., *Nichtlineare operatorgleichungen und operator-differentialgleichungen*. Akademie-Verlag, Berlin (1978)
- [8] Halmos, P.R., *Measure Theory*, Springer-Verlag, New York, 1974.
- [9] Kapustyan O.V., P.O. Kasyanov, J. Valero: Pullback attractors for a class of extremal solutions of the 3D Navier-Stokes system, *J. Math. Anal. Appl.* 373 (2011) 535–547.
- [10] O. V. Kapustyan, V. S. Melnik, and J. Valero: A weak attractor and properties of solutions for the three-dimensional Bénard problem, *Discrete Contin. Dyn. Syst.*, 18, (2007) 449-481.
- [11] Kasyanov P.O., L. Toscano, N.V. Zadoianchuk: Topological Properties of Strong Solutions for the 3D Navier-Stokes Equations, *Solid Mechanics and Its Applications*. 211 (2014) 181–187.
- [12] Kasyanov P.O., L. Toscano, N.V. Zadoianchuk: A criterion for the existence of strong solutions for the 3D Navier-Stokes equations, *Applied Mathematics Letters*. 26 (2013) 15–17.
- [13] Kloeden, P.E., Marin-Rubio, P., Valero, J. The Envelope Attractor of Non-strict Multivalued Dynamical Systems with Application to the 3D Navier-Stokes and Reaction-Diffusion Equations, *Set-Valued and Variational Analysis* 21 (2013) 517–540. doi: 10.1007/s11228-012-0228-x

- [14] Melnik V.S., L. Toscano, On weak extensions of extreme problems for nonlinear operator equations. Part I. Weak solutions: *J. Automat. Inf. Scien.* 38 (2006) 68–78.
- [15] Royden H.L.: *Real Analysis* (Second edition), Macmillan, New York, 1968.
- [16] Serrin J., The initial value problem for the Navier-Stokes equations, in: R.E. Langer (editor), *Nonlinear Problems*, University of Wisconsin Press, Madison, 1963, pp. 69–98.
- [17] Sohr H.: *The Navier-Stokes Equations. An Elementary Functional Analytic Approach*, Verlag, Birkhäuser, 2001.
- [18] Temam R.: *Navier-Stokes equations*, North-Holland, Amsterdam, 1979.
- [19] Zgurovsky M.Z., P.O. Kasyanov, O.V. Kapustyan, J. Valero, N.V. Zadoianchuk: *Evolution Inclusions and Variation Inequalities for Earth Data Processing III*. Springer, Berlin, 2012.