Absolutely split locally free sheaves on proper k-schemes and Brauer-Severi varieties

Saša Novaković

April 2018

Abstract. We classify absolutely split vector bundles on proper k-schemes. More precise, we prove that the closed points of the Picard scheme are in one-to-one correspondence with indecomposable absolutely split vector bundles. Furthermore, we apply the obtained results to study the geometry of (generalized) Brauer–Severi varieties.

Contents

1.	Introduction	1
2.	Generalities on Brauer–Severi varieties and simple algebras	2
3.	Pure vector bundles	4
4.	AS-bundles on proper k -schemes	ϵ
5.	AS-bundles on proper k -schemes with cyclic Picard group	S
6.	AS-bundles on Brauer–Severi varieties	12
7.	AS-bundles on generalized Brauer–Severi varieties	15
References		17

1. Introduction

The goal of this paper is to classify absolutely split vector bundles on arbitrary proper k-schemes X. An absolutely split vector bundle on a proper k-scheme X is a locally free sheaf of finite rank that splits as a direct sum of invertible sheaves after base change to the algebraic closure \bar{k} . For simplicity, we call such sheaves AS-bundles. It is a classical theorem of Grothendieck [16] that the AS-bundles on \mathbb{P}^1 are just the locally free sheaves. Biswas and Nagaraj [7], [8], [9] and the author [27], [28] classified AS-bundles on certain Brauer–Severi varieties. To be precise, AS-bundles were completely classified in dimension one and for Brauer–Severi varieties over \mathbb{R} . Since Brauer–Severi varieties are twisted forms of projective spaces, the results in loc.cit. are generalizations of Grothendieck's theorem for \mathbb{P}^1 . In order to study AS-bundles on arbitrary Brauer–Severi varieties X, we consider the general case where X is an arbitrary proper k-scheme. Our first main result is the following:

Theorem (Theorem 4.6). Let X be a proper k-scheme with $H^0(X, \mathcal{O}_X) = k$. Then the closed points of the Picard scheme $\operatorname{Pic}_{X/k}$ are in one-to-one correspondence with isomorphism classes of indecomposable AS-bundles on X.

Notice that Theorem 4.6 is a direct generalization of the fact that the twisted Picard group equals the set of k-rational points in the Picard scheme (see [24], (7.4)). Since AS-bundles are given by the direct sum of indecomposable AS-bundles, we have a one-to-one correspondence of finite closed subschemes of $\operatorname{Pic}_{X/k}$ and isomorphism classes of AS-bundles. In the case where the proper scheme $X \otimes_k \bar{k}$ has infinite cyclic Picard group, we obtain:

Theorem (Theorem 5.1). Let X be a proper k-scheme with $\operatorname{Pic}(X \otimes_k \overline{k}) \simeq \mathbb{Z}$ and period r and let \mathcal{J} be a generator of $\operatorname{Pic}(X)$. Denote by \mathcal{L} the generator of $\operatorname{Pic}(X \otimes_k \overline{k}) \simeq \mathbb{Z}$ satisfying $\mathcal{J} \otimes_k \overline{k} \simeq \mathcal{L}^{\otimes r}$. Assume there is an indecomposable pure bundle $\mathcal{M}_{\mathcal{L}}$ of type \mathcal{L} . Then all indecomposable AS-bundles \mathcal{E} are of the form

$$\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$$

with unique $a \in \mathbb{Z}$ and $0 \le j \le r - 1$.

See Remark 5.2 for an argument why the indecomposable AS-bundles in Theorem 5.1 do not depend on the choice of the generator of $\operatorname{Pic}(X)$. Notice that for coherent sheaves on proper k-schemes the Krull–Schmidt Theorem holds (see [5]). Therefore, any coherent sheaf can uniquely be decomposed as a direct sum of indecomposables (up to isomorphism and permutation). It is easy to see that all AS-bundles on proper k-schemes are obtained as a direct sum of the indecomposable AS-bundles.

For a more deeper understanding of the AS-bundles it is important to determine the ranks of $\mathcal{M}_{\mathcal{L}^{\otimes j}}$. If $H^0(X, \mathcal{O}_X) = k$, an easy computation shows that the endomorphism algebra $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$ is central simple. One therefore has the notion of the index of $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$. Under the assumption that the Picard group is infinite cyclic, we have the following concrete description of the ranks of $\mathcal{M}_{\mathcal{L}^{\otimes j}}$.

Proposition (Proposition 5.3). Let X be a proper k-scheme with $H^0(X, \mathcal{O}_X) = k$ and $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$. Denote by \mathcal{L} the generator of $\operatorname{Pic}(X \otimes_k \bar{k})$ and assume there is an indecomposable vector bundle $\mathcal{M}_{\mathcal{L}}$ satisfying $\mathcal{M}_{\mathcal{L}} \otimes_k \bar{k} \simeq \mathcal{L}^{\oplus d}$. Then for all $j \in \mathbb{Z}$ one has

$$\operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}}) = \operatorname{ind}(\operatorname{End}(\mathcal{M}_{\mathcal{L}})^{\otimes j}).$$

Applying the above results to the case where X is a (generalized) Brauer–Severi variety we obtain a complete classification of AS-bundles on arbitrary (generalized) Brauer–Severi varieties (see Theorem 6.5 and Corollary 6.6). This gives the results of Biswas and Nagaraj [7], [8], [9] and of the author [27], [28] as corollaries. We also consider the sequence of natural numbers $(d_j)_{j\in\mathbb{Z}}$ with $d_j=\operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes j})$. In fact, for a Brauer–Severi variety of period p it is enough to consider the p+1-tupel $(d_0,d_1,...,d_{p-1},d_p)$ (see Proposition 5.4). This p+1-tupel will be called the AS-type. We determine the AS-type and study the relation between the AS-types of Brauer equivalent and birational Brauer–Severi varieties. Moreover, we will show that the AS-type is a birational invariant (see Proposition 6.12). Finally, as consequences of the Horrocks criterion and a result of Ottaviani [30], we obtain cohomological criteria for a vector bundle on a (generalized) Brauer–Severi variety to be a AS-bundle (see Theorem 6.14 and Theorem 7.7).

Acknowledgement. This paper is an improvement of the first two chapters of my Ph.D. thesis which was supervised by Stefan Schröer whom I would like to thank for a lot of comments and fruitful discussions.

Conventions. Throughout this work k denotes an arbitrary ground field and k^{sep} and \bar{k} a separable respectively algebraic closure. Furthermore, any locally free sheaf is assumed to be of finite rank and will be called vector bundle.

2. Generalities on Brauer-Severi varieties and simple algebras

Throughout the paper k denotes an arbitrary field, unless stated otherwise. We recall the basics of Brauer–Severi varieties and central simple algebras and refer to [4], [15], [33] and [34] for details. For the more general notions of Brauer–Severi schemes and Azumaya algebras we refer the reader to [17] and [18]. A Brauer–Severi variety of dimension n is a scheme X of finite type over k such that $X \otimes_k L \simeq \mathbb{P}^n$ for a finite field extension $k \subset L$. Such a field extension $k \subset L$ is called splitting field of X. Clearly, the algebraic closure \bar{k} is a splitting field for any Brauer–Severi variety. One can show that a Brauer–Severi

variety always splits over a finite separable field extension of k (see [15], Corollary 5.1.4). By embedding the finite separable splitting field into its Galois closure, a Brauer–Severi variety splits over a finite Galois extension of the base field k (see [15], Corollary 5.1.5). It follows from [19], IV, Chapter II, Theorem 2.7.1 that X is projective, integral and smooth over k. There is a well-known one-to-one correspondence between Brauer–Severi varieties and central simple k-algebras. Recall that an associative k-algebra A is called central simple if it is an associative finite-dimensional k-algebra that has no two-sided ideals other than 0 and A and if its center equals k. If the algebra A is a division algebra, it is called central division algebra. Such central simple k-algebras can be characterized by the following well-known fact (see [15], Theorem 2.2.1): A is a central simple k-algebra if and only if there is a finite field extension $k \subset L$ such that $A \otimes_k L \simeq M_n(L)$ if and only if $A \otimes_k \bar{k} \simeq M_n(\bar{k})$.

The degree of a central simple algebra A is now defined to be $\deg(A) := \sqrt{\dim_k A}$. According to the Wedderburn Theorem, for any central simple k-algebra A there is an integer n>0 and a division algebra D such that $A\simeq M_n(D)$. The division algebra D is also central and unique up to isomorphism. Now the degree of the unique central division algebra D is called the *index* of A and is denoted by ind(A). It can be shown that the index is the smallest among the degrees of finite separable splitting fields of A(see [15], Corollary 4.5.9). Two central simple k-algebras $A \simeq M_n(D)$ and $B \simeq M_m(D')$ are called Brauer equivalent if $D \simeq D'$. Brauer equivalence is indeed an equivalence relation and one defines the Brauer group Br(k) of a field k as the group whose elements are equivalence classes of central simple k-algebras and group operation being the tensor product. It is an abelian group with inverse of the equivalence class of A given by the equivalence class of A^{op} . The neutral element is the equivalence class of k. The order of a central simple k-algebra A in Br(k) is called the *period* of A and is denoted by per(A). It can be shown that the period divides the index and that both, period and index, have the same prime factors (see [15], Proposition 4.5.13). Denoting by $BS_n(k)$ the set of all isomorphism classes of Brauer-Severi varieties of dimension n and by $CSA_{n+1}(k)$ the set of all isomorphism classes of central simple k-algebras of degree n+1, there is a canonical identification

$$CSA_{n+1}(k) = BS_n(k)$$

via non-commutative Galois cohomology (see [4], [15] and [35] for details). Hence any n-dimensional Brauer–Severi variety X corresponds to a central simple k-algebra of degree n+1. In view of the one-to-one correspondence between Brauer–Severi varieties and central simple algebras one can also speak about the period of a Brauer–severi variety X. It is defined to be the period of the corresponding central simple k-algebra A.

We also need some facts concerning simple and semisimple rings in general (see [2], [11] for details). Recall that for a ring R with unity, a R-module M is called simple if M has no non-trivial submodules. A R-module M is called simple if M is isomorphic to the direct sum of simple modules (see [2], Chapter 3, § 9). Note that a ring R is called simple (semisimple) if it is a simple (semisimple) left module over itself. Clearly, a simple ring R is of course semisimple. The $Jacobson\ radical$ of a ring R is by definition the intersection of all maximal left ideals in R and is denoted by rad(R). For semisimple rings one has the following well-known characterization: Assume R is left artinian, then R is semisimple if and only if rad(R) = 0 (see [2], Proposition 15.16). In particular R/rad(R) is semisimple. Since central simple k-algebras A are isomorphic to $M_n(D)$, they are simple in the above sense (see [2], § 13, Example 13.1) and one therefore has rad(A) = 0. This fact is needed in the next section (see Proposition 3.2).

3. Pure vector bundles

In this section we study a certain class of vector bundles. We follow the work of Arason, Elman and Jacob [3] and call them *pure*.

Definition 3.1. A vector bundle \mathcal{E} on a proper k-scheme X is called *pure of type* \mathcal{W} if there is an indecomposable vector bundle \mathcal{W} on $X \otimes_k \bar{k}$ such that $\mathcal{E} \otimes_k \bar{k} \simeq \mathcal{W}^{\oplus m}$. If the indecomposable vector bundle \mathcal{W} is invertible, we say \mathcal{E} is *pure of rank one*.

Example 3.2. Let X be a n-dimensional Brauer–Severi variety over k. We consider the Euler sequence on $X \otimes_k \bar{k} \simeq \mathbb{P}^n$ (see [21], II Theorem 8.13):

$$(1) 0 \longrightarrow \Omega^{1}_{\mathbb{P}^{n}} \longrightarrow \mathcal{O}_{\mathbb{P}^{n}}(-1)^{\oplus (n+1)} \longrightarrow \mathcal{O}_{\mathbb{P}^{n}} \longrightarrow 0.$$

This short exact sequence does not split since $\mathcal{O}_{\mathbb{P}^n}(-1)$ has no global sections. Applying $\operatorname{Hom}(\mathcal{O}_{\mathbb{P}^n},-)$ to this sequence gives $\operatorname{Ext}^1(\mathcal{O}_{\mathbb{P}^n},\Omega^1_{\mathbb{P}^n})\simeq \bar{k}$. Therefore, the middle term of the Euler sequence is unique up to isomorphism. Furthermore, since \mathcal{O}_X and $\Omega^1_{X/k}$ exist on X and $\operatorname{Ext}^1(\mathcal{O}_X,\Omega^1_{X/k})\simeq k$, there is also a non-split short exact sequence on X

$$(2) 0 \longrightarrow \Omega^1_{X/k} \longrightarrow \mathcal{V} \longrightarrow \mathcal{O}_X \longrightarrow 0,$$

where the vector bundle \mathcal{V} is unique up to isomorphism. After base change to \bar{k} one gets back the sequence (1) on \mathbb{P}^n . Therefore $\mathcal{V} \otimes_k \bar{k} \simeq \mathcal{O}_{\mathbb{P}^n}(-1)^{\oplus (n+1)}$. This shows that \mathcal{V} is pure of type $\mathcal{O}_{\mathbb{P}^n}(-1)$.

The vector bundle \mathcal{V} from sequence (2) has an interesting property. Consider the endomorphism algebra $\operatorname{End}(\mathcal{V})$ which is a finite-dimensional associative k-algebra. After base change to \bar{k} , we find $\operatorname{End}(\mathcal{V}) \otimes_k \bar{k} \simeq \operatorname{End}(\mathcal{O}_{\mathbb{P}^n}(-1)^{\oplus (n+1)}) \simeq \operatorname{End}(\mathcal{O}_{\mathbb{P}^n}^{\oplus (n+1)}) \simeq M_{n+1}(\bar{k})$. We see that $\operatorname{End}(\mathcal{V})$ is a central simple k-algebra. More generally, we make the following observation.

Proposition 3.3. Let X be a geometrically integral and proper k-scheme and \mathcal{E} pure of rank one. Then $\operatorname{End}(\mathcal{E})$ is a central simple k-algebra. If furthermore \mathcal{E} is indecomposable, $\operatorname{End}(\mathcal{E})$ is a division algebra.

Proof. By definition there exists an invertible sheaf \mathcal{L} on $X \otimes_k \bar{k}$ such that $\mathcal{E} \otimes_k \bar{k} \simeq \bigoplus_{i=0}^n \mathcal{L}$. Let us denote $X_{\bar{k}}$ for the scheme $X \otimes_k \bar{k}$. We have $\mathcal{E} \otimes_k \bar{k} \otimes \mathcal{L}^{\vee} \simeq \mathcal{O}_{X_{\bar{k}}}^{\oplus (n+1)}$. Since X is supposed to be proper, the endomorphism algebra $\operatorname{End}(\mathcal{E})$ is a finite-dimensional k-algebra. Furthermore, since X is geometrically integer we have $\operatorname{End}(\mathcal{O}_{X_{\bar{k}}}) \simeq H^0(X_{\bar{k}}, \mathcal{O}_{X_{\bar{k}}}) \simeq \bar{k}$ and therefore we obtain

$$\operatorname{End}(\mathcal{E}) \otimes_{k} \bar{k} \simeq \operatorname{End}(\bigoplus_{i=0}^{n} \mathcal{L})$$

$$\simeq \operatorname{Hom}(\mathcal{L} \otimes (\bigoplus_{i=0}^{n} \mathcal{O}_{X_{\bar{k}}}), \mathcal{L} \otimes (\bigoplus_{i=0}^{n} \mathcal{O}_{X_{\bar{k}}}))$$

$$\simeq \operatorname{Hom}((\bigoplus_{i=0}^{n} \mathcal{O}_{X_{\bar{k}}}), \mathcal{L}^{\vee} \otimes \mathcal{L} \otimes (\bigoplus_{i=0}^{n} \mathcal{O}_{X_{\bar{k}}}))$$

$$\simeq \operatorname{End}(\mathcal{O}_{X_{\bar{k}}}^{\oplus (n+1)})$$

$$\simeq M_{n+1}(\bar{k}).$$

This implies that $\operatorname{End}(\mathcal{E})$ has to be a central simple k-algebra. Now suppose that \mathcal{E} is indecomposable. According to [3], p.1324 on a proper k-scheme X a vector bundle \mathcal{E} is indecomposable if and only if $\operatorname{End}(\mathcal{E})/\operatorname{rad}(\operatorname{End}(\mathcal{E}))$ is a division algebra over k. Since $\operatorname{End}(\mathcal{E})$ is a central simple k-algebra we have $\operatorname{rad}(\operatorname{End}(\mathcal{E})) = 0$. Hence $\operatorname{End}(\mathcal{E})$ is a central division algebra.

Note that the rank of a pure sheaf as in Proposition 3.3 is equal to the degree of the central simple k-algebra $\operatorname{End}(\mathcal{E})$. Indeed, the above proof shows

(3)
$$\operatorname{rk}(\mathcal{E}) = n + 1 = \operatorname{deg}(\operatorname{End}(\mathcal{E})).$$

Now let X be a geometrically integral and proper k-scheme. Assume $\mathcal E$ is pure of rank one. Suppose furthermore, we are given an indecomposable direct summand $\mathcal F$ of $\mathcal E$. We want to understand the relation between the central simple k-algebras $\operatorname{End}(\mathcal E)$ and $\operatorname{End}(\mathcal F)$. The following result is very useful and will be needed later on quite frequently.

Proposition 3.4. Let X be a proper k-scheme and \mathcal{F} and \mathcal{G} two coherent sheaves. If $\mathcal{F} \otimes_k \bar{k} \simeq \mathcal{G} \otimes_k \bar{k}$, then \mathcal{F} is isomorphic to \mathcal{G} .

Proof. Wiegand [37], Lemma 2.3 proved this in the case X is projective over k, but the proof remains valid for proper k-schemes. For convenience to the reader we reproduce the proof and follow exactly the lines of the proof given in [37]. Since the extension $k \subset \overline{k}$ is a direct limit of finite extensions, it suffices to prove the statement for finite field extensions. Now suppose $k \subset L$ is a finite field extension and $\pi: X \otimes_k L \to X$ the projection. Choose a basis $\{\alpha_1, ..., \alpha_d\}$ for L over k. By assumption we have $\pi^* \mathcal{F} \simeq \pi^* \mathcal{G}$. For the coherent sheaf $\mathcal{A} = \pi_* \pi^* \mathcal{F}$ we have over any affine open set $U \subset X$, $\mathcal{A}(U) = \mathcal{F}(U) \otimes_k L$ and there is a unique $\mathcal{O}_X(U)$ -module isomorphism from $\mathcal{A}(U)$ to $\mathcal{F}(U)^{\oplus d}$, assigning $m \otimes \alpha_i$ to (0,0,...,m,...,0), where m is located at the i^{th} entry. This yields $\pi_* \pi^* \mathcal{F} \simeq \mathcal{F}^{\oplus d}$ and obviously the same holds for $\pi_* \pi^* \mathcal{G}$. Hence $\pi_* \pi^* \mathcal{F} \simeq \mathcal{F}^{\oplus d} \simeq \pi_* \pi^* \mathcal{G} \simeq \mathcal{G}^{\oplus d}$ and we conclude from Krull–Schmidt Theorem that $\mathcal{F} \simeq \mathcal{G}$.

Notice that the statement of Proposition 3.4 also holds if one considers the base change to the separable closure k^{sep} instead of the base change to \bar{k} (see [3], p.1325). A simple consequence of Proposition 3.4 is the following:

Proposition 3.5. Let X be a proper and geometrically integral k-scheme. If \mathcal{E} and \mathcal{E}' are two indecomposable pure sheaves of same type, then $\mathcal{E} \simeq \mathcal{E}'$.

We are now able to understand the relation between the central simple k-algebras $\operatorname{End}(\mathcal{E})$ and $\operatorname{End}(\mathcal{F})$ from above.

Proposition 3.6. Let X be a proper and geometrically integral k-scheme. Let \mathcal{E} be indecomposable and pure of rank one and \mathcal{F} an indecomposable direct summand of \mathcal{E} . Then $\operatorname{End}(\mathcal{E}) \simeq M_n(\operatorname{End}(\mathcal{F}))$ and hence $\operatorname{End}(\mathcal{E})$ and $\operatorname{End}(\mathcal{F})$ are Brauer equivalent.

Proof. Since X is geometrically integral and proper, Proposition 3.3 yields that $\operatorname{End}(\mathcal{E})$ is a central simple k-algebra. Furthermore, since X is proper we can decompose \mathcal{E} as a direct sum of indecomposable vector bundles according to the Krull–Schmidt Theorem. Now let $\mathcal{E} = \bigoplus_{i=1}^n \mathcal{E}_i$ be the Krull–Schmidt decomposition of \mathcal{E} . Since \mathcal{E} is pure of rank one, there is an invertible sheaf \mathcal{L} such that $\mathcal{E} \otimes_k \bar{k} \simeq \bigoplus_{j=1}^r \mathcal{L}$. In view of $\mathcal{E} \otimes_k \bar{k} \simeq \bigoplus_{i=1}^n (\mathcal{E}_i \otimes_k \bar{k})$, we have an isomorphism

$$\bigoplus_{j=1}^{r} \mathcal{L} \simeq \mathcal{E} \otimes_{k} \bar{k} \simeq \bigoplus_{i=1}^{n} (\mathcal{E}_{i} \otimes_{k} \bar{k}).$$

Therefore, by the Krull–Schmidt Theorem for vector bundles on $X \otimes_k \bar{k}$, we obtain that \mathcal{E}_i is also pure of type \mathcal{L} . Since all the bundles \mathcal{E}_i are indecomposable, Proposition 3.5 yields that they are all isomorphic. Hence $\mathcal{E} \simeq \bigoplus_{i=1}^n \mathcal{E}_1$ and thus $\mathcal{F} \simeq \mathcal{E}_1$ by Krull–Schmidt Theorem. By Proposition 3.3, the endomorphism algebra $\operatorname{End}(\mathcal{F})$ is a central division algebra and therefore $\operatorname{End}(\mathcal{E}) \simeq \operatorname{End}(\bigoplus_{i=1}^n \mathcal{F}) \simeq M_n(\operatorname{End}(\mathcal{F}))$. This shows that $\operatorname{End}(\mathcal{F})$ and $\operatorname{End}(\mathcal{E})$ are Brauer equivalent.

Remark 3.7. The proof of the above proposition shows that all the indecomposable direct summands in the Krull–Schmidt decomposition of a vector bundle which is pure

of rank one are isomorphic. In particular, $\operatorname{End}(\mathcal{F})$ is isomorphic to the unique central division algebra D with $M_n(D) \simeq \operatorname{End}(\mathcal{E})$.

4. AS-BUNDLES ON PROPER k-SCHEMES

In this section we prove the first main result of the present paper. We start with the main definition.

Definition 4.1. Let X be a k-scheme. A vector bundle \mathcal{E} on X is called absolutely split (separably split) if it splits after base change as a direct sum of invertible sheaves on $X \otimes_k \bar{k}$ (resp. $X \otimes_k k^{sep}$). For an absolutely split locally free sheaf we shortly write AS-bundle.

Proposition 4.2. Let X be a proper k-scheme and \mathcal{E} a vector bundle on X. Then \mathcal{E} is absolutely split if and only if it is separably split.

Proof. If \mathcal{E} is separably split, then it is also absolutely split. For the other implication, assume \mathcal{E} is not separably split. Then after base change to k^{sep} we obtain from the Krull–Schmidt decomposition

$$\mathcal{E} \otimes_k k^{sep} \simeq \bigoplus_{i=1}^r \mathcal{E}_i$$

that at least one of the indecomposable \mathcal{E}_i on $X \otimes_k k^{sep}$ has rank > 1. According to [31], Proposition 3.1, the locally free sheaves \mathcal{E}_i remain indecomposable after base change to \bar{k} . Therefore, there exists at least one $\mathcal{E}_i \otimes_{k^{sep}} \bar{k}$ with $\mathrm{rk}(\mathcal{E}_i \otimes_{k^{sep}} \bar{k}) > 1$. This implies that

$$\mathcal{E} \otimes_k \bar{k} \simeq (\mathcal{E} \otimes_k k^{sep}) \otimes_{k^{sep}} \bar{k} \simeq \bigoplus_{i=1}^r (\mathcal{E}_i \otimes_{k^{sep}} \bar{k})$$

is not absolutely split. This contradicts the assumption and completes the proof. \Box

For the proof of Theorem 4.5 we need the following generalization of [3], Proposition 3.4.

Proposition 4.3. Let X be a smooth projective variety over k. Let \mathcal{E} be an indecomposable vector bundle on $X \otimes_k k^s$ and suppose $\{\mathcal{E}_1, ..., \mathcal{E}_r\}$ is the $Gal(k^s|k)$ -orbit of \mathcal{E} . Then there is an up to isomorphism unique indecomposable vector bundle \mathcal{F} on X such that $\mathcal{F} \otimes_k k^s \simeq \bigoplus_{i=1}^r \mathcal{E}_i^{\oplus d}$ for a unique positive integer d > 0.

Proof. The vector bundle $\mathcal{V} := \mathcal{E}_1 \oplus \mathcal{E}_2 \oplus ... \oplus \mathcal{E}_r$ is by assumption Galois invariant. Since \mathcal{E} is indecomposable, it follows that any \mathcal{E}_i is indecomposable, too. To get our assertion, we proceed as in the proof of Proposition 3.4 in [3] to obtain a vector bundle \mathcal{M} on X satisfying $\mathcal{M} \otimes_k k^s \simeq \mathcal{V}^{\oplus m} = \bigoplus_{i=1}^r \mathcal{E}_i^{\oplus m}$ for a suitable positive integer m > 0. Take any direct summand \mathcal{W} of \mathcal{M} and observe that $\mathcal{W} \otimes_k k^s \simeq \mathcal{V}^{\oplus r}$ for some positive integer $r \leq m$. In fact this follows from the assumption that $\{\mathcal{E}_1,...,\mathcal{E}_r\}$ is the Gal $(k^s|k)$ -orbit of the indecomposable bundle \mathcal{E} and because $\bigoplus_{i=1}^r \mathcal{E}_i^{\oplus m}$ is the Krull–Schmidt decomposition of $\mathcal{M} \otimes_k k^s$. Now choose among the direct summands of \mathcal{M} a bundle \mathcal{F} with smallest rank. Denote this rank by d. Now one proceeds as in the proof of Proposition 3.4 in [3] to conclude with Krull–Schmidt Theorem and Proposition 3.4 that \mathcal{F} is unique up to isomorphism.

For $G = \operatorname{Gal}(k^{sep}|k)$ we denote by $\operatorname{Pic}^G(X \otimes_k k^{sep}) \subset \operatorname{Pic}(X \otimes_k k^{sep})$ the subgroup of isomorphism classes of G-invariant invertible sheaves. An immediate consequence of Proposition 4.3 is:

Corollary 4.4. Let X be a proper k-scheme. For all $\mathcal{L} \in \operatorname{Pic}^G(X \otimes_k k^{sep})$ there is an up to isomorphism unique indecomposable $\mathcal{M}_{\mathcal{L}}$ on X such that $\mathcal{M}_{\mathcal{L}} \otimes_k \bar{k} \simeq \mathcal{L}^{\oplus n}$.

More general, we have:

Theorem 4.5. Let X be a proper k-scheme and \mathcal{M} an indecomposable separably split vector bundle. Then there is a unique positive integer d > 0 such that $\mathcal{M} \otimes_k k^{sep} \simeq \mathcal{L}_1^{\oplus d} \oplus \cdots \oplus \mathcal{L}_m^{\oplus d}$, where $\{\mathcal{L}_1, ..., \mathcal{L}_l\}$ is the $\operatorname{Gal}(k^{sep}|k)$ -orbit of a unique $\mathcal{L} \in \operatorname{Pic}^G(X \otimes_k k^{sep})$.

Proof. Let \mathcal{M} be an indecomposable separably split vector bundle. By definition we have

$$\mathcal{M} \otimes_k k^{sep} \simeq \bigoplus_{i=1}^n \mathcal{L}_i^{\oplus m_i},$$

for suitable invertible sheaves \mathcal{L}_i on $X \otimes_k k^{sep}$. Since $\mathcal{M} \otimes_k k^{sep}$ is $\operatorname{Gal}(k^{sep}|k)$ invariant, the set $\{\mathcal{L}_1, ..., \mathcal{L}_n\}$ decomposes as the disjoint union of $\operatorname{Gal}(k^{sep}|k)$ -orbits. Let us denote by $\mathbb{B}_l = \{\mathcal{M}(l)_1, ..., \mathcal{M}(l)_{s_l}\}$ the respective orbits. Without loss of generality, $\mathbb{B}_1 = \{\mathcal{L}_1, ..., \mathcal{L}_r\} = \{\sigma^*(\mathcal{L}_1) \mid \sigma \in \operatorname{Gal}(k^{sep}|k)\}$. We claim that $m_1 = m_2 = ... = m_r$. Assume by contradiction that there is a j > 0 such that $m_1 \neq m_j$, j = 1, ..., r. Again, without loss of generality j = 2. Now there is a $\sigma \in \operatorname{Gal}(k^{sep}|k)$ with $\sigma^*(\mathcal{L}_1) \simeq \mathcal{L}_2$. We then must have

$$\sigma^*(\mathcal{L}_1)^{\oplus m_1} \oplus \sigma^*(\mathcal{L}_2)^{\oplus m_2} \oplus \cdots \oplus \sigma^*(\mathcal{L}_r)^{\oplus m_r} \simeq \mathcal{L}_1^{\oplus m_1} \oplus \mathcal{L}_2^{\oplus m_2} \oplus \cdots \oplus \mathcal{L}_r^{\oplus m_r}.$$

It follows from the fact that all \mathcal{L}_i are indecomposable and Krull-Schmidt Theorem that $m_1 = m_2$. This contradicts the assumption $m_1 \neq m_2$. The corresponding statement applies for any \mathbb{B}_l .

Now by Proposition 4.3 it follows that for any \mathbb{B}_l there is a unique d_l and a unique \mathcal{F}_l such that

$$\mathcal{F}_l \otimes_k k^{sep} \simeq \mathcal{M}(l)_1^{\oplus d_l} \oplus \cdots \oplus \mathcal{M}(l)_{s_l}^{\oplus d_l}.$$

We denote by d the least common multiple of all the d_l . By definition, there are integers $n_l \in \mathbb{Z}$ such that $n_l \cdot d_l = d$. We find

$$(\mathcal{M}^{\oplus d}) \otimes_k k^{sep} \simeq \bigoplus_{l=1}^r \bigoplus_{j=1}^{s_l} (\mathcal{M}(l)_j^{\oplus b_l})^{\oplus d}$$

$$\simeq \bigoplus_{l=1}^r \bigoplus_{j=1}^{s_l} (\mathcal{M}(l)_j^{\oplus b_l \cdot d})$$

$$\simeq \bigoplus_{l=1}^r \bigoplus_{j=1}^{s_l} (\mathcal{M}(l)_j^{\oplus d_l})^{\oplus n_l \cdot b_l}$$

$$\simeq (\bigoplus_{l=1}^r \mathcal{F}_l^{\oplus n_l \cdot b_l}) \otimes_k k^{sep}.$$

Now Proposition 3.3 shows

$$\mathcal{M}^{\oplus d} \simeq \bigoplus_{l=1}^r \mathcal{F}_l^{\oplus n_l \cdot b_l}.$$

The Krull–Schmidt Theorem implies that \mathcal{M} must be isomorphic to some \mathcal{F}_l . Hence the assertion.

Theorem 4.5 has a interesting geometric consequence which is the content of the main theorem of this section. To prove this theorem, we first have to look more closely at the Hochschild–Serre spectral sequence for Galois coverings (see [26] for details). In general, for a proper k-scheme X with $H^0(X, \mathcal{O}_X) = k$ one has the four-term exact sequence for the Galois covering $\operatorname{Spec}(k^{sep}) \to \operatorname{Spec}(k)$ with $X' = X \otimes_k k^{sep}$:

$$0 \longrightarrow H^{1}(G, H^{0}(X'_{et}, \mathbb{G}_{m})) \longrightarrow H^{1}(X_{et}, \mathbb{G}_{m}) \longrightarrow H^{0}(G, H^{1}(X'_{et}, \mathbb{G}_{m})) \longrightarrow$$
$$\longrightarrow H^{2}(G, H^{0}(X'_{et}, \mathbb{G}_{m})).$$

Assuming $H^0(X, \mathcal{O}_X) = k$ one has $H^2(G, H^0(X'_{et}, \mathbb{G}_m)) = \operatorname{Br}(k)$ and the above exact sequence becomes

(4)
$$0 \longrightarrow \operatorname{Pic}(X) \longrightarrow \operatorname{Pic}^{G}(X') \longrightarrow \operatorname{Br}(k).$$

In order to get a more geometric interpretation of the group $\operatorname{Pic}^G(X')$ we recall the basics of the Picard scheme. The main references for the Picard scheme are [19], [20] and [23]. For a scheme X, the Picard group $\operatorname{Pic}(X)$ is the same as $H^1(X, \mathcal{O}_X^*)$ (see [21], p.224). This group is also called the absolute Picard group. To get some relative version of this group we fix a S-scheme X with structural morphism $f: X \to S$. Now for a S-scheme T one has the following base change diagram

$$X_T = X \times_S T \longrightarrow X$$

$$\downarrow_{f_T} \qquad \qquad \downarrow_{f}$$

$$T \longrightarrow S$$

and one can form the presheaf $T \mapsto H^1(X_T, \mathcal{O}_{X_T}^*)$ whose associated sheaf is $\mathbb{R}^1 f_{T_*} \mathcal{O}_{X_T}^*$ (see [21], Proposition 8.1, p.250). In the Zariski topology one then defines the relative Picard functor simply as $\operatorname{Pic}_{(X/S)(Zar)}(T) = H^0(T, \mathbb{R}^1 f_{T_*} \mathcal{O}_{X_T}^*)$. One now can imitate this for the fppf-topology and define $\operatorname{Pic}_{(X/S)(fppf)}(T) = H^0(T, \mathbb{R}^1 f_{T_*} \mathbb{G}_m)$. Again the Leray spectral sequence

$$E_2^{pq} = H^p(T, \mathbb{R}^q f_{T_*} \mathcal{F}_{X_T}) \Longrightarrow H^{p+q}(X_T, \mathcal{F}_{X_T}),$$

provides us with the exact sequence of low degree

$$0 \longrightarrow H^1(T, f_{T_*}\mathcal{F}) \longrightarrow H^1(X_T, \mathcal{F}) \longrightarrow H^0(T, \mathbb{R}^1 f_{T_*}\mathcal{F}) \longrightarrow H^2(T, f_{T_*}\mathcal{F}).$$

The cohomology groups occurring in the exact sequence are meant with respect to the fppf-topology. Now for the sheaf \mathbb{G}_m in the fppf-topology, that is also a sheaf in the étale-topology, the above exact sequence becomes:

$$0 \longrightarrow H^1(T, f_{T_*}\mathbb{G}_m) \longrightarrow H^1(X_T, \mathbb{G}_m) \longrightarrow H^0(T, \mathbb{R}^1 f_{T_*}\mathbb{G}_m) \longrightarrow H^2(T, f_{T_*}\mathbb{G}_m).$$

Assuming $f_*\mathcal{O}_X \simeq \mathcal{O}_S$ one obtains $f_{T_*}\mathbb{G}_m = \mathbb{G}_m$ and hence $H^1(T, f_{T_*}\mathbb{G}_m) = H^1(T, \mathbb{G}_m)$, what by [20], p.190-216 implies $H^1(T, f_{T_*}\mathbb{G}_m) = \operatorname{Pic}(T)$. Thus the above exact sequence becomes

$$0 \longrightarrow \operatorname{Pic}(T) \longrightarrow \operatorname{Pic}(X_T) \longrightarrow H^0(T, \mathbb{R}^1 f_{T_*} \mathbb{G}_m) \longrightarrow H^2(T, f_{T_*} \mathbb{G}_m).$$

One defines the relative Picard functor $\operatorname{Pic}_{(X/S)(et)}(T)$ in the étale-topology simply as $H^0(T,\mathbb{R}^1f_{T_*}\mathbb{G}_m)$ since the above exact sequence also exists in this topology. It is a very delicate problem under what kind of assumptions the Picard functor is representable in dependence of the given topology. We refer to [20] and [23] for details. The main theorem of Grothendieck about the Picard functor is that under the assumption that $f: X \to S$ is projective and flat and its geometric fibers are integral, the Picard functor $\operatorname{Pic}_{(X/S)(et)}$ is representable (see [23], Theorem 4.8). The object that represents the Picard functor is a separated scheme locally of finite type over S, called the Picard scheme of X and is denoted by $\operatorname{Pic}_{X/S}$. In particular, for a proper k-scheme X the Picard functor $\operatorname{Pic}_{(X/k)(et)}$ is also representable (see [23], Theorem 4.18.2, Corollary 4.18.3 or [20] p.236 ff.). Specializing further, the above exact sequence becomes for $S = \operatorname{Spec}(k) = T$ and $H^0(X, \mathcal{O}_X) = k$:

$$0 \longrightarrow \operatorname{Pic}(\operatorname{Spec}(k)) \longrightarrow \operatorname{Pic}(X) \longrightarrow \operatorname{Pic}_{(X/k)(fppf)}(k) \longrightarrow \operatorname{Br}(k).$$

As mentioned above, if X is supposed to be proper over $S = \operatorname{Spec}(k) = T$ and $H^0(X, \mathcal{O}_X) = k$, the above sequence also holds in the étale-topology and we consider the Picard functor $H^0(T, \mathbb{R}^1 f_{T_*} \mathbb{G}_m)$ as given in this topology. This gives us the following exact sequence:

$$(5) 0 \longrightarrow \operatorname{Pic}(\operatorname{Spec}(k)) \longrightarrow \operatorname{Pic}(X) \longrightarrow \operatorname{Pic}_{(X/k)(et)}(k) \longrightarrow \operatorname{Br}(k).$$

This means that we can represent elements of $\operatorname{Pic}_{(X/k)(et)}(k)$ by invertible sheaves on X that become zero in the Brauer group $\operatorname{Br}(k)$. If X has a k-rational point, one can show

that $\operatorname{Pic}(X) = \operatorname{Pic}_{(X/k)(et)}(k)$ and therefore the k-rational points of $\operatorname{Pic}_{X/k}$ are in one-to-one correspondence with invertible sheaves on X. The question arises what happens if X does not admit a k-rational point. It is a well known fact that elements of $\operatorname{Pic}_{(X/k)(et)}(k)$ are in one-to-one correspondence with so called twisted line bundles on X (see [24], (7.4)). A generalization of this fact is the following theorem.

Theorem 4.6. Let X be a proper k-scheme with $H^0(X, \mathcal{O}_X) = k$. Then the closed points of $\operatorname{Pic}_{X/k}$ are in one-to-one correspondence with isomorphism classes of indecomposable AS-bundles on X.

Proof. We denote by AS_X the set of isomorphism classes of indecomposable AS-bundles on X. In view of Proposition 4.2, we can restrict ourselves to consider separably split bundles. Let $y \in \operatorname{Pic}_{X/k}$ be a closed point. Then k(y) is a finite extension of k. The closed point y corresponds to a $\operatorname{Gal}(k^{sep}|k)$ -orbit $\{y_1,...,y_r\}$ of points in $\operatorname{Pic}_{X/k}(k^{sep})$. Recall the sequence (5) for $X^{sep} := X \otimes_k k^{sep}$

$$0 \longrightarrow \operatorname{Pic}(\operatorname{Spec}(k^{sep})) \longrightarrow \operatorname{Pic}(X^{sep}) \longrightarrow \operatorname{Pic}_{(X^{sep}/k^{sep})(et)}(k^{sep}) \longrightarrow \operatorname{Br}(k^{sep}).$$

Since $\operatorname{Pic}_{X/k}(k^{sep}) = \operatorname{Pic}_{(X^{sep}/k^{sep})}(k^{sep})$ and $\operatorname{Br}(k^{sep}) = 0$, we see that $\operatorname{Pic}(X^{sep})$ is isomorphic to $\operatorname{Pic}_{X/k}(k^{sep})$. Notice that this isomorphism is compatible with the action of $\operatorname{Gal}(k^{sep}|k)$. So for any $\operatorname{Gal}(k^{sep}|k)$ -orbit $\{y_1,...,y_r\}$ of points in $\operatorname{Pic}_{X/k}(k^{sep})$, there is a unique $\operatorname{Gal}(k^{sep}|k)$ -orbit $\{\mathcal{L}_1,...,\mathcal{L}_r\}$ of line bundles in $\operatorname{Pic}(X^{sep})$. By Proposition 4.2 and Proposition 4.3, there is a uniquely determined indecomposable AS-bundle \mathcal{M}_y satisfying

$$\mathcal{M}_y \otimes_k k^{sep} \simeq \mathcal{L}_1^{\oplus d_y} \oplus \cdots \oplus \mathcal{L}_r^{\oplus d_y}.$$

So there is a well defined map

$$\phi \colon \{ \text{closed points in } \operatorname{Pic}_{X/k} \} \longrightarrow \operatorname{AS}_X,$$

assigning to each closed point y the AS-bundle \mathcal{M}_y as described above. From Theorem 4.5 it follows that ϕ is surjective. It remains to show that ϕ is injective. For this, let y and y' be closed points and \mathcal{M}_y and $\mathcal{M}_{y'}$ the corresponding indecomposable AS-bundles. Assume $\mathcal{M}_y \simeq \mathcal{M}_{y'}$. From Theorem 4.5 we conclude

$$\mathcal{M}_{y} \otimes_{k} k^{sep} \simeq \mathcal{L}_{1}^{\oplus d} \oplus \cdots \oplus \mathcal{L}_{r}^{\oplus d} \simeq \mathcal{N}_{1}^{\oplus d'} \oplus \cdots \oplus \mathcal{N}_{s}^{\oplus d'} \simeq \mathcal{M}_{y'} \otimes_{k} k^{sep}$$

The Krull–Schmidt Theorem implies r=s and d=d'. Moreover, we have $\mathcal{L}_i \simeq \mathcal{N}_j$. Therefore, the Galois orbits $\{y_1,...,y_r\}$ and $\{y_1',...,y_s'\}$ corresponding to the orbits $\{\mathcal{L}_1,...,\mathcal{L}_r\}$ and $\{\mathcal{N}_1,...,\mathcal{N}_s\}$ are the same. By descent, this gives y=y'.

5. AS-BUNDLES ON PROPER k-SCHEMES WITH CYCLIC PICARD GROUP

In this section we prove the second main theorem of the present paper. The results of this section will be applied in Section 6 to study Brauer–Severi varieties.

We start by fixing some notation and stating some facts. From Proposition 3.4 we conclude that $\operatorname{Pic}(X)$ is a subgroup of $\operatorname{Pic}(X \otimes_k \bar{k})$. In particular, if $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$, we have $\operatorname{Pic}(X) \simeq r\mathbb{Z}$. We call the integer |r| the period of X. Let us fix a generator \mathcal{J} of $\operatorname{Pic}(X)$. Now let $\mathcal{L} \in \operatorname{Pic}(X \otimes_k \bar{k})$ be the generator satisfying $\mathcal{J} \otimes_k \bar{k} \simeq \mathcal{L}^{\otimes |r|}$. Assume there is a pure bundle \mathcal{M} of type $\mathcal{L} \in \operatorname{Pic}(X \otimes_k \bar{k})$. We know from Proposition 3.5 that the bundle \mathcal{M} is unique up to isomorphism. We use the notation $\mathcal{M}_{\mathcal{L}}$. It is easy to see that for any invertible sheaf $\mathcal{L}^{\otimes j} \in \operatorname{Pic}(X \otimes_k \bar{k})$ there is an indecomposable pure bundle of type $\mathcal{L}^{\otimes j}$. This is due to the following fact: Let $s = \operatorname{rk}(\mathcal{M}_{\mathcal{L}})$ and consider $(\mathcal{L}^{\oplus s})^{\otimes j} \simeq (\mathcal{L}^{\otimes j})^{\oplus s^j}$. From this one obtains $\mathcal{M}_{\mathcal{L}}^{\otimes j} \otimes_k \bar{k} \simeq (\mathcal{L}^{\oplus s})^{\otimes j} \simeq (\mathcal{L}^{\otimes j})^{\oplus s^j}$. Considering the Krull–Schmidt decomposition of $\mathcal{M}_{\mathcal{L}}^{\otimes j}$ and taking into account that all indecomposable direct summands are isomorphic (see proof of Proposition 3.6 and Remark 3.7), we get a up to isomorphism unique indecomposable locally free sheaf $\mathcal{M}_{\mathcal{L}^{\otimes j}}$ such that $\mathcal{M}_{\mathcal{L}^{\otimes j}} \otimes_k \bar{k} \simeq (\mathcal{L}^{\otimes j})^{\oplus s_j}$, where

 s_j is the rank of $\mathcal{M}_{\mathcal{L}^{\otimes j}}$.

With the above notation we can prove the following result.

Theorem 5.1. Let X be a proper k-scheme with $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$ and period r and let \mathcal{J} be a generator of $\operatorname{Pic}(X)$. Denote by \mathcal{L} the generator of $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$ satisfying $\mathcal{J} \otimes_k \bar{k} \simeq \mathcal{L}^{\otimes r}$. Assume there is an indecomposable pure bundle $\mathcal{M}_{\mathcal{L}}$ of type \mathcal{L} . Then all indecomposable AS-bundles \mathcal{E} are of the form

$$\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$$

with unique $a \in \mathbb{Z}$ and $0 \le j \le r - 1$.

Proof. Let \mathcal{E} be an arbitrary, not necessarily indecomposable AS-bundle and $\pi: X \otimes_k \bar{k} \to X$ the projection. By assumption, there is an indecomposable pure vector bundle $\mathcal{M}_{\mathcal{L}}$ of type \mathcal{L} . We have shown above that there exist up to isomorphism unique indecomposable pure vector bundles of type $\mathcal{L}^{\otimes j}$ for all $j \in \mathbb{Z}$. We denote them by $\mathcal{M}_{\mathcal{L}^{\otimes j}}$. Now consider $\mathcal{M}_{\mathcal{L}^{\otimes j}}$ only for j=0,...,r-1. Let $d=\operatorname{lcm}(\operatorname{rk}(\mathcal{M}_{\mathcal{O}_{X_{\bar{k}}}}),\operatorname{rk}(\mathcal{M}_{\mathcal{L}}),...,\operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes (r-1)}}))$ be the least common multiple and consider the vector bundle $\pi^*(\mathcal{E}^{\oplus d})$. Since \mathcal{E} is an AS-bundle, the vector bundle $\mathcal{E}^{\oplus d}$ is an AS-bundle, too. Therefore $\pi^*(\mathcal{E}^{\oplus d})$ decomposes into a direct sum of invertible sheaves and we find after reordering mod r that $\pi^*(\mathcal{E}^{\oplus d})$ is isomorphic to

$$\left(\bigoplus_{i=0}^{s_0} (\mathcal{L}^{\otimes a_{i_0} \cdot r})^{\oplus d}\right) \oplus \left(\bigoplus_{i=0}^{s_1} (\mathcal{L}^{\otimes a_{i_1} \cdot r+1})^{\oplus d}\right) \oplus \cdots$$
$$\oplus \left(\bigoplus_{i=0}^{s_{r-1}} ((\mathcal{L}^{\otimes a_{i_{r-1}} \cdot r+(r-1)}))^{\oplus d}\right).$$

By definition of d, there are h_j such that $h_j \cdot \operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}}) = d$ for $0 \leq j \leq r-1$. Furthermore, the sheaves $\mathcal{M}_{\mathcal{L}^{\otimes j}}$ satisfy $\pi^* \mathcal{M}_{\mathcal{L}^{\otimes j}} \simeq (\mathcal{L}^{\otimes j})^{\oplus d_j}$, where $d_j = \operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}})$. Now for the direct summands $(\mathcal{L}^{\otimes a_{i_j} \cdot r+j})^{\oplus d}$ we have

$$\left(\mathcal{L}^{\otimes a_{i_j}\cdot r+j}\right)^{\oplus d} = \left(\left(\mathcal{L}^{\otimes a_{i_j}\cdot r+j}\right)^{\oplus d_j}\right)^{\oplus h_j}.$$

By considering the vector bundle $(\mathcal{J}^{\otimes a_{i_j}} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}})^{\oplus h_j}$ on X, we find

$$\pi^* \left(\mathcal{J}^{\otimes a_{i_j}} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}} \right)^{\oplus h_j} \simeq \left(\mathcal{L}^{\otimes a_{i_j} \cdot r + j} \right)^{\oplus d}.$$

For the vector bundle

$$\mathcal{R} = \left(\bigoplus_{i=0}^{r_0} (\mathcal{J}^{\otimes a_{i_0}})^{\oplus d} \right) \oplus \left(\bigoplus_{i=0}^{r_1} (\mathcal{J}^{\otimes a_{i_1}}) \otimes \mathcal{M}_{\mathcal{L}}^{\oplus h_1} \right) \oplus \cdots$$
$$\oplus \left(\bigoplus_{i=0}^{r_{p-1}} (\mathcal{J}^{\otimes a_{i_{p-1}}}) \otimes \mathcal{M}_{\mathcal{L}}^{\oplus h_{r-1}} \right)$$

we therefore have $\pi^*\mathcal{R} \simeq \pi^*(\mathcal{E}^{\oplus d})$. Applying Proposition 3.4 shows that $\mathcal{E}^{\oplus d}$ is isomorphic to \mathcal{R} . Because Krull–Schmidt Theorem holds for vector bundles on X, we conclude that \mathcal{E} is isomorphic to the direct sum vector bundles of the form $\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$ with unique $a \in \mathbb{Z}$ and $0 \leq j \leq r-1$. Furthermore, since all these bundles are indecomposable by definition, we finally get that all the indecomposable AS-bundles are of the form $\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$ with unique $a \in \mathbb{Z}$ and $0 \leq j \leq r-1$. This completes the proof.

Remark 5.2. If we take the other generator \mathcal{J}^{\vee} of $\operatorname{Pic}(X)$, we must take \mathcal{L}^{\vee} as the generator of $\operatorname{Pic}(X \otimes_k \bar{k})$ to have $\mathcal{J}^{\vee} \otimes_k \bar{k} \simeq (\mathcal{L}^{\vee})^{\otimes r}$. Imitating the proof of Theorem 5.1 get that all indecomposable AS-bundles are of the form $(\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}})^{\vee}$ with unique $a \in \mathbb{Z}$ and $-r+1 \leq j \leq 0$. Now Proposition 5.4 below shows that these bundles are isomorphic

to the indecomposable AS-bundles from Theorem 5.1. Therefore, the isomorphism classes of indecomposable AS-bundles do not depend on the choice of the generator of Pic(X).

Now we want to determine the ranks of the sheaves $\mathcal{M}_{\mathcal{L}^{\otimes j}}$. If $H^0(X, \mathcal{O}_X) = k$, we obtain from Proposition 3.3 that $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$ is central simple. One therefore has the notion of the index of $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$.

Proposition 5.3. Let X be a proper k-scheme with $H^0(X, \mathcal{O}_X) = k$ and $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$. Denote by \mathcal{L} a generator of $\operatorname{Pic}(X \otimes_k \overline{k})$ and assume there is an indecomposable pure vector bundle $\mathcal{M}_{\mathcal{L}}$ of type \mathcal{L} . Then for all $j \in \mathbb{Z}$

$$\operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}}) = \operatorname{ind}(\operatorname{End}(\mathcal{M}_{\mathcal{L}})^{\otimes j}).$$

Proof. Let $\pi: X \otimes_k \bar{k} \to X$ denote the projection. For the bundle $\mathcal{M}_{\mathcal{L}}^{\otimes j}$ we have

$$\pi^* \mathcal{M}_{\mathcal{L}}^{\otimes j} \simeq (\mathcal{L}^{\oplus \mathrm{rk}(\mathcal{M}_{\mathcal{L}})})^{\otimes j} \simeq (\mathcal{L}^{\otimes j})^{\oplus \mathrm{rk}(\mathcal{M}_{\mathcal{L}})^j}.$$

This shows that $\mathcal{M}_{\mathcal{L}}^{\otimes j}$ is pure of type $\mathcal{L}^{\otimes j}$. In general, $\mathcal{M}_{\mathcal{L}}^{\otimes j}$ need not to be indecomposable. Since $H^0(X, \mathcal{O}_X) = k$, Proposition 3.3 implies that the endomorphism algebra $\operatorname{End}(\mathcal{M}_{\mathcal{L}^{\otimes j}})$ is a central division algebra. Applying the Krull-Schmidt Theorem for $\mathcal{M}_{\mathcal{L}}^{\otimes j}$ on X we get a decomposition

$$\mathcal{M}_{\mathcal{L}}^{\otimes j} \simeq \bigoplus_{i=1}^{m} \mathcal{E}_{i},$$

where all \mathcal{E}_i are indecomposable. After base change to the algebraic closure we have

$$(\mathcal{L}^{\oplus \operatorname{rk}(\mathcal{M}_{\mathcal{L}})})^{\otimes j} \simeq (\mathcal{M}_{\mathcal{L}}^{\otimes j}) \otimes_{k} \bar{k} \simeq \bigoplus_{i=1}^{m} \mathcal{E}_{i} \otimes_{k} \bar{k}.$$

Applying Krull-Schmidt Theorem for any \mathcal{E}_i on $X \otimes_k \bar{k}$ shows that \mathcal{E}_i are pure of type $\mathcal{L}^{\otimes j}$. According to Proposition 3.5 and Remark 3.7, all \mathcal{E}_i are isomorphic to $\mathcal{M}_{\mathcal{L}^{\otimes j}}$. Proposition 3.6 implies that $\operatorname{End}(\mathcal{M}_{\mathcal{L}}^{\otimes j})$ is a matrix algebra over the central division algebra $\operatorname{End}(\mathcal{M}_{\mathcal{L}^{\otimes j}}). \ \operatorname{By} \ (3), \text{ the rank of } \mathcal{M}_{\mathcal{L}^{\otimes j}} \text{ equals } \operatorname{deg}(\operatorname{End}(\mathcal{M}_{\mathcal{L}^{\otimes j}})). \ \operatorname{But } \operatorname{deg}(\operatorname{End}(\mathcal{M}_{\mathcal{L}^{\otimes j}})) =$ $\operatorname{ind}(\operatorname{End}(\mathcal{M}_{\mathcal{L}}^{\otimes j}))$. Now it is an easy exercise to show $\operatorname{End}(\mathcal{M}_{\mathcal{L}}^{\otimes j}) \simeq \operatorname{End}(\mathcal{M}_{\mathcal{L}})^{\otimes j}$ as central simple algebras. Finally, it follows $\operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}}) = \operatorname{deg}(\operatorname{End}(\mathcal{M}_{\mathcal{L}^{\otimes j}})) = \operatorname{ind}(\operatorname{End}(\mathcal{M}_{\mathcal{L}})^{\otimes j}).$

Let X be as in Theorem 5.1 and consider the sequence $(\operatorname{rk}(\mathcal{M}_{\mathcal{L}^{\otimes j}}))_{j \in \mathbb{Z}}$. Since $\operatorname{Pic}(X \otimes_k$ \bar{k}) is infinite cyclic one has a periodicity in the above sequence with respect to the period r of X.

Proposition 5.4. Let X be a proper k-scheme of period r with $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$ and $H^0(X, \mathcal{O}_X) = k$ and let \mathcal{L} , \mathcal{J} and $\mathcal{M}_{\mathcal{L}^{\otimes j}}$ be as in Theorem 5.1. Then for all $j \in \mathbb{Z}$ the following hold:

- $\begin{array}{ll} \textbf{(i)} \ \ \mathcal{M}_{\mathcal{L}^{\otimes j}}^{\vee} \simeq \mathcal{M}_{\mathcal{L}^{\otimes (-j)}}. \\ \textbf{(ii)} \ \ \mathcal{M}_{\mathcal{L}^{\otimes (j+ar)}} \simeq \mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}. \end{array}$

In particular one has $\operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes j}) = \operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes (-j)})$ and $\operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes (j+ar)}) = \operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes j})$.

Proof. For the invertible sheaf \mathcal{L} there is are indecomposable vector bundles $\mathcal{M}_{\mathcal{L}}$ and $\mathcal{M}_{\mathcal{L}^{\vee}}$. Since $\mathcal{M}_{\mathcal{L}}$ is pure of type \mathcal{L} , we conclude that $\mathcal{M}_{\mathcal{L}}^{\vee}$ is pure of type \mathcal{L}^{\vee} . According to Proposition 3.5, $\mathcal{M}_{\mathcal{L}}^{\vee}$ is isomorphic to $\mathcal{M}_{\mathcal{L}^{\vee}}$. The same argument shows $\mathcal{M}_{\mathcal{L}^{\otimes j}}^{\vee} \simeq$ $\mathcal{M}_{\mathcal{L}^{\otimes (-j)}}$. This proves (i). To prove (ii), we apply the same argument. For this, let \mathcal{J} be the generator of $\operatorname{Pic}(X)$ and note that by the definition of the period of X we have $\mathcal{J} \otimes_k \bar{k} \simeq \mathcal{L}^{\otimes r}$. Now consider the bundle $\mathcal{M}_{\mathcal{L}^{\otimes (j+ar)}}$ and note that it is pure of type $\mathcal{L}^{\otimes (j+ar)}$. The vector bundle $\mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$ is indecomposable and also pure of type $\mathcal{L}^{\otimes (j+ar)}$. Again with Proposition 3.5 we conclude $\mathcal{M}_{\mathcal{L}^{\otimes (j+ar)}} \simeq \mathcal{J}^{\otimes a} \otimes \mathcal{M}_{\mathcal{L}^{\otimes j}}$. This completes the proof.

Remark 5.5. Under the assumption of the above proposition we see that the sequence $(\operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes j}))_{j\in\mathbb{Z}}$ is completely determined by the tupel $(1,\operatorname{rk}(\mathcal{M}_{\mathcal{L}}),...,\operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes r-1}),1)$. Note that $\operatorname{rk}(\mathcal{M}_{\mathcal{O}_{X\otimes_k\bar{k}}}) = \operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes r}) = 1$, since both, the structure sheaf and $\mathcal{L}^{\otimes r}$, descent.

In view of Proposition 5.4 and Remark 5.5 we define the AS-type of X to be the r+1-tupel $(1, \operatorname{rk}(\mathcal{M}_{\mathcal{L}}), ..., \operatorname{rk}(\mathcal{M}_{\mathcal{L}\otimes r-1}), 1)$. It is interesting to study the relation between the AS-types of proper k-schemes X and Y which are related by certain morphisms $f: X \to Y$. Moreover, for the central simple k-algebras $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$ one also has the notion of period as invariant and it is reasonable to study which geometric properties of X are reflected by the AS-type and the periods of $\operatorname{End}(\mathcal{M}_{\mathcal{L}})$.

6. AS-BUNDLES ON BRAUER-SEVERI VARIETIES

In this section we describe the structure of AS-bundles on Brauer–Severi varieties. Furthermore, we study the AS-type and find a relation between the AS-types of Brauer equivalent and birational Brauer–Severi varieties.

As mentioned in Section 2, the n-dimensional Brauer–Severi varieties over k are in one-to-one correspondence with central simple k-algebras of degree n+1. As mentioned above, the period of a Brauer–Severi variety X is the order of the corresponding central simple k-algebra A in $\operatorname{Br}(k)$. But there is also a geometric interpretation of the period. Hochschild–Serre spectral sequence for Galois coverings applied to a Brauer–Severi variety X yields the following exact sequence (see [4]):

$$0 \longrightarrow \operatorname{Pic}(X) \xrightarrow{deg} \mathbb{Z} \longrightarrow \operatorname{Br}(k).$$

A Theorem due to Lichtenbaum (see [15], Theorem 5.4.10.) states that the boundary map $\delta: \mathbb{Z} \to \operatorname{Br}(k)$ is given by sending 1 to the class of X in $\operatorname{Br}(k)$. Here by class of X we mean that of the corresponding central simple k-algebra. It follows that $\operatorname{Pic}(X)$ is identified with some subgroup $r\mathbb{Z}$ of \mathbb{Z} . So |r| is the order of A in $\operatorname{Br}(k)$, where A is the central simple k-algebra corresponding to X. Hence the period of X can also be thought of as the smallest positive integer s such that $\mathcal{O}_X(s)$ exists in $\operatorname{Pic}(X)$. Therefore, the period of a Brauer–Severi variety equals the period as defined in Section 5. For further investigations we need the following result contained in [33], Theorem 5.5. Denote by (m,n) the greatest common divisor of the natural numbers m and n.

Theorem 6.1. Let A be a central simple k-algebra of index i. Then for $r \geq 0$ one has:

- (i) The index of $A^{\otimes r}$ divides $(\binom{i}{r}, i)$.
- (ii) Suppose i and r are coprime. Then $A^{\otimes r}$ has index i.
- (iii) Let e be (r,i). Then $A^{\otimes r}$ has index dividing i/e.

Notice that for Brauer–Severi varieties X over k one has $H^0(X, \mathcal{O}_X) = k$ and $\operatorname{Pic}(X) \simeq \mathbb{Z}$. Since $X \otimes_k \bar{k} \simeq \mathbb{P}^n$, one also has $\operatorname{Pic}(X \otimes_k \bar{k}) \simeq \mathbb{Z}$. To apply Theorem 5.1 we have to show that for the generator $\mathcal{O}_{\mathbb{P}^n}(1) \in \operatorname{Pic}(X \otimes_k \bar{k})$ there exists an indecomposable pure vector bundle of type $\mathcal{O}_{\mathbb{P}^n}(1)$.

Proposition 6.2. Let X be a n-dimensional Brauer–Severi variety over k and $\mathcal{O}_{\mathbb{P}^n}(1)$ the generator of $\operatorname{Pic}(X \otimes_k \overline{k}) \simeq \mathbb{Z}$. Then there exists an indecomposable pure vector bundle \mathcal{W}_1 on X of type $\mathcal{O}_{\mathbb{P}^n}(1)$.

Proof. We have seen that the vector bundle \mathcal{V} of Example 3.2 satisfies $\pi^*\mathcal{V} \simeq \mathcal{O}_{\mathbb{P}^n}(-1)^{\oplus (n+1)}$, where π denotes the projection $\pi: X \otimes_k \bar{k} \to X$. Hence $\pi^*\mathcal{V}^\vee$ is isomorphic to $\mathcal{O}_{\mathbb{P}^n}(1)^{\oplus (n+1)}$ and therefore \mathcal{V}^\vee is pure of type $\mathcal{O}_{\mathbb{P}^n}(1)$. Let us denote the vector bundle \mathcal{V}^\vee by \mathcal{W} . We decompose \mathcal{W} according to the Krull–Schmidt Theorem and write $\mathcal{W} \simeq \bigoplus_{i=1}^m \mathcal{W}_i$. Since all the \mathcal{W}_i have to be isomorphic (see Remark 3.7), it follows $\mathcal{W} \simeq \mathcal{W}_1^{\oplus m}$, where \mathcal{W}_1 is

unique up to isomorphism. From Krull-Schmidt Theorem it follows that W_1 is also pure of type $\mathcal{O}_{\mathbb{P}^n}(1)$. This completes the proof.

We saw that the period p of a Brauer–Severi variety X can be thought of as the smallest positive integer r such that $\mathcal{O}_X(r)$ exists in $\operatorname{Pic}(X)$. Hence the period defined for Brauer–Severi varieties equals the period defined for arbitrary proper k-schemes in Section 5. Now we write \mathcal{W}_j for the vector bundles $\mathcal{M}_{\mathcal{L}^{\otimes j}}$ of Section 5 with $\mathcal{L} = \mathcal{O}_{\mathbb{P}^n}(1)$. Then Theorem 5.1 applies and we obtain:

Corollary 6.3. Let X be a Brauer-Severi variety over a field k of period p. Then all AS-bundles \mathcal{E} are of the form

$$\mathcal{E} \simeq igoplus_{j=0}^{p-1} \left(igoplus_{i=0}^{r_j} \mathcal{O}_X(a_{i_j}p) \otimes \mathcal{W}_j
ight)$$

with unique $a_{i_j} \in \mathbb{Z}$ and $r_j > 0$.

As mentioned in Section 5, for a complete understanding of the AS-bundles on Brauer–Severi varieties one has to determine the ranks of the \mathcal{W}_j . So consider the sequence of natural numbers $(d_j)_{j\in\mathbb{Z}}$, with $d_j=\operatorname{rk}(\mathcal{W}_j)$. Proposition 5.4 and Remark 5.5 show that we do not have to consider the hole sequence $(d_j)_{j\in\mathbb{Z}}$. Furthermore, we note that $\mathcal{W}_0=\mathcal{O}_X$ and $\mathcal{W}_p=\mathcal{O}_X(p)$, where p is the period of X. This implies that $\operatorname{rk}(\mathcal{W}_0)=1=\operatorname{rk}(\mathcal{W}_p)$. In fact, for the vector bundle \mathcal{W} of the proof of Proposition 6.2 one can show $\operatorname{End}(\mathcal{W})\simeq A$, where A is the central simple algebra corresponding to X (see [32], p.144 or [36], §3, 3.6). Applying Proposition 5.3 implies:

Corollary 6.4. Let X be a n-dimensional Brauer–Severi variety over a field k corresponding to a central simple k-algebra A. Then for every $j \in \mathbb{Z}$ one has

$$\operatorname{rk}(\mathcal{W}_i) = \operatorname{ind}(A^{\otimes j}).$$

Theorem 6.5. Let X be a Brauer–Severi variety over k and A the corresponding central simple k-algebra of period p. Then the AS-type of X is given by $d_j = \operatorname{ind}(A^{\otimes j})$ for $0 \leq j \leq p$.

Corollary 6.4 together with Theorem 6.5 now give a complete classification of AS-bundles on Brauer–Severi varieties and thus the results of Biswas and Nagaraj [7], [8], [9] and the author [27], [28] as corollary.

Corollary 6.6. ([28], Theorem 5.1 and [8], Theorem 1.1) Let X be a n-dimensional Brauer–Severi variety over k of index two. Then the AS-bundles are of the form

$$\mathcal{E} \simeq \left(igoplus_{i=1}^r \mathcal{O}_X(2a_i)
ight) \oplus \left(igoplus_{j=1}^s \mathcal{O}_X(2b_j) \otimes \mathcal{W}_1
ight)$$

with unique r, s, a_i and b_j . The vector bundle W_1 satisfies $W_1 \otimes_k L \simeq \mathcal{O}_{\mathbb{P}^n}(1)^{\oplus 2}$, where $k \subset L$ is a degree two Galois extension that splits X.

Proof. Since the index of X is two and the period divides the index (see [15], Proposition 4.5.13), we conclude that the period is also two. Note that the period cannot be one, since this would imply that X is the projective space, contradicting the fact that the index of X is two. Hence the AS-type of X is (1,2,1) according to Theorem 6.5. Now the assertion follows from Corollary 6.3 and [15], Corollary 4.5.9.

Note that the results in [7], [9] and [27] concern the case of non-trivial one-dimensional Brauer–Severi varieties and therefore follow directly from Corollary 6.6 since non-split Brauer–Severi varieties of dimension are of index two.

In the case where the Brauer–Severi variety corresponds to a central simple k-algebra with period equals index, the AS-type can be stated very explicitly. We first need the following facts.

Lemma 6.7. Let A be a central simple k-algebra of period p. Then one has $p/(p,r) = per(A^{\otimes r})$.

Proposition 6.8. Let A be a central simple k-algebra with period p and index i. Then for all $r \geq 0$ one has that p/(p,r) divides $\operatorname{ind}(A^{\otimes r})$ and $\operatorname{ind}(A^{\otimes r})$ divides i/(i,r). In particular one has $p/(p,r) \leq \operatorname{ind}(A^{\otimes r}) \leq i/(i,r)$.

Proof. By Lemma 6.7 we have $p/(p,r) = \operatorname{per}(A^{\otimes r})$, and since the period always divides the index (see [15], Proposition 4.5.13) we find that p/(p,r) divides $\operatorname{ind}(A^{\otimes r})$. The second inequality $\operatorname{ind}(A^{\otimes r}) \leq i/(i,r)$ and the fact that $\operatorname{ind}(A^{\otimes r})$ divides i/(i,r) is (iii) of Theorem 6.1.

The last proposition enables us to determine the AS-type of Brauer–Severi varieties with same period and index.

Proposition 6.9. Let X be a Brauer-Severi variety over k corresponding to a central simple k-algebra A such that the period p equals the index i. Then the AS-type of X is given by

$$\operatorname{rk}(\mathcal{W}_j) = p/(p,j)$$

for $0 \le j \le p$.

Proof. Since the period p equals the index i, we conclude from Proposition 6.8 that $p/(p,j)=\operatorname{ind}(A^{\otimes j})$ for $0\leq j\leq p$. The assertion then follows from Corollary 6.4.

Remark 6.10. The problem for which fields k the period equals the index is called *period-index problem* and is highly non-trivial. For further discussion on this problem we refer the reader to [6] and to the work of de Jong [12].

If k is local or global, any central division algebra over k is cyclic and the period equals the index (see [33], Theorem 10.7). In this case Proposition 6.9 applies and we can determine the AS-type . The next observation shows that it is enough to consider minimal linear subvarieties to determine the AS-type.

Proposition 6.11. Let X and Y be Brauer-Severi varieties over k which are Brauer equivalent. Then X and Y have the same AS-type.

Proof. Let A be the central simple k-algebra corresponding to X and B the central simple k-algebra corresponding to Y. Since A and B are Brauer equivalent, there is a unique central division algebra D such that $A \simeq M_n(D)$ and $B \simeq M_m(D)$ for suitable n and m. Hence $\operatorname{ind}(A^{\otimes j}) = \operatorname{ind}(D^{\otimes j}) = \operatorname{ind}(B^{\otimes j})$ for all $j \in \mathbb{Z}$. Since Brauer equivalent Brauer–Severi varieties have the same period, Theorem 6.5 yields the assertion.

Interestingly, it turns out that the AS-type is also a birational invariant for Brauer–Severi varieties.

Proposition 6.12. Let X and Y be two birational Brauer–Severi varieties. Then they have the same AS-type.

Proof. Let A be the central simple k-algebra corresponding to X and B the algebra corresponding to Y. Since X and Y are supposed to be birational, [15], Corollary 5.4.2 implies that A and B generate the same cyclic subgroup in Br(k). Denoting by i the index of A, Proposition 6.8 shows that $\operatorname{ind}(A^{\otimes r})$ divides i/(i,r). Since i/(i,r) divides i, we conclude that $\operatorname{ind}(A^{\otimes r})$ divides $i = \operatorname{ind}(A)$. The same holds for B and we see that $\operatorname{ind}(B^{\otimes s})$ divides $\operatorname{ind}(B)$. In what follows we show that $\operatorname{ind}(A^{\otimes r}) = \operatorname{ind}(B^{\otimes r})$ for all r. Since A and B generate the same cyclic subgroup in $\operatorname{Br}(k)$, we know that A is Brauer equivalent to $B^{\otimes l}$

and B to $A^{\otimes m}$ for suitable l and m. Therefore $\operatorname{ind}(A^{\otimes m}) = \operatorname{ind}(B)$ divides $\operatorname{ind}(A)$ and $\operatorname{ind}(B^{\otimes l}) = \operatorname{ind}(A)$ divides $\operatorname{ind}(B)$. This shows that $\operatorname{ind}(B)$ divides $\operatorname{ind}(A)$ and vice verse. Thus $\operatorname{ind}(B) = \operatorname{ind}(A)$. The same argument applied to $\operatorname{ind}(A^{\otimes r})$ and $\operatorname{ind}(B^{\otimes r})$ yields that $\operatorname{ind}(A^{\otimes rm}) = \operatorname{ind}(B^{\otimes r})$ divides $\operatorname{ind}(A^{\otimes r}) = \operatorname{ind}(B^{\otimes r})$. This shows $\operatorname{ind}(A^{\otimes r}) = \operatorname{ind}(B^{\otimes r})$ for all r. As mentioned above, A and B generate the same cyclic subgroup and therefore have the same period. Now Theorem 6.5 implies that X and Y have the same AS-type.

Remark 6.13. Note that the other implication in Proposition 6.11 and 6.12 does not hold. Indeed, by [15], Theorem 1.4.2, two non-split one-dimensional Brauer–Severi varieties X and Y are birational if and only if they are isomorphic. But according to Theorem 6.5 and Corollary 6.6 any one-dimensional Brauer–Severi variety has the same AS-type. So indeed there are Brauer–Severi varieties which are not Brauer equivalent (respectively not birational) but have the same AS-type.

As a consequence of the Horrocks criterion on \mathbb{P}^n (see [29]) we can state a cohomological criterion for a vector bundle on a Brauer–Severi variety to be a AS-bundle.

Theorem 6.14 (AS-criterion). Let X be a n-dimensional Brauer–Severi variety over k and period p. A vector bundle \mathcal{E} is a AS-bundle if and only if for 0 < i < n one has

$$H^i(X, \mathcal{E} \otimes \mathcal{O}_X(ap) \otimes \mathcal{W}_i) = 0$$

for every $a \in \mathbb{Z}$ and every $0 \le j \le p-1$.

We end up this section mentioning that the duals of the indecomposable AS-bundles W_j with $j \geq 0$ on a Brauer–Severi variety X generate the Grothendieck group $K_0(X)$. This follows directly from [32], Section 8, Theorem 4.1 or [22], Theorem 3.1.

7. AS-BUNDLES ON GENERALIZED BRAUER-SEVERI VARIETIES

In this section we will see that nearly all results stated in the last section for Brauer–Severi varieties also hold for generalized Brauer–Severi varieties.

Let A be a central simple k-algebra of degree n and $1 \le d \le n$. Now consider the subset of $\operatorname{Grass}_k(d \cdot n, A)$ consisting of those subspaces of A that are left ideals L of dimension $d \cdot n$. This subset of $\operatorname{Grass}_k(d \cdot n, A)$ can be given a structure of a projective scheme over k, defined by the relations stating that the L are left ideals (see [10], p.100 for details). This scheme is denoted by $\operatorname{BS}(d,A)$ and is called generalized Brauer–Severi variety. It is a closed subscheme of the Grassmannian $\operatorname{Grass}_k(dn,A)$. For a field extension $k \subset E$ one has $\operatorname{BS}(d,A\otimes_k E) \cong \operatorname{BS}(d,A)\otimes_k E$ (see [10]). As in the case of Brauer–Severi varieties, there is always a finite separable and therefore a finite Galois extension $k \subset K$ such that $\operatorname{BS}(d,A)\otimes_k K \cong \operatorname{Grass}_K(d,n)$. Clearly, the generalized Brauer–Severi variety becomes isomorphic to the Grassmannian after base change to the algebraic closure \bar{k} . Note that for d=1 one obtains the usual Brauer–Severi variety as defined in Section 2. Unfortunately, there is no one-to-one correspondence between generalized Brauer–Severi varieties and central simple k-algebras (see [10], Theorem 1). Nonetheless, it is easy to classify all AS-bundles on generalized Brauer–Severi varieties (see Theorem 7. and Proposition 7.2 below). Recall the tautological exact sequence on $X = \operatorname{Grass}_k(d,n)$

$$0 \longrightarrow \mathcal{S} \longrightarrow \mathcal{O}_X^{\oplus n} \longrightarrow \mathcal{Q} \longrightarrow 0$$

with tautological sheaf S which is a vector bundle of rank d. On the generalized Brauer–Severi variety BS(d, A) one has also a tautological short exact sequence

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{O}_{\mathrm{BS}(d,A)}^{\oplus n^2} \longrightarrow \mathcal{R} \longrightarrow 0.$$

This short becomes after base change $BS(d, A) \otimes_k L \simeq Grass_L(d, n)$ to some splitting field L of A (see [25], Section 4):

$$0 \longrightarrow \mathcal{S}^{\oplus n} \longrightarrow \mathcal{O}_{\operatorname{Grass}_L(d,n)}^{\oplus n^2} \longrightarrow \mathcal{Q}^{\oplus n} \longrightarrow 0.$$

The vector bundle \mathcal{I} is called $tautological\ sheaf$ on $Y=\mathrm{BS}(d,A)$. Recall that for $\mathrm{Grass}(d,n)$ the Picard group coincides with its first Chow group (see [14], Example 15.3.6). By Proposition 14.6.6 in [14] we have $\mathrm{Pic}(\mathrm{Grass}(d,n))\simeq \mathbb{Z}$, where the ample generator can be taken to be $\mathcal{L}=\det(\mathcal{S}^\vee)$. Proposition 3.3 now implies $\mathrm{Pic}(\mathrm{BS}(d,A))\simeq \mathbb{Z}$. In order to apply Theorem 5.1, we have to show that there is a up to isomorphism unique pure vector bundle of type \mathcal{L} on Y. For this, let Σ^λ be the Schur functor associated with the partition $\lambda=(\lambda_1,\lambda_2,...,\lambda_d)$, where $0\leq\lambda_i\leq n-d$ (see [1], [13] for details). Levine, Srinivas and Weyman [25], Section 4 proved that for the tautological sheaf \mathcal{S} on $\mathrm{Grass}(d,n)$ the vector bundles $\Sigma^\lambda(\mathcal{S})^{\oplus n\cdot|\lambda|}$ descent to vector bundles \mathcal{N}_λ on $\mathrm{BS}(d,A)$. Note that these vector bundles are unique up to isomorphism by Proposition 3.3. In particular, for the ample generator $\mathcal{L}=\det(\mathcal{S}^\vee)$ of $\mathrm{Pic}(\mathrm{Grass}(d,n))$ there is a up to isomorphism unique indecomposable vector bundle $\mathcal{W}_\mathcal{L}$ satisfying $\mathcal{W}_\mathcal{L}\otimes_k k^{sep}\simeq \mathcal{L}^{\oplus \mathrm{rk}(\mathcal{W}_\mathcal{L})}$. As explained in Section 5, for all $\mathcal{L}^{\otimes j}\in\mathrm{Pic}(\mathrm{Grass}(d,n))$ there exist up to isomorphism unique indecomposable vector bundles $\mathcal{W}_{\mathcal{L}\otimes j}$ with $\mathcal{W}_{\mathcal{L}\otimes j}\otimes_k k^{sep}\simeq (\mathcal{L}^{\otimes j})^{\oplus \mathrm{rk}(\mathcal{W}_\mathcal{L}\otimes j)}$. Let \mathcal{M} be the generator of $\mathrm{Pic}(\mathrm{BS}(d,A))$ satisfying $\mathcal{M}\otimes_k k^{sep}\simeq \mathcal{L}^{\otimes r}$, where r>0 denotes the period of $\mathrm{BS}(d,A)$. Now Theorem 5.1 immediately implies:

Theorem 7.1. Let X = BS(d, A) be a generalized Brauer–Severi variety of period r for the central simple k-algebra A and M the generator of Pic(BS(d, A)). Then all indecomposable AS-bundles of finite rank are of the form:

$$\mathcal{M}^{\otimes a} \otimes \mathcal{W}_{\mathcal{L}^{\otimes j}}$$

with unique $a \in \mathbb{Z}$ and unique $0 \le j \le r - 1$.

An immediate consequence of Proposition 5.3 is the following:

Proposition 7.2. Let X = BS(d, A) be a generalized Brauer–Severi variety over k for the central simple k-algebra A of degree n. Then for all $j \in \mathbb{Z}$ one has:

$$\operatorname{rk}(\mathcal{W}_{\mathcal{L}\otimes j}) = \operatorname{ind}(A^{\otimes j \cdot d}).$$

Proof. Let $\lambda = (\lambda_1, \lambda_2, ..., \lambda_d)$ be a partition with $0 \leq \lambda_i \leq n - d$ and \mathcal{S} the tautological sheaf on $\mathrm{BS}(d,A) \otimes_k k^{sep} \simeq \mathrm{Grass}(d,n)$. As mentioned above, in [25], Section 4 it is proved that the vector bundles $\Sigma^{\lambda}(\mathcal{S})^{n\cdot |\lambda|}$ descent to vector bundles \mathcal{N}_{λ} which are unique up to isomorphism according to Proposition 3.4. In particular, $\det(\mathcal{S})^{\oplus n \cdot d}$ descents to a vector bundle which we denote by \mathcal{N} . For the endomorphism algebra of \mathcal{N}^{\vee} one has $\mathrm{End}(\mathcal{N}^{\vee}) \simeq A^{\otimes d}$ (see also [25]). Since \mathcal{N}^{\vee} is pure of type $\det(\mathcal{S}^{\vee})$, but in general not indecomposable, we conclude that $A^{\otimes d}$ is a matrix algebra over $\mathrm{End}(\mathcal{W}_{\mathcal{L}})$ (see Proposition 3.6). Proposition 5.3 gives $\mathrm{rk}(\mathcal{W}_{\mathcal{L}\otimes j}) = \mathrm{ind}(\mathrm{End}(\mathcal{W}_{\mathcal{L}})^{\otimes j})$. But the index of $\mathrm{End}(\mathcal{W}_{\mathcal{L}})^{\otimes j}$ is the same as the index of $A^{\otimes j \cdot d}$. This completes the proof.

Remark 7.3. Let $G = \text{Gal}(k^{sep}|k)$ be the absolute Galois group and Y the generalized Brauer–Severi variety BS(d, A). With a result of Blanchet [10], Theorem 7, the exact sequence

$$0 \longrightarrow \operatorname{Pic}(Y) \longrightarrow \operatorname{Pic}^{G}(Y \otimes_{k} k^{sep}) \stackrel{\delta}{\longrightarrow} \operatorname{Br}(k) \stackrel{res}{\longrightarrow} \operatorname{Br}(F(Y))$$

and the arguments of the proof of Theorem 5.4.1 of [15] one directly see that the period of Y (as defined in Section 5) is equal to the period of $A^{\otimes d}$.

Now we prove Propositions 6.11 and 6.12 for generalized Brauer–Severi varieties.

Proposition 7.4. Let D be a central division algebra over k of degree n and $A = M_m(D)$. Then BS(d, D) and BS(d, A) have the same AS-type.

Proof. Since D and A are Brauer equivalent, we conclude that $\operatorname{ind}(D^{\otimes i}) = \operatorname{ind}(A^{\otimes i})$ for all $i \in \mathbb{Z}$. Thus $\operatorname{ind}(D^{\otimes d \cdot i}) = \operatorname{ind}(A^{\otimes d \cdot i})$ for all $i \in \mathbb{Z}$. Furthermore, the period of $D^{\otimes d}$ equals the period of $A^{\otimes d}$. Applying Proposition 7.2 and Theorem 7.1 yields the assertion.

Proposition 7.5. Let X = BS(d, A) and Y = BS(d, B) be two birational generalized Brauer–Severi varieties over k, then they have the same AS-type.

Proof. As a consequence of [10], Theorem 7, the period of $A^{\otimes d}$ equals the period of $B^{\otimes d}$ and hence the period of X equals the period of Y. Now the same arguments as in the proof of Proposition 6.12 show that $\operatorname{ind}(A^{\otimes d \cdot j}) = \operatorname{ind}(B^{\otimes d \cdot j})$. From Theorem 7.1 and Proposition 7.2 we conclude that both have the same AS-type.

Ottaviani [30], Theorem 2.1 proved a splitting criterion for vector bundles on Grass(d, n), at least if char(k) = 0. As a consequence of this result we obtain a criterion for a vector undle on BS(d, A) to be a AS-bundle. We first cite Ottaviani's result.

Theorem 7.6. Let k be a field of characteristic zero and $n \ge 3$. Then a locally free sheaf \mathcal{E} of finite rank on $X = \operatorname{Grass}_k(d,n)$ splits as a direct sum of invertible sheaves if and only if for $0 < r < \dim(X)$ and all $t \in \mathbb{Z}$ one has

$$H^{r}(X, \bigwedge^{i_1}(\mathcal{Q}^{\vee}) \otimes \cdots \otimes \bigwedge^{i_s}(\mathcal{Q}^{\vee}) \otimes \mathcal{E}(t)) = 0$$

for all $i_1, ..., i_s$ such that $0 \le i_1, ..., i_s \le n - d$ and $s \le d$.

Now denote by \mathcal{M} the (ample) generator of $\operatorname{Pic}(\operatorname{BS}(d,A))$. Considering the above $i_1,...,i_s$, we note that for a fixed s-tupel $i_1,...,i_s$ the i_s can be ordered in a weakly decreasing way. We denote the reordering by $\lambda_1 \geq \lambda_2 \geq ... \geq \lambda_s$. So in this way we get a partition $\lambda = (\lambda_1,...,\lambda_s)$ and we can associate a Young diagram to it with at most d rows and n-d columns. Now let μ' be the conjugate of the partition $\mu = (\lambda_i)$ (we have exactly λ_i boxes). Then we have $\Sigma^{\mu'}(\mathcal{Q}^\vee) = \bigwedge^{\lambda_i}(\mathcal{Q}^\vee)$ on $\operatorname{BS}(d,A) \otimes_k \bar{k} \simeq \operatorname{Grass}_{\bar{k}}(d,n)$. In [25], Section 4 it is shown that $(\Sigma^{\mu'}(\mathcal{Q}^\vee))^{\oplus n\cdot |\mu'|}$ descents to a vector bundle \mathcal{P}_{λ_i} on $\operatorname{BS}(d,A)$. From Proposition 3.4 we know that these vector bundles are unique up to isomorphism. We simply write $\mathcal{F}(m)$ for $\mathcal{F} \otimes \mathcal{M}^{\otimes m}$, where $\mathcal{M} \in \operatorname{Pic}(\operatorname{BS}(d,A))$ denotes the generator. With this notation we have the following result:

Theorem 7.7 (AS-criterion). Let BS(d, A) be the generalized Brauer–Severi variety of period r for the central simple k-algebra A of degree $n \geq 3$ and \mathcal{P}_{λ_i} the vector bundles from above. A vector bundle \mathcal{E} is a AS-bundle if and only if for $0 < r < \dim(\mathrm{BS}(d,A))$ and all $t \in \mathbb{Z}$ one has

$$H^r(\mathrm{BS}(d,A),\mathcal{P}_{\lambda_1}\otimes\cdots\otimes\mathcal{P}_{\lambda_n}\otimes\mathcal{E}(t))=0$$

for all $\lambda_1 > \lambda_2 > ... > \lambda_s$ such that $0 < \lambda_1, ..., \lambda_s < n - d$ and s < d.

We note that Levine, Srinivas and Weyman [25] calculated the K-theory of generalized Brauer–Severi varieties and showed that the Grothendieck group is generated by \mathcal{N}_{λ} , where \mathcal{N}_{λ} are the vector bundles obtained by descent from $\Sigma^{\lambda}(\mathcal{S})^{\oplus n \cdot |\lambda|}$. Therefore, as distinguished from the case of Brauer–Severi varieties, the AS-bundles on generalized Brauer–Severi varieties do not generate the Grothendieck group of $\mathrm{BS}(d,A)$ for d>1.

References

- K. Akin, D.A. Buchsbaum and J. Weyman: Schur Functors and Schur Complexes. Adv. in Math. Vol. 44 (1982), 207-278.
- [2] F.W. Anderson and K.R. Fuller: Rings and Categories of modules. Springer-Verlag, New York (1974).
- [3] J.K. Arason, R. Elman and B. Jacob: On indecomposable vector bundles. Comm. in Alg. Vol. 20 (1992), 1323-1351.

- [4] M. Artin: Brauer-Severi varieties. Brauer groups in ring theory and algebraic geometry, Lecture Notes in Math. 917, Notes by A. Verschoren, Berlin, New York: Springer-Verlag (1982), 194?10.
- [5] M. Atiyah: On the Krull-Schmidt Theorem with application to sheaves. Bull. Soc. Math. France Vol. 84 (1956), 307-317.
- [6] A. Auel, E. Brussel, S. Garibaldi and U. Vishne: Open Problems on Central Simple Algebras. arXiv:1006.3304 [math.RA] (2010).
- [7] I. Biswas and D. Nagaraj: Classification of real algebraic vector bundles over the real anisotropic conic. Int. J. Math. Vol. 16 (2005), 1207-1220.
- [8] I. Biswas and D. Nagaraj: Absolutely split real algebraic vector bundles over a real form of projective space. Bull. Sci. Math. Vol. 131 (2007), 686-696.
- [9] I. Biswas and D. Nagaraj: Vector bundles over a nondegenerate conic. J. Aust. Math. Soc. Vol. 86 (2009), 145-154.
- [10] A. Blanchet: Function Fields of Generalized Brauer-Severi Varieties. Communications in Algebra. Vol. 19 (1991), 97-118.
- [11] N. Bourbaki: Algébre, Chapitre X, Masson. (1980).
- [12] A.D. de Jong: The period-index problem for the Brauer group of an algebraic surface. Duke Math. J. 123 No. 1 (2004), 71-94.
- [13] W. Fulton: Young Tableaux, with Applications to Representation Theory and Geometry. Cambridge University Press (1997).
- [14] W. Fulton: Intersection Theory. Ergebnisse der Mathematik und ihre Grenzgebiete, Springer, second edition (1998).
- [15] P. Gille and T. Szamuely: Central Simple Algebras and Galois Cohomology. Cambridge Studies in advanced Mathematics. 101. Cambridge University Press. (2006)
- [16] A. Grothendieck: Sur la classification des fibré holomorphes sur la sphère de Riemann. Amer. J. Math. Vol. 79 (1957), 121-138.
- [17] A. Grothendieck: Le group de Brauer I: Algebras d Azumaya et interpretations diverses, Seminaire Bourbaki. No. 290 (1964).
- [18] A. Grothendieck: Le group de Brauer II: Theorie cohomologique, Seminaire Bourbaki. No. 297 (1965).
- [19] A. Grothendieck et J. Dieudonne: Elements de Géométrie Algébrique. Publ. Math. IHES. (1960-68).
- [20] A. Grothendieck: Fondements de la Géométrie Algébrique [Extraits du Séminaire Bourbaki 1957-1962], Secrétariat mathématique (1962).
- [21] R. Hartshorne: Algebraic Geometry. Springer-Verlag, New York, Berlin, Heidelberg (1977).
- [22] N. Karpenko: Codimension 2 Cycles on Severi–Brauer Varieties. K-Theory 13 No. 4 (1998), 305-330.
- [23] S. L. Kleiman: The Picard scheme: arXiv:math/0504020 [math.AG] (2005).
- [24] J. Kollár: Severi-Brauer varieties: A geometric treatment.
- [25] M. Levine, V. Srinivas and J. Weyman: K-Theory of Twisted Grassmannians. K-Theory Vol. 3 (1989), 99-121.
- [26] J. Milne: Étale Cohomology. Princeton Mathematical Series. Vol. 33, Princeton University Press (1980).
- [27] S. Novaković: Der Spaltungssatz für Quadriken. Diploma Thesis, Heinrich-Heine Universität Düsseldorf (2009).
- [28] S. Novaković: Absolutely split locally free sheaves on Brauer-Severi varieties of index two. Bull. d. Sci. Math. Vol. 136 (2012), 413-422.
- [29] C. Okonek, M. Schneider and H. Spindler: Vector Bundles on Complex Projective Space. Progress in Mathematics. 3. Birkhäuser, Boston (1980).
- [30] G. Ottaviani: Some extensions of the Horrocks criterion to vector bundles on grassmannians and quadrics. Annali di Matem. Vol. 155 (1989), 317-341.
- [31] S. Pumplün: Vector bundles and symmetric bilinear forms over curves of genus 1 and arbitrary index. Mathematische Zeitschrift Vol. 246 (2004), 563-602
- [32] D. Quillen: Higher algebraic K-theory. Algebraic K-theory I, Lecture Notes in Math. 341, Springer (1979), 85-147.
- [33] D. Saltman: Lectures on division algebras. Vol. 94 of CBMS Regional conference Series in Mathematics. American Math. Soc. (1999).
- [34] J.-P. Serre: Local Fields. Springer-Verlag, New York, Berlin (1980).

- [35] J.-P. Serre: Cohomologie Galoisienne. Lecure Notes in Mathematics 5, Springer-Verlag Berlin (1994); English translation: Galois Cohomology, Springer-Verlag Berlin (2002).
- [36] V.E. Voskresenskii: Algebraic Groups and Their Birational Invariants. Translated by Boris Kunyavski, Translations of Math. Monographs AMS Vol. 179 (2000).
- [37] R. Wiegand: Torsion in Picard Groups of Affine Rings. Contemp. Math. Vol 159 (1994), 433-444.

MATHEMATISCHES INSTITUT, HEINRICH–HEINE–UNIVERSITÄT 40225 DÜSSELDORF, GERMANY

 $\hbox{E-mail adress: } novakovic@math.uni-duesseldorf.de$