

# CABLING, CONNECTED SUM OPERATIONS AND THE SLOPE CONJECTURE

KIMIHIKO MOTEGI AND TOSHIE TAKATA

ABSTRACT. The slope conjecture proposed by Garoufalidis asserts that the Jones slopes given by the sequence of degrees of the colored Jones polynomials are boundary slopes. We prove that operations of cabling and connected sum keep the positivity of the slope conjecture. As an application, we verify the slope conjecture for graph knots.

## 1. INTRODUCTION

Let  $K$  be a knot in the 3–sphere  $S^3$  and  $E(K)$  the exterior  $S^3 - \text{int}N(K)$ . Denote by  $(\mu, \lambda)$  the preferred meridian-longitude pair of  $K$ . Then any homotopically nontrivial simple closed curves in  $\partial E(K)$  represents  $p[\mu] + q[\lambda] \in H_1(\partial E(K))$  for some relatively prime integers  $p$  and  $q$ . We call  $p/q \in \mathbb{Q} \cup \{\infty\}$  a *boundary slope* of  $K$  if there exists a connected, orientable, incompressible and boundary-incompressible surface  $F$  such that a component of  $\partial F$  represents  $p[\mu] + q[\lambda] \in H_1(\partial E(K))$ . Let us define:

$$bs(K) = \left\{ r \in \mathbb{Q} \cup \{\infty\} \mid r \text{ is a boundary slope of } K \right\}.$$

Following Hatcher [8]  $bs(K)$  is a finite subset of  $\mathbb{Q} \cup \{\infty\}$  for every knot  $K$ .

The *colored Jones function* of  $K$  is a sequence of Laurent polynomials  $J_{K,n}(q) \in \mathbb{Z}[q^{\pm 1}]$  for  $n \in \mathbb{N}$ , where  $J_{K,2}(q)$  is the ordinary Jones polynomial of  $K$ . Let  $\delta_K(n)$  be the maximal degree of  $J_{K,n}(q) \in \mathbb{Z}[q^{\pm 1}]$ . We call  $x \in \mathbb{R}$  a *cluster point* of a sequence  $\{x_n\}$  if  $x$  is a limit point of a subsequence of  $\{x_n\}$ . We define  $js(K)$  as follows:

$$js(K) = \left\{ \text{cluster points of the sequence } \left\{ \frac{4\delta_K(n)}{n^2} \right\}_{n \in \mathbb{N}} \right\}.$$

Since the colored Jones function is  $q$ -holonomic [4, Theorem 1], Theorem 1 in [3] shows:

$$\delta_K(n) = c_2(n)n^2 + c_1(n)n + c_0(n)$$

for rational valued periodic functions  $c_i(n)$  with an integral period. By Lemma 1.8 in [3],  $js(K)$  is the finite set of 4 times the rational values of the periodic function  $c_2(n)$ . Using the minimal degree  $\delta_K^*(q)$  of  $J_{K,n}(q)$  instead of  $\delta_K(n)$ , we can define:

$$js^*(K) = \left\{ \text{cluster points of the sequence } \left\{ \frac{4\delta_K^*(n)}{n^2} \right\}_{n \in \mathbb{N}} \right\}.$$

---

2010 *Mathematics Subject Classification*. Primary 57M25, 57M27

As noted in [3, 1.4],  $\delta_K^*(n) = -\delta_{K^*}(n)$  and thus  $js^*(K) = -js(K^*)$ , where  $K^*$  is the mirror image of  $K$  and  $-X := \{-x_1, \dots, -x_m\}$  if  $X = \{x_1, \dots, x_m\}$ .

We call an element in  $js(K) \cup js^*(K)$  a *Jones slope* of  $K$ .

In [3], Garoufalidis proposed the following conjecture which relates Jones slopes and boundary slopes.

**Conjecture 1.1 (Slope conjecture).** *For any knot  $K$ , every Jones slope is a boundary slope, i.e.  $js(K) \cup js^*(K) \subset bs(K)$ .*

The conjecture was verified for torus knots, some non-alternating knots including the  $(-2, 3, 7)$ -pretzel knot [3], adequate knots [2] and a 2-parameter family of 2-fusion knots [1, 5]. Note that the class of adequate knots includes all alternating knots and most Montesinos knots. In the present note we prove the following results which give further supporting evidence for the slope conjecture.

**Theorem 1.2.** *Suppose that  $K_1$  and  $K_2$  satisfy the slope conjecture, then the connected sum  $K_1 \sharp K_2$  also satisfies the slope conjecture.*

**Theorem 1.3.** *Suppose that  $K$  satisfies the slope conjecture. Then a  $(p, q)$ -cable  $C_{p,q}(K)$  of  $K$  also satisfies the slope conjecture.*

A knot  $K$  is called a *graph knot* if its exterior  $E(K)$  is a graph manifold, i.e. there is a family of tori which decomposes  $E(K)$  into Seifert fiber spaces. This implies that any graph knot is obtained from unknots by a finite sequence of operations of cabling and connected sum; see [6, Corollary 4.2]. A graph knot can be also characterized as a knot whose Gromov volume vanishes [7, 12, 13]. The trivial knot  $O$  obviously satisfies the slope conjecture ( $js(O) = bs(O) = \{0\}$ ), Theorems 1.2 and 1.3 establish:

**Corollary 1.4.** *Every graph knot satisfies the slope conjecture.*

**Acknowledgments.** K.M. has been partially supported by JSPS Grants-in-Aid for Scientific Research (C), 26400099, The Ministry of Education, Culture, Sports, Science and Technology, Japan and Joint Research Grant of Institute of Natural Sciences at Nihon University for 2014. T.T. has been partially supported by JSPS Grants-in-Aid for Scientific Research (C), 25400094, The Ministry of Education, Culture, Sports, Science and Technology, Japan.

## 2. THE SLOPE CONJECTURE AND CONNECTED SUM OPERATION

In this section we establish Theorem 1.2.

**Lemma 2.1.** *If  $p/q \in js(K_1 \sharp K_2)$ , then there exist  $p_1/q_1 \in js(K_1)$  and  $p_2/q_2 \in js(K_2)$  such that  $p/q = p_1/q_1 + p_2/q_2$ .*

*Proof of Lemma 2.1.* Let us put  $\delta_{K_i}(n) = \alpha_i(n)n^2 + \beta_i(n)n + \gamma_i(n)$  for  $i = 1, 2$ , where  $\alpha_i(n)$ ,  $\beta_i(n)$  and  $\gamma_i(n)$  are periodic functions with integral periods. It is known that

$$J_{K_1 \# K_2, n}(q) = J_{K_1, n}(q)J_{K_2, n}(q)$$

and so we have:

$$\begin{aligned} \delta_{K_1 \# K_2}(n) &= \delta_{K_1}(n) + \delta_{K_2}(n) \\ &= (\alpha_1(n) + \alpha_2(n))n^2 + (\beta_1(n) + \beta_2(n))n + (\gamma_1(n) + \gamma_2(n)). \end{aligned}$$

Since  $\alpha_i(n)$ ,  $\beta_i(n)$  and  $\gamma_i(n)$  are periodic functions with integral periods, so are  $\alpha_1(n) + \alpha_2(n)$ ,  $\beta_1(n) + \beta_2(n)$ , and  $\gamma_1(n) + \gamma_2(n)$ . Therefore, the Jones slope  $p/q$  of  $K_1 \# K_2$  is an element of the finite set of the rational values of  $4(\alpha_1(n) + \alpha_2(n)) = 4\alpha_1(n) + 4\alpha_2(n)$ , and hence  $p/q = p_1/q_1 + p_2/q_2$  for some Jones slopes  $p_1/q_1 \in js(K_1), p_2/q_2 \in js(K_2)$ .  $\square$ (Lemma 2.1)

Since every composite knot has an essential meridional annulus, in the following we may assume  $q_i > 0$  for  $i = 1, 2$ .

**Lemma 2.2.** *If  $p_1/q_1 \in bs(K_1)$  and  $p_2/q_2 \in bs(K_2)$ , then  $p_1/q_1 + p_2/q_2 \in bs(K_1 \# K_2)$ .*

*Proof of Lemma 2.2.* Let  $A$  be an essential annulus in  $E(K_1 \# K_2)$  which decomposes  $E(K_1 \# K_2)$  into  $E(K_1)$  and  $E(K_2)$ . Let  $F_i$  be an essential surface in  $E(K_i)$  and  $m_i$  the number of boundary components of  $F_i$ . (If  $p_i \neq 0$ , then for homological reason  $m_i$  is an even integer.) Then  $\partial F_i$  consists of  $m_i$  mutually parallel loops each of which has slope  $p_i/q_i$  ( $q_i > 0$ ). Note that the core of  $A$  is a meridian of  $K_i$  and choose  $F_i$  so that  $A \cap F_i$  consists of  $m_i q_i$  spanning arcs in  $A$ . See Figure 2.1.




FIGURE 2.1. Essential surfaces  $F_1 \subset E(K_1)$  and  $F_2 \subset E(K_2)$  with  $m_1 = 2$ ,  $q_1 = 2$ ,  $m_2 = 2$ ,  $q_2 = 1$ ;  $\partial F_1 = \gamma_{1,1} \cup \gamma_{1,2}$ ,  $\partial F_2 = \gamma_{2,1} \cup \gamma_{2,2}$ .

Orient  $\partial F_i$  so that they run the same direction on  $\partial E(K_i)$  (independent of an orientation induced from  $F_i$ ) and a component of  $\partial F_1$  and that of  $\partial F_2$  has opposite orientations on  $A$  as in Figure 2.1. In general,  $m_1q_1 \neq m_2q_2$ , i.e. the number of components of  $A \cap F_1$  does not coincide with that of  $A \cap F_2$ , so we take  $m_2q_2$  parallel copies of  $F_1$  and  $m_1q_1$  parallel copies of  $F_2$ . Let us denote these (disconnected) surfaces by  $m_2q_2F_1 \subset E(K_1)$  and  $m_1q_1F_2 \subset E(K_2)$ , respectively. We give an orientation on the boundary of  $m_2q_2F_1$  (resp.  $m_1q_1F_2$ ) so that it coincides with that of  $\partial F_1$  (resp.  $\partial F_2$ ). Since both  $A \cap m_2q_2F_1$  and  $A \cap m_1q_1F_2$  consist of  $m_1m_2q_1q_2$  spanning arcs in  $A$ , we can connect  $m_2q_2F_1$  and  $m_1q_1F_2$  along the annulus  $A$  to obtain a possibly disconnected surface  $F'$  in  $E(K_1 \# K_2)$ . Note that all the components of  $\partial F'$  run the same direction on  $\partial E(K_1 \# K_2)$  with respect to the orientation given in the above.

**Claim 2.3.** *Each component of  $F' \cap \partial E(K_1 \# K_2)$  has slope  $p_1/q_1 + p_2/q_2$ .*

*Proof of Claim 2.3.* Let  $(\mu_i, \lambda_i)$  and  $(\mu, \lambda)$  be preferred meridian-longitude pairs of  $K_i$  and  $K_1 \# K_2$ ; we take  $\mu_1 = \mu_2 = \mu \subset \partial A$ . Orient them so that  $\langle \mu_i, \lambda_i \rangle = \langle \mu, \lambda \rangle = 1$  and  $\langle \mu, \partial F' \rangle = \langle \mu_1, \partial(m_2q_2F_1) \rangle = \langle \mu_2, \partial(m_1q_1F_2) \rangle > 0$ , where  $\langle \alpha, \beta \rangle$  denotes the algebraic intersection number between  $\alpha$  and  $\beta$ . Then

$$\langle \mu, \partial F' \rangle = \langle \mu_1, \partial(m_2q_2F_1) \rangle = m_2q_2 \langle \mu_1, \partial F_1 \rangle = m_2q_2(m_1q_1) = m_1m_2q_1q_2$$

and

$$\begin{aligned} \langle \partial F', \lambda \rangle &= \langle \partial(m_2q_2F_1), \lambda_1 \rangle + \langle \partial(m_1q_1F_2), \lambda_2 \rangle \\ &= m_2q_2 \langle \partial F_1, \lambda_1 \rangle + m_1q_1 \langle \partial F_2, \lambda_2 \rangle \\ &= m_2q_2(m_1p_1) + m_1q_1(m_2p_2) \\ &= m_1m_2p_1q_2 + m_1m_2q_1p_2. \end{aligned}$$

Thus  $F' \cap \partial E(K_1 \# K_2)$  represents

$$\begin{aligned} &(m_1m_2p_1q_2 + m_1m_2q_1p_2)[\mu] + m_1m_2q_1q_2[\lambda] \\ &= m_1m_2((p_1q_2 + q_1p_2)[\mu] + q_1q_2[\lambda]) \in H_1(\partial E(K_1 \# K_2)). \end{aligned}$$

Let  $k$  be the greatest common divisor of  $p_1q_2 + q_1p_2$  and  $q_1q_2$ . Then  $F' \cap \partial E(K_1 \# K_2)$  consists of  $m_1m_2k$  parallel loops each of which has slope

$$\frac{(p_1q_2 + q_1p_2)/k}{(q_1q_2)/k} = (p_1q_2 + q_1p_2)/q_1q_2 = p_1/q_1 + p_2/q_2.$$

□(Claim 2.3)

Let  $F$  be a connected component of  $F'$ . If  $F$  is non-orientable, then we take a tubular neighborhood  $N(F)$  of  $F$  in  $E(K_1 \# K_2)$  and we replace  $F$  by  $\partial N(F)$ , which is an orientable double cover of  $F$  and each component of  $\partial N(F)$  has slope  $p_1/q_1 + p_2/q_2$ ; for simplicity we continue to use the same symbol  $F$  to denote  $\partial N(F)$ . Since  $F_i \subset E(K_i)$  is orientable,  $F \cap E(K_i)$  consists of parallel copies of  $F_i$  for  $i = 1, 2$ . Note also that for each component of  $F \cap E(K_i)$ , its boundary component across  $A$  in the same direction.

**Claim 2.4.** *The surface  $F$  is essential in  $E(K_1 \# K_2)$ .*

*Proof of Claim 2.4.* Suppose for a contradiction that  $F$  is compressible. Let  $D$  be a compressing disk of  $F$ . If  $A \cap D = \emptyset$ , then  $D$  is entirely contained in  $E(K_i)$  and  $F_i$  is compressible, contradicting the assumption. So in the following we assume  $A \cap D \neq \emptyset$ . Recall that  $F \cap E(K_i)$  consists of parallel copies of  $F_i$ . Note that  $A \cap F$  consists of spanning arcs in  $A$  in minimal number of components (Figure 2.2). We may assume that  $D$  intersects  $A$  transversely and the number of components of  $A \cap D$  is minimal. Then  $A \cap D$  consists of circles and arcs whose endpoints belong to  $A \cap F$ . Since  $A$  is incompressible, we eliminate the circle components, and thus  $A \cap D$  consists of arcs; see Figure 2.2.




FIGURE 2.2.  $A \cap F$  and  $A \cap D$

Then  $A \cap D$  consists of properly embedded arcs in  $D$ . Let  $\gamma$  be an outermost arc of  $A \cap D$  in  $D$ ;  $\gamma$  cuts off an outermost disk  $\Delta$ . There are two possibilities: (i)  $\partial\gamma$  is contained in a single arc  $\tau$  of  $A \cap F$  (Figure 2.3(i)), or (ii)  $\gamma$  is an arc connecting two spanning arcs  $\tau_1$  and  $\tau_2$  of  $A \cap F$  (Figure 2.3(ii)).




FIGURE 2.3. An outermost arc  $\gamma$  of  $A \cap D$  in  $D$  and its possible situation in  $A$

Suppose that  $\partial\gamma$  is contained in a single arc  $\tau$  of  $A \cap F$  (Figure 2.3(i)). Then  $\gamma$  is parallel to  $\tau$ ;  $\gamma$  and  $\tau$  cobound a disk  $\Delta' \subset A$  (If  $A \cap F = \tau$ , although there would be a possibility that  $\gamma$  starts from one side of  $\tau$  and ends in the other side of  $\tau$ , this cannot happen for homological reason.) Let  $F_\Delta$  be a unique component of  $F \cap E(K_1)$  or  $F \cap E(K_2)$  intersecting  $\partial\Delta$ ;  $F_\Delta$  is a parallel copy of  $F_1$  or  $F_2$ . Then by the incompressibility of  $F_i$  in  $E(K_i)$  and the irreducibility of  $E(K_i)$ , the

disk  $\Delta \cup \Delta'$  is parallel to a disk in  $\partial E(K_i)$ . Thus we can isotope  $D$  so that  $\gamma$  is removed from  $A \cap D$ . This contradicts the minimality of the number of components of  $A \cap D$ .

Next assume that  $\gamma$  is an arc connecting two spanning arcs  $\tau_1$  and  $\tau_2$  of  $A \cap F$  (Figure 2.3(ii)). As above we take a unique component  $F_\Delta$  of  $F \cap E(K_1)$  or  $F \cap E(K_2)$  intersecting  $\partial\Delta$ ;  $F_\Delta$  is a parallel copy of  $F_1$  or  $F_2$ . Since  $F_\Delta$  is boundary-incompressible,  $\tau_1$  and  $\tau_2$  are contained in a single component of  $\partial F_\Delta$  and run in opposite directions in  $A$ , a contradiction. It follows that  $F$  is incompressible in  $E(K_1 \# K_2)$ .

Since  $K_i$  is non-trivial and  $F_i$  is not a disk for  $i = 1, 2$ ,  $F$  is not an annulus. Hence [9, Lemma 1.10] shows that  $F$  is boundary-incompressible as well. Thus  $F$  is a desired essential surface in  $E(K_1 \# K_2)$  with boundary slope  $p_1/q_1 + p_2/q_2$ .

This completes a proof of Lemma 2.2. □(Lemma 2.2)

**Remark 2.5.** *In the above construction of the surface  $F'$ , we can slide or twist several times  $m_1 q_1 F_2$  along the annulus  $A$  before connecting with  $m_2 q_2 F_1$  without changing its boundary slope, so  $F'$  is not unique.*

Let us turn to a proof of Theorem 1.2. Before proving the theorem, we note the following general fact.

**Claim 2.6.** *Let  $K$  be a knot in  $S^3$ . If  $js^*(K) \subset bs(K)$ , then  $js(K^*) \subset bs(K^*)$ .*

*Proof of Claim 2.6.* If  $r \in js(K^*)$ , then  $-r \in -js(K^*) = js^*(K) \subset bs(K)$ . Thus  $r \in -bs(K) = bs(K^*)$ . □(Claim 2.6)

*Proof of Theorem 1.2.* Assume first that  $p/q \in js(K_1 \# K_2)$ . Then as shown in Lemma 2.1,  $p/q = p_1/q_1 + p_2/q_2$  for some Jones slopes  $p_1/q_1 \in js(K_1)$  and  $p_2/q_2 \in js(K_2)$ . Since  $p_1/q_1 \in js(K_1) \subset bs(K_1)$  and  $p_2/q_2 \in js(K_2) \subset bs(K_2)$  by the initial assumption, Lemma 2.2 shows that  $p/q = p_1/q_1 + p_2/q_2 \in bs(K_1 \# K_2)$ .

Next assume that  $p/q \in js^*(K_1 \# K_2)$ . Then  $-p/q \in js((K_1 \# K_2)^*) = js(K_1^* \# K_2^*)$ . Since  $K_i$  satisfies the slope conjecture,  $js^*(K_i) \subset bs(K_i)$ , thus by Claim 2.6  $js(K_i^*) \subset bs(K_i^*)$ . Apply the above argument to  $K_1^*$  and  $K_2^*$  to conclude that  $-p/q \in bs(K_1^* \# K_2^*) = bs((K_1 \# K_2)^*)$ . Hence  $p/q \in bs(K_1 \# K_2)$ . This completes a proof of Theorem 1.2. □(Theorem 1.2)

### 3. THE SLOPE CONJECTURE AND CABLING OPERATION

Let  $V$  be a standardly embedded solid torus in  $S^3$  and  $k$  a 0-bridge braid in  $V$  which wraps  $p$  times in meridional direction and  $q$  times in longitudinal direction;  $k$  is a  $(p, q)$ -torus knot in  $S^3$ . In the following, we assume  $q > 0$ . Given a nontrivial knot  $K$ , take an orientation preserving embedding  $f : V \rightarrow S^3$  such that the core of  $f(V)$  is  $K$  and  $f$  sends a preferred longitude of  $V$  to that of  $K$ . Then the image  $f(k)$  is called the  $(p, q)$ -cable of  $K$  and denoted by  $C_{p,q}(K)$ . We begin by describing Jones slopes of  $C_{p,q}(K)$ .

**Lemma 3.1.** *If  $r \in js(C_{p,q}(K))$ , then  $r = pq$  or  $aq^2/b$  for some  $a/b \in js(K)$ .*

*Proof of Lemma 3.1.* Let us put  $J'_{\bullet,n}(t) := J_{\bullet,n}(t^4) \frac{t^{2n} - t^{-2n}}{t^2 - t^{-2}} \in \mathbb{Z}[t^{\pm 1}]$  and denote its maximal degree by  $\delta'_{\bullet}(n)$ . For simplicity, we write  $C := C_{p,q}(K)$ . From [11] (see also [14, Theorem 1]), we have:

$$J'_{C,n}(t) = t^{pq(n^2-1)} \sum_{k=-\frac{n-1}{2}}^{k=\frac{n-1}{2}} t^{-4pk(kq+1)} J'_{K,2kq+1}(t),$$

where we use the convention that  $J'_{K,2kq+1}(t) = -J'_{K,-2kq-1}(t)$  if  $2kq+1 \leq 0$ . Then,

$$\begin{aligned} \delta'_C(n) &= pq(n^2 - 1) + \max\{\delta'_K(|2kq+1|) - 4pk(kq+1) \mid -\frac{n-1}{2} \leq k \leq \frac{n-1}{2}\} \\ &= pq(n^2 - 1) + \max\{\delta'_K(|kq+1|) - pk(kq+2) \mid -(n-1) \leq k \leq n-1\}, \end{aligned}$$

where we transform  $2k$  to  $k$  in the second equality.

Let us write  $\delta_K(n) = c_2(n)n^2 + c_1(n)n + c_0(n)$ , where  $c_i(n)$  is a periodic function with an integral period. Then  $\delta'_K(n) = 4\delta_K(n) + 2(n-1) = 4c_2(n)n^2 + (4c_1(n) + 2)n + 4c_0(n) - 2$ . Thus we can write

$$\delta'_K(n) = \alpha(n)n^2 + \beta(n)n + \gamma(n),$$

where  $\alpha(n), \beta(n)$  and  $\gamma(n)$  are periodic functions  $4c_2(n), 4c_1(n) + 2$  and  $4c_0(n) - 2$ , respectively. Then,

$$\begin{aligned} &\delta'_K(|kq+1|) - pk(kq+2) \\ &= \alpha(|kq+1|)(kq+1)^2 + \beta(|kq+1|)|kq+1| + \gamma(|kq+1|) - pk(kq+2) \\ &= \begin{cases} \alpha'_+(k)k^2 + \beta'_+(k)k + \gamma'_+(k) & \text{if } 0 \leq k \leq n-1, \\ \alpha'_-(k)k^2 + \beta'_-(k)k + \gamma'_-(k) & \text{if } -(n-1) \leq k < 0, \end{cases} \end{aligned}$$

where we put

$$\begin{aligned} \alpha'_\pm(k) &:= q(\alpha(\pm qk \pm 1)q - p) \\ \beta'_\pm(k) &:= 2\alpha(\pm qk \pm 1)q \pm \beta(\pm qk \pm 1)q - 2p \\ \gamma'_\pm(k) &:= \alpha(\pm qk \pm 1) \pm \beta(\pm qk \pm 1) + \gamma(\pm qk \pm 1). \end{aligned}$$

Let  $\pi_{\alpha'_\pm}$  be the period of  $\alpha'_\pm(n)$  and

$$\begin{aligned} A_\pm &:= \max\{\alpha'_\pm(k) \mid 1 \leq k \leq \pi_{\alpha'_\pm}\} \cup \{0\} \\ &= \max\{q(\alpha(\pm qk \pm 1)q - p) \mid 1 \leq k \leq \pi_{\alpha'_\pm}\} \cup \{0\}. \end{aligned}$$

From Lemma 3.2 below, for large  $n$  we have:

$$\begin{aligned} \max\{\alpha'_+(k)k^2 + \beta'_+(k)k + \gamma'_+(k) \mid 0 \leq k \leq n-1\} &= A_+n^2 + O(n), \\ \max\{\alpha'_-(k)k^2 + \beta'_-(k)k + \gamma'_-(k) \mid -(n-1) \leq k < 0\} &= A_-n^2 + O(n), \end{aligned}$$

where  $O(n)$  denotes a term that is at most linear in  $n$ .

We put  $A := \max\{A_+, A_-\}$ , which is a nonnegative rational number. If  $A > 0$ , then there exists  $j_A \in \mathbb{N}$  such that  $A = q(\alpha(j_A)q - p) = \alpha(j_A)q^2 - pq$ .

Therefore, we obtain:

$$\begin{aligned}\delta'_C(n) &= pq(n^2 - 1) + An^2 + O(n) \\ &= \begin{cases} \alpha(j_A)q^2n^2 + O(n) & \text{if } A > 0, \\ pqn^2 + O(n) & \text{if } A = 0. \end{cases}\end{aligned}$$

Hence,

$$4\delta_C(n) = \begin{cases} \alpha(j_A)q^2n^2 + O(n) & \text{if } A > 0, \\ pqn^2 + O(n) & \text{if } A = 0. \end{cases}$$

Since  $js(K) = \{4c_2(n) \mid n \in \mathbb{N}\} = \{\alpha(n) \mid n \in \mathbb{N}\}$  and  $js(C) \subset \{\alpha(j_A)q^2, pq\}$ , if  $r \in js(C)$ , then  $r = pq$  or  $aq^2/b$ , where  $a/b = \alpha(j_A) \in js(K)$ . This complete a proof of Lemma 3.1.  $\square$ (Lemma 3.1)

**Lemma 3.2.** *Let  $f(k) = \alpha(k)k^2 + \beta(k)k + \gamma(k)$  be a quadratic quasi polynomial and  $\pi_\alpha$  the period of  $\alpha(k)$ . We put  $A := \max(\{\alpha(j) \mid 1 \leq j \leq \pi_\alpha\} \cup \{0\})$ . Then, for large  $n$ ,*

$$\max\{f(k) \mid 0 \leq k \leq n-1\} = An^2 + O(n).$$

*Proof of Lemma 3.2.* For  $1 \leq j \leq \pi_\alpha$ , we compute

$$\begin{aligned}M_j(n) &:= \max\{f(k) \mid k \equiv j \pmod{\pi_\alpha}, 0 \leq k \leq n-1\} \\ &= \max\{\alpha(j)k^2 + \beta(k)k + \gamma(k) \mid k \equiv j \pmod{\pi_\alpha}, 0 \leq k \leq n-1\}.\end{aligned}$$

We note that there exist polynomials  $f_0, \dots, f_{\pi-1}$  such that  $\alpha(j)k^2 + \beta(k)k + \gamma(k) = f_i(k)$  when  $k \equiv i \pmod{\pi}$ , where  $\pi$  is a period of  $\alpha(j)k^2 + \beta(k)k + \gamma(k)$ , i.e.  $\pi$  is the common period of  $\beta(k)$  and  $\gamma(k)$ .

If  $\alpha(j) > 0$ , then we see that  $M_j(n) = \alpha(j)n^2 + O(n)$  at  $k = n-1$  for large  $n$ . If  $\alpha(j) = 0$ , it is clear that  $M_j(n) = O(n)$ . If  $\alpha(j) < 0$ , then  $M_j(n) = f(c)$  for some constant  $c$  independent of  $n$ , and so  $M_j(n) = O(n)$ . Moreover, if  $0 \leq b < a$ , then  $bn^2 + O(n) < an^2 + O(n)$  for large  $n$ . Hence we obtain:

$$\max\{f(k) \mid 0 \leq k \leq n-1\} = \max_{1 \leq j \leq \pi_\alpha} M_j(n) = An^2 + O(n).$$

$\square$ (Lemma 3.2)

Since the Jones slope  $pq$  is the boundary slope of the cabling annulus of  $C_{p,q}(K)$ ,  $pq \in bs(C_{p,q}(K))$ . The next result was essentially shown by Klaff and Shalen [10], but we give a modified proof here for completeness.

**Lemma 3.3.** *If  $a/b \in bs(K)$ , then  $aq^2/b \in bs(C_{p,q}(K))$ .*

*Proof of Lemma 3.3.* Let  $M_{p,q} = V - \text{int}N(k)$ , the standard  $(p, q)$ -cable space. We denote preferred meridian-longitude pairs of  $N(k)$  and  $V$  by  $(\mu, \lambda)$  and  $(\mu_V, \lambda_V)$ , respectively. Then  $H_1(M_{p,q}) \cong \mathbb{Z} \oplus \mathbb{Z}$  is generated by  $[\mu]$  and  $[\lambda_V]$ . Let  $D$  be a  $q$ -th punctured meridian disk of  $V$  and  $A$  an obvious annulus connecting  $\partial V$  and  $\partial N(k)$ . With suitable orientations we have  $[D \cap \partial N(k)] = q[\mu]$ ,  $[D \cap \partial V] = [\mu_V]$ ,  $[A \cap \partial N(k)] = pq[\mu] + [\lambda]$ ,  $[A \cap \partial V] = p[\mu_V] + q[\lambda_V]$ , and thus  $[\mu_V] = q[\mu]$ ,  $[\lambda] = q[\lambda_V]$  in  $H_1(M_{p,q})$ .

Let  $S$  be an oriented surface in  $M_{p,q}$  representing the nontrivial homology class  $(aq - bp)[D] + b[A] \in H_2(M_{p,q}, \partial M_{p,q})$ . We can construct  $S$  by the “double-curve sum” of  $(aq - bp)$  parallel copies of  $D$  and  $b$  parallel copies of  $A$  (i.e. cut and paste along their intersection arcs to get an embedded surface representing the desired homology class); see Figure 3.1.




FIGURE 3.1. Double-curve sum

Then it is easy to see that each component of  $S \cap \partial V$  has slope  $a/b$ , and that of  $S \cap \partial N(k)$  has slope  $aq^2/b$ . If  $S$  is compressible, then after compression, we take a connected component  $S_0$  of  $S$  which represents nontrivial homology class in  $H_2(M_{p,q}, \partial M_{p,q})$ . Since  $S_0$  represents a nontrivial homology class, it is not a boundary-parallel annulus. Thus the incompressible surface  $S_0$  is also boundary-incompressible [9, Lemma 1.10], i.e.  $S_0$  is essential in  $M_{p,q}$ . Following [9, Proposition 1.11], we may assume (up to isotopy) that  $S_0$  is horizontal, i.e. transverse to all Seifert fibers of  $M_{p,q}$ . This implies  $S_0 \cap \partial V$  and  $S_0 \cap \partial N(k)$  are not empty. In particular, each component of  $S_0 \cap \partial V$  has slope  $a/b$ , and that of  $S_0 \cap \partial N(k)$  has slope  $aq^2/b$ . Let  $m_0$  be the number of components of  $\partial S_0$  on  $\partial V$ .

Now recall that  $C_{p,q}(K) = f(V)$ , where  $f : V \rightarrow S^3$  is the orientation preserving embedding such that the core of  $f(V)$  is  $K$ . Then  $E(C_{p,q}(K)) = E(K) \cup f(M_{p,q})$  in which  $T = \partial E(K) = \partial f(V)$  is an essential torus. Since  $a/b \in bs(K)$ , we have an essential surface  $S_1 \subset E(K)$  which has  $m_1$  boundary components each of which has slope  $a/b$ . Let us take  $m_1$  parallel copies of  $f(S_0)$  and  $m_0$  parallel copies of  $S_1$ . Connecting them, we obtain a possibly disconnected surface  $F'$  in  $E(C_{p,q}(K))$ . Let  $F$  be a connected component of  $F'$ . If  $F$  is non-orientable, then as in the proof of Lemma 2.2 we replace  $F$  by  $\partial N(F)$ , where  $N(F)$  is a tubular neighborhood of  $F$  in  $E(C_{p,q}(K))$ . In the latter case, we continue to use the same symbol  $F$  to denote  $\partial N(F)$ . Since  $S_0$  and  $S_1$  are orientable,  $F \cap E(K)$  consists of parallel copies of  $S_1$ , and  $F \cap f(M_{p,q})$  consists of parallel copies of  $f(S_0)$ . Applying the proof of Claim 2.4, where we use the essentiality of  $T$  instead of that of  $A$ , we see that  $F$  is an essential surface in  $E(C_{p,q}(K))$  with boundary slope  $aq^2/b$ .  $\square$ (Lemma 3.3)

*Proof of Theorem 1.3.* Assume first that  $r \in js(C_{p,q}(K))$ . Then by Lemma 3.1  $r = aq^2/b$  or  $pq$ , where  $a/b \in js(K)$ . If  $r = aq^2/b$ , then Lemma 3.3 shows that

$r \in bs(C_{p,q}(K))$ . If  $r = pq$ , then each boundary component of the cabling annulus has slope  $pq$ , and thus  $r \in bs(C_{p,q}(K))$ .

Next assume that  $r \in js^*(C_{p,q}(K))$ . Then  $-r \in js((C_{p,q}(K))^*) = js(C_{-p,q}(K^*))$ . Since  $K$  satisfies the slope conjecture,  $js^*(K) \subset bs(K)$ . Hence by Claim 2.6  $js(K^*) \subset bs(K^*)$ . Let us apply the above argument to  $K^*$  and its  $(-p, q)$ -cable to conclude that  $-r \in bs(C_{-p,q}(K^*)) = bs((C_{p,q}(K))^*)$ . This then implies  $r \in bs(C_{p,q}(K))$ , and completes a proof.  $\square$ (Theorem 1.3)

#### REFERENCES

- [1] N.M. Dunfield and S. Garoufalidis; Incompressibility criteria for spun-normal surfaces, *Trans. Amer. Math. Soc.*, **364** (2012), 6109–6137.
- [2] D. Futer, E. Kalfagianni and J. Purcell; Slopes and colored Jones polynomials of adequate knots, *Proc. Amer. Math. Soc.*, **139** (2011), 1889–1896.
- [3] S. Garoufalidis; The Jones slopes of a knot, *Quantum Topology* **2** (2011), 43–69.
- [4] S. Garoufalidis and T.T. Le; The colored Jones function is  $q$ -holonomic, *Geom. Topol.* **9** (2005), 1253–1293.
- [5] S. Garoufalidis and R. van der Veen; Quadratic integer programming and the slope conjecture, arXiv:1405.5088.
- [6] C.McA. Gordon; Dehn surgery and satellite knots, *Trans. Amer. Math. Soc.*, **275** (1983), 687–708.
- [7] M. Gromov; Volume and bounded cohomology, *Inst. Hautes Études Sci. Publ. Math.* **56** (1983), 213–307.
- [8] A.E. Hatcher; On the boundary curves of incompressible surfaces, *Pacific J. Math.* **99** (1982), 373–377.
- [9] A.E. Hatcher; Notes on basic 3-manifold topology (2000). Freely available at <http://www.math.cornell.edu/hatcher>
- [10] B. Klaff and P.B. Shalen; The diameter of the set of boundary slopes of a knot, *Algebr. Geom. Topol.* **6** (2006), 1095–1112.
- [11] H. Morton; The coloured Jones function and Alexander polynomial for torus knots, *Proc. Camb. Philos. Soc.* **117** (1995), 129–135.
- [12] T. Soma; The Gromov invariant of links, *Invent. Math.* **64** (1981), 445–454.
- [13] W.P. Thurston; The geometry and topology of 3-manifolds, Lecture notes, Princeton University, 1979.
- [14] R. van der Veen; A Cabling formula for the colored Jones polynomial, arXiv:0807.2679.

DEPARTMENT OF MATHEMATICS, NIHON UNIVERSITY, 3-25-40 SAKURAJOSUI, SETAGAYA-KU, TOKYO 156-8550, JAPAN

*E-mail address:* motegi@math.chs.nihon-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU, FUKUOKA 819-0395, JAPAN

*E-mail address:* ttakata@math.kyushu-u.ac.jp