

Seifert surgery on knots via Reidemeister torsion and Casson-Walker-Lescop invariant II

Teruhisa Kadokami, Noriko Maruyama and Tsuyoshi Sakai

June 3, 2015

Abstract

For a knot K with $\Delta_K(t) \doteq t^2 - 3t + 1$ in a homology 3-sphere, let M be the result of $2/q$ -surgery on K . We show that an appropriate assumption on the Reidemeister torsion of the universal abelian covering of M implies $q = \pm 1$, if M is a Seifert fibered space.

1 Introduction

The first author [Kd1] studied the Reidemeister torsion of Seifert fibered homology lens spaces, and showed the following:

Theorem 1.1 ([Kd1, Theorem 1.4]) *Let K be a knot in a homology 3-sphere Σ such that the Alexander polynomial of K is $t^2 - 3t + 1$. The only surgeries on K that may produce a Seifert fibered space with base S^2 and with $H_1 \neq \{0\}, \mathbb{Z}$ have coefficients $2/q$ and $3/q$, and produce Seifert fibered space with three singular fibers. Moreover (1) if the coefficient is $2/q$, then the set of multiplicities is $\{2\alpha, 2\beta, 5\}$ where $\gcd(\alpha, \beta) = 1$, and (2) if the coefficient is $3/q$, then the set of multiplicities is $\{3\alpha, 3\beta, 4\}$ where $\gcd(\alpha, \beta) = 1$.*

It is conjectured that Seifert surgeries on non-trivial knots are integral (except some cases). We [KMS] have studied the $2/q$ -Seifert surgery, one of the remaining cases of the above theorem, by applying the Reidemeister torsion and the Casson-Walker-Lescop invariant, and have given sufficient conditions to determine the integrality of $2/q$ ([KMS, Theorems 2.1, 2.3]).

In this paper, we give another condition for the integrality of $2/q$ (Theorem 2.1). Like as in [KMS], the condition is also suggested by computations for the figure eight knot ([KMS, Example 2.2]).

We note two differences of this paper from [KMS]; one is that the surgery coefficient appears in the condition instead of the Casson-Walker-Lescop invariant, and another is that we need more delicate estimation for the Dedekind sum to prove the result.

(1) Let Σ be a homology 3-sphere, and let K be a knot in Σ . Then $\Delta_K(t)$ denotes the Alexander polynomial of K , and $\Sigma(K; p/r)$ denotes the result of p/r -surgery on K .

(2) The first author [Kd2] introduced the norm of polynomials and homology lens spaces: Let ζ_d be a primitive d -th root of unity. For an element α of $\mathbb{Q}(\zeta_d)$, $N_d(\alpha)$ denotes the norm of α

⁰2010 *Mathematics Subject Classification*: 11R04, 11R27, 57M25, 57M27.

Keywords: Reidemeister torsion, Casson-Walker-Lescop invariant, Seifert fibered space.

associated to the algebraic extension $\mathbb{Q}(\zeta_d)$ over \mathbb{Q} . Let $f(t)$ be a Laurent polynomial over \mathbb{Z} . We define $|f(t)|_d$ by

$$|f(t)|_d = |N_d(f(\zeta_d))| = \left| \prod_{i \in (\mathbb{Z}/d\mathbb{Z})^\times} f(\zeta_d^i) \right|.$$

Let X be a homology lens space with $H_1(X) \cong \mathbb{Z}/p\mathbb{Z}$. Then there exists a knot K in a homology 3-sphere Σ such that $X = \Sigma(K; p/r)$ ([BL, Lemma 2.1]). We define $|X|_d$ by

$$|X|_d = |\Delta_K(t)|_d,$$

where d is a divisor of p . Then $|X|_d$ is a topological invariant of X (Refer to [Kd2] for details).

(3) Let X be a closed oriented 3-manifold. Then $\lambda(X)$ denotes the Lescop invariant of X ([Le]). Note that $\lambda(S^3) = 0$.

2 Result

Let K be a knot in a homology 3-sphere Σ . Let M be the result of $2/q$ -surgery on K : $M = \Sigma(K; 2/q)$. Let $\pi : X \rightarrow M$ be the universal abelian covering of M (i.e. the covering associated to $\text{Ker}(\pi_1(M) \rightarrow H_1(M))$). Since $H_1(M) \cong \mathbb{Z}/2\mathbb{Z}$, π is the 2-fold unbranched covering.

In [KMS], we have defined $|K|_{(q,d)}$ by the following formula, if $|X|_d$ is defined:

$$|K|_{(q,d)} := |X|_d.$$

Assume that the Alexander polynomial of K is $t^2 - 3t + 1$. Then, as noted in [KMS], $H_1(X) \cong \mathbb{Z}/5\mathbb{Z}$ and $|K|_{(q,5)}$ is defined.

We then have the following.

Theorem 2.1 *Let K be a knot in a homology 3-sphere Σ . We assume the following.*

(2.1) $\lambda(\Sigma) = 0$,

(2.2) $\Delta_K(t) \doteq t^2 - 3t + 1$,

(2.3) $|q| \geq 3$,

(2.4) $\sqrt{|K|_{(q,5)}} > 4q^2$.

Then $M = \Sigma(K; 2/q)$ is not a Seifert fibered space.

Remark 2.2 Let K be the figure eight knot in S^3 . Note that $\Delta_K(t) \doteq t^2 - 3t + 1$. Then $|K|_{(q,5)} = (5q^2 - 1)^2$ by [KMS, Example 2.2]. Hence (2.4) holds if $|q| \geq 3$.

Remark 2.3 Theorem 2.1 seems to suggest studying the asymptotic behavior of $|K|_{(q,d)}$ as a function of q .

3 An inequality for the Dedekind sum

To prove Theorem 2.1, we need the following inequality for the Dedekind sum $s(\cdot, \cdot)$ ([RG]):

Proposition 3.1 ([Ma, Lemma 3]) *For an even integer $p \geq 8$ and for an odd integer q such that $3 \leq q \leq p - 3$ and $\gcd(p, q) = 1$, we have*

$$|s(q, p)| < f(2, p)$$

where $f(2, p) = \frac{(p-1)(p-5)}{24p}$.

By this proposition, we immediately have the following.

Lemma 3.2 *For an even integer $p \geq 8$ and for an integer q_* such that $q_* \not\equiv \pm 1 \pmod{p}$ and $\gcd(p, q_*) = 1$, we have*

$$|s(q_*, p)| < \frac{p}{24}.$$

Proof. By assumptions, there exists q such that $q_* \equiv q \pmod{p}$ and $3 \leq q \leq p - 3$. Hence by Proposition 3.1, we have

$$|s(q_*, p)| = |s(q, p)| < \frac{(p-1)(p-5)}{24p} < \frac{p}{24}.$$

□

Remark 3.3 The estimation given in Proposition 3.1 has a natural application ([Ma]).

4 Proof of Theorem 2.1

Suppose that $M = \Sigma(K; 2/q)$ is a Seifert fibered space. Then, as shown in [KMS], we may assume that

(*) : M has a framed link presentation as in Figure 1,

where $1 \leq \alpha < \beta$ and $\gcd(\alpha, \beta) = 1$.




Figure 1: A framed link presentation of $M = \Sigma(K; 2/q)$

Also as shown in [KMS], $\sqrt{|K|_{(q,5)}} = (\alpha\beta)^2$. Hence by **(2.4)**,

$$(\alpha\beta)^2 > 4q^2 \quad (4.1)$$

By **(2.1)**, **(2.2)** and [Le, 1.5 T2], we have $\lambda(M) = -q$. Hence $(\alpha\beta)^2 > 4\{\lambda(M)\}^2$, and hence

$$|\lambda(M)| < \frac{\alpha\beta}{2} \quad (4.2)$$

We now consider e defined as follows:

$$e := \frac{q_1}{2\alpha} + \frac{q_2}{2\beta} + \frac{q_3}{5}.$$

According to the sign of e , we treat two cases separately: We first consider the case $e > 0$. Then the order of $H_1(M)$ is $20\alpha\beta e$. Since $H_1(M) \cong \mathbb{Z}/2\mathbb{Z}$, $20\alpha\beta e = 2$, and $e = 1/(10\alpha\beta)$. Hence by (*) and [Le, Proposition 6.1.1], we have

$$\lambda(M) = \left(-\frac{4}{5}\right)\alpha\beta + \frac{5\beta}{24\alpha} + \frac{5\alpha}{24\beta} + \frac{1}{120\alpha\beta} - \frac{1}{4} - T \quad (4.3)$$

where $T = s(q_1, 2\alpha) + s(q_2, 2\beta) + s(q_3, 5)$.

By (4.2), we have

$$-\frac{\alpha\beta}{2} < \lambda(M).$$

Hence by (4.3),

$$-\frac{\alpha\beta}{2} < \left(-\frac{4}{5}\right)\alpha\beta + \frac{5\beta}{24\alpha} + \frac{5\alpha}{24\beta} + \frac{1}{120\alpha\beta} - \frac{1}{4} + |T|.$$

Consequently

$$\frac{3}{10}\alpha\beta < -\frac{1}{4} + \frac{5}{24\alpha}\beta + \frac{5}{24}\left(\frac{\alpha}{\beta}\right) + \frac{1}{120\alpha\beta} + |T| \quad (4.4)$$

As in [KMS], we show that $\alpha \geq 2$ implies a contradiction: Suppose that $\alpha \geq 2$. Since $\alpha < \beta$, we have $\beta \geq 3$ and $\alpha/\beta < 1$. Hence

$$\frac{3}{5}\beta < -\frac{1}{4} + \frac{5}{24 \cdot 2}\beta + \frac{5}{24} + \frac{1}{120 \cdot 2 \cdot 3} + |T|.$$

Since $|s(q_1, 2\alpha)| \leq \frac{2\alpha}{12} < \frac{2\beta}{12}$, $|s(q_2, 2\beta)| \leq \frac{2\beta}{12}$, and $|s(q_3, 5)| \leq \frac{1}{5}$ as in [KMS], we have

$$|T| \leq |s(q_1, 2\alpha)| + |s(q_2, 2\beta)| + |s(q_3, 5)| \leq \frac{\beta}{3} + \frac{1}{5}.$$

Hence

$$\frac{3}{5}\beta < -\frac{1}{4} + \frac{5}{48}\beta + \frac{5}{24} + \frac{1}{120 \cdot 6} + \left(\frac{\beta}{3} + \frac{1}{5}\right).$$

Thus

$$\left(\frac{3}{5} - \frac{5}{48} - \frac{1}{3}\right)\beta < -\frac{1}{4} + \frac{5}{24} + \frac{1}{120 \cdot 6} + \frac{1}{5}.$$

Therefore

$$\frac{39}{240}\beta < \frac{1}{240} \left(38 + \frac{1}{3}\right) < \frac{39}{240}.$$

This contradicts $\beta \geq 3$.

We next show that $\alpha = 1$ implies a contradiction: Suppose that $\alpha = 1$. By (4.1), $\beta^2 > 4q^2$. Since $|q| \geq 3$, $\beta^2 > 4 \cdot 3^2 = 36$. Hence $\beta > 6$. Since $\alpha = 1$, $e = \frac{1}{10\beta}$. Hence

$$\frac{q_1}{2} + \frac{q_2}{2\beta} + \frac{q_3}{5} = \frac{1}{10\beta}$$

and hence we have the following equation.

$$(5\beta)q_1 + 5q_2 + (2\beta)q_3 = 1 \tag{4.5}$$

Since q_1 and q_2 are odd (see Figure 1), β must be even. Since $\beta > 6$, we have $\beta \geq 8$. We then have

$$(\#) : \quad q_2 \not\equiv \pm 1 \pmod{2\beta}.$$

In fact, since q_1 is odd, $(5\beta)q_1 \equiv \beta \pmod{2\beta}$. Hence by (4.5),

$$\beta + 5q_2 \equiv 1 \pmod{2\beta}.$$

Now suppose that $q_2 \equiv 1 \pmod{2\beta}$. Then $\beta + 5 \equiv 1 \pmod{2\beta}$. This is impossible since $\beta \geq 8$. Next suppose that $q_2 \equiv -1 \pmod{2\beta}$. Then $\beta - 5 \equiv 1 \pmod{2\beta}$. This is also impossible since $\beta \geq 8$. Thus $(\#)$ holds.

Substituting $\alpha = 1$ in (4.4),

$$\frac{3}{10}\beta < -\frac{1}{4} + \frac{5}{24}\beta + \frac{5}{24\beta} + \frac{1}{120\beta} + |T|$$

where $T = s(q_2, 2\beta) + s(q_3, 5)$ (since $s(q_1, 2) = 0$). By $(\#)$ and Lemma 3.2,

$$|s(q_2, 2\beta)| < \frac{2\beta}{24} = \frac{\beta}{12}.$$

Hence

$$|T| \leq |s(q_2, 2\beta)| + |s(q_3, 5)| < \frac{\beta}{12} + \frac{1}{5}.$$

Since $\beta \geq 8$,

$$\frac{3}{10}\beta < -\frac{1}{4} + \frac{5}{24}\beta + \frac{5}{24 \cdot 8} + \frac{1}{120 \cdot 8} + \left(\frac{\beta}{12} + \frac{1}{5}\right).$$

Thus

$$\left(\frac{3}{10} - \frac{5}{24} - \frac{1}{12}\right)\beta < -\frac{1}{4} + \frac{5}{24 \cdot 8} + \frac{1}{120 \cdot 8} + \frac{1}{5}$$

and hence $\frac{1}{120}\beta < 0$. This is a contradiction, and ends the proof in the case $e > 0$.

We finally consider the case $e < 0$. Then $e = -\frac{1}{10\alpha\beta}$. By $(*)$ and [Le, Proposition 6.1.1], we have

$$\lambda(M) = -\left\{\left(-\frac{4}{5}\right)\alpha\beta + \frac{5\beta}{24\alpha} + \frac{5\alpha}{24\beta} + \frac{1}{120\alpha\beta} - \frac{1}{4} + T\right\}.$$

Remaining part of the proof is similar to that in the case $e > 0$.

This completes the proof of Theorem 2.1. □

References

- [BL] S. Boyer and D. Lines, *Surgery formulae for Casson's invariant and extensions to homology lens spaces*, J. Reine Angew. Math., **45** (1990), 181–220.
- [Kd1] T. Kadokami, *Reidemeister torsion of Seifert fibered homology lens spaces and Dehn surgery*, Algebr. Geom. Topol., **7** (2007), 1509–1529.
- [Kd2] T. Kadokami, *Reidemeister torsion and lens surgeries on knots in homology 3-spheres II*, Top. Appl., **155**, no.15 (2008), 1699–1707.
- [KMS] T. Kadokami, N.Maruyama and T. Sakai, *Seifert surgery on knots via Reidemeister torsion and Casson-Walker-Lescop invariant*, Top. Appl., **188** (2015), 64–73.
- [Le] C. Lescop, *Global surgery formula for the Casson-Walker invariant*, Ann. of Math. Studies, Princeton Univ. Press., **140** (1996).
- [Ma] N. Maruyama, *On a distribution of rational homology 3-spheres captured by the CWL invariant*, Preprint (2009).
- [RG] H. Rademacher and E. Grosswald, *Dedekind sums*, The Carus Mathematical Monograph, **16** (1972).

Teruhisa Kadokami
Department of Mathematics, East China Normal University,
Dongchuan-lu 500, Shanghai, 200241, China
mshj@math.ecnu.edu.cn, kadokami2007@yahoo.co.jp

Noriko Maruyama
Musashino Art University,
Ogawa 1-736, Kodaira, Tokyo 187-8505, Japan
maruyama@musabi.ac.jp

Tsuyoshi Sakai
Department of Mathematics, Nihon University,
3-25-40, Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan