Blowing-up of locally monomially foliated space

Aymen Braghtha

January 27, 2022

Abstract

In this paper, we prove that the blowing-up preserve the local monomiality of foliated space.

1 Locally monomial foliations

Let M be an analytic manifold of dimension n and $D \subset M$ be a divisor with normal crossings. We denote respectively by \mathcal{O}_M and $\Theta_M[\log D]$ the sheaf of holomorphic functions and the sheaf of vector fields on M which are tangent to D.

A singular foliation on (M, D) is coherent subsheaf \mathcal{F} of $\Theta_M[\log D]$ which is reduced and integrable (see [1] and [2]). The dimension (or the rank) of \mathcal{F} is given by

$$s = \max_{p \in M} \dim \mathcal{F}(p)$$

where $\mathcal{F}(p) \subset T_p M$ denote the vector subspace generated by the evaluation of \mathcal{F} at p.

Let \mathbb{F} be a field (we usually take $\mathbb{F} = \mathbb{Q}, \mathbb{R}$ or \mathbb{C}). We shall say that \mathcal{F} is \mathbb{F} -locally monomial if for each point $p \in M$ there exists

1. a local system of coordinates $x = (x_1, \ldots, x_n)$ at p

2. an s-dimensional vector subspace $V \subset \mathbb{F}^n$

such that D is locally given by

$$D_p = \{x_i = 0 : i \in I\}, \text{ for some } I \subset \{1, \dots, n\}$$

and \mathcal{F}_p is the $\mathcal{O}_{M,p}$ -module generated by the abelian Lie algebra

$$\mathcal{L}(V) = \{\sum_{i=1}^{n} a_i x_i \frac{\partial}{\partial x_i} : a \in V\} \bigoplus_{i \overline{I}: e_i \in M} \mathbb{F} \frac{\partial}{\partial x_i}$$

where $\overline{I} = \{1, \ldots, n\} \setminus I$. We shall say that the triple (M, D, \mathcal{F}) is locally monomially foliated space and that (x, I, V) is a local presentation at p.

Lemma 1.1. Let (x, I, V) be a local presentation for (M, D, \mathcal{F}) at a point p. Then, for each vector $m \in V^{\perp}$, the (possibly multivalued) function $f(x) = x^m$ is a first integral of \mathcal{F} .

2 Blowing-up

Let $Y \subset M$ be a smooth submanifold of codimension r. We shall say that Y has normal crossings with (M, D, \mathcal{F}) if for each point $p \in Y$ there exists a local presentation (x, I, V) at p such that Y is given by

$$Y = \{x_1 = x_2 = \dots = x_r = 0\}.$$

Proposition 1.2. Let $\Phi : \widetilde{M} \to M$ be the blowing-up with a center Y which has normal crossings with (M, D, \mathcal{F}) . Let

 $\widetilde{D} = \Phi^{-1}(D)$ and $\widetilde{\mathcal{F}} \subset \Theta_{\widetilde{M}}[\log \widetilde{D}]$

denote the total transform of D and the strict transform of \mathcal{F} respectively. Then, the triple $(\widetilde{M}, \widetilde{D}, \widetilde{\mathcal{F}})$ is a locally monomially foliated space.

The proof is based on the following result on linear algebra.

3 Some linear algebra

Let \mathbb{F} be a field and let $V \subset \mathbb{F}^n$ be a vector subspace of dimension s. Let us fix a disjoint partition of indices $\{1, \ldots, n\} = I_1 \sqcup I_2$ write $\mathbb{F}^n = \mathbb{F}^{I_1} \oplus \mathbb{F}^{I_2}$ and let $\pi_I : \mathbb{F}^n \to \mathbb{F}^I$ denote the projection in the corresponding subspace \mathbb{F}^I generated by $\{e_i : i \in I\}$.

Lemma 3.1. There exists a basis for V such that for each vector v in this basis, either

$$v = \pi_{I_2}(v)$$
 or $v = \pi_{I_2}(v) + e_i$

for some $i \in I_1$

Proof. Up to a permutation of coordinates, we can suppose that $I_1 = \{1, \ldots, n_1\}$ and $I_1 = \{n_1 + 1, \ldots, n\}$ (with the convention that $n_1 = 0$ if $I_1 = \emptyset$)

Let $M = [m_1, \ldots, m_s]$ be a $s \times n$ matrix whose rows are an arbitrary basis of V. By a finite number of elementary row operations and permutations of columns (which leave invariant the subsets I_1 and I_2), we can suppose that the matrix M has the form

$$M = \begin{pmatrix} Id_{k_1 \times k_1} & A_{k_1 \times l_1} & 0_{k_1 \times k_2} & B_{k_1 \times l_2} \\ \hline 0_{k_2 \times k_1} & 0_{k_2 \times l_1} & Id_{k_2 \times k_2} & C_{k_2 \times l_2} \end{pmatrix}$$

where $k_i + l_i = |I_i|$ for $i = 1, 2, k_1 + k_2 = s$ and $Id_{k,l}$ and $0_{k,l}$ denote the $k \times l$ identity and zero matrix respectively.

respectively. Now, it suffices to define $v_i = e_i + \sum_{n_1+k_2+1 \leq j} m_{i,j}e_j$, for $i = 1, \ldots, k_1$ and $v_i = e_{i+n_1} + \sum_{n_1+k_2+1 \leq j} m_{i,j}e_j$ for $i = k_1 + 1, \ldots, s$.

References

- Paul Baum and Raoul Bott, Singularities of holomorphic foliations. J. Differential Geometry, 7: 279-342, 1972.
- Yoshiki Mitera and Junya Yoshizaki. The local analytical triviality of a complex analytic singular foliation. Hokkaido Math. J., 33(2):275-297, 2004

Burgundy University, Burgundy Institue of Mathematics, U.M.R. 5584 du C.N.R.S., B.P. 47870, 21078 Dijon Cedex - France. E-mail adress: aymenbraghtha@yahoo.fr