

On acylindrical hyperbolicity of groups with positive first ℓ^2 -Betti number

D. Osin

Abstract

We prove that every finitely presented group with positive first ℓ^2 -Betti number that surjects onto \mathbb{Z} is acylindrically hyperbolic. In particular, this implies acylindrical hyperbolicity of groups of deficiency at least 2.

1 Introduction

Recall that a group G is *acylindrically hyperbolic* if it admits a non-elementary acylindrical action on a hyperbolic space; in this context, the action is non-elementary if and only if G is not virtually cyclic and has unbounded orbits [13]. The class of acylindrically hyperbolic groups is very broad and includes non-elementary hyperbolic and relatively hyperbolic groups, all but finitely many mapping class groups of orientable closed surfaced (possibly with punctures), $Out(F_n)$ for $n \geq 2$, most 3-manifold groups, non-virtually cyclic groups acting properly on proper $CAT(0)$ spaces and containing rank-1 elements, and many other examples.

On the other hand, being acylindrically hyperbolic is a rather strong property. In particular, many aspects of the theory of hyperbolic and relatively hyperbolic groups can be generalized in the context of acylindrical hyperbolicity. These include various algebraic, model-theoretic, and analytic properties (see [13] for a more detailed survey), small cancellation theory [7], and group theoretic Dehn surgery [4].

The first ℓ^2 -Betti number, denoted $\beta_1^{(2)}(G)$, is a numerical invariant of G ranging in the interval $[0, \infty]$. The study of ℓ^2 -Betti numbers of manifolds was started by Atiyah. The definition was extended to groups in the groundbreaking work of Cheeger and Gromov [2], and since then the theory has been further developed by Gaboriau, Lück and others (see [5, 9] and references therein). This work and more recent papers of Peterson-Thom [15] and Chifan-Sinclair-Udrea [3] show that groups with positive first ℓ^2 -Betti numbers share many algebraic and analytic properties with groups defined by various geometric “negative curvature” type conditions.

For instance, if G is a countable group with $\beta_1^{(2)}(G) > 0$, then G has finite amenable radical, is not a direct product of two infinite groups, is not boundedly generated, and is

not inner amenable (hence the von Neumann algebra of G does not have property Γ of Murray and von Neumann) provided all non-trivial conjugacy classes of G are infinite; if, in addition, G satisfies a weak version of the Atiyah Conjecture, then it contains non-cyclic free subgroups and its reduced C^* -algebra is simple and has unique trace [3, 15]. All these properties, or their stronger analogues, are also known for acylindrically hyperbolic groups.

At first glance, this might seem to be a plain coincidence. However, it turns out that the classes of acylindrically hyperbolic groups and groups with positive first ℓ^2 -Betti numbers are related more closely than one might think.

Theorem 1.1. *Let G be a finitely presented group. Suppose that $\beta_1^{(2)}(G) > 0$ and G maps onto \mathbb{Z} . Then G is acylindrically hyperbolic.*

By the Lück Approximation Theorem, every finitely presented residually finite group G with $\beta_1^{(2)}(G) > 0$ virtually maps onto \mathbb{Z} . Thus we obtain the following.

Corollary 1.2. *Let G be a finitely presented residually finite group with $\beta_1^{(2)}(G) > 0$. Then G is virtually acylindrically hyperbolic.*

Note that finite presentability of G is essential here. Indeed there exist finitely generated (but not finitely presented) residually finite torsion groups with positive first ℓ^2 -Betti number [10], while acylindrically hyperbolic groups are never torsion [4].

The next application of Theorem 1.1 is to groups of positive deficiency. It answers a question asked in [11, Problem 8.3]. In the particular case of groups with one relation and at least 3 generators this result was proved in [11]. Recall that a group G has deficiency at least k if there exists a finite presentation of G with k more generators than relations.

Corollary 1.3. *Every finitely presented group of deficiency at least 2 is acylindrically hyperbolic.*

Recall that, by the Baumslag-Pride theorem, every finitely presented group of deficiency at least 2 is large, i.e., has a finite index subgroup that admits an epimorphism onto a non-cyclic free group [1]. However, this does not guarantee acylindrical hyperbolicity. For example, $F_2 \times \mathbb{Z}$ is large, but not acylindrically hyperbolic.

We list here few immediate consequences of our main theorem. Note that all of them but (a) are new even for groups of deficiency 2 and, in general, fail for large groups.

Corollary 1.4. *Let G be a finitely presented group with $\beta_1^{(2)}(G) > 0$ that surjects onto \mathbb{Z} . Then the following hold.*

- (a) G is SQ-universal.
- (b) $\dim H_b^{(2)}(G, \ell^p(G)) = \infty$ for all $1 \leq p < \infty$.

If, in addition, G has no non-trivial finite normal subgroups, then:

(c) (cf. [15, Cor. 4.6]) The reduced C^* -algebra of G is simple and has unique trace.

(d) G does not satisfy any non-trivial mixed identity.

Recall that G is *SQ-universal* if every countable group embeds in a quotient of G . In particular, (a) implies that G is “algebraically large”; for example, it contains non-cyclic free subgroups and has uncountably many normal subgroups. By a *mixed identity* of a group G we mean any element $w \in G * \langle x_1, \dots, x_k \rangle$ such that every homomorphism $G * \langle x_1, \dots, x_k \rangle \rightarrow G$ that restricts to the identity map on G takes w to 1. Not satisfying any non-trivial mixed identity is a rather restrictive conditions. It implies several interesting model-theoretic, geometric, and algebraic properties of G . For details see [8].

2 Preliminaries

In this section we collect results about ℓ^2 -Betti numbers and acylindrically hyperbolic groups used in the proof of Theorem 1.1 and its corollaries.

An isometric action of a group G on a metric space S is called *acylindrical* if for every $\varepsilon > 0$ there exist $R, N > 0$ such that for every two points $x, y \in S$ satisfying $d(x, y) \geq R$, there are at most N elements $g \in G$ such that

$$d(x, gx) \leq \varepsilon \quad \text{and} \quad d(y, gy) \leq \varepsilon.$$

By [13, Theorem 1.2], a group G is acylindrically hyperbolic if and only if there exists a (possibly infinite) generating set X of G such that the Cayley graph of G with respect to X , denoted $\Gamma(G, X)$, is hyperbolic, its Gromov boundary consists of more than 2 points, and the action of G on $\Gamma(G, X)$ is acylindrical.

For the definition of the first ℓ^2 -Betti number of a group we refer to [9]. Readers who are not familiar with this theory may think of the following Lück’s Approximation Theorem as a definition in the particular case of finitely presented residually finite groups.

Theorem 2.1 (Lück). *Let G be a finitely presented group and let $G_1 \geq G_2 \geq \dots$ be a chain of finite index normal subgroups of G such that $\bigcap_{i=1}^{\infty} G_i = \{1\}$. Then*

$$\beta_1^{(2)}(G) = \lim_{i \rightarrow \infty} \frac{b_1(G_i)}{[G : G_i]},$$

where $b_1(G_i) = \dim(G_i/[G_i, G_i] \otimes \mathbb{Q})$ is the ordinary first Betti number of G_i .

Recall that a subgroup H of a group G is *s-normal* if the intersection $H^g \cap H$ is infinite for every $g \in G$. The following theorem was proved by Peterson and Thom in [15, Theorem

5.12]. In fact, they prove this result under a slightly weaker assumption that H is weakly s -normal, which we do not need (and do not explain) here. Note also that the assumption that H is infinite in the formulation of [15, Theorem 5.12] is redundant since s -normal subgroups are infinite by the definition.

Theorem 2.2 (Peterson-Thom). *Let G be a countable group, H an infinite index s -normal subgroup of G . Suppose that $\beta_1^{(2)}(H) < \infty$ (e.g., this is the case if H is finitely generated). Then $\beta_1^{(2)}(G) = 0$.*

We also need the following theorem, which provides a sufficient condition for an HNN-extension to be acylindrically hyperbolic.

Theorem 2.3 (Minasyan-Osin, [11, Crollary 2.2]). *Let G be an HNN-extension of a group A with associated subgroups C and D . Suppose that $C \neq A \neq D$ and there exists $g \in G$ such that $C^g \cap C$ is finite. Then G is acylindrically hyperbolic.*

3 Proofs

Our proof of Theorem 1.1 is based on the following elementary lemma.

Lemma 3.1. *Let G be a finitely presented group admitting an epimorphism $\varepsilon: G \rightarrow \mathbb{Z}$. Then G is an HNN-extension of a finitely generated group with finitely generated associated subgroups.*

Proof. We fix a finite presentation

$$G = \langle t, a_1, \dots, a_k \mid R_1, \dots, R_m \rangle \quad (1)$$

of the group G such that $\varepsilon(t)$ generates \mathbb{Z} . Note that the total sum of exponents of all occurrences of $t^{\pm 1}$ in each R_i is 0. Let N be the maximal number of occurrences of t in a relator in (1). Then every R_i can be rewritten (possibly after conjugation by a suitable power of t) as a product of $b_{\alpha\beta}^{\pm 1} = t^\beta a_\alpha^{\pm 1} t^{-\beta}$, where the indices range as follows: $1 \leq \alpha \leq k$, $0 \leq \beta \leq N$. Let S_i be the word in the alphabet $B^{\pm 1}$, where

$$B = \{b_{\alpha\beta} \mid 1 \leq \alpha \leq k, 0 \leq \beta \leq N\},$$

obtained from R_i after such a rewriting. By using Tietze transformations, we can rewrite (1) in the form

$$G = \langle t, B \mid S_1, \dots, S_m, \mathcal{T} \rangle,$$

where \mathcal{T} is the set of all relations of the form

$$tb_{\alpha,\beta}t^{-1} = b_{\alpha,\beta+1}$$

for $1 \leq \alpha \leq k$ and $0 \leq \beta \leq N - 1$. In particular, G is an HNN-extension of the subgroup generated by B with finitely generated associated subgroups. \square

Proof of Theorem 1.1. Let G be a group satisfying the assumptions of Theorem 1.1. By Lemma 3.1, G splits as an HNN-extension of a finitely generated group A with finitely generated associated subgroups C, D with a stable letter t .

We first observe that $C \neq A \neq D$. Indeed, assume $A = C$ or $A = D$. Let $\varepsilon: G \rightarrow \mathbb{Z}$ denote the natural homomorphism sending t to a generator of \mathbb{Z} and all elements of A to 0. Let K_n denote the preimage of $n\mathbb{Z}$ under ε . Then $\beta_1^{(2)}(K_n) = n\beta_1^{(2)}(G) \rightarrow \infty$ as $n \rightarrow \infty$. On the other hand, it is straightforward to check that K_n is generated by A and t^n , which yields a uniform upper bound on the number of generators of G . Since the first ℓ^2 -Betti number of any group is bounded above by the number of generators minus 1, we get a contradiction.

Thus $C \neq A \neq D$. By Theorem 2.2, a countable group with positive first ℓ^2 -Betti number cannot contain finitely generated s -normal subgroups of infinite index. In particular, C is not s -normal in G , i.e., there exists $g \in G$ such that $|g^{-1}Cg \cap C| < \infty$. Now applying Theorem 2.3 we conclude that G is acylindrically hyperbolic. \square

Proof of Corollary 1.2. By Theorem 2.1 every finitely presented residually finite group G with $\beta_1^{(2)}(G) > 0$ has a finite index subgroup that surjects onto \mathbb{Z} . Hence we can apply Theorem 1.1. \square

Proof of Corollary 1.3. Let G be a group of deficiency $\text{def}(G) \geq 2$. Recall that $\beta_1^{(2)}(G) \geq \text{def}(G) - 1$. Hence G has positive first ℓ^2 -Betti number. Note also that G obviously surjects onto \mathbb{Z} . It remains to apply Theorem 1.1. \square

Proof of Corollary 1.4. Parts (a)–(d) are well-known for acylindrically hyperbolic groups. For (a) and (c) see Theorem 2.33 and Theorem 2.35 in [4]. Claim (b) is essentially due to Hamenstädt who proved it in different terms (see [6] and the discussion before Theorem 8.3 in [13]). For (d) we refer to [8]. \square

We mention one more corollary. Recall that a group G belongs to the class \mathcal{C}_{reg} if $H_b^2(G, \ell^2(G)) \neq 0$; further G belongs to the class \mathcal{D}_{reg} if G is non-amenable and there exists an unbounded quasi-cocycle $G \rightarrow \ell^2(G)$. These classes were introduced by Monod-Shalom [12] and Thom [16] in their work on rigidity of group actions. Currently no relation between these classes is known, although they seem likely to coincide. For a more detailed discussion we refer to [16].

It is well-known and fairly easy to see that every quasi-cocycle $G \rightarrow \ell^2(G)$ that is not a sum of a true cocycle and a bounded map gives rise to a non-trivial element of $H_b^2(G, \ell^2(G))$. Since $\beta_1^{(2)}(G) > 0$ if and only if G is non-amenable and there exists a non-zero cocycle $G \rightarrow \ell^2(G)$, we obtain the following alternative first proved by Thom [16]: If $G \in \mathcal{D}_{reg}$, then either $G \in \mathcal{C}_{reg}$ or $\beta_1^{(2)}(G) > 0$. From part (a) of the previous corollary and the Lück approximation, we immediately obtain the following.

Corollary 3.2. *Let $G \in \mathcal{D}_{reg}$ be a finitely presented group. Suppose that G surjects onto \mathbb{Z} . Then $G \in \mathcal{C}_{reg}$.*

We conclude with two open questions.

Problem 3.3. *Is every finitely presented group G with positive first ℓ^2 -Betti number acylindrically hyperbolic?*

Note that there exist finitely presented groups with positive first ℓ^2 -Betti numbers that do not even virtually surject on \mathbb{Z} (e.g., one can take $G = Q * Q$, where Q is a finitely presented infinite simple group). Thus there is no way to reduce the general case of this problem to Theorem 1.1.

The second question is motivated by the fact that the use of ℓ^2 -Betti numbers and the Peterson-Thom theorem in our proof is unavoidable, even if we only want to prove our main theorem for groups of deficiency at least 2. It would be interesting to find an alternative (group theoretic) proof in this particular case. The main difficulty here is the following.

Problem 3.4. *Find a group theoretic argument showing that every finitely generated s -normal subgroup of a finitely presented group of deficiency at least 2 is of finite index.*

References

- [1] B. Baumslag, S. Pride, Groups with two more generators than relators, *J. London Math. Soc.* (2) **17** (1978), no. 3, 425–426.
- [2] J. Cheeger, M. Gromov, L_2 -cohomology and group cohomology, *Topology* **25** (1986), no. 2, 189–215.
- [3] I. Chifan, T. Sinclair, B. Udrea, Inner amenability for groups and central sequences in factors, *arXiv:1307.5002*.
- [4] F. Dahmani, V. Guirardel, D. Osin, Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces, *arXiv:1111.7048*; *Memoirs AMS*, to appear.
- [5] D. Gaboriau, Orbit equivalence and measured group theory. Proceedings of the International Congress of Mathematicians. Volume III, 1501–1527, Hindustan Book Agency, New Delhi, 2010.
- [6] U. Hamenstädt, Bounded cohomology and isometry groups of hyperbolic spaces, *J. Eur. Math. Soc.* **10** (2008), no. 2, 315–349.
- [7] M. Hull, Small cancellation in acylindrically hyperbolic groups, *arXiv:1308.4345*.

- [8] M. Hull, D. Osin, Transitivity degrees of countable groups and acylindrical hyperbolicity, prep., 2015.
- [9] W. Lück, L^2 -invariants: theory and applications to geometry and K -theory. A Series of Modern Surveys in Mathematics, 44. Springer-Verlag, Berlin, 2002.
- [10] W. Lück, D. Osin, Approximating the first L^2 -betti number of residually finite groups, *J. Topol. Anal.* **3** (2011), no. 2, 153–160.
- [11] A. Minasyan, D. Osin, Acylindrical hyperbolicity of groups acting on trees, *arXiv:1310.6289*; *Math. Ann.*, to appear.
- [12] N. Monod, Y. Shalom, Orbit equivalence rigidity and bounded cohomology. *Ann. Math.* **164** (2006), no. 3, 825–878.
- [13] D. Osin, Acylindrically hyperbolic groups, *arXiv:1304.1246*; *Trans. AMS*, to appear.
- [14] D. Osin, A. Thom, Normal generation and ℓ^2 -beti numbers of groups, *Math. Ann.*, 355 (2013), no. 4, 1331–1347.
- [15] J. Peterson, A. Thom, Group cocycles and the ring of affiliated operators, *Invent. Math.*, **185**, (2011), no. 3, 561–592.
- [16] A. Thom, Low degree bounded cohomology and L^2 -invariants for negatively curved groups. *Groups Geom. Dyn.* **3** (2009), no. 2, 343–358.

D. Osin: Stevenson Center 1326, Department of Mathematics, Vanderbilt University,
Nashville, TN 37240, USA
email: *denis.osin@gmail.com*