

Single shot simulations of dynamic quantum many-body systems

Kaspar Sakmann¹ and Mark Kasevich¹

¹ *Department of Physics, Stanford University, Stanford, California 94305, USA*

The single-particle density is the most basic quantity that can be calculated from a given many-body wave function. It provides the probability to find a particle at a given position when the average over many realizations of an experiment is taken. However, the outcome of single experimental shots of ultracold atom experiments is determined by the N -particle probability density. This difference can lead to surprising results. For example, independent Bose-Einstein condensates (BECs) with definite particle numbers form interference fringes even though no fringes would be expected based on the single-particle density [1–4]. By drawing random deviates from the N -particle probability density single experimental shots can be simulated from first principles [1, 3, 5]. However, obtaining expressions for the N -particle probability density of realistic time-dependent many-body systems has so far been elusive. Here, we show how single experimental shots of general ultracold bosonic systems can be simulated based on numerical solutions of the many-body Schrödinger equation. We show how full counting distributions of observables involving any number of particles can be obtained and how correlation functions of any order can be evaluated. As examples we show the appearance of interference fringes in interacting independent BECs, fluctuations in the collisions of strongly attractive BECs, the appearance of randomly fluctuating vortices in rotating systems and the center of mass fluctuations of attractive BECs in a harmonic trap. The method described is broadly applicable to bosonic many-body systems whose phenomenology is driven by information beyond what is typically available in low-order correlation functions.

Let us briefly outline how single experimental shots can be simulated from a general many-body wave function Ψ . The probability to find N particles at positions $\mathbf{r}_1, \dots, \mathbf{r}_N$ in a many-body system is determined by the N -particle probability distribution $P(\mathbf{r}_1, \dots, \mathbf{r}_N) = |\Psi(\mathbf{r}_1, \dots, \mathbf{r}_N)|^2$. In experiments on ultracold bosons snapshots of the positions of the particles are taken and single experimental shots sample the N -particle probability distribution. This distribution is high-dimensional and sampling it directly from a given N -boson wave function is hopeless. However, it can be rewritten as a product of conditional probabilities

$$P(\mathbf{r}_1, \dots, \mathbf{r}_N) = P(\mathbf{r}_1)P(\mathbf{r}_2|\mathbf{r}_1) \times \dots \times P(\mathbf{r}_N|\mathbf{r}_{N-1}, \dots, \mathbf{r}_1), \quad (1)$$

where e.g. $P(\mathbf{r}_2|\mathbf{r}_1)$ denotes the conditional probability to find a particle at \mathbf{r}_2 given that another particle is at \mathbf{r}_1 . By drawing \mathbf{r}_1 from $P(\mathbf{r})$, \mathbf{r}_2 from $P(\mathbf{r}|\mathbf{r}_1)$, \mathbf{r}_3 from $P(\mathbf{r}|\mathbf{r}_2, \mathbf{r}_1)$, etc., one random deviate of $P(\mathbf{r}_1, \dots, \mathbf{r}_N)$ is generated. Obtaining the conditional probabilities in (1) is a formidable combinatorial problem though, even for special cases [1, 5]. Here, we provide a general algorithm to simulate single shots from any given N -boson wave function $|\Psi\rangle = \sum_{\vec{n}} C_{\vec{n}} |\vec{n}\rangle$, where $|\vec{n}\rangle = |n_1, \dots, n_M\rangle$ are configurations constructed by distributing N bosons over M orbitals ϕ_i . We apply this algorithm to many-body states obtained by solving the time-dependent many-body Schrödinger equation numerically using the multiconfigurational time-dependent Hartree for bosons method (MCTDHB) [6–8]. This combination of many-body Schrödinger dynamics and sampling of the N -particle probability allows us to simulate single experimental shots from first principles in realistic settings, see

Methods for the algorithm and details.




Figure 1. Interference of independent interacting condensates. Two independent, repulsively interacting condensates collide in an elongated trap. Shown is the single-particle density (left column) and random deviates of the N -particle density (right column) at different times. In the overlap region interference fringes show up in the N -particle density, but not in the single-particle density. The results are obtained by solving the many-body Schrödinger equation in two spatial dimensions. Parameter values: $N = 10000$ bosons. Interaction strength $\lambda = 4.95$. See text for details. All quantities shown are dimensionless.

It is instructive to briefly review Bose-Einstein condensation. A many-boson state is condensed if its reduced single-particle density matrix has exactly one nonzero eigenvalue ρ_i of order N [9]. The eigenvalues ρ_i are

known as natural occupations, the eigenvectors as natural orbitals. The BEC is fragmented if more than one eigenvalue ρ_i is of order N [10, 11], see Methods for details. Fully condensed states, i.e. states with $\rho_1 = N$, are of the form $\phi(\mathbf{r}_1)\phi(\mathbf{r}_2) \times \dots \times \phi(\mathbf{r}_N)$. (1) then becomes a trivial product of independent, identical probability distributions, and there are no correlations between particles. For instance, Gross-Pitaevskii (GP) mean-field states are of this form. Any other state, in particular fragmented states, exhibit correlations and vice versa any correlated state is to some degree fragmented. We will now show how fragmented BECs lead to macroscopically fluctuating outcomes in single shots.

In the following we use dimensionless units $\hbar = m = 1$ and solve the time-dependent many-body Schrödinger equation $i\frac{\partial}{\partial t}|\Psi\rangle = \hat{H}|\Psi\rangle$ using the MCTDHB method [6–8]. Here,

$$H = \sum_{i=1}^N -\frac{1}{2} \frac{\partial^2}{\partial \mathbf{r}_i^2} + V(\mathbf{r}_i) + \lambda_0 \sum_{i<j} \delta_\epsilon(\mathbf{r}_i - \mathbf{r}_j) \quad (2)$$

denotes a general many-body Hamiltonian in D dimensions with an external potential $V(\mathbf{r})$ and a regularized contact interaction $\delta_\epsilon(\mathbf{r}) = (2\pi\epsilon^2)^{-D/2} e^{-\mathbf{r}^2/2\epsilon^2}$. We parameterize the interaction strength by the mean-field parameter $\lambda = \lambda_0(N-1)$, see Methods for details.

Let us begin with an example of two interfering, independent condensates of $N = 10000$ bosons in an elongated trap with tight harmonic confinement along the z direction such that we can work in $D = 2$ dimensions and $\mathbf{r} = (x, y)$. We use $V(\mathbf{r}) = V_x(x) + V_y(y) + V_g(x)$ as an external potential, where $V_x(x)$ and $V_y(y)$ are harmonic traps and $V_g(x)$ is an additional Gaussian potential that flattens the bottom of the trap along the x -direction. As an initial state we use two independent condensates, each of which is the mean-field ground state (corresponding to $M = 1$ in the MCTDHB formalism) of $N/2$ bosons of the displaced traps $V_\pm(\mathbf{r}) = V(x \pm d, y)$ with $d = 18.6$ in harmonic oscillator units of the y -direction at an interaction strength $\lambda = 4.95$. The initial state $|\Psi(0)\rangle = |N/2, N/2\rangle$ is fragmented with $\rho_1 = \rho_2 = N/2$. We then solve the time-dependent many-body Schrödinger equation for $|\Psi(0)\rangle$ using $M = 2$ orbitals. Fig. 1 shows the single-particle density as well as random deviates of the N -particle density at different times. The two condensates accelerate towards each other, collide and separate again. During the collision interference fringes appear in deviates of the N -particle density at locations that fluctuate randomly from shot to shot, but not in the single-particle density. This is also expected based on simplified models [1, 3]. However, here this result follows directly from the solution of the many-body Schrödinger equation. The interparticle interaction is weak here; interaction effects only become visible as ripples in the density after the collision and the natural occupations remain practically constant all along.

We now go one step further and investigate collisions between strongly attractive independent condensates in




Figure 2. Collision of independent attractively interacting condensates. Two independent attractively interacting condensates collide in an elongated trap in two spatial dimensions. (a) Single-particle density at different times. The condensates approach each other without spreading and bounce off one another. (b) Random deviates of the N -particle density at the time of the collision. Correlations lead to either a single strongly localized density maximum containing practically all particles or two smaller maxima containing about half the particles each. (c) Fragmentation of the condensate as a function of time. The initial state is two fold fragmented with $\rho_1/N = \rho_N/N = 49.4\%$. During the collision two additional natural occupations become significantly occupied and the system can no longer be separated into two independent condensates. Parameter values: $N = 100$ bosons. Interaction strength $\lambda = -5.94$. See text for details. All quantities shown are dimensionless.

the same trap. For this purpose we use $N = 100$ bosons at an interaction strength $\lambda = -5.94$ which is about 2% above the threshold for collapse of the GP mean-field ground state in this trap. For the initial state we first compute the many-body ground state of fifty bosons using two orbitals and imaginary time-propagation. This ground state is highly condensed, $\rho_1/N = 98.7\%$. The initial state is then taken as the symmetrized product of the ground state and a displaced copy of it located at $\mathbf{r} = (-d, 0)$. Thus, the initial state has natural occupations $\rho_1/N = \rho_2/N = 49.4\%$ and $\rho_3/N = \rho_4/N = 0.6\%$. We then propagate this initial state using $M = 4$ orbitals.

Fig. 2 (a) shows the single particle-density at different times. The condensates approach each other without spreading significantly, collide and separate again. During the collision the single-particle density exhibits two maxima, the condensates seem to bounce off each other. However, single shots at the time of the collision reveal a different result, see Fig. 2 (b). In about half of all shots a strongly localized density maximum is visible, whereas in the other half two smaller well separated maxima appear. We stress that at no point any type of (possibly random) phase relationship between the colliding parts was assumed. In fact, for independent condensates the assumption of a preexisting, but random relative phase




Figure 3. Fluctuating vortices. A repulsive condensate in the ground state of a harmonic trap is stirred by a rotating potential in two spatial dimensions. Over the course of time the system fragments and vortices appear at random positions in single shots. (a) First column: single-particle density at different times. Second to fourth column: single shots at the same times. (b) Fragmentation of the condensate as a function of time. Starting from a condensed state, the system of bosons fragments as it is stirred. While the system is condensed single shots and the single-particle density look alike. When the system is fragmented vortices appear at random positions. Parameter values: $N = 10000$. Interaction strength: $\lambda = 17$. See text for details. All quantities shown are dimensionless.

is at variance with quantum mechanics [12]. The macroscopic fluctuations in the outcomes follow directly from the intrinsic correlations of the many-body state. Fig. 2 (c) shows the natural occupations of the system. As long as the condensates are far apart, the natural occupations remain close to their initial values. However, during the collision two additional natural orbitals become occupied indicating a buildup of even stronger correlations. As a consequence after the collision the system can no longer be separated into two independent condensates.

In the previous two examples already the initial states were fragmented. We now turn to a system where fragmentation builds up dynamically. Stirring a BEC can lead to fragmentation and vortex nucleation that cannot be explained within the mean-field framework of quantized vortices [5, 13]. Consider the ground state of a repulsively interacting BEC of $N = 10000$ bosons in a pancake shaped trap with $\omega_x = \omega_y = 1$ at an interaction strength $\lambda = 17$. We compute the many-body ground state using $M = 2$ orbitals which is practically fully condensed with $\rho_1/N = 99.98\%$. We then switch on a time-dependent stirring potential $V_s(\mathbf{r}, t) = \frac{1}{2}\eta(t)[x(t)^2 - y(t)^2]$ that imparts angular momentum onto the BEC. Here $x(t)$ and $y(t)$ vary harmonically and the amplitude $\eta(t)$ is linearly ramped up from zero to a finite value and back down, see Methods for details. Fig. 3 (a) shows the density together with single shots at different times. The evolution of the natural occupations is shown in Fig. 3




Figure 4. Full counting distribution of the center of mass operator. Shown are 10000 random deviates of the center of mass operator of the ground state of an attractively interacting condensate in one spatial dimension. The center of mass fluctuations of the mean-field result (blue) are significantly smaller than those of the many-body results where the bosons are allowed to occupy $M = 2, 3, 10$ (green, magenta, red) orbitals. The $M = 10$ result coincides with the exact analytical one (black). Parameter values: $N = 10$ bosons; interaction strength $\lambda = -0.423$, trap frequency $\omega_x = 1/100$. All quantities shown are dimensionless.

(b). While the system is condensed, single shots reproduce the single-particle density. Over the course of time an additional natural orbital becomes occupied and the BEC becomes correlated. As correlations build up the outcome of single shots fluctuates more and more and vortices appear at random locations in every single shot. This is in stark contrast to mean-field theory, where due to the lack of correlations vortices always appear at the same location.

As a last example let us show how full distribution functions of N -body operators can be evaluated by simulating single shots. Consider the ground state of N attractively interacting bosons in a harmonic trap, $\omega_x = 1/100$, in one dimension, i.e. $D = 1$ and $\mathbf{r} = x$. The exact wave function of the center of mass coordinate $X = \frac{1}{N} \sum_i x_i$ of the many-body ground state is given by a Gaussian $\Psi_{mb}(X) = (\sqrt{\pi}X_{mb})^{-1/2}e^{-X^2/2X_{mb}^2}$ with $X_{mb} = 1/\sqrt{N\omega_x}$ [14]. On the other hand, the mean-field ground state is uncorrelated and hence its center of mass width is given by $X_{mf} = \sigma_{mf}/\sqrt{N}$, where $\sigma_{mf}^2 = \langle \phi_{mf}|x^2|\phi_{mf} \rangle$ is the variance of the mean-field orbital ϕ_{mf} , see Methods. In the limit of a weak trap, $\omega_x \rightarrow 0$, the mean-field solution approaches a soliton with $\sigma_{mf} = \pi/(\sqrt{3}|\lambda|)$. Thus, for sufficiently strong attractive interaction X_{mb} exceeds X_{mf} . We compute the ground state of $N = 10$ bosons at an interaction strength $\lambda = -0.423$ using imaginary time-propagation for different numbers of orbitals. From the obtained ground states we generate 10000 random deviates of the center of mass coordinate. Fig. 4 shows fits to the obtained histograms

of the center of mass deviates together with the exact center of mass distribution. The many-body result for $M = 10$ orbitals is indistinguishable from the exact one and significantly broader than the mean-field ($M = 1$) result. In the present example the many-body correlations are the cause for the onset of the delocalization of the ground state.

METHODS

Bose-Einstein condensation.

For an N -boson state $|\Psi\rangle = \sum_{\vec{n}} C_{\vec{n}}(t)|\vec{n}\rangle$ and a bosonic field operator $\hat{\Psi}(\mathbf{r}) = \sum_j \hat{b}_j \phi_j(\mathbf{r})$ the reduced single-particle density matrix is defined as

$$\rho^{(1)}(\mathbf{r}|\mathbf{r}') = \langle \Psi | \hat{\Psi}^\dagger(\mathbf{r}') \hat{\Psi}(\mathbf{r}) | \Psi \rangle = \sum_{i,j} \rho_{ij} \phi_i^*(\mathbf{r}') \phi_j(\mathbf{r}) \quad (3)$$

with $\rho_{ij} = \langle \Psi | \hat{b}_i^\dagger \hat{b}_j | \Psi \rangle$. By diagonalizing ρ_{ij} one obtains $\rho^{(1)}(\mathbf{r}|\mathbf{r}') = \sum_i \rho_i \phi_i^{NO}(\mathbf{r}) \phi_i^{NO*}(\mathbf{r}')$. The eigenvalues $\rho_1 \geq \rho_2 \geq \dots$ are known as natural occupations, the eigenvectors $\phi_i^{NO}(\mathbf{r})$ as natural orbitals. If there is only one eigenvalue $\rho_1 = \mathcal{O}(N)$ the BEC is condensed [9], if more than one $\rho_i = \mathcal{O}(N)$ the BEC is fragmented [10, 11]. The diagonal $\rho(\mathbf{r}) \equiv \rho^{(1)}(\mathbf{r}|\mathbf{r}' = \mathbf{r})$ is the single-particle density of the N -boson wave function.

Single Shot Algorithm.

Here we show how single shots can be simulated from a general N -boson wave function expanded in M orbitals $|\Psi\rangle = \sum_{\vec{n}} C_{\vec{n}}|\vec{n}\rangle$, where $|\vec{n}\rangle = |n_1, \dots, n_M\rangle$ and $\sum_{i=1}^M n_i = N$. Special cases (for $M = 2$) have been treated in earlier works [1, 5]. The goal is to draw the positions $\mathbf{r}_1, \dots, \mathbf{r}_N$ of N bosons from the probability distribution $P(\mathbf{r}_1, \dots, \mathbf{r}_N)$. We achieve this by evaluating the conditional probabilities in (1). For this purpose we define reduced wave functions

$$|\Psi^{(k)}\rangle = \begin{cases} |\Psi\rangle, & \text{if } k = 0 \\ \mathcal{N}_k \hat{\Psi}(\mathbf{r}_k) |\Psi^{(k-1)}\rangle, & \text{if } k = 1, \dots, N-1 \end{cases} \quad (4)$$

of $n = N - k$ bosons with normalization constants \mathcal{N}_k . The respective single-particle densities are given by $\rho_k(\mathbf{r}) = \langle \Psi^{(k)} | \hat{\Psi}^\dagger(\mathbf{r}) \hat{\Psi}(\mathbf{r}) | \Psi^{(k)} \rangle$ and $\mathcal{N}_k = \rho_{k-1}(\mathbf{r}_k)^{-1/2}$. The first position \mathbf{r}_1 is drawn from $P(\mathbf{r}) = \rho_0(\mathbf{r})/N$. Assuming that positions $\mathbf{r}_k, \dots, \mathbf{r}_1$ have already been drawn, the conditional probability density for the next particle $P(\mathbf{r}|\mathbf{r}_k, \dots, \mathbf{r}_1) = P(\mathbf{r}, \mathbf{r}_k, \dots, \mathbf{r}_1)/P(\mathbf{r}_k, \dots, \mathbf{r}_1)$ is given by

$$P(\mathbf{r}|\mathbf{r}_k, \dots, \mathbf{r}_1) \propto \rho_k(\mathbf{r}), \quad (5)$$

since $P(\mathbf{r}_k, \dots, \mathbf{r}_1)$ is a constant. The problem is thus reduced to obtaining the wave function $|\Psi^{(k)}\rangle =$

$\sum_{\vec{n}} C_{\vec{n}}^{(k)} |\vec{n}\rangle$ from the wave function $|\Psi^{(k-1)}\rangle = \sum_{\vec{n}} C_{\vec{n}}^{(k-1)} |\vec{n}\rangle$, where the sums over run over all configurations of n and $n+1$ bosons, respectively. Defining $\vec{n}^q = (n_1, \dots, n_q + 1, \dots, n_M)$ one finds from (4)

$$C_{\vec{n}}^{(k)} = \mathcal{N}_k \sum_{q=1}^M \phi_q(\mathbf{r}) C_{\vec{n}^q}^{(k-1)} \sqrt{n_q + 1} \quad (6)$$

Using (6) in a general M orbital algorithm requires an ordering of the $\binom{n+M-1}{n}$ configurations $|\vec{n}\rangle$ for all particle numbers $n = 1, \dots, N$. Combinadics [7] provide such an ordering by associating the index

$$J(n_1, \dots, n_M) = 1 + \sum_{i=1}^{M-1} \binom{n+M-1-i-\sum_{j=1}^i n_j}{M-i} \quad (7)$$

with each configuration $|\vec{n}\rangle$. Using (7) all coefficients $C_{\vec{n}}^{(k)}$ can then be obtained by evaluating the sums in (6) and \mathcal{N}_k is determined by normalization. Using the coefficients $C_{\vec{n}}^{(k)}$ we evaluate $\rho_k(\mathbf{r})$ and by means of (5) we then draw \mathbf{r}_{k+1} from $P(\mathbf{r}|\mathbf{r}_k, \dots, \mathbf{r}_1)$. This concludes the algorithm to simulate single shots. It is now easy to see that also correlation functions of arbitrary order can be evaluated. By realizing that

$$\langle \Psi | \hat{\Psi}^\dagger(\mathbf{r}_1) \dots \hat{\Psi}^\dagger(\mathbf{r}_k) \hat{\Psi}(\mathbf{r}_k) \dots \hat{\Psi}(\mathbf{r}_1) | \Psi \rangle = \prod_{j=1}^k \rho_{j-1}(\mathbf{r}_j) \quad (8)$$

the k -th order correlation function is evaluated at $\mathbf{r}_1, \dots, \mathbf{r}_k$ as the product of the reduced densities $\rho_{j-1}(\mathbf{r}_j)$. To evaluate the correlation function $\langle \Psi | \hat{\Psi}^\dagger(\mathbf{r}_1) \dots \hat{\Psi}^\dagger(\mathbf{r}_k) \hat{\Psi}(\mathbf{r}_k) \dots \hat{\Psi}(\mathbf{r}_1) | \Psi \rangle$ the only modification to the single shot algorithm above consists in choosing the positions $\mathbf{r}_1, \dots, \mathbf{r}_k$ rather than drawing them randomly.

MCTDHB.

In the MCTDHB [6–8] method the many-boson wave function is expanded in all configurations that can be constructed by distributing N bosons over M time-dependent orbitals $\phi_i(\mathbf{r}, t)$. The ansatz for the time-dependent many-boson wave function reads:

$$|\Psi(t)\rangle = \sum_{\vec{n}} C_{\vec{n}}(t) |\vec{n}; t\rangle \quad (9)$$

In (9) the $C_{\vec{n}}(t)$ are time-dependent expansion coefficients and the $|\vec{n}; t\rangle$ are time-dependent permanents built from the orbitals $\phi_i(\mathbf{r}, t)$. The MCTDHB equations of motion are derived by requiring stationarity of the many-body Schrödinger action functional

$$S[\{C_{\vec{n}}(t)\}, \{\phi_j(x, t)\}] = \int dt \{ \langle \Psi(t) | H - i \frac{\partial}{\partial t} | \Psi(t) \rangle - \sum_{k,j=1}^M \mu_{kj}(t) [\langle \phi_k | \phi_j \rangle - \delta_{kj}] \}, \quad (10)$$

with respect to variations of the coefficients and the orbitals. The $\mu_{kj}(t)$ are time-dependent Lagrange multipliers that ensure the orthonormality of the orbitals. For bosons interacting via a delta-function interaction and $M = 1$ the MCTDHB equations of motion reduce to the time-dependent Gross-Pitaevskii equation. For more information see the literature [6–8].

Parameters.

For the $D = 2$ dimensional simulations in this work we assume tight harmonic confinement with a frequency ω_z and a harmonic oscillator length $l_z = \sqrt{\hbar/(m\omega_z)}$ along the z -direction. The bosons interact via a regularized contact interaction potential $\frac{\hbar^2\lambda_0}{m}\delta_\epsilon(\mathbf{r})$, with $\delta_\epsilon(\mathbf{r}) = (2\pi\epsilon^2)^{-1}e^{-\mathbf{r}^2/2\epsilon^2}$ and a dimensionless interaction strength $\lambda_0 = \sqrt{8\pi}a/l_z$, where a is the scattering length and m the mass of boson. We note that it is important to regularize contact interaction potentials for $D > 1$ [15, 16]. The contributions to the external potential are given by $V_x(x) = \frac{1}{2}m\omega_x^2x^2$, $V_y(y) = \frac{1}{2}m\omega_y^2y^2$, and $V_g(x) = Ce^{-x^2/2\sigma^2}$, with $C = m\sigma^2\omega_x^2$. We obtain dimensionless units $\hbar = m = 1$ and the Hamiltonian (2) by measuring energy in units of $\hbar\omega_y$, length in units of $l_y = \sqrt{\hbar/(m\omega_y)}$ and time in units of $1/\omega_y$. We use a plane wave discrete variable representation to represent all orbitals and operators. The width of the contact interaction is $\epsilon = 0.15$ and the grid spacing is $\Delta x = \Delta y = \epsilon/2$ for all simulations in this work. For the elongated trap the parameter values are $\omega_x = 0.07$, $\omega_y = 1$ and $\sigma = 10$

on a grid $[-43.2, 43.2] \times [-3.6, 3.6]$. For the rotating BEC the parameter values are $\omega_x = \omega_y = 1$ and $\eta(t)$ is linearly ramped up from zero to $\eta_{max} = 0.1$ over a time span $t_r = 80$. $\eta(t)$ is then kept constant for $t_{up} = 220$ and ramped back down to zero over a time span t_r . The potential $V_s(\mathbf{r}, t) = \frac{1}{2}\eta(t)[x(t)^2 - y(t)^2]$ rotates harmonically with $x(t) = x \cos(\Omega t) + y \sin(\Omega t)$ and $y(t) = -x \sin(\Omega t) + y \cos(\Omega t)$, where $\Omega = \pi/4$. The grid size is $[-8, 8] \times [-8, 8]$.

For the $D = 1$ dimensional simulations we assume tight harmonic confinement along the y - and z -directions with a radial frequency $\omega_\perp = \omega_y = \omega_z$ and an oscillator length $l_\perp = \sqrt{\hbar/(m\omega_\perp)}$. The contact interaction potential is then given by $\frac{2\hbar^2a}{ml_\perp}\delta_\epsilon(x)$, with $\delta_\epsilon(x) = (2\pi\epsilon^2)^{-1/2}e^{-x^2/2\epsilon^2}$. We use $\hbar\omega_\perp$ as the unit of energy and l_\perp as the unit length. The dimensionless interaction strength is then given by $\lambda_0 = 2a/l_\perp$. The harmonic potential along the x -direction $\omega_x = 1/100$ is much weaker than the radial confinement $\omega_\perp = 1$. The grid size is $[-90, 90]$. The Gross-Pitaevskii soliton solution on an infinite line takes on the form $\phi_{mf}(x) = \sqrt{\lambda/4} \text{sech}(\lambda x/2)$.

A. Image processing.

The histograms of the positions of particles obtained using the single shot algorithm have a resolution that is determined by the grid spacing. For better visibility and in analogy to a realistic imaging system we convoluted the data points of each histogram with a point-spread function (PSF). As a PSF we used a Gaussian of width 3×3 pixels.

-
- [1] Javanainen, J. & Yoo, S. M. Quantum Phase of a Bose-Einstein Condensate with an Arbitrary Number of Atoms. *Phys. Rev. Lett.* **76**, 161 (1996)
 - [2] Andrews, M. R. *et al.* Observation of Interference Between Two Bose Condensates. *Science* **275**, 637 (1997)
 - [3] Castin, Y. & Dalibard, J. Relative phase of two Bose-Einstein condensates. *Phys. Rev. A* **55**, 4330 (1997)
 - [4] Hofferberth, S. Lesanovsky, I. Fischer, B. Verdu, J. & Schmiedmayer J. Radiofrequency-dressed-state potentials for neutral atoms. *Nat. Phys.* **2**, 710 (2006)
 - [5] Dagnino, D. Barberán N. & Lewenstein, M. Vortex nucleation in a mesoscopic Bose superfluid and breaking of the parity symmetry. *Phys. Rev. A* **80**, 053611 (2009)
 - [6] Alon, O. E. Streltsov, A. I. & Cederbaum, L. S. Multiconfigurational time-dependent Hartree method for bosons: Many-body dynamics of bosonic systems. *Phys. Rev. A* **77**, 033613 (2008)
 - [7] Streltsov, A. I. Alon, O. E. & Cederbaum, L. S. General mapping for bosonic and fermionic operators in Fock space. *Phys. Rev. A* **81**, 022124 (2010)
 - [8] Streltsov, A. I. Sakmann, K. Lode, A. U. J. Alon, O. E. & Cederbaum L. S., *The Multiconfigurational Time-Dependent Hartree for Bosons Package*, version 2.3, Heidelberg (2013)
 - [9] Penrose, O. & Onsager, L. Bose-Einstein Condensation and Liquid Helium. *Phys. Rev.* **104**, 576 (1956)
 - [10] Nozières, P. & James D. S. Particle vs. pair condensation in attractive Bose-liquids. *J. Phys. France* **43** 1133 (1982)
 - [11] Streltsov, A. I. Alon, O. E. & Cederbaum, L. S. General variational many-body theory with complete self-consistency for trapped bosonic systems. *Phys. Rev. A* **73**, 063626 (2006)
 - [12] Mullin, W. J. & Laloë, F. Interference of Bose-Einstein condensates: Quantum nonlocal effects. *Phys. Rev. A* **78**, 061605(R) (2008).
 - [13] Weiner, S. E. Tsatsos, M. C. Cederbaum, L. S. & Lode, A. U. J. Angular momentum in interacting many-body systems hides in phantom vortices. *arXiv:1409.7670* (2014)
 - [14] Brey, L. Johnson, N. F. & Halperin, B. I. Optical and magneto-optical absorption in parabolic quantum wells. *Phys. Rev. B* **40**, 10647 (1989)
 - [15] Esry, B. D. & Greene, C. H. Validity of the shape-independent approximation for Bose-Einstein condensates. *Phys. Rev. A* **60**, 1451 (1999)
 - [16] Doganov, R. A. Klaiman, S. Alon, O. E. Streltsov, A. I. & Cederbaum, L. S. Two trapped particles interacting by a

finite-range two-body potential in two spatial dimensions. Phys. Rev. A **87**, 033631 (2013)

ACKNOWLEDGEMENTS

Financial support through the Karel Urbanek Postdoctoral Research Fellowship is gratefully acknowledged by K. S. Computing time was provided by the High Performance Computing Center (HLRS) in Stuttgart, Germany.

CONTRIBUTIONS

K. S. and M. K. conceived the ideas and designed the study. K. S. developed the algorithm and carried out the

simulations. K. S. and M. K. wrote the paper.

COMPETING INTERESTS

The authors declare that they have no competing financial interests.

CORRESPONDING AUTHOR

Correspondence and requests for materials should be addressed to Kaspar Sakmann (email: kaspar.sakmann@gmail.com)