

SYMMETRIES, HOPF FIBRATIONS AND SUPERCRITICAL ELLIPTIC PROBLEMS

MÓNICA CLAPP AND ANGELA PISTOIA

ABSTRACT. We consider the semilinear elliptic boundary value problem

$$-\Delta u = |u|^{p-2} u \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

in a bounded smooth domain Ω of \mathbb{R}^N for supercritical exponents $p > \frac{2N}{N-2}$.

Until recently, only few existence results were known. An approach which has been successfully applied to study this problem, consists in reducing it to a more general critical or subcritical problem, either by considering rotational symmetries, or by means of maps which preserve the Laplace operator, or by a combination of both.

The aim of this paper is to illustrate this approach by presenting a selection of recent results where it is used to establish existence and multiplicity or to study the concentration behavior of solutions at supercritical exponents.

KEY WORDS: Nonlinear elliptic boundary value problem; supercritical nonlinearity; nonautonomous critical problem.

2010 MSC: 35J61 (35J20, 35J25).

1. INTRODUCTION

Consider the model problem

$$(\wp_p) \quad \begin{cases} -\Delta u = |u|^{p-2} u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Δ is the Laplace operator, Ω is a bounded domain in \mathbb{R}^N with smooth boundary, $N \geq 3$, and $p > 2$.

Despite its simple form, this problem has been an amazing source of open problems, and the process of understanding it has helped develop new and interesting techniques which can be applied to a wide variety of problems.

The behavior of this problem depends strongly on the exponent p . It is called subcritical, critical or supercritical depending on whether $p \in (2, 2_N^*)$, $p = 2_N^*$ or $p \in (2_N^*, \infty)$, where $2_N^* := \frac{2N}{N-2}$ is the so-called critical Sobolev exponent.

In the subcritical case, standard variational methods yield the existence of a positive solution and infinitely many sign changing solutions. But if p is critical or supercritical the existence of solutions becomes a delicate issue. It depends on the domain. An identity obtained by Pohozaev [26] implies that (\wp_p) does not have a nontrivial solution if Ω is strictly starshaped and $p \in [2_N^*, \infty)$. On the other hand, Kazdan and Warner [17] showed that infinitely many radial solutions exist for every $p \in (2, \infty)$ if Ω is an annulus.

Date: March 1, 2022.

Research supported by CONACYT grant 129847 and PAPIIT grant IN106612 (Mexico) and Università degli Studi di Roma "La Sapienza" Accordi Bilaterali "Esistenza e proprietà geometriche di soluzioni di equazioni ellittiche non lineari" (Italy).

The critical problem has received much attention during the last thirty years, partly due to the fact that it is a simple model for equations which arise in some fundamental questions in differential geometry, like the Yamabe problem or the prescribed scalar curvature problem. Still, many questions remain open in this case.

Until quite recently, only few existence results were known for $p \in (2_N^*, \infty)$. A fruitful approach which has been applied in recent years to treat supercritical problems consists in reducing problem (φ_p) to a more general elliptic critical or subcritical problem, either by considering rotational symmetries, or by means of maps which preserve the Laplace operator, or by a combination of both.

The aim of this paper is to illustrate this approach by presenting a selection of recent results where it is used to establish existence and multiplicity or to study the concentration behavior of solutions at certain supercritical exponents. To put these results into perspective, we first present some nonexistence results.

2. NONEXISTENCE RESULTS

When $p = 2_N^*$ a remarkable result obtained by Bahri and Coron [2] establishes the existence of at least one positive solution to problem (φ_p) in every domain Ω having nontrivial reduced homology with $\mathbb{Z}/2$ -coefficients. Moreover, if Ω is invariant under the action of a closed subgroup G of the group $O(N)$ of linear isometries of \mathbb{R}^N and every G -orbit in Ω is infinite¹, the critical problem is known to have infinitely many solutions [6].

Passaseo showed in [22, 23] that neither of these conditions is enough to guarantee existence in the supercritical case. He proved the following result.

Theorem 2.1. *For each $1 \leq k \leq N - 3$ there is a domain Ω such that*

- (a) Ω has the homotopy type of \mathbb{S}^k ,
- (b) Ω is $O(k+1)$ -invariant with infinite $O(k+1)$ -orbits,
- (c) (φ_p) has no solution for $p \geq 2_{N,k}^* := \frac{2(N-k)}{N-k-2}$,
- (d) (φ_p) has infinitely many solutions for $p < 2_{N,k}^*$.

Here $O(k+1)$ is the group of all linear isometries of \mathbb{R}^{k+1} acting on the first $k+1$ coordinates of a point in \mathbb{R}^N . Note that

$$2_{N,k}^* := \frac{2(N-k)}{N-k-2} = 2_{N-k}^*$$


is the critical Sobolev exponent in dimension $N-k$. It is called the $(k+1)$ -st critical exponent in dimension N . Note also that $2_N^* < 2_{N,1}^* < 2_{N,2}^* < \dots < 2_{N,N-3}^* = 6$, so $2_{N,k}^*$ is supercritical in dimension N if $k \geq 1$.

Passaseo's domains are defined as

$$\Omega := \{(y, z) \in \mathbb{R}^{k+1} \times \mathbb{R}^{N-k-1} : (|y|, z) \in B\},$$

¹Recall that the G -orbit of a point $x \in \mathbb{R}^n$ is the set $Gx := \{gx : g \in G\}$. A subset X of \mathbb{R}^n is said to be G -invariant if $Gx \subset X$ for every $x \in X$, and a function $f : X \rightarrow \mathbb{R}$ is G -invariant if it is constant on every G -orbit of X .

where B is any open ball, centered in $(0, \infty) \times \{0\}$, whose closure is contained in the halfspace $(0, \infty) \times \mathbb{R}^{N-k-1}$.



Passaseo's result was extended to more general domains by Faya and the authors of this paper in [8]. They also showed that existence may fail even in domains with richer topology. More precisely, they proved the following result.

Theorem 2.2. *For every $\varepsilon > 0$ there is a domain Ω such that*


- (a) Ω has the homotopy type of $\mathbb{S}^1 \times \dots \times \mathbb{S}^1$ (k factors),
- (b) Ω is $O(2) \times \dots \times O(2)$ -invariant with infinite orbits,
- (c) (φ_p) does not have a nontrivial solution for $p \geq 2_{N,k}^* + \varepsilon$,
- (d) (φ_p) has infinitely many solutions for $p < 2_{N,k}^*$.

Here \mathbb{S}^1 stands for the unit circle in \mathbb{R}^2 , and the group $O(2) \times \dots \times O(2)$ with k factors acts on $\mathbb{S}^1 \times \dots \times \mathbb{S}^1$ in the obvious way. Note that there are k cohomology classes in $H^1(\Omega; \mathbb{Z})$ whose cup-product is nonzero. In fact, the cup-length of Ω is $k + 1$.

These domains are of the form

$$(2.1) \quad \Omega := \{(y^1, \dots, y^k, z) \in \mathbb{R}^2 \times \dots \times \mathbb{R}^2 \times \mathbb{R}^{N-2k} : (|y^1|, \dots, |y^k|, z) \in B\},$$

where B is an open ball centered in $(0, \infty)^k \times \{0\}$, whose closure is contained in $(0, \infty)^k \times \mathbb{R}^{N-2k}$ and whose radius decreases as $\varepsilon \rightarrow 0$.



The question whether one can get rid of ε remains open.

Problem 1. *Is it true that (φ_p) does not have a nontrivial solution in the domain Ω defined in (2.1) if $p \geq 2_{N,k}^*$ and B is an open ball of arbitrary radius?*

3. SOLUTIONS FOR HIGHER CRITICAL EXPONENTS VIA HOPF MAPS

A fruitful approach to produce solutions to a supercritical problem (φ_p) is to reduce it to some problem of the form

$$(3.1) \quad -\operatorname{div}(a(x)\nabla v) = b(x)|v|^{p-2}v \text{ in } \Theta, \quad v = 0 \text{ on } \partial\Theta,$$

in a bounded smooth domain Θ in \mathbb{R}^n , with $n := \dim \Theta < \dim \Omega = N$ and the same exponent p . Thus, if $p \in (2_N^*, 2_n^*]$, then p is subcritical or critical for problem (3.1) but it is supercritical for (\wp_p) .

3.1. A reduction via Hopf maps. Hopf maps provide a way to obtain such a reduction.

For $N = 2, 4, 8, 16$ we write $\mathbb{R}^N = \mathbb{K} \times \mathbb{K}$, where \mathbb{K} is either the real numbers \mathbb{R} , or the complex numbers \mathbb{C} , or the quaternions \mathbb{H} , or the Cayley numbers \mathbb{O} .

The Hopf map $\mathfrak{h}_{\mathbb{K}} : \mathbb{R}^N = \mathbb{K} \times \mathbb{K} \rightarrow \mathbb{R} \times \mathbb{K} = \mathbb{R}^{(N/2)+1}$ is given by

$$\mathfrak{h}_{\mathbb{K}}(z_1, z_2) = (|z_1|^2 - |z_2|^2, 2\overline{z_1}z_2).$$

Topologically, it is just the quotient map of $\mathbb{K} \times \mathbb{K}$ onto its orbit space under the action of $\mathbb{S}_{\mathbb{K}} := \{\zeta \in \mathbb{K} : |\zeta| = 1\}$ given by multiplication on each coordinate, i.e. $\zeta(y, z) := (\zeta y, \zeta z)$ for $\zeta \in \mathbb{S}_{\mathbb{K}}$, $(y, z) \in \mathbb{K} \times \mathbb{K}$.

But, regarding our problem, the most relevant property of $\mathfrak{h}_{\mathbb{K}}$ is of geometric nature. It is the fact that $\mathfrak{h}_{\mathbb{K}}$ preserves the Laplace operator. Maps with this property are called harmonic morphisms [3, 29]. The following statement can be derived by straightforward computation or from the general theory of harmonic morphisms.

Proposition 3.1. *Let $\Omega \subset \mathbb{K}^2$ be an $\mathbb{S}_{\mathbb{K}}$ -invariant domain such that $0 \notin \overline{\Omega}$. Set $\Theta := \mathfrak{h}_{\mathbb{K}}(\Omega)$. Then u is an $\mathbb{S}_{\mathbb{K}}$ -invariant solution to problem (\wp_p) iff the function v given by $u = v \circ \mathfrak{h}_{\mathbb{K}}$ is a solution to problem*

$$(3.2) \quad -\Delta v = \frac{1}{2|x|} |v|^{p-2} v \quad \text{in } \Theta, \quad v = 0 \quad \text{on } \partial\Theta.$$

Note that, if $\mathbb{K} = \mathbb{C}, \mathbb{H}$ or \mathbb{O} , then $\dim \Theta = \dim \mathbb{K} + 1 < 2 \dim \mathbb{K} = \dim \Omega$. Therefore, $p := 2_{\dim \mathbb{K}+1}^* = 2_{N, \dim \mathbb{K}-1}^*$ is critical for (3.2) and supercritical for (\wp_p) .

Recently Pacella and Srikanth showed that the real Hopf map provides a one-to-one correspondence between $[O(m) \times O(m)]$ -invariant solutions of (\wp_p) in a domain Ω in \mathbb{R}^{2m} and $O(m)$ -invariant solutions of (3.2) in some domain Θ in \mathbb{R}^{m+1} , where $O(m)$ acts on the last m coordinates of $\mathbb{R}^{m+1} \cong \mathbb{R} \times \mathbb{R}^m$. In [21] they proved the following result.

Proposition 3.2. *Let $N = 2m$ and Ω be an $[O(m) \times O(m)]$ -invariant bounded smooth domain in \mathbb{R}^{2m} such that $0 \notin \overline{\Omega}$. Set*

$$\Theta := \{(t, \zeta) \in \mathbb{R} \times \mathbb{R}^m : \mathfrak{h}_{\mathbb{R}}(|y_1|, |y_2|) = (t, |\zeta|) \text{ for some } (y_1, y_2) \in \Omega\}.$$

Then $v(t, \zeta) = w(t, |\zeta|)$ is an $O(m)$ -invariant solution of problem (3.2) iff $u(y_1, y_2) = w(\mathfrak{h}_{\mathbb{R}}(|y_1|, |y_2|))$ is an $[O(m) \times O(m)]$ -invariant solution of problem (\wp_p) .

3.2. Multiplicity results in symmetric domains. The previous propositions suggest to study the critical problem

$$(\wp_b^*) \quad \begin{cases} -\Delta v = b(x) |v|^{2_n^*-2} v & \text{in } \Theta, \\ v = 0 & \text{on } \partial\Theta, \end{cases}$$

in a bounded smooth domain Θ in \mathbb{R}^n , $n \geq 3$, where $b : \overline{\Theta} \rightarrow \mathbb{R}$ is a positive continuous function.

This problem is variational but, due to the lack of compactness of the associated energy functional, classical variational methods cannot be applied to establish existence of solutions.

Under suitable symmetry assumptions compactness is restored: if G is a closed subgroup of the group $O(n)$ of linear isometries of \mathbb{R}^n , Θ and b are G -invariant, and every G -orbit in $\overline{\Theta}$ has infinite cardinality, problem (φ_b^*) is known to have infinitely many G -invariant solutions [6]. This fact, together with Proposition 3.1, provides examples of domains in which problem (φ_p) has infinitely many solutions for some higher critical exponents. For example, one has the following result.

Theorem 3.3. *Let Θ be a solid of revolution around the z -axis in \mathbb{R}^3 whose closure does not intersect the z -axis and set $\Omega := \mathfrak{h}_{\mathbb{C}}^{-1}(\Theta)$. Then the supercritical problem*

$$(\varphi_{2_{4,1}}^*) \quad -\Delta u = |u|^4 u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

has infinitely many solutions which are constant on $\mathfrak{h}_{\mathbb{C}}^{-1}\{(r \cos \vartheta, r \sin \vartheta, t) : \vartheta \in [0, 2\pi]\}$ for each $(r, 0, t) \in \Theta$.

Note that Ω is homeomorphic to $\Theta \times \mathbb{S}^1$. Similar results for problems $(\varphi_{2_{8,3}}^*)$ and $(\varphi_{2_{16,7}}^*)$ can be derived using the Hopf maps $\mathfrak{h}_{\mathbb{H}}$ and $\mathfrak{h}_{\mathbb{O}}$.

On the other hand, if Θ contains a finite orbit, problem (φ_b^*) might not have a nontrivial solution, as occurs when Θ is a ball centered at the origin in \mathbb{R}^n and $b \equiv 1$. Conditions which guarantee that problem (φ_b^*) has a prescribed number of solutions in domains having finite orbits were recently obtained by Faya and the authors. They proved the following result in [8]. Special cases of it were previously established in [7, 10].

Theorem 3.4. *Fix a closed subgroup Γ of $O(n)$ and a bounded smooth Γ -invariant domain D in \mathbb{R}^n such that every Γ -orbit in D has infinite cardinality. Assume that b is Γ -invariant. Then there exists a sequence of real numbers (ℓ_m) with the following property: if $\Theta \supset D$ and Θ is invariant under the action of a subgroup G of Γ such that*


$$(3.3) \quad \min_{x \in \Theta} \frac{\#Gx}{b(x)^{\frac{n-2}{2}}} > \ell_m,$$

then problem (φ_b^) has at least m pairs $\pm u$ of G -invariant solutions in Θ ; one pair does not change sign and the rest are sign changing.*

Here $\#Gx$ denotes the cardinality of the G -orbit of x .

We illustrate this result with an example. Let D be a torus of revolution around the z -axis in \mathbb{R}^3 , Γ be the group of all rotations around the z -axis, and $b(x) := \frac{1}{2|x|}$. Fix $\varepsilon > 0$ smaller than the distance of D to the z -axis. Let G_k be the group generated by the rotation of angle $\frac{2\pi}{k}$. If $k\sqrt{2}\varepsilon > \ell_m$ and Θ is a G_k -invariant domain which contains D whose distance to the z -axis is at least ε , then every G_k -orbit in Θ has cardinality k and the inequality (3.3) holds true. We call a G_k -invariant domain which contains D and does not intersect the z -axis a *k-teething*

toy.



Applying Proposition 3.1 to this example we obtain the following result.

Theorem 3.5. *If Θ is a k -teething toy whose distance to the z -axis is at least ε and $k\sqrt{2\varepsilon} > \ell_m$, then the supercritical problem*

$$(\wp_{2_{4,1}^*}) \quad -\Delta u = |u|^4 u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

has m pairs of \mathbb{S}_C -invariant solutions in $\Omega := \mathfrak{h}_C^{-1}(\Theta)$; one of them does not change sign and the rest are sign changing.

A more general result can be derived from Theorem 3.4, as stated in [8].

3.3. Existence in domains with thin spherical holes. Next we consider problem (\wp_b^*) in a punctured domain

$$\Theta_\varepsilon := \{x \in \Theta : |x - \xi| > \varepsilon\},$$

where Θ is a bounded smooth domain in \mathbb{R}^n , $n \geq 3$, $\xi \in \Theta$, $\varepsilon > 0$ is small, and $b : \overline{\Theta} \rightarrow \mathbb{R}$ is a positive \mathcal{C}^2 -function. Additionally, we assume that Θ and b are invariant under the action of some closed subgroup G of $O(n)$ and that ξ is a fixed point, i.e. $g\xi = \xi$ for all $g \in G$.

If $b \equiv 1$ Coron showed in [12] that problem $(\wp_{2_n^*})$ has a positive solution in Θ_ε for ε small enough. Coron's proof takes advantage of the fact that the variational functional associated to problem $(\wp_{2_n^*})$ satisfies the Palais-Smale condition between the ground state level and twice that level. This is not true anymore when $b \not\equiv 1$, so Coron's argument does not carry over to the nonautonomous case.

A method which has proved to be very successful in dealing with critical problems which involve small perturbations of the domain is the Lyapunov-Schmidt reduction method, see e.g. [16] and the references therein. This method was used by Faya and the authors in [9] to prove the following result.

Theorem 3.6. *If $\nabla b(\xi) \neq 0$ then, for ε small enough,*

$$(\wp_b^*) \quad -\Delta v = b(x) |v|^{2_n^*-2} v \quad \text{in } \Theta_\varepsilon, \quad v = 0 \quad \text{on } \partial\Theta_\varepsilon,$$

has a positive G -invariant solution v_ε in Θ_ε which concentrates at the boundary of the hole and blows up at ξ as $\varepsilon \rightarrow 0$.

Note that G may be the trivial group, so this result is true in a non-symmetric setting and, combined with Proposition 3.1, yields solutions to supercritical problems concentrating around a spherical hole, see [9].

But we may also combine it with Proposition 3.2 as follows: Let $N = 2m$, $m \geq 2$, Ω be an $[O(m) \times O(m)]$ -invariant bounded smooth domain in $\mathbb{R}^N \equiv \mathbb{R}^m \times \mathbb{R}^m$ such that $0 \notin \bar{\Omega}$, and $\xi \in \Omega \cap (\mathbb{R}^m \times \{0\})$. For $\varepsilon > 0$ small enough set

$$\Omega_\varepsilon := \{x \in \Omega : \text{dist}(x, S_\xi) > \varepsilon\},$$

where $S_\xi := \{(x, 0) \in \mathbb{R}^m \times \{0\} : |x| = |\xi|\}$. The following result, obtained by combining Theorem 3.6 with Proposition 3.2, was established in [9].

Theorem 3.7. *For each ε small enough the supercritical problem*

$$(\mathcal{P}_{2m, m-1}^*) \quad -\Delta u = |u|^{4/(m-1)} u \quad \text{in } \Omega_\varepsilon, \quad v = 0 \quad \text{on } \partial\Omega_\varepsilon,$$

has a positive $[O(m) \times O(m)]$ -invariant solution u_ε which concentrates along the set $\{x \in \Omega : \text{dist}(x, S_\xi) = \varepsilon\}$ and blows up at the $(m-1)$ -dimensional sphere S_ξ as $\varepsilon \rightarrow 0$.


4. SOLUTIONS FOR HIGHER CRITICAL EXPONENTS VIA ROTATIONS

Supercritical problems in domains obtained through rotations can be reduced to subcritical or critical problems as follows.

4.1. A reduction via rotations. Fix $k_1, \dots, k_m \in \mathbb{N}$ and set $k := k_1 + \dots + k_m$. If $N \geq k + m$ let

$$\Omega := \{(y^1, \dots, y^m, z) \in \mathbb{R}^{k_1+1} \times \dots \times \mathbb{R}^{k_m+1} \times \mathbb{R}^{N-k-m} : (|y^1|, \dots, |y^m|, z) \in \Theta\},$$

where Θ is a bounded smooth domain in \mathbb{R}^{N-k} whose closure is contained in $(0, \infty)^m \times \mathbb{R}^{N-k-m}$.



Each point $\xi \in \Theta$ gives rise to a subset

$$(4.1) \quad T_\xi := \{(y^1, \dots, y^m, z) \in \Omega : |y^i| = \xi_i, z = (\xi_{m+1}, \dots, \xi_{N-k})\}$$

of Ω which is homeomorphic to the product of spheres $\mathbb{S}^{k_1} \times \dots \times \mathbb{S}^{k_m}$. A straightforward computation yields the following result.

Proposition 4.1. *A function u of the form $u(y^1, \dots, y^m, z) = v(|y^1|, \dots, |y^m|, z)$ is a solution of problem (\mathcal{P}_p) iff v is a solution of*

$$(4.2) \quad -\text{div}(a(x)\nabla v) = a(x)|v|^{p-2}v \quad \text{in } \Theta, \quad v = 0 \quad \text{on } \partial\Theta,$$

with $a(x_1, \dots, x_{N-k}) := x_1^{k_1} \dots x_m^{k_m}$.

4.2. Multiplicity results in domains obtained by rotation. Wei and Yan considered domains as above, with $m = 1$, where Θ is invariant under the action of the group $O(2) \times O(1)^{N-k-3}$ on the last $N - k - 1$ coordinates of \mathbb{R}^{N-k} , i.e.

$$(s, r \cos \theta, r \sin \theta, x_3, \dots, x_{N-k}) \in \Theta \text{ for all } \theta \in (0, 2\pi) \text{ if } (s, r, 0, \dots, x_{N-k}) \in \Theta, \\ (x_1, \dots, -x_i, \dots, x_{N-k}) \in \Theta \text{ if } (x_1, \dots, x_i, \dots, x_{N-k}) \in \Theta \text{ and } i = 4, \dots, N - k.$$

They proved the following result in [28].

Theorem 4.2. *Let $N \geq 5$. Assume that Θ is $[O(2) \times O(1)^{N-k-3}]$ -invariant and that there is a point*

$$(s^*, r^*) \in \mathcal{S} := \{(s, r) \in \mathbb{R}^2 : (s, r, 0, \dots, 0) \in \partial\Theta\},$$

which is a strict local minimum (or a strict local maximum) of the distance of \mathcal{S} to $\{0\} \times \mathbb{R}$. Set $\xi_j := (s^, r^* \cos(2\pi j/\ell), r^* \sin(2\pi j/\ell), 0, \dots, 0)$. Then, for large enough $\ell \in \mathbb{N}$, problem $(\wp_{2_{N,k}^*})$ has a solution u_ℓ with ℓ positive layers; which concentrate along each of the k -dimensional spheres $T_{\xi_j} \subset \partial\Omega$, $j = 0, \dots, \ell - 1$, and blow up at $\partial\Omega$ as $\ell \rightarrow \infty$.*

They derived this result from Proposition 4.1 after proving the existence of ℓ -multibubble solutions for the critical problem

$$(\wp_{a,a}^*) \quad -\operatorname{div}(a(x)\nabla v) = a(x)|v|^{2_{N-k}^*-2}v \text{ in } \Theta, \quad v = 0 \text{ on } \partial\Theta,$$

which concentrate at the points $\xi_j \in \partial\Theta$. A precise description of the solutions is given in [28].

In [19] Kim and Pistoia considered domains with thin k -dimensional holes. More precisely, for some fixed $\xi \in \Theta$, they considered

$$\Omega_\varepsilon := \{x \in \Omega : \operatorname{dist}(x, T_\xi) > \varepsilon\},$$

with T_ξ as in (4.1) and $\varepsilon > 0$ sufficiently small. Note that Ω_ε is obtained by rotating the punctured domain $\Theta_\varepsilon := \{x \in \Theta : |x - \xi| > \varepsilon\}$ as described in subsection 4.1. For $N - k \geq 4$ they proved the existence of towers of bubbles with alternating signs around ξ for the anisotropic problem $(\wp_{a,a}^*)$ in Θ_ε , thus extending a previous result by Ge, Musso and Pistoia [16] for the autonomous problem $a \equiv 1$. The number of bubbles increases as $\varepsilon \rightarrow 0$. Combining this result with Proposition 4.1 they obtained the following one.

Theorem 4.3. *Let $N \geq k + 4$. Then, for every $\ell \in \mathbb{N}$ there exists $\varepsilon_\ell > 0$ such that, for each $\varepsilon \in (0, \varepsilon_\ell)$, the supercritical problem*

$$-\Delta u = |u|^{2_{N,k}^*-2}u \text{ in } \Omega_\varepsilon, \quad v = 0 \text{ on } \partial\Omega_\varepsilon,$$

has a solution u_ε with ℓ layers of alternating signs, which concentrate with different rates along the boundary of the tubular neighborhood of radius ε of T_ξ and blow up at T_ξ as $\varepsilon \rightarrow 0$.

A precise description of the solutions can be found in [19].

5. CONCENTRATION ALONG MANIFOLDS AT THE HIGHER CRITICAL EXPONENTS

5.1. Approaching the higher critical exponents from below. For domains as those described in subsection 4.1 and $p \in (2, 2_{N,k}^*)$ problem (\wp_p) has infinitely many nontrivial solutions of the form $u(y^1, \dots, y^m, z) = v(|y^1|, \dots, |y^m|, z)$. This

follows immediately from Proposition 4.1 using standard variational methods because problem (4.2) is subcritical and the domain Θ is bounded. Existence and nonexistence results in some unbounded domains are also available [11].

So the question is whether one can establish existence of solutions to (\wp_p) which exhibit a certain concentration behavior as $p \rightarrow 2_{N,k}^*$.

For the slightly subcritical problem $(\wp_{2_{N,k}^* - \varepsilon})$ positive and sign changing solutions u_ε which blow up at one or several points in Ω as $\varepsilon \rightarrow 0$ have been obtained e.g. in [4, 25, 27]. Recently, del Pino, Musso and Pacard [15] considered the case in which p approaches the second critical exponent from below. They showed that, if $N \geq 8$ and $\partial\Omega$ contains a nondegenerate closed geodesic Γ with negative inner normal curvature then, for every $\varepsilon > 0$ small enough, away from an explicit discrete set of values, problem $(\wp_{2_{N,1}^* - \varepsilon})$ has a positive solution u_ε which concentrates and blows up at Γ as $\varepsilon \rightarrow 0$.

It is natural to ask whether similar concentration phenomena can be observed as p approaches the $(k+1)$ -st critical exponent $2_{N,k}^*$ from below, i.e. whether there are domains in which problem $(\wp_{2_{N,k}^* - \varepsilon})$ has a solution u_ε which concentrates and blows up at a k -dimensional submanifold of Ω as $\varepsilon \rightarrow 0$. Ackermann, Kim and the authors have given positive answers to this question for domains Ω as in subsection 4.1.

Let \mathcal{K} be the set of all nondegenerate critical points ξ of the restriction of the function $a(x_1, \dots, x_{N-k}) := x_1^{k_1} \dots x_m^{k_m}$ to $\partial\Theta$ such that $\nabla a(\xi)$ points into the interior of Θ , and let T_ξ be the set defined in (4.1). The following result was proved in [1].

Theorem 5.1. *For any subset $\{\xi_1, \dots, \xi_\ell\}$ of \mathcal{K} and $1 \leq m \leq \ell$ there exists $\varepsilon_0 > 0$ such that, for each $\varepsilon \in (0, \varepsilon_0)$, problem $(\wp_{2_{N,k}^* - \varepsilon})$ has a solution u_ε with m positive layers and $\ell - m$ negative layers; which concentrate at the same rate and blows up along one of the sets T_{ξ_i} as $\varepsilon \rightarrow 0$.*

Sign changing solutions are also available. Statement (a) in the following theorem was proved in [1] and statement (b) was proved in [18].

Theorem 5.2. *Assume there exist $\xi_0 \in \mathcal{K}$ and $\tau_1, \dots, \tau_{N-k-1} \in \mathbb{R}^{N-k}$ such that the set $\{\nabla a(\xi_0), \tau_1, \dots, \tau_{N-k-1}\}$ is orthogonal and Θ and a are invariant with respect to the reflection ρ_i on the hyperplane through ξ_0 which is orthogonal to τ_i , for each $i = 1, \dots, N-k-1$. Then the following statements hold true:*

- (a) *For each $\varepsilon > 0$ small enough problem $(\wp_{2_{N,k}^* - \varepsilon})$ has a sign changing solution u_ε with one positive and one negative layer, which concentrate at the same rate along T_{ξ_0} and blow up at T_{ξ_0} as $\varepsilon \rightarrow 0$.*
- (b) *If $k \leq N-4$ then, for any integer $\ell \geq 2$, there exists $\varepsilon_\ell > 0$ such that, for $\varepsilon \in (0, \varepsilon_\ell)$, problem $(\wp_{2_{N,k}^* - \varepsilon})$ has a solution u_ε with ℓ layers of alternating signs which concentrate at different rates along T_{ξ_0} and blow up at T_{ξ_0} as $\varepsilon \rightarrow 0$.*

These results follow from Proposition 4.1 once the corresponding statements for the slightly subcritical problems

$$-\operatorname{div}(a(x)\nabla v) = a(x)|v|^{2_{N-k}^* - \varepsilon - 2}v \quad \text{in } \Theta, \quad v = 0 \quad \text{on } \partial\Theta,$$

have been established. This is done in [1, 18], where a precise description of the solutions is given. The following questions were raised in [1]:

Problem 2. *Can statement (a) in Theorem 5.2 be improved to establish existence of solutions with ℓ layers of alternating signs which concentrate at the same rate along T_{ξ_0} and blow up at T_{ξ_0} as $\varepsilon \rightarrow 0$, for any $\ell \geq 2$?*

Problem 3. *Does Theorem 5.2 hold true without the symmetry assumption?*

Recently, Pacella and Pistoia considered the case in which $N = 2m$ and Ω is the annulus $\Omega := \{x \in \mathbb{R}^N : 0 < a < |x| < b\}$. They used Proposition 3.2 to establish the existence of positive and sign changing solutions to problem $(\wp_{2_{N,k}^* - \varepsilon})$ concentrating along the $(m - 1)$ -dimensional spheres

$$S_1 := \{(x, 0) \in \mathbb{R}^m \times \{0\} : |x| = a\}, \quad S_2 := \{(0, y) \in \{0\} \times \mathbb{R}^m : |y| = a\}$$

as $\varepsilon \rightarrow 0$. Their results can be found in [20].

5.2. Approaching the higher critical exponents from above. For the slightly supercritical problem $(\wp_{2_{N,k}^* + \varepsilon})$, $\varepsilon > 0$, in a domain Ω with nontrivial topology del Pino, Felmer and Musso established the existence of a positive solution u_ε with two bubbles which concentrate at two different points $\xi_1, \xi_2 \in \Omega$ as $\varepsilon \rightarrow 0$ [13]. Solutions with more than two bubbles are also available, see e.g. [4, 14, 24]. For ε sufficiently small solutions with only one bubble do not exist [5].

The following problem is fully open to investigation.

Problem 4. *Are there domains Ω in which $(\wp_{2_{N,k}^* + \varepsilon})$ has positive or sign changing solutions u_ε which concentrate and blow up at k -dimensional manifolds as $\varepsilon \rightarrow 0$?*

REFERENCES

- [1] N. ACKERMANN, M. CLAPP, A. PISTOIA, Boundary clustered layers for some supercritical problems. *J. Differential Equations* **254** (2013), 4168-4183.
- [2] A. BAHRI, J.M. CORON, On a nonlinear elliptic equation involving the critical Sobolev exponent: The effect of the topology of the domain. *Comm. Pure Appl. Math.* **41** (1988), 253-294.
- [3] P. BAIRD, J.C. WOOD, *Harmonic morphisms between Riemannian manifolds*. London Mathematical Society Monographs. New Series, 29. The Clarendon Press, Oxford University Press, Oxford, 2003.
- [4] T. BARTSCH, A.M. MICHELETTI, AND A. PISTOIA, On the existence and the profile of nodal solutions of elliptic equations involving critical growth. *Calc. Var. Partial Differential Equations* **26** (2006), 265-282.
- [5] M. BEN AYED, K. EL MEHDI, O. REY, M. GROSSI, A nonexistence result of single peaked solutions to a supercritical nonlinear problem. *Commun. Contemp. Math.* **5** (2003), 179-195.
- [6] M. CLAPP, A global compactness result for elliptic problems with critical nonlinearity on symmetric domains, in *Nonlinear Equations: Methods, Models and Applications*, 117-126, *Progr. Nonlinear Differential Equations Appl.* **54** Birkhauser, Boston, 2003.
- [7] M. CLAPP, J. FAYA, Multiple solutions to the Bahri-Coron problem in some domains with nontrivial topology, *Proc. Amer. Math. Soc.* **141** (2013), 4339-4344.
- [8] M. CLAPP, J. FAYA, AND A. PISTOIA, Nonexistence and multiplicity of solutions to elliptic problems with supercritical exponents. *Calc. Var. Partial Differential Equations* **48** (2013), 611-623.
- [9] M. CLAPP, J. FAYA, AND A. PISTOIA, Positive solutions to a supercritical elliptic problem which concentrate along a thin spherical hole. *J. Anal. Math.*, to appear.
- [10] M. CLAPP, F. PACELLA, Multiple solutions to the pure critical exponent problem in domains with a hole of arbitrary size, *Math. Z.* **259** (2008), 575-589.
- [11] M. CLAPP, A. SZULKIN, A supercritical elliptic problem in a cylindrical shell. Preprint.
- [12] J.M. CORON, Topologie et cas limite des injections de Sobolev, *C.R. Acad. Sc. Paris* **299**, Ser. I (1984), 209-212.

- [13] M. DEL PINO, P. FELMER, M. MUSSO, Two-bubble solutions in the super-critical Bahri-Coron's problem. *Calc. Var. Partial Differential Equations* **16** (2003), 113–145.
- [14] M. DEL PINO, P. FELMER, M. MUSSO, Multi-bubble solutions for slightly super-critical elliptic problems in domains with symmetries. *Bull. London Math. Soc.* **35** (2003), 513–521.
- [15] M. DEL PINO, M. MUSSO, F. PACARD, Bubbling along boundary geodesics near the second critical exponent. *J. Eur. Math. Soc. (JEMS)* **12** (2010), 1553–1605.
- [16] Y. GE, M. MUSSO, A. PISTOIA, Sign changing tower of bubbles for an elliptic problem at the critical exponent in pierced non-symmetric domains. *Comm. Partial Differential Equations* **35** (2010), 1419–1457.
- [17] J. KAZDAN, F. WARNER, Remarks on some quasilinear elliptic equations. *Comm. Pure Appl. Math.* **38** (1975), 557–569.
- [18] S. KIM, A. PISTOIA, Boundary towers of layers for some supercritical problems. *J. Differential Equations* **255** (2013), 2302–2339.
- [19] S. KIM, A. PISTOIA, Supercritical problems in domains with thin toroidal holes. Preprint.
- [20] F. PACELLA, A. PISTOIA, Bubble concentration on spheres for supercritical elliptic problems. Preprint.
- [21] F. PACELLA, P.N. SRIKANTH, A reduction method for semilinear elliptic equations and solutions concentrating on spheres. Preprint. arXiv:1210.0782.
- [22] D. PASSASEO, Nonexistence results for elliptic problems with supercritical nonlinearity in nontrivial domains. *J. Funct. Anal.* **114** (1993), 97–105.
- [23] D. PASSASEO, New nonexistence results for elliptic equations with supercritical nonlinearity. *Differential Integral Equations* **8** (1995), 577–586.
- [24] A. PISTOIA, O. REY, Multiplicity of solutions to the supercritical Bahri-Coron's problem in pierced domains. *Adv. Differential Equations* **11** (2006), 647–666.
- [25] A. PISTOIA AND T. WETH, Sign changing bubble tower solutions in a slightly subcritical semilinear Dirichlet problem. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **24** (2007), 325–340.
- [26] S.I. POHOZHAEV, Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$. *Soviet Math. Dokl.* **6** (1965), 1408–1411.
- [27] O. REY, Blow-up points of solutions to elliptic equations with limiting nonlinearity. *Differential Integral Equations* **4** (1991), 1155–1167.
- [28] J. WEI, S. YAN, Infinitely many positive solutions for an elliptic problem with critical or supercritical growth. *J. Math. Pures Appl.* **96** (2011), 307–333.
- [29] J.C. WOOD, *Harmonic morphisms between Riemannian manifolds*. Modern trends in geometry and topology, 397–414, Cluj Univ. Press, Cluj-Napoca, 2006.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIRCUITO EXTERIOR, C.U., 04510 MÉXICO D.F., MEXICO
E-mail address: monica.clapp@im.unam.mx

DIPARTIMENTO DI METODI E MODELLI MATEMATICI, UNIVERSITÀ DI ROMA "LA SAPIENZA", VIA ANTONIO SCARPA 16, 00161 ROMA, ITALY
E-mail address: pistoia@dmmm.uniroma1.it