

# ON A SMOOTH COMPACTIFICATION OF $\mathrm{PSL}(n, \mathbb{C})/T$

INDRANIL BISWAS, S. SENTHAMARAI KANNAN, AND D. S. NAGARAJ

**ABSTRACT.** Let  $T$  be a maximal torus of  $\mathrm{PSL}(n, \mathbb{C})$ . For  $n \geq 4$ , we construct a smooth compactification of  $\mathrm{PSL}(n, \mathbb{C})/T$  as a geometric invariant theoretic quotient of the wonderful compactification  $\overline{\mathrm{PSL}(n, \mathbb{C})}$  for a suitable choice of  $T$ -linearized ample line bundle on  $\overline{\mathrm{PSL}(n, \mathbb{C})}$ . We also prove that the connected component, containing the identity element, of the automorphism group of this compactification of  $\mathrm{PSL}(n, \mathbb{C})/T$  is  $\mathrm{PSL}(n, \mathbb{C})$  itself.

## 1. INTRODUCTION

Let  $G$  be a semisimple group of adjoint type over the field  $\mathbb{C}$  of complex numbers. De Concini and Procesi in [DP] constructed a smooth projective variety  $\overline{G}$  with an action of  $G \times G$  such that

- the variety  $G$  equipped with the action of  $G \times G$  given by the left and right translations is an open dense orbit of it, and
- the boundary  $\overline{G} \setminus G$  is a union of  $G \times G$  stable normal crossing divisors.

This variety  $\overline{G}$  is known as the wonderful compactification of  $G$ .

Fix a maximal torus  $T$  of  $G$ . Consider the right action of  $T$  on  $\overline{G}$ , meaning the action of the subgroup  $1 \times T \subset G \times G$ . For a  $T$ -linearized ample line bundle  $\mathcal{L}$  on  $\overline{G}$ , let  $\overline{G}_T^{ss}(\mathcal{L})$  and  $\overline{G}_T^s(\mathcal{L})$  denote respectively the loci of semistable and stable points of  $\overline{G}$  (see [MFK, p. 30, p. 40]).

Our first main result (Proposition 3.3) says that there is a  $T$ -linearized ample line bundle  $\mathcal{L}$  on  $\overline{G}$  such that  $\overline{G}_T^{ss}(\mathcal{L}) = \overline{G}_T^s(\mathcal{L})$ .

For  $G = \mathrm{PSL}(n, \mathbb{C})$ , we show that there is a  $T$ -linearized ample line bundle  $\mathcal{L}$  on  $\overline{\mathrm{PSL}(n, \mathbb{C})}$  such that

- the GIT quotient  $\overline{\mathrm{PSL}(n, \mathbb{C})}_T^{ss}(\mathcal{L})//T$  is smooth, and
- the boundary  $(\overline{\mathrm{PSL}(n, \mathbb{C})}_T^{ss}(\mathcal{L})//T) \setminus (\mathrm{PSL}(n, \mathbb{C})/T)$  is a union of  $\mathrm{PSL}(n, \mathbb{C})$  stable normal crossing divisors.

We further show that for  $n \geq 4$ , the connected component of the automorphism group of  $\overline{\mathrm{PSL}(n, \mathbb{C})}_T^{ss}//T$  containing the identity automorphism is  $\mathrm{PSL}(n, \mathbb{C})$  (Theorem 4.1).

---

2000 *Mathematics Subject Classification.* 14F17.

*Key words and phrases.* Wonderful compactification, GIT quotients, automorphism group, Frobenius splitting.

## 2. PRELIMINARIES AND NOTATION

In this section we recall some preliminaries and notation about Lie algebras and algebraic groups; see for example [Hu1] and [Hu2] for the details. Let  $G$  be a simple group of adjoint type of rank  $n$  over the field of complex numbers. Let  $T$  be a maximal torus of  $G$  and  $B \supset T$  a Borel subgroup of  $G$ . Let  $N_G(T)$  denote the normalizer of  $T$  in  $G$ . So  $W := N_G(T)/T$  is the Weyl group of  $G$  with respect to  $T$ .

The Lie algebra of  $G$  will be denoted by  $\mathfrak{g}$ . Let  $\mathfrak{h} \subset \mathfrak{g}$  be the Lie algebra of  $T$ . The set of roots of  $G$  with respect to  $T$  will be denoted by  $R$ . Let  $R^+ \subset R$  be the set of positive roots with respect to  $B$ . Let

$$S = \{\alpha_1, \alpha_2, \dots, \alpha_n\} \subset R^+$$

be the set of simple roots with respect to  $B$ . The group of characters of  $T$  will be denoted by  $X(T)$ , while the group of one-parameter subgroups of  $T$  will be denoted by  $Y(T)$ . Let

$$\{\lambda_i \mid 1 \leq i \leq n\}$$

be the ordered set of one-parameter subgroups of  $T$  satisfying the condition that  $\langle \alpha_i, \lambda_j \rangle = \delta_{ij}$ , where

$$\langle -, - \rangle : X(T) \times Y(T) \longrightarrow \mathbb{Z}$$

is the natural pairing, and  $\delta_{ij}$  is the Kronecker delta function. Let  $\leq$  (respectively,  $\geq$ ) be the partial order on  $X(T)$  defined as follows:  $\chi_1 \leq \chi_2$ , (respectively,  $\chi_1 \geq \chi_2$ ) if  $\chi_2 - \chi_1$  (respectively,  $\chi_1 - \chi_2$ ) is a linear combination of simple roots with non-negative integers as coefficients.

Let  $(-, -)$  denote the restriction of the Killing form of  $\mathfrak{g}$  to  $\mathfrak{h}$ . Let

$$\{\omega_j \mid 1 \leq j \leq n\}$$

be the ordered set of fundamental weights corresponding to  $S$ , in other words,

$$\frac{2(\omega_i, \alpha_j)}{(\alpha_j, \alpha_j)} = \delta_{ij}, \quad 1 \leq i, j \leq n.$$

For  $1 \leq i \leq n$ , let  $s_{\alpha_i}$  denote the simple reflection corresponding to  $\alpha_i$ .

The longest element of  $W$  corresponding to  $B$  will be denoted by  $w_0$ . Let

$$B^- = w_0 B w_0^{-1}$$

be the Borel subgroup of  $G$  opposite to  $B$  with respect to  $T$ .

For the notion of a  $G$ -linearization, and the GIT quotients, we refer to [MFK, p. 30, p. 40].

Consider the flag variety  $G/B$  that parametrizes all Borel subgroups of  $G$ . For a character  $\chi$  of  $B$ , let

$$L_\chi = G \times_B \mathbb{C} \longrightarrow G/B$$

be the  $G$ -linearized line bundle associated to the action of  $B$  on  $G \times \mathbb{C}$  given by  $b.(g, z) = (gb, \chi(b^{-1})z)$  for  $b \in B$  and  $(g, z) \in G \times \mathbb{C}$ . So, in particular,  $L_\chi$  is  $T$ -linearized. When  $L_\chi$  is

ample, we denote by  $(G/B)_T^{ss}(L_\chi)$  (respectively,  $(G/B)_T^s(L_\chi)$ ) the semistable (respectively, stable) locus in  $G/B$  for the  $T$ -linearized ample line bundle  $L_\chi$ .

Next we recall some facts about the wonderful compactification of  $G$ . Let  $\chi$  be a regular dominant weight of  $G$  with respect to  $T$  and  $B$ , and let  $V(\chi)$  be the irreducible representation of  $\widehat{G}$  with highest weight  $\chi$ , where  $\widehat{G}$  is the simply connected covering of  $G$ . By [DP, p. 16, 3.4], the wonderful compactification  $\overline{G}$ , which we denote by  $X$ , is the closure of the  $G \times G$ -orbit of the point

$$[1] \in \mathbb{P}(V(\chi) \otimes V(\chi)^*)$$

corresponding to the identity element 1 of  $V(\chi) \otimes V(\chi)^* = \mathrm{End}(V(\chi)^*)$ . We denote by  $\mathcal{L}_\chi$  the ample line bundle on  $X$  induced by this projective embedding. Since the regular dominant weights generate the weight lattice, given a weight  $\chi$ , we have the line bundle  $\mathcal{L}_\chi$  on  $X$  associated to  $\chi$ .

By [DP, Theorem, p. 14, Section 3.1], there is a unique closed  $G \times G$ -orbit  $Z$  in  $X$ . Note that

$$Z = \bigcap_{i=1}^n D_i,$$

where  $D_i$  is the  $G \times G$  stable irreducible component of  $\overline{G} \setminus G$  such that  $\mathcal{O}(D_i) = \mathcal{L}_{\alpha_i}$  [DP, p. 29, Section 8.2, Corollary]. Further,  $Z$  is isomorphic to  $G/B \times G/B^-$  as a  $G \times G$  variety. By [DP, p. 26, 8.1], the pullback homomorphism

$$i^* : \mathrm{Pic}(X) \longrightarrow \mathrm{Pic}(Z),$$

for the inclusion map  $i : Z \hookrightarrow X$  is injective and is given by

$$i^*(\mathcal{L}_\chi) = p_1^*(L_\chi) \otimes p_2^*(L_{-\chi}),$$

where  $L_\chi$  (respectively,  $L_{-\chi}$ ) is the line bundle on  $G/B$  (respectively,  $G/B^-$ ) associated  $\chi$  (respectively,  $-\chi$ ) and  $p_j$  is the projection to the  $j$ -th factor of  $G/B \times G/B^-$  for  $j = 1, 2$ .

### 3. CHOICE OF A POLARIZATION ON $\overline{G}$

We continue with the notation of Section 2. Let  $G$  be a simple algebraic group of adjoint type of rank  $n \geq 2$ , such that its root system  $R$  is different from  $A_2$ . Let

$$\mathrm{NS} := \left\{ \sum_{i=1}^n m_i \alpha_i : m_i \in \mathbb{N} \right\}.$$

Then, we have the following:

**Lemma 3.1.** *The above defined  $\mathrm{NS}$  contains a regular dominant character  $\chi$  of  $T$  such that  $s_{\alpha_i}(\chi) \geq 0$  and  $\langle \chi, w(\lambda_i) \rangle \neq 0$  for every  $w \in W$  and  $1 \leq i \leq n$ .*

*Proof.* Denote by  $X(T)_\mathbb{Q}$  the rational vector space generated by  $X(T)$ , and also denote by  $X(T)^+$  the semi-group of it given by the dominant characters of  $T$ . Let  $\rho \in X(T)_\mathbb{Q}$  be the half sum of positive roots of  $R$ . Then,  $2\rho = 2(\sum_{i=1}^n \omega_i) \in X(T)^+$  is a regular dominant character of  $T$ , and we have  $2\rho \in \mathrm{NS}$ .

Since  $R$  is irreducible of rank at-least 2 and different from  $A_2$ , we see that for every simple root  $\alpha_i$ , there are at-least 3 positive roots  $\beta$  satisfying  $\alpha_i \leq \beta$ . Hence, the coefficient of every simple root  $\alpha_j$  in the expression of  $s_{\alpha_i}(2\rho) = 2\rho - 2\alpha_i$  (as a non-negative integral linear combination of simple roots) is positive. Hence, we have  $s_{\alpha_i}(2\rho) \in \mathbb{N}S$ . Thus, we have

$$2\rho \in X(T)^+ \cap \left( \bigcap_{i=1}^n s_{\alpha_i}(\mathbb{N}S) \right).$$

Denote by  $N$  the determinant of the Cartan matrix of  $R$ . Then we have  $N\omega_i \in \mathbb{N}S$  for every  $i = 1, 2, \dots, n$ . By the previous discussion, there exists  $m \in \mathbb{N}$  such that  $ms_{\alpha_i}(2\rho) - N\alpha_i \in \mathbb{N}S$  for every  $1 \leq i \leq n$ . Hence, we get

$$s_{\alpha_i}(2m\rho + N\omega_i) = ms_{\alpha_i}(2\rho) - N\alpha_i + N\omega_i \in \mathbb{N}S, \quad 1 \leq i \leq n,$$

and from this it follows that

$$2m\rho + N\omega_i \in X(T)^+ \cap \left( \bigcap_{j=1}^n s_{\alpha_j}(\mathbb{N}S) \right), \quad 1 \leq i \leq n.$$

Consider the characters  $2m\rho, 2m\rho + N\omega_2, \dots, 2m\rho + N\omega_n$  of  $T$ . These are linearly independent in  $X(T)$  and by construction they all lie in the rational cone

$$\mathcal{C} \subset X(T)_{\mathbb{Q}}^+$$

generated by the semi-group  $X(T)^+ \cap \left( \bigcap_{i=1}^n s_{\alpha_i}(\mathbb{N}S) \right)$ . It follows that  $\mathcal{C}$  has a maximal dimension in  $X(T)_{\mathbb{Q}}$ , hence it is not contained in any hyperplane of  $X(T)_{\mathbb{Q}}$ . Therefore, there exists a regular dominant character  $\chi \in \mathcal{C} \cap \mathbb{N}S$  such that  $\langle \chi, w(\lambda_i) \rangle \neq 0$  for all  $1 \leq i \leq n$  and every  $w \in W$ , and hence the lemma follows.  $\square$

**Lemma 3.2.** *Let  $\chi \in \mathbb{N}S$  be a regular dominant character of  $T$  satisfying the properties stated in Lemma 3.1. Then we have*

- (1)  $(G/B)_T^{ss}(L_\chi) = (G/B)_T^s(L_\chi)$ , and
- (2) the set of all unstable points

$$(G/B) \setminus (G/B)_T^{ss}(L_\chi)$$

*is contained in the union of  $W$ -translates of all Schubert varieties of codimension at least two.*

*Proof.* Set  $L := L_\chi$ . Since  $\langle \chi, w(\lambda_i) \rangle \neq 0$  for every  $w \in W$  and  $1 \leq i \leq n$ , by [Ka1, p. 38, Lemma 4.1] we have

$$(G/B)_T^{ss}(L) = (G/B)_T^s(L).$$

This proves (1).

To prove (2), take an unstable point  $x \in G/B$  for the polarization  $L$ . Then, there is a one-parameter subgroup  $\lambda$  of  $T$  such that  $\mu^L(x, \lambda) < 0$ . Let  $\phi \in W$  be such that  $\phi(\lambda)$  is in the fundamental chamber, say

$$\phi(\lambda) = \sum_{i=1}^n c_i \lambda_i,$$

where  $\{c_i\}$  are non-negative integers. Consequently, we have

$$\mu^L(n_\phi(x), \phi(\lambda)) = \mu^L(x, \lambda) < 0,$$

where  $n_\phi$  is a representative of  $\phi$  in  $N_G(T)$ . Now, let  $n_\phi(x)$  be in the Schubert cell  $BwB/B$  for some  $w \in W$ . By [Se, Lemma 5.1], we have

$$\mu^L(n_\phi(x), \phi(\lambda)) = \left(-\sum_{i=1}^n c_i \langle w(\chi), \lambda_i \rangle\right) < 0.$$

(The sign here is negative because we are using left action of  $B$  on  $G/B$  while in [Se, Lemma 5.1] the action of  $B$  on  $B \backslash G$  is on the right.) Therefore we have  $w(\chi) \not\leq 0$ . For every  $1 \leq i \leq n$  we have  $s_{\alpha_i}(\chi) \geq 0$ , and hence  $w_0 s_{\alpha_i}(\chi) \leq 0$ . Hence we have  $l(w_0) - l(w) \geq 2$ . This completes the proof of (2).  $\square$

**Proposition 3.3.** *Let  $X = \overline{G}$  be the wonderful compactification of  $G$ . Let  $\chi$  be as in Lemma 3.2, and let  $X_T^{ss}(\mathcal{L}_\chi)$  (respectively,  $X_T^s(\mathcal{L}_\chi)$ ) be the semi-stable (respectively, stable) locus of  $X$  for the action of  $1 \times T$  and the polarization  $\mathcal{L}_\chi$  on  $X$ . Then we have*

- (1)  $X_T^{ss}(\mathcal{L}_\chi) = X_T^s(\mathcal{L}_\chi)$ , and
- (2) *the set of unstable points  $X \setminus (X_T^{ss}(\mathcal{L}_\chi))$  is a union of irreducible closed subvarieties of codimension at least three.*

*Proof.* Let  $Z$  be the unique closed  $G \times G$ -orbit in  $X$ . Let  $Z_T^{ss}(\mathcal{L}_\chi)$  (respectively,  $Z_T^s(\mathcal{L}_\chi)$ ) be the semi-stable (respectively, stable) locus of  $Z$  for the action of  $1 \times T$  and the polarization  $i^*(\mathcal{L}_\chi)$ , where  $i : Z \hookrightarrow X$  is the inclusion map. Since  $Z$  is isomorphic to  $G/B \times G/B^-$  and  $i^*(\mathcal{L}_\chi) = p_1^*(L_\chi) \otimes p_2^*(L_{-\chi})$ , we see that

$$Z_T^{ss}(\mathcal{L}_\chi) \simeq (G/B) \times ((G/B^-)_T^{ss}(L_{-\chi}))$$

and  $Z_T^s(\mathcal{L}_\chi) \simeq (G/B) \times ((G/B^-)_T^s(L_{-\chi}))$ . Set  $Z^{ss} = Z_T^{ss}(\mathcal{L}_\chi)$  and  $Z^s = Z_T^s(\mathcal{L}_\chi)$ . By Lemma 3.2 and above discussion, we have  $Z^{ss} = Z^s$ .

For convenience, we will denote  $X_T^{ss}(\mathcal{L}_\chi)$  and  $X_T^s(\mathcal{L}_\chi)$  by  $X^{ss}$  and  $X^s$  respectively. If  $X^{ss} \neq X^s$ , then the complement  $X^{ss} \setminus X^s$  is a non-empty  $G \times T$  invariant closed subset of  $X^{ss}$ . Hence, the complement  $(X^{ss} // T) \setminus (X^s // T)$  is a non-empty  $G \times \{1\}$ -invariant closed subset of  $X^{ss} // T$ . In particular,  $(X^{ss} // T) \setminus (X^s // T)$  is a finite union of non-empty  $G \times \{1\}$ -invariant projective varieties. Therefore, there is a  $B \times \{1\}$ -fixed point in  $(X^{ss} // T) \setminus (X^s // T)$ . Let

$$p \in (X^{ss} // T) \setminus (X^s // T)$$

be a  $B \times \{1\}$ -fixed point. Let  $Y$  be the closed  $\{1\} \times T$ -orbit in the fiber  $\pi^{-1}(\{p\})$  over  $p$  for the geometric invariant theoretic quotient map  $\pi : X^{ss} \rightarrow X^{ss} // T$ . Since this map  $\pi$  is  $G \times \{1\}$  equivariant, we conclude that  $\pi^{-1}(\{p\})$  is  $B \times \{1\}$ -invariant. Hence, for any  $b \in B$ , the translation  $(b, 1) \cdot Y$  lies in  $\pi^{-1}(\{p\})$ . Since the actions of  $B \times \{1\}$  and  $\{1\} \times T$  on  $X$  commute with each other, we see that  $(b, 1) \cdot Y$  is also a closed  $\{1\} \times T$ -orbit in  $\pi^{-1}(\{p\})$ . By the uniqueness of the closed  $\{1\} \times T$ -orbit in  $\pi^{-1}(\{p\})$  we conclude that  $(b, 1) \cdot Y = Y$ . Hence  $Y$  is preserved by the action of  $B \times \{1\}$ . In particular,  $Y$  is  $U \times \{1\}$ -invariant, where  $U \subset B$  is the unipotent radical. The action of  $U \times \{1\}$  on

$Y$  induces a homomorphism from  $U$  to  $T/S$  of algebraic groups, where  $\{1\} \times S$  is the stabilizer in  $\{1\} \times T$  of some point  $q$  in  $Y$ . Since there is no nontrivial homomorphism from an unipotent group to a torus, we conclude that  $U \times \{1\}$  fixes the point  $q$ .

By [DP, p. 32, Proposition], for any regular dominant character  $\chi$  of  $T$  with respect to  $B$ , the morphism  $X \hookrightarrow \mathbb{P}(V(\chi) \otimes V(\chi)^*)$  is a  $G \times G$  equivariant embedding, where  $V(\chi)$  is the irreducible representation of  $G$  with highest weight  $\chi$ , and  $V(\chi)^*$  is its dual. Hence, the  $U \times 1$ -fixed point set of  $X$  is equal to  $X \cap \mathbb{P}(\mathbb{C}_\chi \otimes V(\chi)^*)$ , where  $\mathbb{C}_\chi$  is the one dimensional  $B$ -module associated to the character  $\chi$ . Therefore, by the above discussion, we have  $q \in X \cap \mathbb{P}(\mathbb{C}_\chi \otimes V(\chi)^*)$ .

Further, by [DP, Theorem, p. 30] we have

$$H^0(X, \mathcal{L}_\chi) = \bigoplus_{\nu \leq \chi} V(\nu)^* \otimes V(\nu),$$

where the sum runs over all dominant characters  $\nu$  of  $T$  satisfying  $\nu \leq \chi$ . By [DP, p. 29, Corollary] and [DP, p. 30, Theorem], the zero locus of

$$\bigoplus_{\nu < \chi} V(\nu)^* \otimes V(\nu) \subset H^0(X, \mathcal{L}_\chi)$$

in  $X$  is the unique closed  $G \times G$ -orbit  $Z = G/B \times G/B^-$ . Hence, by the discussion in the previous paragraph, we have  $q \in Z$ . This contradicts the choice of the polarization  $\mathcal{L}_\chi$ . Therefore, the proof of (1) is complete.

To prove (2), note that  $X \setminus X^{ss}$  is a closed subset of  $X$ , and

$$Z \setminus Z^{ss} = (X \setminus X^{ss}) \cap Z.$$

Also, by Lemma 3.2, the complement  $Z \setminus Z^{ss} \subset Z$  is of codimension at least two. Since we have  $Z = \bigcap_{i=1}^n D_i$ , the complement  $D_i \setminus D_i^{ss}$  is of codimension at least two for all  $1 \leq i \leq n$ . Further, every point in the open subset  $G \subset X$  is semistable. Hence,  $X \setminus X^{ss}$  is of codimension at least three.  $\square$

The following lemma will be used in the proof of Corollary 3.5.

**Lemma 3.4.** *Let  $H$  be a reductive algebraic group acting linearly on a polarized projective variety  $V$ . Assume that  $V^{ss} = V^s$ , where  $V^{ss}$  (respectively,  $V^s$ ) is the set of semi-stable (respectively, stable) points of  $V$  for the action of  $H$ . Then the set of all points in  $V^{ss}$  whose stabilizer in  $H$  is trivial is actually a Zariski open subset (it may be possibly empty).*

*Proof.* Consider the morphism

$$f : H \times V^{ss} \longrightarrow V^{ss} \times V^{ss}, \quad (h, v) \longmapsto (h \cdot v, v).$$

Since  $V^{ss} = V^s$ , this map  $f$  is proper [MFK, p. 55, Corollary 2.5]. Hence the image

$$M := f(H \times V^{ss}) \subset V^{ss} \times V^{ss}$$

is a closed subvariety. Now, let

$$U' \subset V^{ss}$$

be the locus of points with trivial stabilizer (for the action of  $H$ ). Take any

$$v_0 \in U',$$

and set  $z_0 := f((1, v_0)) = (v_0, v_0)$ . Then,  $(f_*\mathcal{O}_{H \times V^{ss}})_{z_0}$  is a free  $\mathcal{O}_{M, z_0}$ -module of rank one. Hence by [Mu, p. 152, Souped-up version II of Nakayama's Lemma], the locus of points  $x \in M$  such that  $(f_*\mathcal{O}_{H \times V^{ss}})_x$  is a free  $\mathcal{O}_{M, x}$ -module of rank at most one is a non-empty Zariski open subset. Since  $(f_*(\mathcal{O}_{H \times V^{ss}}))_z$  is nonzero for all  $z \in M$ , the set of all points  $x \in M$  such that  $(f_*(\mathcal{O}_{H \times V^{ss}}))_x$  is a free  $\mathcal{O}_{M, x}$ -module of rank one is a Zariski open subset of  $M$ ; this Zariski open subset of  $M$  will be denoted by  $U$ . Note that

$$f^{-1}(U) = p_2^{-1}(U'),$$

where  $p_2 : H \times V^{ss} \rightarrow V^{ss}$  is the second projection. Since  $p_2$  is flat of finite type over  $\mathbb{C}$ , it is an open map (see [Ha, p. 266, Exercise 9.1]). Hence  $U' = p_2(f^{-1}(U))$  is a Zariski open subset. This finishes the proof of the lemma.  $\square$

**Corollary 3.5.** *Let  $X = \overline{\mathrm{PSL}(n+1, \mathbb{C})}$  be the wonderful compactification of  $\mathrm{PSL}(n+1, \mathbb{C})$ ,  $n \geq 3$ . For the choice of the regular dominant character  $\chi$  of  $T$  as in Proposition 3.3,*

- (1) *the action of  $\{1\} \times T$  on  $X_T^{ss}(\mathcal{L}_\chi)//T$  is free,*
- (2)  *$X_T^{ss}(\mathcal{L}_\chi)//T$  is a smooth projective embedding of  $G/T$ , and*
- (3) *the set of unstable points  $X \setminus (X_T^{ss}(\mathcal{L}_\chi))$  is a union of irreducible closed subvarieties of codimension at least three.*

*Proof.* Let  $\chi$  be a regular dominant character of  $T$  as in Proposition 3.3. As in the proof of Proposition 3.3, let  $Z$  denote the unique closed  $G \times G$  orbit in  $X$ . Also, let  $X^{ss}$ ,  $X^s$ ,  $Z^{ss}$  and  $Z^s$  be as in the proof of Proposition 3.3. By Proposition 3.3 we have  $X^{ss} = X^s$ . Hence by Lemma 3.4, the locus  $V$  of points in  $X^{ss}$  with trivial stabilizer (for the action of  $\{1\} \times T$ ) is a Zariski open subset of  $X^{ss}$ . Therefore,  $X^{ss} \setminus V$  is a  $G \times \{1\}$  stable closed subvariety of  $X^{ss}$ . By using the arguments in the the proof of Proposition 3.3 we see that the set of  $B \times \{1\}$ -fixed points in  $Z \cap (X^{ss} \setminus V)$  is non-empty. But on the other hand by the proof of [Ka2, p. 194, Example 3.3] we see that given any point  $z \in Z^{ss}$ , its stabilizer subgroup in  $\{1\} \times T$  is trivial. This is a contradiction. Hence we conclude that the action of  $\{1\} \times T$  on  $X^{ss}$  is free. This proves Part (1) and Part (2).

Part (3) follows immediately from the corresponding statement in Proposition 3.3.  $\square$

#### 4. AUTOMORPHISM GROUP OF $\overline{\mathrm{PSL}(n+1, \mathbb{C})}_T^{ss}(\mathcal{L})//T$

Let  $G = \mathrm{PSL}(n+1, \mathbb{C})$ , with  $n \geq 3$ , and define

$$Y := \overline{\mathrm{PSL}(n+1, \mathbb{C})}_T^{ss}(\mathcal{L}_\chi)//T,$$

where  $\chi$  is as in Proposition 3.3.

**Theorem 4.1.** *Let  $A$  denote the connected component, containing the identity element, of the group of holomorphic (= algebraic) automorphisms of  $Y$ . Then*

- (1)  $A$  is isomorphic to  $G$ , and
- (2) the Picard group of  $Y$  is a free Abelian group of rank  $2n$ .

*Proof.* Let  $TY$  denote the algebraic tangent bundle of  $Y$ . From [MO, Theorem 3.7] we know that  $A$  is an algebraic group. The Lie algebra of  $A$  is  $H^0(Y, TY)$  equipped with the Lie bracket operation of vector fields.

The Lie algebra of  $G$  will be denoted by  $\mathfrak{g}$ . Define  $X := \overline{\text{PSL}(n+1, \mathbb{C})}$  and  $U := \overline{\text{PSL}(n+1, \mathbb{C})}_T^{\text{ss}}(\mathcal{L}_X)$ . The connected component, containing the identity element, of the automorphism group of  $X$  is  $G \times G$  [Br, Example 2.4.5]. From this, and the fact that the complement  $X \setminus U \subset X$  is of codimension at least three (see Corollary 3.5), we conclude that

$$H^0(U, TU) = H^0(X, TX) = \mathfrak{g} \oplus \mathfrak{g}.$$

Let  $\phi : U \rightarrow Y$  be the geometric invariant theoretic quotient map. Let

$$T_U \supset T_\phi \rightarrow U$$

be the relative tangent bundle for  $\phi$ . Since  $\phi$  makes  $U$  a principal  $T$ -bundle over  $Y$  (see Corollary 3.5(1)), we have the following short exact sequence of vector bundles on  $U$

$$(4.1) \quad 0 \rightarrow T_\phi \rightarrow TU \rightarrow \phi^*(TY) \rightarrow 0,$$

and the relative tangent bundle  $T_\phi$  is identified with the trivial vector bundle  $\mathcal{O}_U \otimes_{\mathbb{C}} \mathfrak{h}$ , where  $\mathfrak{h}$  is the Lie algebra of  $T$ .

Set  $Z = X \setminus U$ . Since  $\text{codim}(Z) \geq 3$  (Corollary 3.5), we have

$$H^0(U, T_\phi) = H^0(X, \mathcal{O}_X \otimes \mathfrak{h}) = \mathfrak{h}.$$

Note that  $H^1(U, T_\phi) = H_Z^2(X, \mathcal{O}_X \otimes \mathfrak{h})$ . Indeed, this follows from the following cohomology exact sequence (see [Gr, Corollary 1.9])

$$H^1(X, \mathcal{O}_X \otimes \mathfrak{h}) \rightarrow H^1(U, \mathcal{O}_X \otimes \mathfrak{h}) \rightarrow H_Z^2(X, \mathcal{O}_X \otimes \mathfrak{h}) \rightarrow H^2(X, \mathcal{O}_X \otimes \mathfrak{h})$$

combined with the fact that  $H^i(X, \mathcal{O}_X) = 0$  for all  $i > 0$  [DP, p. 30, Theorem]. As  $X$  is smooth and  $\text{codim}(Z) \geq 3$ , it follows from [Gr, Theorem 3.8 and Proposition 1.4] that

$$H_Z^2(X, \mathcal{O}_X) = 0,$$

and hence  $H^1(U, T_\phi) = 0$ . Now, using this fact in the long exact sequence of cohomologies corresponding to the short exact sequence in (4.1) we obtain the following short exact sequence:

$$0 \rightarrow 0 \oplus \mathfrak{h} \rightarrow \mathfrak{g} \oplus \mathfrak{g} \rightarrow H^0(U, \phi^*TY) \rightarrow 0.$$

Hence, we have

$$H^0(U, \phi^*TY) = \mathfrak{g} \oplus (\mathfrak{g}/\mathfrak{h}).$$

By using geometric invariant theory,  $H^0(Y, TY)$  is the invariant part

$$H^0(Y, TY) = H^0(U, \phi^*TY)^{\{1\} \times T} \subset H^0(U, \phi^*TY).$$

Thus we have  $H^0(Y, TY) = \mathfrak{g}$ . This proves (1).



To prove (2), let  $\{D_i \mid 1 \leq i \leq n\}$  be the  $(G \times G)$ -stable irreducible closed subvarieties of  $\overline{G}$  of codimension one such that

$$G = \overline{G} \setminus \left( \bigcup_{i=1}^n D_i \right).$$

Let  $D_i^{ss} = D_i \cap X^{ss} \subset D_i$  be the semistable locus of  $D_i$ . Set  $Z := Y \setminus (G/T)$ , and write it as a union

$$Z = \bigcup_{i=1}^n Z_i,$$

where each  $Z_i = D_i^{ss} // T$  is an irreducible closed subvariety of  $Y$  of codimension one. As  $Y$  is smooth, each  $Z_i$  produces a line bundle  $L_i \rightarrow Y$  whose pullback to  $X^{ss}$  is  $\mathcal{O}_{X^{ss}}(D_i^{ss})$ . Since  $\mathrm{Pic}(X^{ss}) = \mathrm{Pic}(X)$  and  $\{\mathcal{O}_X(D_i)\}_{1 \leq i \leq n}$  are linearly independent in  $\mathrm{Pic}(X)$  (see [DP, p. 26, 8.1]), we get that  $L_i$ ,  $1 \leq i \leq n$ , are linearly independent in  $\mathrm{Pic}(Y)$ . The Picard group of  $G/T$  is isomorphic to the group of characters of the inverse image  $\widehat{T}$  of  $T$  inside the simply connected covering  $\widehat{G}$  of  $G$  (see [KKV]). Now it follows from the exact sequence in [Fu, Proposition 1.8] that  $\mathrm{Pic}(Y)$  is a free Abelian group of rank  $2n$ , thus completing the proof of (2).  $\square$

**Remark 4.2.** The compactification  $Y$  of  $G/T$  constructed here is an example of a non-spherical variety for the action of  $G$  whose connected component of the automorphism group is  $G$ .

**Remark 4.3.** Note that both  $Y$  and  $G/B \times G/B$  are smooth compactifications of  $G/T$  with isomorphic Picard groups. Further both are Fano varieties, i.e., the anti-canonical line bundle is ample. The fact that  $G/B \times G/B$  is Fano is well known. That the variety  $Y$  is Fano follows as a consequence of the exact sequence in (4.1) together with the facts that  $X$  is Fano (see [DP]) and the codimension of  $X \setminus U$  is greater than or equal to 3, where  $X$  and  $U$  are as in the proof of Theorem 4.1. But  $Y$  and  $G/B \times G/B$  are not isomorphic, as  $\mathrm{Aut}^0(Y) \simeq G$  and  $\mathrm{Aut}^0(G/B \times G/B) \simeq G \times G$ , where  $\mathrm{Aut}^0(M)$  denotes the connected component of the group of algebraic automorphisms of a smooth projective variety  $M$  containing the identity element.

**Remark 4.4.** In [St], Strickland extended the construction of  $\overline{G}$  to any arbitrary algebraically closed field. Also,  $\overline{G}$  is a Frobenius split variety in positive characteristic [St, p. 169, Theorem 3.1] (see [MR] for the definition of Frobenius splitting). Since  $T$  is linearly reductive, using Reynolds operator, one can see that the geometric invariant theoretic quotient of  $\overline{G}$  for the action of  $T$  is also Frobenius split for any polarization on  $\overline{G}$ .

#### ACKNOWLEDGEMENTS

We are grateful to the referee for comments to improve the exposition. The first-named author thanks the Institute of Mathematical Sciences for hospitality while this work was carried out. He also acknowledges the support of the J. C. Bose Fellowship. The second named author would like to thank the Infosys Foundation for the partial support.

## REFERENCES

- [Br] M. Brion, The total coordinate ring of a wonderful variety, *Jour. Alg.* **313** (2007), 61–99.
- [DP] C. De Concini and C. Procesi, Complete symmetric varieties, *Invariant theory* (Montecatini, 1982), 1–44, Lecture Notes in Math., 996, Springer, Berlin, 1983.
- [Fu] W. Fulton, *Intersection Theory*, Second edition, Springer, Berlin, 1998.
- [Gr] A. Grothendieck, *Local cohomology*, Lecture Notes in Math., 41, Springer, Berlin, 1967.
- [Ha] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, Springer-Verlag, Berlin, Heidelberg, 1977.
- [Hu1] J. E. Humphreys, *Introduction to Lie algebras and Representation theory*, Springer-Verlag, Berlin Heidelberg, New York, 1972.
- [Hu2] J. E. Humphreys, *Linear Algebraic Groups Representation theory*, GTM 21, Springer-Verlag, Berlin Heidelberg, New York, 1975.
- [Ka1] S.S. Kannan, Torus quotients of homogeneous spaces-II, *Proc. Indian Acad. Sci.(Math. Sci)* **109** (1999), 23–39.
- [Ka2] S.S. Kannan, GIT related problems of the flag variety for the action of a maximal torus, Springer Proceedings in Mathematics & Statistics, **82** (2014), 189–203.
- [KKV] F. Knop, H. Kraft and T. Vust, The Picard group of a  $G$ -variety, *Algebraische Transformationsgruppen und Invariantentheorie*, 77–87, DMV Sem., **13**, Birkhäuser, Basel, 1989.
- [MO] H. Matsumura and F. Oort, Representability of group functors, and automorphisms of algebraic schemes, *Invent. Math.* **4** (1967), 1–25.
- [MR] V.B. Mehta and A. Ramanathan, Frobenius splitting and cohomology vanishing for Schubert varieties, *Ann. Math.* **122** (1985), 27–40.
- [Mu] D. Mumford, *The red book of varieties and schemes*, Second expanded edition, Springer, 1999.
- [MFK] D. Mumford, J. Fogarty and F. Kirwan, *Geometric Invariant Theory*, Springer-Verlag, Berlin Heidelberg, New York, 1994.
- [Se] C.S. Seshadri, Quotient Spaces modulo reductive algebraic groups, *Ann. Math.* **95** (1972), 511–556.
- [St] E. Strickland, A vanishing theorem for group compactifications, *Math. Ann.* **277** (1987), 165–171.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

*E-mail address:* `indranil@math.tifr.res.in`

CHENNAI MATHEMATICAL INSTITUTE, H1, SIPCOT IT PARK, SIRUSERI, KELAMBAKKAM 603103, INDIA

*E-mail address:* `kannan@cmi.ac.in`

THE INSTITUTE OF MATHEMATICAL SCIENCES, CIT CAMPUS, TARAMANI, CHENNAI 600113, INDIA

*E-mail address:* `dsn@imsc.res.in`