

# An Improvement of the Cipolla-Lehmer Type Algorithms

Namhun Koo<sup>1</sup>, Gook Hwa Cho<sup>2</sup>, Byeonghwan Go<sup>2</sup>, and Soonhak Kwon<sup>2</sup>

Email: nhkoo@nims.re.kr, achimheasal@nate.com, kobhh@skku.edu, shkwon@skku.edu

National Institute for Mathematical Sciences, Daejeon, Republic of Korea<sup>1</sup>

Sungkyunkwan University, Suwon, Republic of Korea<sup>2</sup>

## Abstract

Let  $\mathbb{F}_q$  be a finite field with  $q$  elements with prime power  $q$  and let  $r > 1$  be an integer with  $q \equiv 1 \pmod{r}$ . In this paper, we present a refinement of the Cipolla-Lehmer type algorithm given by H. C. Williams, and subsequently improved by K. S. Williams and K. Hardy. For a given  $r$ -th power residue  $c \in \mathbb{F}_q$  where  $r$  is an odd prime, the algorithm of H. C. Williams determines a solution of  $X^r = c$  in  $O(r^3 \log q)$  multiplications in  $\mathbb{F}_q$ , and the algorithm of K. S. Williams and K. Hardy finds a solution in  $O(r^4 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ . Our refinement finds a solution in  $O(r^3 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ . Therefore our new method is better than the previously proposed algorithms independent of the size of  $r$ , and the implementation result via SAGE shows a substantial speed-up compared with the existing algorithms.

**Keywords** : finite field,  $r$ -th root, Cipolla-Lehmer algorithm, Adleman-Manders-Miller algorithm, primitive root

**MSC 2010 Codes** : 11T06, 11Y16, 68W40

## 1 Introduction

Let  $r > 1$  be an integer and  $q$  be a power of a prime. Finding  $r$ -th root (or finding a root of  $X^r = c$ ) in finite field  $\mathbb{F}_q$  has many applications in computational number theory and in many other related topics. Some such examples include point halving and point compression on elliptic curves [15], where square root computations are needed. Similar applications for high genus curves require  $r$ -th root computations also.

Among several available root extraction methods of the equation  $X^r - c = 0$ , there are two well known algorithms applicable for arbitrary integer  $r > 1$ ; the Adleman-Manders-Miller algorithm [1], a straightforward generalization of the Tonelli-Shanks square root algorithm [16, 18] to the case of  $r$ -th root extraction, and the Cipolla-Lehmer algorithms [7, 11]. Due to the cumbersome extension field arithmetic needed for the Cipolla-Lehmer algorithm, one usually prefers the Tonelli-Shanks or the Adleman-Manders-Miller, and other related researches [2, 3, 10] exist to improve the Tonelli-Shanks.

The efficiency of the Adleman-Manders-Miller algorithm heavily depends on the exponent  $\nu$  of  $r$  satisfying  $r^\nu | q - 1$  and  $r^{\nu+1} \nmid q - 1$ , which becomes quite slow if  $\nu \approx \log q$ . Even in the case of  $r = 2$ , it had been observed in [14] that, for a prime  $p = 9 \times 2^{3354} + 1$ , running the Tonelli-Shanks algorithm using various software such as Magma, Mathematica and Maple cost roughly 5 minutes, 45 minutes, 390 minutes, respectively while the Cipolla-Lehmer costs under 1 minute in any of the above softwares. It should be mentioned that such extreme cases (of  $p$  with  $p - 1$  divisible by high powers of 2) may happen in some cryptographic applications.

For example, one of the NIST suggested curve [15] P-224 :  $y^2 = x^3 - 3x + b$  over  $\mathbb{F}_p$  uses a prime  $p = 2^{224} - 2^{96} + 1$ .

A generalization to  $r$ -th root extraction of the Cipolla-Lehmer square root algorithm is proposed by H. C. Williams [19] and the complexity of the proposed algorithm is  $O(r^3 \log q)$  multiplications in  $\mathbb{F}_q$ . A refinement of the algorithm in [19] was given by K. S. Williams and K. Hardy [20] where the complexity is reduced to  $O(r^4 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ . For the case of the square root, a new Cipolla-Lehmer type algorithm based on the Lucas sequence was given by Müller [14]. A similar result for the case  $r = 3$  was also obtained by Cho et al. [5], and a possible generalization to the  $r$ -th root extraction of Müller's square root algorithm was given in [6].

In this paper, we present a new Cipolla-Lehmer type algorithm for  $r$ -th root extractions in  $\mathbb{F}_q$  whose complexity is  $O(r^3 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ , which improves previously proposed results in [19, 20]. We also compare our algorithm with those in [19, 20] using the software SAGE, and show that our algorithm performs consistently better than those in [19, 20] as is expected from the theoretical complexity estimation. In [19] and [20], only the case where  $r$  is an odd prime was considered but we will give the general arguments (i.e., no restriction on  $r$ ) here.

The remainder of this paper is organized as follows: In Section 2, we briefly summarize the Cipolla-Lehmer algorithm, and introduce the works of H. C. Williams [19] and K. S. Williams and K. Hardy [20]. In Section 3, we present our refinement of the Cipolla-Lehmer algorithm. In Section 4, we give the complexity analysis of our algorithm and show the result of SAGE implementations of the three algorithms (in [19], [20], and ours). Finally, in Section 5, we give the concluding remarks.

## 2 Cipolla-Lehmer Algorithm in $\mathbb{F}_q$

Let  $q$  be a prime power and  $\mathbb{F}_q$  be a finite field with  $q$  elements. Let  $c \neq 0 \in \mathbb{F}_q$  be an  $r$ -th power residue in  $\mathbb{F}_q$  for an integer  $r > 1$  with  $q \equiv 1 \pmod{r}$ . We restrict  $r$  as an odd prime in this section.

### 2.1 H. C. Williams' algorithm

Let  $b \in \mathbb{F}_q$  be an element such that  $b^r - c$  is not an  $r$ -th power residue in  $\mathbb{F}_q$ . Such  $b$  can be found after  $r$  random trials of  $b$ . (See pp.479-480 in [20] for further explanation.) Then the polynomial  $X^r - (b^r - c)$  is irreducible over  $\mathbb{F}_q$  and there exists  $\theta \in \mathbb{F}_{q^r} - \mathbb{F}_q$  such that  $\theta^r = b^r - c$ . Let  $\omega = \theta^{q-1} = (b^r - c)^{\frac{q-1}{r}}$ . Then we have  $\omega^r = 1$  where  $\omega$  is a primitive  $r$ -th root because  $b^r - c$  is not an  $r$ -th power in  $\mathbb{F}_q$ .

For all  $0 \leq i \leq r-1$ , using  $q \equiv 1 \pmod{r}$ , one has  $\theta^{q^i} = \theta \cdot \theta^{q^i-1} = \theta \cdot (\theta^{q-1})^{1+q+\dots+q^{i-1}} = \theta \omega^i$ , which implies  $(b - \theta)^{q^i} = b - \theta^{q^i} = b - \omega^i \theta$ . Letting  $\alpha = b - \theta$ , one has

$$\alpha^{\sum_{j=0}^{r-1} q^j} = (b - \theta)^{1+q+q^2+\dots+q^{r-1}} = \prod_{i=0}^{r-1} (b - \omega^i \theta) = b^r - \theta^r = c. \quad (1)$$

Thus one may find an  $r$ -th root of  $c$  by computing  $\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}} \in \mathbb{F}_q[\theta] = \mathbb{F}_q[X]/\langle X^r - (b^r - c) \rangle$ .

**Proposition 1.** [H. C. Williams]

Suppose that  $c \neq 0$  is an  $r$ -th power in  $\mathbb{F}_q$ . Let  $\theta^r = b^r - c$  with  $\theta \in \mathbb{F}_{q^r}$  and  $b \in \mathbb{F}_q$  such that

$b^r - c$  is not an  $r$ -th power in  $\mathbb{F}_q$ . Then letting  $\alpha = b - \theta$ ,

$$\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}} \in \mathbb{F}_q$$

is an  $r$ -th root of  $c$ .

The usual ‘square and multiply method’ (or ‘double and add method’ if one uses a linear recurrence relation) requires roughly  $\log \frac{\sum_{j=0}^{r-1} q^j}{r} \approx r \log q$  steps for the evaluation of  $\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}}$ , and therefore the complexity of the algorithm of H. C. Williams is  $O(r^3 \log q)$  multiplications in  $\mathbb{F}_q$ . H. C. Williams’ result can be expressed in Algorithm 1 using the recurrence relation technique of Section 2.2.

---

**Algorithm 1** H. C. Williams’  $r$ -th root algorithm [19]

---

**Input :** An  $r$ -th power residue  $c$  in  $\mathbb{F}_q$

**Output :**  $x \in \mathbb{F}_q$  satisfying  $x^r = c$

- 1: **do** Choose a random  $b \in \mathbb{F}_q$  **until**  $b^r - c$  is not an  $r$ -th power residue.
  - 2:  $M \leftarrow \frac{1+q+\dots+q^{r-1}}{r}$
  - 3:  $A \leftarrow (b, -1, 0, \dots, 0)$  //  $A$  is a coefficient vector of  $\alpha = b - \theta$ . //
  - 4:  $A \leftarrow \text{RecurrenceRelation}(A, M)$  //  $A$  is a coefficient vector of  $\alpha^M$ . //
  - 5:  $x \leftarrow$  corresponding element of  $A$  //  $x = \alpha^M$  //
  - 6: **return**  $x$
- 

Note that  $\alpha = b + \theta$  is used in the original paper [19], while our presentation is based on [20] where it uses  $\alpha = b - \theta$ . We followed [20] because it is more convenient to deal with general  $r$  which is not necessarily odd prime. For example, if one uses  $\alpha = b + \theta$  as in [19], then the case of even  $r$  (such as  $r = 2$ ) cannot be covered. Detailed explanations will be given in Section 3.

## 2.2 Recurrence relation

Given  $\sum_{i=0}^{r-1} a_i \theta^i \in \mathbb{F}_q[\theta]$ , define  $a_i(j) \in \mathbb{F}_q$  ( $0 \leq i \leq r-1$ ,  $1 \leq j$ ) as

$$\sum_{i=0}^{r-1} a_i(j) \theta^i = \left( \sum_{i=0}^{r-1} a_i \theta^i \right)^j. \quad (2)$$

In particular, one has  $a_i(1) = a_i$  for all  $0 \leq i \leq r-1$ . Then one has

$$\begin{aligned} \sum_{i=0}^{r-1} a_i(m+n) \theta^i &= \left( \sum_{i=0}^{r-1} a_i(m) \theta^i \right) \left( \sum_{j=0}^{r-1} a_j(n) \theta^j \right) \\ &= \sum_{l=0}^{r-1} \left( \sum_{j=0}^l a_j(m) a_{l-j}(n) \right) \theta^l + (b^r - c) \sum_{l=0}^{r-2} \left( \sum_{j=l+1}^{r-1} a_j(m) a_{l+r-j}(n) \right) \theta^l, \end{aligned}$$

which implies

$$a_l(m+n) = \sum_{j=0}^l a_j(m) a_{l-j}(n) + (b^r - c) \sum_{j=l+1}^{r-1} a_j(m) a_{l+r-j}(n) \quad (3)$$

for all  $0 \leq l \leq r-1$ . When  $l = r-1$ , the second summation in the equation (3) does not happen so that one has  $a_{r-1}(m+n) = \sum_{j=0}^{r-1} a_j(m)a_{r-1-j}(n)$ . This recurrence relation is summarized in Algorithm 2.

---

**Algorithm 2** RecurrenceRelation( $A, M$ )

---

**Input :** A coefficient vector  $A = (a_0, a_1, \dots, a_{r-1})$  of  $a = \sum_{i=0}^{r-1} a_i \theta^i \in \mathbb{F}_q[\theta]$  and  $M \in \mathbb{Z}^+$

**Output :** A coefficient vector of  $a^M \in \mathbb{F}_q[\theta]$

```

1: Write  $M = \sum M_i 2^i$  where  $M_i \in \{0, 1\}$ .
2:  $(B_0, B_1, \dots, B_{r-1}) \leftarrow (a_0, a_1, \dots, a_{r-1})$ 
3: for  $k$  from  $\lfloor \log M \rfloor - 1$  downto 0 do
4:    $(A_0, A_1, \dots, A_{r-1}) \leftarrow (B_0, B_1, \dots, B_{r-1})$ 
5:   for  $i$  from 0 to  $r-1$  do
6:      $B_i \leftarrow \sum_{j=0}^i A_j A_{i-j} + (b^r - c) \sum_{j=i+1}^{r-1} A_j A_{r+i-j}$ 
7:   if  $M_k = 1$  then
8:      $(A_0, A_1, \dots, A_{r-1}) \leftarrow (B_0, B_1, \dots, B_{r-1})$ 
9:     for  $i$  from 0 to  $r-1$  do
10:       $B_i \leftarrow \sum_{j=0}^i A_j a_{i-j} + (b^r - c) \sum_{j=i+1}^{r-1} A_j a_{r+i-j}$ 
11: return  $(B_0, \dots, B_{r-1})$ 

```

---

### 2.3 An improvement of K. S. Williams and K. Hardy

Williams and Hardy [20] improved the algorithm of H. C. Williams by reducing the loop length to  $\log q$  as follows. Write  $\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}}$  (where  $\alpha = b - \theta$ ) as

$$\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}} = E_1^{\frac{q-1}{r}} \cdot E_2, \quad (4)$$

where

$$E_1 = \alpha^{(q-1)^{r-2}}, \quad E_2 = \alpha^{\frac{q^r-1}{r(q-1)} - \frac{(q-1)^{r-1}}{r}}.$$

By noticing that the exponent  $\frac{q^r-1}{r(q-1)} - \frac{(q-1)^{r-1}}{r}$  of  $E_2$  is a polynomial of  $q$  with integer coefficients and using the binomial theorem, one has the following expression of  $E_1$  and  $E_2$  as

$$E_1 = \prod_{i=0}^{r-2} X_i \quad \text{with } X_i = (b - \omega^i \theta)^{(-1)^{r-i} \binom{r-2}{i}}, \quad (5)$$

$$E_2 = \prod_{i=1}^{r-1} Y_i \quad \text{with } Y_i = (b - \omega^{r-i-1} \theta)^{\frac{1 - (-1)^i \binom{r-1}{i}}{r}}. \quad (6)$$

Thus we have the following result of Williams and Hardy.

**Proposition 2.** [Williams-Hardy]

(1) Under same assumption as in Proposition 1,  $E_1^{\frac{q-1}{r}} \cdot E_2$  is an  $r$ -th root of  $c$ , where

$$E_1 = \alpha^{(q-1)^{r-2}}, \quad E_2 = \alpha^{\frac{q^r-1}{r(q-1)} - \frac{(q-1)^{r-1}}{r}}.$$

(2)  $E_1$  and  $E_2$  can be efficiently computed using the relations

$$E_1 = \prod_{i=0}^{r-2} (b - \omega^i \theta)^{(-1)^{r-i} \binom{r-2}{i}}, \quad E_2 = \prod_{i=1}^{r-1} (b - \omega^{r-i-1} \theta)^{\frac{1-(-1)^i \binom{r-1}{i}}{r}}.$$

---

**Algorithm 3** Williams-Hardy  $r$ -th root algorithm [20]

---

**Input :** An  $r$ -th power residue  $c$  in  $\mathbb{F}_q$

**Output :**  $x \in \mathbb{F}_q$  satisfying  $x^r = c$

- 1: **do** Choose a random  $b \in \mathbb{F}_q$  **until**  $b^r - c$  is not an  $r$ -th power residue.
  - 2:  $\omega \leftarrow (b^r - c)^{\frac{q-1}{r}}$ , where  $\theta^r = b^r - c$ .
  - 3:  $E_1 \leftarrow 1$ ,  $E_2 \leftarrow 1$
  - 4: **for**  $i$  from 1 to  $r-1$  **do**
  - 5:  $X_i \leftarrow (b - \omega^{i-1} \theta)^{(-1)^{r-i+1} \binom{r-2}{i-1}}$ ,  $Y_i \leftarrow (b - \omega^{r-i-1} \theta)^{\frac{1-(-1)^i \binom{r-1}{i}}{r}}$
  - 6:  $E_1 \leftarrow E_1 \cdot X_i$ ,  $E_2 \leftarrow E_2 \cdot Y_i$
  - 7:  $A \leftarrow$  coefficient vector of  $E_1$
  - 8:  $A \leftarrow \text{RecurrenceRelation}(A, \frac{q-1}{r})$
  - 9:  $E'_1 \leftarrow$  corresponding element of  $A$  in  $\mathbb{F}_q[\theta]$
  - 10:  $x \leftarrow E'_1 \cdot E_2$
  - 11: **return**  $x$
- 

The complexity of computing each of  $X_i$  in the equation (5) is of  $O(\log q) + O(r) + O(r^2 \log \binom{r-2}{i})$  multiplications in  $\mathbb{F}_q$ . Hence all  $X_i$  can be computed in  $O(r \log q + r^4)$   $\mathbb{F}_q$ -multiplications. Since the  $O(r)$  multiplications of all  $X_i$  ( $0 \leq i \leq r-2$ ) in  $\mathbb{F}_{q^r}$  need  $O(r^3)$  multiplications in  $\mathbb{F}_q$ , the total complexity of computing  $E_1$  (as a polynomial of  $\theta$  degree at most  $r-1$ ) is  $O(r \log q + r^4)$   $\mathbb{F}_q$ -multiplications. Similarly the complexity of computing  $E_2$  is also  $O(r \log q + r^4)$   $\mathbb{F}_q$ -multiplications. For a detailed explanation, see [20]. Since the exponentiation  $E_1^{\frac{q-1}{r}}$  (using the recurrence relation) needs  $O(r^2 \log \frac{q-1}{r}) = O(r^2 \log q)$  multiplications in  $\mathbb{F}_q$  and since the multiplication of two elements  $E_1^{\frac{q-1}{r}}$  and  $E_2$  needs  $O(r)$  multiplications in  $\mathbb{F}_q$  (because only the constant term of the  $\theta$  expansion is needed), the total cost of computing an  $r$ -th root of  $c$  using the algorithm of K. S. Williams and K. Hardy [20] is  $O(r^2 \log q + r^4)$ .

### 3 Our New $r$ -th Root Algorithm

In this section, we give an improved version of the Cipolla-Lehmer type algorithm by generalizing the method of [20]. Our new algorithm is applicable for all  $r > 1$  with  $q \equiv 1 \pmod{r}$ . Throughout this section, we assume that  $r$  is not necessarily a prime. Thus  $\omega = \theta^{q-1} = (b^r - c)^{\frac{q-1}{r}}$  may not be a primitive  $r$ -th root of unity even if  $b^r - c$  is not an  $r$ -th power in  $\mathbb{F}_q$ . Consequently a more stronger condition is needed for the primitivity of  $\omega$ . That is,  $\omega$  is a primitive  $r$ -th root of unity if and only if  $\omega^{\frac{r}{p}} \neq 1$  for every prime  $p|r$ , which holds if and only if  $(b^r - c)^{\frac{q-1}{p}} \neq 1$  for every prime  $p|r$ . From now on, we will assume that  $(b^r - c)^{\frac{q-1}{p}} \neq 1$  for every prime  $p|r$  and therefore  $\omega$  is a primitive  $r$ -th root of unity.

Let  $\alpha \in \mathbb{F}_{q^r}$ . Then, by extracting  $r$ -th roots from the following simple identity

$$\alpha^r \left( 1 \cdot \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+q^2+\dots+q^{r-2}} \right)^q = \left( 1 \cdot \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+q^2+\dots+q^{r-2}} \right) \alpha^{1+q+\dots+q^{r-1}},$$

one may expect that  $\alpha \left(1 \cdot \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+q^2+\dots+q^{r-2}}\right)^{\frac{q-1}{r}}$  equals  $\alpha^{\frac{1+q+\dots+q^{r-1}}{r}}$  up to  $r$ -th roots of unity. In fact, they are exactly the same element in  $\mathbb{F}_q$  and can be verified as follows;

$$\alpha^{\frac{1+q+\dots+q^{r-1}}{r}} = \alpha^{\frac{\sum_{i=0}^{r-1} q^i}{r}} = \alpha \cdot \alpha^{\frac{(\sum_{i=0}^{r-1} q^i) - r}{r}} \quad (7)$$

$$= \alpha \cdot \alpha^{\frac{\sum_{i=0}^{r-1} (q^i - 1)}{r}} = \alpha \cdot \alpha^{\frac{(q-1) \sum_{i=1}^{r-1} \sum_{j=0}^{i-1} q^j}{r}} \quad (8)$$

$$= \alpha \cdot \left( \alpha^{\sum_{i=1}^{r-1} \sum_{j=0}^{i-1} q^j} \right)^{\frac{q-1}{r}} \quad (9)$$

$$= \alpha \cdot \left(1 \cdot \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+q^2+\dots+q^{r-2}}\right)^{\frac{q-1}{r}}. \quad (10)$$

**Proposition 3.** [Main Theorem]

Let  $q \equiv 1 \pmod{r}$  with  $r > 1$  and let  $(b^r - c)^{\frac{q-1}{r}} \neq 1$  for all prime divisors  $p$  of  $r$ . Then letting  $\alpha = b - \theta$  where  $\theta^r = b^r - c$ ,

$$\alpha \cdot \left(1 \cdot \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+q^2+\dots+q^{r-2}}\right)^{\frac{q-1}{r}}$$

is an  $r$ -th root of  $c$ .

Based on the above simple result, we may present a new  $r$ -th root algorithm (Algorithm 4) of complexity  $O(r^2 \log q + r^3)$  with given information of the prime factors of  $r$ . It should be mentioned that our proposed algorithm is general in the sense that  $r$  can be any (composite) positive integer  $> 1$  satisfying  $q \equiv 1 \pmod{r}$ , while  $r$  was assumed to be an odd prime both in [19] and [20].

Both in [19] and [20],  $b$  was chosen so that  $\omega = (b^r - c)^{\frac{q-1}{r}} \neq 1$ , and since  $r$  is prime,  $\omega$  is automatically a primitive  $r$ -th root. This property guarantees the validity of the equation (1), namely

$$(b - \theta)(b - \omega\theta)(b - \omega^2\theta) \dots (b - \omega^{r-1}\theta) = b^r - \theta^r = c. \quad (11)$$

However if  $r$  is composite, then  $\omega = (b^r - c)^{\frac{q-1}{r}}$  is not a primitive  $r$ -th root in general. In fact, letting  $s > 1$  be the least positive integer satisfying  $\omega^s = 1$ , the degree of the irreducible polynomial of  $\theta$  (where  $\theta^r = b^r - c$ ) is  $s$  because

$$\theta^{q^s-1} = (\theta^{q-1})^{q^{s-1}+q^{s-2}+\dots+q+1} = \omega^{q^{s-1}+q^{s-2}+\dots+q+1} = \omega^s,$$

and one has

$$(b - \theta)(b - \omega\theta) \dots (b - \omega^{r-1}\theta) = \{(b - \theta)(b - \omega\theta) \dots (b - \omega^{s-1}\theta)\}_s^{\frac{r}{s}} = (b^s - \theta^s)_s^{\frac{r}{s}} \neq c \quad (12)$$

if  $s < r$ . Therefore the methods of [19] and [20] do not work for a composite  $r$  unless one assumes the primitivity of  $\omega$ .

Also, even if one assumes the primitivity of  $\omega = (b^r - c)^{\frac{q-1}{r}}$ , one still has some problems both in [19] and [20], which will be explained in the following remarks.

**Remark 1.** In [19],  $\alpha = b + \theta$  was used (instead of  $b - \theta$ ) under the assumption of  $\theta^r = c - b^r$  with  $(c - b^r)^{\frac{q-1}{r}} \neq 1$ . If we choose  $\alpha = b + \theta$  following [19], then we get

$$(b + \theta)(b + \omega\theta) \dots (b + \omega^{r-1}\theta) = b^r - (-\theta)^r = b^r + (-1)^{r+1}\theta^r. \quad (13)$$

Therefore if  $r$  is odd prime (as was originally assumed in [19]), one has  $b^r + \theta^r = c$  and the  $r$ -th root algorithm is essentially same to the case  $\alpha = b - \theta$ . However when  $r$  is even (for example, when  $r = 2$ ), the original method in [19] cannot be used because  $b^r + (-1)^{r+1}\theta^r = b^r - \theta^r \neq c$ .

---

**Algorithm 4** Our new  $r$ -th root algorithm

---

**Input :** An  $r$ -th power residue  $c$  in  $\mathbb{F}_q$

**Output :**  $x \in \mathbb{F}_q$  satisfying  $x^r = c$

```
1: do Choose a random  $b \in \mathbb{F}_q$  until  $(b^r - c)^{\frac{q-1}{r}}$  is a primitive  $r$ -th root of unity.
2:  $\omega \leftarrow (b^r - c)^{\frac{q-1}{r}}$ ,  $\alpha \leftarrow b - \theta$  where  $\theta^r = b^r - c$ .
3:  $P \leftarrow \alpha, A \leftarrow \alpha, W \leftarrow 1$ 
4: for  $i = 1$  to  $r - 2$  do           //  $A, P \in \mathbb{F}_q[\theta]$  and  $W \in \mathbb{F}_q$  //
5:    $W \leftarrow W\omega, V \leftarrow b - W\theta$  //  $W = \omega^i, V = b - \omega^i\theta = \alpha^{q^i}$  //
6:    $A \leftarrow AV, P \leftarrow PA$        //  $A = \alpha^{1+q+\dots+q^i}, P = \alpha \cdot \alpha^{1+q} \dots \alpha^{1+q+\dots+q^i}$  //
7:  $B \leftarrow$  coefficient vector of  $P$ 
8:  $B \leftarrow \text{RecurrenceRelation}(B, \frac{q-1}{r})$ 
9:  $P \leftarrow$  corresponding element of  $B$  in  $\mathbb{F}_q[\theta]$ 
10:  $x \leftarrow \alpha \cdot P$            //  $x \in \mathbb{F}_q$  //
11: return  $x$ 
```

---

**Remark 2.** The algorithm in [20] needs  $E_1$  and  $E_2$  satisfying  $\alpha^{\frac{\sum_{j=0}^{r-1} q^j}{r}} = E_1^{\frac{q-1}{r}} \cdot E_2$ . However for composite  $r$ ,  $E_2$  cannot be well-defined in some cases, since the exponent  $\frac{1 - (-1)^i \binom{r-1}{i}}{r}$  in the equation (6) is not an integer in general. That is, the property  $(-1)^i \binom{r-1}{i} \equiv 1 \pmod{r}$  only holds when  $r$  is prime. Therefore the algorithm in [20] fails to give the answer when  $r$  is composite such as  $r = 4, 6, 9, \dots$  (i.e., when  $r = 4$ , one has  $E_2 = \alpha^{q^2 - \frac{1}{2}q}$  so the coefficient  $\frac{1}{2}$  of  $q$  in the exponent is not an integer and one cannot compute  $E_2$ .) The problem of  $E_2$  being undefined is unavoidable even if one assumes the primitivity of  $\omega$ .

## 4 Complexity Analysis and Comparison

### 4.1 Complexity analysis

An initial step of the proposed algorithm requires one to find a primitive  $r$ -th root  $\omega$  in  $\mathbb{F}_q$ . When  $r$  is prime, one only needs to find  $b$  satisfying  $\omega = (b^r - c)^{\frac{q-1}{r}} \neq 0, 1$  and the probability that a random  $b$  satisfies the required property is  $\frac{1}{r} + O(q^{-\frac{1}{4}})$  ([20] pp.480) under the assumption of  $r \leq q^{\frac{1}{4}}$ . When  $r$  is composite, one further needs to check whether  $\omega^{\frac{r}{p}} \neq 1$  for every prime divisor  $p$  of  $r$ . Since the complexity estimation  $O(r^3 \log q)$  in [19] and  $O(r^2 \log q + r^4)$  in [20] still hold if one assumes that a primitive root  $\omega = (b^r - c)^{\frac{q-1}{r}}$  is already given, we will also assume that a primitive root  $\omega$  is given in our estimation for a fair comparison.

At each  $i$ -th step of the for-loop of our proposed algorithm, step 5 needs 1  $\mathbb{F}_q$  multiplication. In step 6, the computation  $AV$  needs 1  $\mathbb{F}_{q^r}$  multiplication which, in fact, can be executed with  $2r$   $\mathbb{F}_q$  multiplications because  $V = b - \omega^i\theta$  is linear in  $\theta$ . The computation  $PA$  needs 1  $\mathbb{F}_{q^r}$  multiplication which can be executed with  $r^2$   $\mathbb{F}_q$  multiplications. Therefore, at the end of the for-loop, one needs at most  $(r-2)(1+2r+r^2) < (r+1)^3$   $\mathbb{F}_q$  multiplications (of order  $O(r^3)$ ). Since the exponentiation  $P^{\frac{q-1}{r}}$  (in steps 7-9) needs  $O(r^2 \log q)$   $\mathbb{F}_q$  multiplications, the total cost of our proposed algorithm is  $O(r^3 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ . On the other hand, the cost of Algorithm 1 [19] is  $O(r^2 \log \frac{q^r-1}{r}) = O(r^3 \log q)$ , and the cost of Algorithm 3 [20] is  $O(r^4 + r^2 \log q)$  where  $O(r^4)$  comes from the cost of computing  $E_1$  and  $E_2$  in steps 4-6 of

Algorithm 3. The theoretical estimation shows that our proposed algorithm is better than Algorithm 3 as  $r$  gets larger.

Finally, when  $r = 2$ , the for-loop can be omitted in our algorithm so that one only needs to compute  $P \cdot P^{\frac{q-1}{2}}$  which is exactly same to the original Cipolla-Lehmer algorithm.

## 4.2 Implementation results

Table 1 shows the implementation results using SAGE of the above mentioned two algorithms and our proposed one. The implementation was performed on Intel Core i7-4770 3.40GHz with 8GB memory.

Table 1: Running time (in seconds) for  $r$ -th root algorithms

$r$	3	4	43	101	211
Algorithm 1 [19]	0.467	fail	2026.962	Interr.	Interr.
Algorithm 3 [20]	0.254	fail	53.849	535.043	3956.433
Our proposed algorithm	0.253	0.355	48.359	256.601	1098.401

For convenience, we used prime fields  $\mathbb{F}_p$  with size about 2000 bits. Average timings of the  $r$ -th root computations for 5 different inputs of  $r$ -th power residue  $c \in \mathbb{F}_p$  are computed for the primes  $r = 3, 43, 101, 211$ . As one can see in the table, our proposed algorithm performs better than the algorithms in [19] and [20]. The table also shows that our algorithm gets dramatically faster than other algorithms as  $r$  gets larger. For example, when  $r = 101$ , our algorithm is roughly 2 times faster than Algorithm 3, and when  $r = 211$ , our algorithm is 4 times faster than Algorithm 3. For  $r = 101, 211$ , the SAGE computation were interrupted after 3 hours for Algorithm 1.

## 5 Conclusions

We proposed a new Cipolla-Lehmer type algorithm for  $r$ -th root extractions in  $\mathbb{F}_q$ . Our algorithm has the complexity of  $O(r^3 + r^2 \log q)$  multiplications in  $\mathbb{F}_q$ , which improves the previous results of  $O(r^3 \log q)$  in [19] and of  $O(r^4 + r^2 \log q)$  in [20]. Our algorithm is applicable for any integer  $r > 1$ , whereas the previous algorithms are effective only for odd prime  $r$ . Software implementations via SAGE also show that our proposed algorithm is consistently faster than the previously proposed algorithms, and becomes much more effective as  $r$  gets larger.

## References

- [1] L. Adleman, K. Manders and G. Miller, *On taking roots in finite fields*, Proceeding of 18th IEEE Symposium on Foundations on Computer Science (FOCS), pp. 175-177, 1977.
- [2] A. O. L. Atkin, *Probabilistic primality testing*, summary by F. Morain, Inria Research Report 1779, pp.159-163, 1992.
- [3] D. Bernstein, *Faster square root in annoying finite field*, preprint, Available from <http://cr.yp.to/papers/sqroot.pdf>, 2001.



- [4] Z. Cao, Q. Sha, and X. Fan, *Adleman-Manders-Miller root extraction method revisited*, preprint, available from <http://arxiv.org/abs/1111.4877>, 2011.
- [5] G. H. Cho, N. Koo, E. Ha, and S. Kwon, *New cube root algorithm based on third order linear recurrence relation in finite field*, to appear in Designs, Codes and Cryptography, available from <http://link.springer.com/article/10.1007%2Fs10623-013-9910-8>.
- [6] G. H. Cho, N. Koo, E. Ha, and S. Kwon, *Trace expression of  $r$ -th root over finite field*, preprint, available from <http://eprint.iacr.org/2013/041.pdf>, 2013.
- [7] M. Cipolla, *Un metodo per la risoluzione della congruenza di secondo grado*, Rendiconto dell'Accademia Scienze Fisiche e Matematiche, Napoli, Ser.3, Vol. IX, pp. 154-163, 1903.
- [8] J. Doliskani and E. Schost, *Taking roots over high extensions of finite fields*, Mathematics of Computation, Vol.83, pp. 435-446, 2014.
- [9] G. Gong and L. Harn, *Public key cryptosystems based on cubic finite field extensions*, IEEE Transactions on Information Theory, Vol.45, pp. 2601-2605, 1999.
- [10] F. Kong, Z. Cai, J. Yu, and D. Li, *Improved generalized Atkin algorithm for computing square roots in finite fields*, Information Processing Letters, Vol. 98, no. 1, pp. 1-5, 2006.
- [11] D. H. Lehmer, *Computer technology applied to the theory of numbers*, Studies in Number Theory, Englewood Cliffs, NJ: Prentice-Hall, pp. 117-151, 1969.
- [12] R. Lidl and H. Niederreiter, *Finite Fields*, Cambridge University Press, 1997.
- [13] A. J. Menezes, I. F. Blake, X. Gao, R. C. Mullin, S. A. Vanstone, and T. Yaghoobian, *Applications of Finite Fields*, Springer, 1992.
- [14] S. Müller, *On the computation of square roots in finite fields*, Designs, Codes and Cryptography, Vol.31, pp. 301-312, 2004.
- [15] NIST, *Digital Signature Standard*, Federal Information Processing Standard 186-3, National Institute of Standards and Technology, Available from <http://csrc.nist.gov/publications/fips/>, 2000.
- [16] D. Shanks, *Five number-theoretic algorithms*, Proceeding of 2nd Manitoba Conference on Numerical Mathematics, Manitoba, Canada, pp. 51-70, 1972.
- [17] I. Shparlinski, *Finite Fields: Theory and Computation*, Springer, 1999.
- [18] A. Tonelli, *Bemerkung über die Auflösung quadratischer Congruenzen*, Göttinger Nachrichten, pp. 344-346, 1891.
- [19] H. C. Williams, *Some algorithm for solving  $x^q \equiv N \pmod{p}$* , Proc. 3rd Southeastern Conf. on Combinatorics, Graph Theory, and Computing (Florida Atlantic University), pp. 451-462, 1972.
- [20] K. S. Williams and K. Hardy, *A refinement of H. C. Williams'  $q$ th root algorithm*, Mathematics of Computation, Vol.61, pp. 475-483, 1993.