

The Effects of Non Standard Neutrino Interactions (NSIs) in $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}, B^+ \rightarrow \pi^+ \nu \bar{\nu}, B^+ \rightarrow K^{*+} \nu \bar{\nu}$ and $B \rightarrow X_s \nu \bar{\nu}$

Shakeel Mahmood⁽¹⁾; Farida Tahir; Azeem Mir
*Comsats Institute of Information Technology,
Department of Physics, Park Road, Chek Shazad, Islamabad*

⁽¹⁾shakeel_mahmood@hotmail.com

Abstract

We study the rare decays $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}, B^+ \rightarrow \pi^+ \nu \bar{\nu}, B^+ \rightarrow K^{*+} \nu \bar{\nu}$ and $B \rightarrow X_s \nu \bar{\nu}$ for the search of NSIs. We want to constraint the NSIs by using these reactions. We show that there is a strong dependence of these reactions on new physics free parameter $\epsilon_{\tau\tau}^{QL}$, where $Q = u, c, t$. We include second and third generation of quarks in the loop for these decays. We show that the $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$ is providing very precise bounds as compared to all other semileptonic decays, having neutrinos in their final state. We further show that the interference between standard model and NSIs is giving dominant contribution for $B^+ \rightarrow \pi^+ \nu \bar{\nu}, B^+ \rightarrow K^{*+} \nu \bar{\nu}$ and $B \rightarrow X_s \nu \bar{\nu}$. We point out that the constraints for $\epsilon_{\tau\tau}^{cL}$ and $\epsilon_{\tau\tau}^{tL}$ are more precise as compared to $\epsilon_{\tau\tau}^{uL}$. The analysis of $B^+ \rightarrow \pi^+ \nu \bar{\nu}, B^+ \rightarrow K^{*+} \nu \bar{\nu}$ and $B \rightarrow X_s \nu \bar{\nu}$, provide us that the u quark induced Br of NSIs are giving very very small contribution. We also compare these decays to the decays of charm and kaons having neutrinos in the final state.

Keywords: NSIs; rare decays; B decays.

PACS numbers: 12.60.-i, 13.15.+g, 13.20.-v

1 Introduction

After the remarkable discovery of Higgs by the ATLAS [1] and CMS [2] collaboration and confirmation in [3] that it is Standard Model (SM) higgs, one important question arises. Is there any room for new physics (NP) beyond SM? No doubt SM predictions have been verified experimentally to the highest level of precision [4][5]. But, along with other limitations, SM lacks any explanation for a possible pattern for particle mass, known as mass hierarchy problem. SM can not predict top quark mass without experimental evidence. The experiments on B meson [6][7][8] are also giving some cracks in standard model.

We are yet unable to explain dark matter and matter anti-matter asymmetry. Gravity is not included in the SM. Theoretically SM is thought to be unsatisfactory and there can be some new particles as well as new interactions. It has been believed that standard model is a low energy approximation of more general theory. So, many theoretical extensions of SM has been presented. But, so far, the only concrete evidence against it has been provided by the neutrino oscillations [9] [10][11][12][13][14][15]. To explore NP the study of mesonic rare decays involving neutrinos in final state, can be interesting. These decays proceeds through flavor changing neutral currents (FCNC), highly suppressed [16] due to GIM mechanism [17] and occur at loop level [18][19]. The discrepancies between experiments and theory (SM) for such reactions provide us an excellent window towards NP. New particles can be added in the loops to improve theory or we can have new interactions. So, FCNC reactions involving neutrinos in the final state can be interesting.

Theoretically, $B^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$ thought to be more clean than $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ because it has only top quark contribution and no contribution from charm sector. Non standard neutrino interactions (NSIs) of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $D_s^+ \rightarrow D^+ \nu \bar{\nu}$ are studied in [20] and [21] respectively and constrained are found for $\epsilon_{\tau\tau}^{uL}$. The NSIs constraints for three generations of quarks are given in [22] As all of these have same loop structure, so, similar thing should happen to $B^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$. Two more loop level processes useful for the search of new physics are inclusive $B \rightarrow X_s \nu \bar{\nu}$ and exclusive $B^+ \rightarrow K^{*+} \nu \bar{\nu}$ due to their theoretical cleanliness [23].

In this paper we the scheme of study is as: first of all we give experimental status of $B^+ \rightarrow \pi^+ \nu \bar{\nu}$, $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$, $B^+ \rightarrow K^{*+} \nu \bar{\nu}$ and $B \rightarrow X_s \nu \bar{\nu}$ then we revise the SM contribution of these reaction. Next, we study these reactions in NSIs with u quark in the loop which is the usual case of NSIs and obtain the constraints $\epsilon_{\tau\tau}^{uL}$. Then we modify the operators for c and t quarks. and find out the constraints $\epsilon_{\tau\tau}^{cL}$ and $\epsilon_{\tau\tau}^{tL}$. We compare these constraints among the three generations. We also compare these constraints to the constraints of same type of reactions from D and K decays. Then discussion and results are provided and conclusion is given at the end.

2 Experimental Status of $B^+ \rightarrow \pi^+ \nu \bar{\nu}$ Decay

B decays are being studied in the detectors like CLEO, CDF, BaBar, Belle, ALEPH collaborations and LHCb but the decays involving neutrinos in the final state will be tested at super B factories in experimentally clean environment. The detection of $B^+ \rightarrow \pi^+ \nu \bar{\nu}$ is a hard task and currently we have only experimental bound for this reaction but this will be in the range of super B factories. The experimental bound is given in the Table 1 along with their SM prediction. Current experimental bounds for this are $< 9.8 \times 10^{-5}$ but our experience with $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ guide us that the experimental value should be of the order of 10^{-7} . It means we are not discarding the SM values but just searching for the small room for new physics. Experimental value for $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$ is

Figure 1: SM b decays to d netrino antinetrino

$< 2.6 \times 10^{-8}$ [36]. $B^+ \rightarrow K^{*+} \nu \bar{\nu}$ is easy to measure experimentally because we have vector particle in the final state, so polarization does matter and its latest bound is $< 4 \times 10^{-5}$.

Inclusive process $B \rightarrow X_s \nu \bar{\nu}$ are very difficult to observe experimentally because we need to tag all the particles involve in X_s along with missing neutrinos. The limit available till to date is $< 64 \times 10^{-5}$, given in table 1 with reference.

2.1 $M \rightarrow M' \nu \bar{\nu}$ Decays in the Standard Model

Here mass of $M >$ mass of M' and both are representing the mesons. The SM calculation can be divided into two categories, short distance and long distance..This can be found from [24] [25] [26] and [27] that the dominant contribution for $M \rightarrow M' \nu \bar{\nu}$ comes from short distance because long distance contribution is 10^{-3} less than short distance, if the quark level process is $b \rightarrow d \nu \bar{\nu}$ or $s \rightarrow d \nu \bar{\nu}$. The quark level process for our decay is $b \rightarrow d \nu \bar{\nu}$ which can be represented by the feynman diagrams shown in figure 1.

In such reactions we can easily separate hadronic interactions from leptonic interaction. For B decays the dominant contribution comes from the short distance just like K decays and we use perturbation theory due to asymptotic freedom. The effective Hamiltonian for such reactions quark level reaction will be

$$H_{eff}^{SM} = \frac{G_F}{\sqrt{2}} \frac{\alpha_{em}}{2\pi \sin^2 \theta_W} \sum_{\alpha, \beta=e, \mu, \tau} V_{tb}^* V_{td} X(x_t) \times (\bar{d}b)_{V-A} (\nu_\alpha \bar{\nu}_\beta)_{V-A}$$

where $V - A$ in the subscript represents the vector and axial vector current

respectively. For such reactions charm quark contribution in the loop is negligible in contrast to K decay due to smallness of off diagonal CKM element and $X(x_t)$ is the loop integral of top-quark exchange [19]. For this reaction we have two penguin and one box diagram [30] and sum of all give the contribution

$$X(x_t) = \eta_X \frac{x_t}{8} \left[\frac{x_t + 2}{x_t - 1} + \frac{3x_t - 6}{(x_t - 1)^2} \ln x_t \right]$$

Here $x_t = \frac{m_t^2}{m_w^2}$ and $\eta_X = 0.985$ is QCD small distance correction. By using above Hamiltonian we can obtain Br as

$$Br(B^+ \rightarrow \pi^+ \nu \bar{\nu})_{SM} = r_{iso} \frac{3\alpha_{em}^2}{|V_{ub}|^2 2\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{td} X(x_t)|^2 Br(B^+ \rightarrow \pi^0 l^+ \nu_l)$$

$r_{iso} \simeq 0.94$ is the isospin breaking effect for B. It is discussed for K mesons in [32] which depends on at least three things (1) mass effect (2) a suppression of about 4% in neutral form factor comes from $\eta - \pi$ mixing and (3) about 2% suppression due to absence of log leading correction.

The reaction $B^+ \rightarrow K^{*+} \bar{\nu} \nu$, proceed through quark level process $\bar{b} \rightarrow \bar{s} \nu \bar{\nu}$ and the effective Hamiltonian is same except to replace d with s . Here we do not have a tree level reaction for normalization so we have to use $B^+ \rightarrow \rho^0 l^+ \nu_l$

$$Br(B^+ \rightarrow K^{*+} \nu \bar{\nu})_{SM} = r_{iso} \frac{3\alpha_{em}^2}{|V_{ub}|^2 2\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{ts} X(x_t)|^2 Br(B^+ \rightarrow \rho^0 l^+ \nu_l)$$

For $B \rightarrow X_s \nu \bar{\nu}$ again the effective Hamiltonian is same except to replace d with s , and we do not have a tree level process like $B^+ \rightarrow \pi^0 l^+ \nu_l$. So we have to normalize with the process $B \rightarrow X_c \nu \bar{\nu}$ and due to different phase spaces for X_s and X_c , we have to include other factors. The Br will be

$$Br(B \rightarrow X_s \nu \bar{\nu})_{SM} = \frac{3\alpha_{em}^2}{4\pi^2 \sin^4 \theta_W} \left| \frac{V_{ts} V_{tb} X(x_t)}{V_{cb}} \right|^2 \frac{\bar{\eta}}{f(z) \kappa(z)} Br(B \rightarrow X_c l \nu_l)$$

where $f(z) = 1 - 8z + 8z^3 - z^4 - 12z^2 \ln(z)$ with $z = \frac{m_c^2}{m_b^2}$

and $\kappa(z) = 0.88$, $\eta = \kappa(0) = 0.83$.

A useful discussion can about the factors can be found in [27] and [30]. With the latest values of the constants we have the Br

$$Br(B \rightarrow X_c \nu \bar{\nu})_{SM} = 3.6 \times 10^{-5}$$

$K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$ along with $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ is thought to be theoretically clean reactions for the search of new physics. The effective Hamiltonian for $K_L^0 \rightarrow \pi^0 \nu \bar{\nu}$ can be written as

$$H_{eff}^{SM} = \frac{G_F}{\sqrt{2}} \frac{\alpha_{em}}{2\pi \sin^2 \theta_W} \sum_{\alpha, \beta=e, \mu, \tau} V_{ts}^* V_{td} X(x_t) \times (\bar{d}s)_{V-A} (\nu_\alpha \bar{\nu}_\beta)_{V-A}$$

Figure 2: NSIs b decays to d neutrino antineutrino

and the Br can be extracted by normalizing with tree level reaction, so that all the hadronic uncertainties are absorbed

$$Br(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})_{SM} = R_{iso} \frac{3\alpha_{em}^2}{|V_{ub}|^2 16\pi^2 \sin^4 \theta_W} \frac{\tau(K_L^0)}{\tau(K^+)} |V_{ts}^* V_{td} X(x_t)|^2 Br(K^+ \rightarrow \pi^0 e^+ \nu_e)$$

Here isospin symmetry is exploited as

$$\langle \pi^0 | (\bar{d}s)_{V-A} | \bar{K}^0 \rangle = \langle \pi^0 | (\bar{s}u)_{V-A} | K^+ \rangle$$

and isospin breaking effect is $R_{iso} = 0.94$, and other values from [36] are used to obtained the Br

$$Br(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})_{SM} = 2.06 \times 10^{-11}$$

2.2 Model Independent Approach

The NSI for the process is shown by the Fig 2 and represented by

$$H_{eff}^{NSI} = \frac{G_F}{\sqrt{2}} (V_{tb}^* V_{tq}) \frac{\alpha_{em}}{4\pi \sin^2 \theta_W} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w} \times (\nu_\alpha \bar{\nu}_\beta)_{V-A} (\bar{q}b)_{V-A}$$

We use $V_{ub} = (4.15 \pm 0.49) \times 10^{-3}$ and $Br(B^+ \rightarrow \pi^0 l^+ \nu_l) = (7.78 \pm 0.28) \times 10^{-5}$ to find out Br

$$Br(B^+ \rightarrow \pi^+ \bar{\nu} \nu)_{NSI} = r_{iso} \frac{\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{ub}^* V_{ud} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}|^2 Br(B^+ \rightarrow \pi^0 l^+ \nu_l)$$

Although the current experimental results of B decays are narrowing the gap between theory and experiments but when we will get more precise experimental data than we will need more accurate theoretical results. With the assumption that experiments will give us the value of 10^{-7} , we can constraint the NSIs from this reaction. As α and β can be any lepton we take them as τ , because for other leptons we have already more precise constraints [33].

Similarly NSIs Br of $B^+ \rightarrow K^{*+} \bar{\nu} \nu$ can be found as

$$Br(B^+ \rightarrow K^{*+} \bar{\nu} \nu)_{NSI} = \frac{\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{ub}^* V_{us} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}|^2 Br(B^+ \rightarrow \rho^0 l^+ \nu_l)$$

For $B \rightarrow X_s \nu \bar{\nu}$ NSIs Br

$$Br(B \rightarrow X_c \nu \bar{\nu})_{NSIs} = \frac{\alpha_{em}^2}{16\pi^2 \sin^4 \theta_W} \left| \frac{V_{us} V_{ub}}{V_{cb}} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w} \right|^2 \frac{\bar{\eta}}{f(z)\kappa(z)} Br(B \rightarrow X_c l \nu_l)$$

$$Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{NSI} = r_{iso} \frac{\alpha_{em}^2}{|V_{ub}|^2 16\pi^2 \sin^4 \theta_W} |V_{ub}^* V_{ud} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}|^2 Br(K^+ \rightarrow \pi^0 e^+ \nu_e)$$

2.3 Inteferece between Standard Model and NSIs

For all above decays the dominant contribution is coming from the interference between the SM and NSIs. The Br of interference are obtained as

$$Br(B^+ \rightarrow \pi^+ \bar{\nu} \nu)_{Int\,er\,f\,e\,r\,e\,n\,c\,e} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{td} X(x_t) V_{ub}^* V_{ud} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \pi^0 l^+ \nu_l)$$

$$Br(B^+ \rightarrow K^{*+} \bar{\nu} \nu)_{Int\,er\,f\,e\,r\,e\,n\,c\,e} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{ts} X(x_t) V_{ub}^* V_{us} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \rho^0 l^+ \nu_l)$$

$$Br(B \rightarrow X_c \nu \bar{\nu})_{Int\,er\,f\,e\,r\,e\,n\,c\,e} = \frac{\alpha_{em}^2}{16\pi^2 \sin^4 \theta_W |V_{cb}|} |V_{ts} V_{tb} X(x_t) V_{us} V_{ub} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| \frac{\bar{\eta}}{f(z)\kappa(z)} Br(B \rightarrow X_c l \nu_l)$$

$$Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{Int\,er\,f\,e\,r\,e\,n\,c\,e} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 16\pi^2 \sin^4 \theta_W} |V_{ts}^* V_{td} X(x_t) V_{ub}^* V_{ud} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(K^+ \rightarrow \pi^0 e^+ \nu_e)$$

The contribution from interference for B decays is is so large as compared to the NSIs that NSIs effects can easily be ignored. So that the bounds on the constraints are obtained from the interference only. But for the K_L^0 the $Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{Int\,er\,f\,e\,r\,e\,n\,c\,e}$ is 10^{-3} less than the $Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{NSIs}$, so the contribution is ignored. The numerical values are given in table 1 and 2.

2.3.1 c-quark in the Loop

For c-quark we have to modify the operators as and obtain the constraints

$$Br(B^+ \rightarrow \pi^+ \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{td} X(x_t) V_{cb}^* V_{cd} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \pi^0 l^+ \nu_l)$$

$$\epsilon_{\tau\tau}^{cL} \leq 1.5$$

$$Br(B^+ \rightarrow K^{*+} \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{ts} X(x_t) V_{cb}^* V_{cs} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \rho^0 l^+ \nu_l)$$

$$\epsilon_{\tau\tau}^{cL} \leq 1.5$$

For $B \rightarrow X_s \nu \bar{\nu}$ the Br and constraints are

$$Br(B \rightarrow X_c \nu \bar{\nu})_{Interference} = \frac{\alpha_{em}^2}{16\pi^2 \sin^4 \theta_W |V_{cb}|} |V_{ts} V_{tb} X(x_t) V_{cs} V_{cb} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| \frac{\bar{\eta}}{f(z)\kappa(z)} Br(B \rightarrow X_c l \nu_l)$$

$$\epsilon_{\tau\tau}^{cL} \leq 0.8$$

$$Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 16\pi^2 \sin^4 \theta_W} |V_{ts}^* V_{td} X(x_t) V_{cb}^* V_{cd} \epsilon_{\alpha\beta}^{cL} \ln \frac{\Lambda}{m_w}| Br(K^+ \rightarrow \pi^0 e^+ \nu_e)$$

2.3.2 t-quark in the Loop

With t-quark we have following operator and constraint

$$Br(B^+ \rightarrow \pi^+ \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{td} X(x_t) V_{tb}^* V_{td} \epsilon_{\alpha\beta}^{tL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \pi^0 l^+ \nu_l)$$

$$\epsilon_{\tau\tau}^{tL} \leq 1.5$$

$$Br(B^+ \rightarrow K^{*+} \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 8\pi^2 \sin^4 \theta_W} |V_{tb}^* V_{ts} X(x_t) V_{tb}^* V_{ts} \epsilon_{\alpha\beta}^{tL} \ln \frac{\Lambda}{m_w}| Br(B^+ \rightarrow \rho^0 l^+ \nu_l)$$

$$\epsilon_{\tau\tau}^{tL} \leq 1.5$$

For $B \rightarrow X_s \nu \bar{\nu}$ the Br and constraints are

$$Br(B \rightarrow X_c \nu \bar{\nu})_{Interference} = \frac{\alpha_{em}^2}{16\pi^2 \sin^4 \theta_W |V_{cb}|} |V_{ts} V_{tb} X(x_t) V_{ts} V_{tb} \epsilon_{\alpha\beta}^{tL} \ln \frac{\Lambda}{m_w}| \frac{\bar{\eta}}{f(z)\kappa(z)} Br(B \rightarrow X_c l \nu_l)$$

$$\epsilon_{\tau\tau}^{tL} \leq 0.8$$

$$Br(K_L^0 \rightarrow \pi^0 \bar{\nu} \nu)_{Interference} = r_{iso} \frac{2\alpha_{em}^2}{|V_{ub}|^2 16\pi^2 \sin^4 \theta_W} |V_{ts}^* V_{td} X(x_t) V_{tb}^* V_{td} \epsilon_{\alpha\beta}^{uL} \ln \frac{\Lambda}{m_w}| Br(K^+ \rightarrow \pi^0 e^+ \nu_e)$$

Reaction	Theoretical	Experimental	NSIs with u	NSIs with c	NSIs with t
$B^+ \rightarrow \pi^+ \bar{\nu} \nu$ $(u\bar{b}) \rightarrow (\bar{d}u) \bar{\nu} \nu$	1.5×10^{-7} [35]	$< 9.8 \times 10^{-5}$ [36]	3×10^{-8} $\epsilon_{\tau\tau}^{uL} \leq 1.5$	1×10^{-7} $\epsilon_{\tau\tau}^{cL} \leq 1.5$	1×10^{-7} $\epsilon_{\tau\tau}^{tL} \leq 1.5$
$B \rightarrow X_s \bar{\nu} \nu$ $b \rightarrow s \bar{\nu} \nu$	3.6×10^{-5}	$< 64 \times 10^{-5}$ [37]	$\sim 10^{-8}$ $\epsilon_{\tau\tau}^{uL} \leq 0.8$	1×10^{-5} $\epsilon_{\tau\tau}^{cL} \leq 0.8$	1×10^{-5} $\epsilon_{\tau\tau}^{tL} \leq 0.8$
$B^+ \rightarrow K^{*+} \bar{\nu} \nu$ $u\bar{b} \rightarrow (u\bar{s}) \bar{\nu} \nu$	3.56×10^{-6}	$< 4 \times 10^{-5}$	$\sim 10^{-8}$ $\epsilon_{\tau\tau}^{uL} \leq 1.5$	1.5×10^{-6} $\epsilon_{\tau\tau}^{cL} \leq 1.5$	1.5×10^{-6} $\epsilon_{\tau\tau}^{tL} \leq 1.5$

Table 1: Comparison of the constraints for B decays

3 Discussion and results

We study three processes $B^+ \rightarrow \pi^+ \bar{\nu} \nu$, $B^+ \rightarrow K^{*+} \bar{\nu} \nu$ (exclusive) and $B \rightarrow X_s \bar{\nu} \nu$ (inclusive). which are theoretically clean processes. So, these are ideal for the search of new physics. Very high Br of $B^+ \rightarrow \pi^+ \bar{\nu} \nu$ and $B^+ \rightarrow K^{*+} \bar{\nu} \nu$ making it very attractive for the experimentalists too. Although, $B \rightarrow X_s \bar{\nu} \nu$ is very difficult to detect but it is much clean as compared to any other rare decay that's why it is studied for the search of new physics. The results are summarized in table 1 and plots are provided in figures 3, 4,5,6,7 and 8 to make the comparison more clear. The constraints on NSIs with the decays of charm and kaon rare decays involving neutrinos in their final states are calculated in [21][22][34], which give very precise constraints $O(10^{-2})$ for the up type quarks. For the charm and kaons the interference between the standard model and NSIs is very small, $O(10^{-3})$ less than $NSIs$. But, for the case of B rare decays having two neutrinos in their final state the dominant contribution is coming from the interference. One more difference is that for charm and kaons the Br of first and second generation are giving higher values, but, for B the contribution from second and third generation is providing leading contribution as compared to the first generation. As for as the comparison with in the generation is concerned we are getting more precise constraints with second generation of quarks (c quark) for B , D and K rare decays having neutrino in final state. But, the constraints from B decays are less precise for u -quark than D and K decays. It is contrary to the usual NSIs, in which the dominate contribution comes from u -quark induced processes. The rare B decays will be in the range of much clean environment of the B factories. That's why the analysis of such reaction is very important for the new physics. We analysis $K_L^0 \rightarrow \pi^0 \bar{\nu} \nu$ for the study of NSIs with three generations of quarks which provides even be more precise constraints as compared to D and K decays of same type. The constraint for $K_L^0 \rightarrow \pi^0 \bar{\nu} \nu$ are $O(10^{-3})$ for all the generations but the Br for third generations is very small and can be ignored. This is shown in figures 9,10 and 11.

Figure 3: u quark induced NSIs of B+ Decays to Pi+, neutrino and antineutrino

Figure 4: c quark induced NSIs of B+ Decays to Pi+, neutrino and antineutrino

Reaction	Theoretical	Experimental	NSIs with u	NSIs with c	NSIs with t
$K_L^0 \longrightarrow \pi^0 \bar{\nu} \nu$ $(u\bar{b}) \longrightarrow (\bar{d}u) \bar{\nu} \nu$	2.06×10^{-11}	$< 2.6 \times 10^{-8}$ [36]	1.4×10^{-11} $\epsilon_{\tau\tau}^{uL} \leq O(10^{-3})$	1.5×10^{-11} $\epsilon_{\tau\tau}^{cL} \leq O(10^{-3})$	3×10^{-17} $\epsilon_{\tau\tau}^{tL} \leq O(10^{-3})$

Table 2: Comparison of the constraints for KL

Figure 5: t quark induced NSIs of B^+ Decays to Π^+ , neutrino and antineutrino

Figure 6: u quark induced NSIs of B Decays to X_c , neutrino and antineutrino

Figure 7: c quark induced NSIs of B Decays to Xc , neutrino and antineutrino

Figure 8: t quark induced NSIs of B Decays to Xc , neutrino and antineutrino

Figure 9: u quark induced NSIs of KL Zero Decay

Figure 10: c quark induced NSIs of KL zero Decay

Figure 11: t quark induced NSIs of KL Zero Decay

4 Conclusion

We have studied $B^+ \rightarrow \pi^+ \bar{\nu} \nu$, $B^+ \rightarrow K^{*+} \bar{\nu} \nu$, $B \rightarrow X_s \bar{\nu} \nu$ and $K_L^0 \rightarrow \pi^0 \bar{\nu} \nu$ for the search of new physics in the form of "Non Standard Neutrino Interactions" (NSIs). We have calculated the Branching ratio (Br) of these reactions and constrained NSIs by using the mismatch between standard Model and the experiments. NSIs are giving very small contributions in rare decays of B mesons as compared to the D and K decays involving neutrinos in their final state. But, the interference effects for B are giving dominant contribution and it is used to find out constraint on NSIs. We found the contrarians for three generations of up type quarks as; $\epsilon_{\tau\tau}^{uL}$, $\epsilon_{\tau\tau}^{cL}$ and $\epsilon_{\tau\tau}^{tL}$ using interference. Charm quark and top quark induced constraint, $\epsilon_{\tau\tau}^{cL}$, $\epsilon_{\tau\tau}^{tL}$ are much more precise as compared to u quark. For $B^+ \rightarrow K^{*+} \bar{\nu} \nu$ and $B \rightarrow X_s \bar{\nu} \nu$, the $\epsilon_{\tau\tau}^{uL}$ constraint is same but the Br is very small. This shows that NSIs will have effects on the rare decays of B meson both inclusive and exclusive. $K_L^0 \rightarrow \pi^0 \bar{\nu} \nu$ is showing exactly the same behavior as of the D and K decays but providing more precise bounds. Here the interference effects are small and the bounds from $K_L^0 \rightarrow \pi^0 \bar{\nu} \nu$ are $O(10^{-3})$ which are $O(10^{-1})$ smaller than D and K .

References

- [1] G. Aad et al. (ATLAS Collaboration), Phys. Lett. B 716, 1 (2012).
- [2] S. Chatrchyan et al. (CMS Collaboration), Phys. Lett. B 716, 30 (2012).
- [3] CMS Collaboration, NPHYS3005 22 JUNE 2014.

- [4] N. Polonsky: Supersymmetry: Structure and Phenomena, Lecture Notes in Physics Monographs Springer-Verlag, Heidelberg, 2001; hep-ph/0108236.
- [5] J. Ellis: Supersymmetry for Alp Hikers, CERN-TH/2002-052, hep-ph/0203114.
- [6] M. Tanaka and R. Watanabe, Phys. Rev. D 82 (2010) 034027; arXiv:1302.2171v1 [hep-ex].
- [7] J. P. Lees et al. (The BABAR Collaboration), Phys. Rev. Lett. 109 (2012) 211801.
- [8] J. P. Lees et al. (The BABAR Collaboration), Phys. Rev. D 86 (2012) 051105.
- [9] SuperKamiokande, Y. Fukuda, et al., Phys. Rev. Lett. 82 (1999) 2644.
- [10] Super-Kamiokande, S. Fukuda, et al., Phys. Rev. Lett. 85 (2000) 3999.
- [11] SuperKamiokande, S. Fukuda, et al., Phys. Rev. Lett. 86 (2001) 5651.
- [12] Super-Kamiokande, S. Fukuda, et al., Phys. Lett. B 539 (2002) 179.
- [13] SNO, Q.R. Ahmad, et al., Phys. Rev. Lett. 87 (2001) 071301.
- [14] SNO, Q.R. Ahmad, et al., Phys. Rev. Lett. 89 (2002) 011301.
- [15] SNO, Q.R. Ahmad, et al., Phys. Rev. Lett. 89 (2002) 011302; KamLAND, K. Eguchi et al., (2002).
- [16] Alejandro Jaramillo and Luis A. Sanchez, Phys. Rev. D 84, 115001 (2011); Alakabha Datta, Phys. Rev. D 78, 095004 (2008).
- [17] S. L. Glashow, J. Iliopoulos, L. Maiani, Phys. Rev. D 2, 7 (1970).
- [18] F. Tahir, A. Mir, S. Mahmood, Chinese Physics C Vol. 38, No. 12 (2014) 123101.
- [19] Takeo Inami and C.S. Lim, Prog. Theor. Phys, 65, 1 (1981).
- [20] C. H. Chen, C. Q. Geng, T. C. Yuan, Phys. Rev. D 75, 077301 (2007).
- [21] S. Mahmood, F. Tahir and A. Mir, Mod. Phys. Lett. A 30, 1 (2015) 1550004.
- [22] S. Mahmood, F. Tahir and A. Mir, Int. J. Mod. Phys A Vol. 30, No. 2 (2015) 1550013.
- [23] W. Altmannshofer, A. J. Buras, D. M. Strauba and M. Wick, JHEP 04 (2009) 022.
- [24] D. Rein and L. M. Sehgal, Phys. Rev. D 39, 11 (1989).
- [25] Lu, Ming et al. Phys. Lett. B 324 (1994) 461-464 hep-ph/9401204 CALT-68-1911.

- [26] Rein, D. et al. Phys.Rev. D39 (1989) 3325 PITHA-89/03.
- [27] Andrzej J. Buras, arXiv:hep-ph/9806471v1 24 Jun 1998.
- [28] J. Boger et al. (SNO Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A 449 (2000) 172.
- [29] M. L. Mangano et al., hep-ph/0105155.
- [30] G. Buchalla, A. J. Buras, and M. E. Lautenbacher, Rev. Mod. Phys. 68 (1996) 1125.
- [31] arXiv:1303.3719v2 [hep-ex] 29 Apr 2013.
- [32] W. Marciano and Z. Parsa, Phys. Rev. D 53, R1 (1996).
- [33] S. Davidson, C. Pena-Garay, N. Rius, and A. Santamaria, J. High Energy Phys. 03 (2003) 011.
- [34] S. Mahmood, F. Tahir and A. Mir, Int. J. Mod. Phys A Vol. 30, Nos. 4 & 5 (2015) 1550024.
- [35] Zhen-Jun Xiao, Ying-Ying Fan, Wen-Fei Wang and Shan Cheng, arXiv:1401.0571v2 [hep-ph] 28 Feb 2014.
- [36] K.A. Olive et al., Chinese Physics C Vol. 38, No. 9 (2014) 090001.
- [37] ALEPH collaboration, R. Barate et al., Eur. Phys. J.C 19 (2001) 213.