## HARMONIC AND SPECTRAL ANALYSIS OF ABSTRACT PARABOLIC OPERATORS IN HOMOGENEOUS FUNCTION SPACES

#### ANATOLY G. BASKAKOV AND ILYA A. KRISHTAL

ABSTRACT. We use methods of harmonic analysis and group representation theory to study the spectral properties of the abstract parabolic operator  $\mathcal{L} = -d/dt + A$  in homogeneous function spaces. We provide sufficient conditions for invertibility of such operators in terms of the spectral properties of the operator A and the semi-group generated by A. We introduce a homogeneous space of functions with absolutely summable spectrum and prove a generalization of the Gearhart-Prüss Theorem for such spaces. We use the results to prove existence and uniqueness of solutions of a certain class of non-linear equations.

#### 1. Introduction

In this paper we study the spectral properties of a differential operator

$$(1.1) \mathcal{L} = -d/dt + A: D(\mathcal{L}) \subset \mathcal{F}(\mathbb{R}, X) \to \mathcal{F}(\mathbb{R}, X)$$

in homogeneous Banach spaces  $\mathcal{F}(\mathbb{R},X)$  of functions with values in a complex Banach space X. The operator  $A:D(A)\subset X\to X$  in (1.1) is assumed to be the infinitesimal generator of a  $C_0$ -semigroup  $T:\mathbb{R}_+=[0,\infty)\to B(X)$ . The homogeneous spaces  $\mathcal{F}=\mathcal{F}(\mathbb{R},X)$  and the operator  $\mathscr{L}$  are identified precisely in Definitions 2.1 and 3.1, respectively.

The properties of the homogeneous spaces  $\mathcal{F}$  allow us to correctly define the *Howland semigroup*  $\mathcal{T}: \mathbb{R}^+ \to \mathcal{B}(\mathcal{F})$  by

$$(\mathscr{T}(t)x)(s) = T(t)x(s-t), \ s \in \mathbb{R}, x \in \mathcal{F}, t \in \mathbb{R}^+.$$

In some homogeneous spaces such as  $C_0(\mathbb{R}, X)$  and  $L^p(\mathbb{R}, X)$ , see Example 2.1 for the definitions, it was proved [4, 5] that the operator

Date: January 27, 2022.

Key words and phrases. Abstract parabolic operators, Homogeneous function spaces, Green functions, Beurling spectrum.

The first author is supported in part by RFBR grant 13-01-00378. The second author is supported in part by NSF grant DMS-1322127.

 $\mathscr{L}$  is the infinitesimal generator of the semigroup  $\mathscr{T}$ . Moreover, in a large class of homogeneous spaces the following result holds.

**Theorem 1.1.** The following are equivalent:

ullet The operator  $\mathcal L$  is invertible, that is, the equation

$$\frac{dx}{dt} = Ax + y$$

has a unique (mild) solution  $x \in \mathcal{F}(\mathbb{R}, X)$  for any  $y \in \mathcal{F}(\mathbb{R}, X)$ ;

- The semigroup T is hyperbolic, that is, the spectrum  $\sigma(T)$  of the semigroup T satisfies  $\sigma(T(1)) \cap \mathbb{T} = \emptyset$ ,  $\mathbb{T} = \{\lambda \in \mathbb{C} : |\lambda| = 1\}$ ;
- The Howland semigroup  $\mathcal{T}$  is hyperbolic.

For  $\mathcal{F} \in \{C_0, L^p\}$ ,  $1 \leq p < \infty$ , the above result was proved in [30, 31] and [17, Theorem 2.39]. For a larger class of spaces, including  $L^{\infty}$  and  $C_b$ , the theorem appears in [4, 5, 9]. We remark that in many homogeneous spaces the result still holds even though the Howland semigroup  $\mathcal{T}$  is not strongly continuous (see [9]).

In this paper we prove that one of the implications in the above theorem holds for all homogeneous spaces in Definition 2.1:

**Theorem 1.2.** If T is a hyperbolic semigroup, then the operator  $\mathscr{L}$  is invertible.

If X is a Hilbert space, classical results of L. Gearhart and J. Prüss [24, 35, 20, 34] provide another equivalence to the statements in Theorem 1.1.

**Theorem 1.3.** The operator  $\mathcal{L}$  is invertible if and only if the imaginary axis  $i\mathbb{R}$  is a subset of the resolvent set  $\rho(A)$  of the generator A, i.e.

(1.3) 
$$\sigma(A) \cap (i\mathbb{R}) = \emptyset,$$

and the resolvent operator  $R(\lambda, A) = (A - \lambda I)^{-1}$  satisfies

(1.4) 
$$M = \sup_{\lambda \in i\mathbb{R}} ||R(\lambda, A)|| < \infty.$$

If X is not a Hilbert space, this equivalence does not hold in general (see [25] and [20, Counterexample IV.2.7]). Typical generalizations of the above result to Banach spaces [32, and references therein] would impose additional restrictions on the resolvent  $R(\lambda, A)$  or the Banach space X. In this paper, we pursue a different kind of generalization, where we deal with an arbitrary (complex) Banach space X and impose no additional restrictions on the resolvent. Instead, we define a class  $\mathcal{F}_{as}$  of functions in  $\mathcal{F}(\mathbb{R}, X)$  which is a Banach function space where the differential operator  $\mathcal{L}$  is invertible provided that (1.3) and (1.4) hold.

The space  $\mathcal{F}_{as}$  of functions with the absolutely summable spectrum (see Definition 2.5) is defined using the spectral theory of Banach modules [7, 28].

We refer to [1, 17, 20] for more information on the background and history of research related to this paper.

The remainder of the paper is organized as follows. In Section 2 we define homogeneous function spaces and introduce the space  $\mathcal{F}_{as}$  of functions with the absolutely summable spectrum. We do the latter by means of the spectral theory of Banach  $L^1(\mathbb{R})$ -modules, basic notions of which are also discussed in Section 2. In Section 3 we study the basic properties of the differential operator  $\mathcal{L}_{\mathcal{F}}$  and prove the first of our two main results: a sufficient condition for invertibility of  $\mathcal{L}_{\mathcal{F}}$  in a homogeneous Banach function space, which appears in Theorem 3.3. Our other main result, Theorem 4.1, appears in Section 4 and establishes sufficient conditions for invertibility of the operator  $\mathcal{L}$  in any homogeneous space of functions with the absolutely summable spectrum. We also provide an estimate for  $\|\mathcal{L}_{\mathcal{F}_{as}}^{-1}\|$ . In the final section of the paper we illustrate our results with a counterexample to the Gearhart-Prüss Theorem and an application to a special kind of non-linear differential equations.

#### 2. Preliminaries

In this section we introduce the notation, define homogeneous function spaces, and survey the necessary tools from the spectral theory of Banach  $L^1(\mathbb{R})$ -modules. We also introduce the space of functions with the absolutely summable spectrum.

The symbol X will denote a complex Banach space and B(X) will be the Banach algebra of all bounded linear operators on X. By T:  $\mathbb{R}_+ \to B(X)$  we shall denote a  $C_0$ -semigroup of operators in B(X) and  $A: D(A) \subset X \to X$  will be its infinitesimal generator [20].

By  $L^1(\mathbb{R}, X)$  we shall denote the Banach space of all (equivalence classes) of Bochner integrable X-valued functions with the standard  $L^1$ -norm:

$$||f|| = ||f||_1 = \int_{\mathbb{R}} ||f(t)||_X dt, \quad f \in L^1(\mathbb{R}, X).$$

If  $X = \mathbb{C}$ , we shall use the notation  $L^1 = L^1(\mathbb{R})$  for the standard group algebra of Lebesgue integrable functions. For  $f \in L^1(\mathbb{R})$ , we shall denote by  $\hat{f}$  the Fourier transform of f given by

$$\hat{f}(\lambda) = \int_{\mathbb{R}} f(t)e^{-it\lambda}dt, \ \lambda \in \mathbb{R}.$$

The space  $L^1(\mathbb{R}, B(X))$  defined as above, however, may be too small for our purposes. Occasionally, we will use the space  $L^1_s(\mathbb{R}, B(X))$  that consists of all functions  $F: \mathbb{R} \to B(X)$  with the following properties:

- (1) For all  $x \in X$  the function  $s \mapsto F(s)x : \mathbb{R} \to X$  is measurable;
- (2) There is  $f \in L^1(\mathbb{R})$  such that

(2.1) 
$$||F(s)|| \le f(s)$$
 a.e.

For  $F \in L^1_s(\mathbb{R}, B(X))$  we let

$$||F|| = ||F||_1 = \inf ||f||,$$

where the infimum is taken over all functions f that satisfy (2.1).

The space  $L_s^1(\mathbb{R}, B(X))$  is a Banach algebra with the multiplication given by

$$(F_1 * F_2)(t) = \int_{\mathbb{R}} F_1(s) F_2(t-s) ds, \quad F_1, F_2 \in L_s^1(\mathbb{R}, B(X)).$$

In particular,  $||F_1 * F_2|| \le ||F||_1 ||F||_2$ .

We shall also use the space of locally integrable X-valued functions  $L^1_{loc}(\mathbb{R},X)$ , which consists of all measurable functions  $f:\mathbb{R}\to X$  such that

$$\int_{K} \|f(t)\|_{X} dt < \infty$$

for any compact set  $K \subset \mathbb{R}$ .

For  $p \in [1, \infty)$ , the Stepanov space  $S^p = S^p(\mathbb{R}, X)$  consists of all functions  $x \in L^1_{loc}(\mathbb{R}, X)$  such that

$$||x||_{\mathcal{S}^p} = \sup_{t \in \mathbb{R}} \left( \int_0^1 ||x(s+t)||^p ds \right)^{1/p} < \infty.$$

## 2.1. Homogeneous Banach function spaces.

In this paper we consider Banach function spaces  $\mathcal{F}(\mathbb{R},X)$  that are homogeneous according to the following definition.

**Definition 2.1.** A Banach function space  $\mathcal{F} = \mathcal{F}(\mathbb{R}, X)$  is *homogeneous* if it has the following properties:

- (1)  $\mathcal{F}$  is continuously embedded into  $\mathcal{S}^1$ ;
- (2) For all  $t \in \mathbb{R}$  and  $x \in \mathcal{F}$  we have  $S(t)x \in \mathcal{F}$ , where

$$(2.2) S(t)x(s) = x(t+s),$$

and the translation operator S(t) is an isometry in  $B(\mathcal{F})$ .

(3) Given  $x \in \mathcal{F}$  and  $C \in B(X)$ , the function

$$y(t) = C(x(t))$$

belongs to  $\mathcal{F}$  and  $||y|| \le ||C|| ||x||$ .

(4) Given  $x \in \mathcal{F}$  and  $F \in L_s^1(\mathbb{R}, B(X))$ , the convolution

$$(F * x)(t) = \int_{\mathbb{R}} F(s)x(t - s)ds$$

belongs to  $\mathcal{F}$  and  $||F * x|| \le ||F||_1 ||x||$ .

(5) If  $x \in \mathcal{F}$  is such that f \* x = 0 for all  $f \in L^1(\mathbb{R})$  then x = 0.

Example 2.1. The following Banach spaces are homogeneous or have an equivalent norm that makes them homogeneous:

(1) The spaces  $L^p = L^p(\mathbb{R}, X), p \in [1, \infty]$ , of functions  $x \in L^1_{loc}(\mathbb{R}, X)$  such that

$$||x||_{L^p} = \left(\int_{\mathbb{R}} ||x(s)||^p ds\right)^{1/p} < \infty, \ p \in [1, \infty),$$

or  $||x||_{\infty} = \operatorname{ess\,sup}_{t \in \mathbb{R}} |x(t)| < \infty;$ 

- (2) Stepanov spaces  $S^p = S^p(\mathbb{R}, X), p \in [1, \infty);$
- (3) Wiener amalgam spaces  $L^{p,q} = L^{p,q}(\mathbb{R}, X), p, q \in [1, \infty),$  of functions  $x \in L^1_{loc}(\mathbb{R}, X)$  such that

$$||x||_{L^{p,q}} = \left(\sum_{k \in \mathbb{Z}} \left(\int_0^1 ||x(s+k)||^p ds\right)^{q/p}\right)^{1/q} < \infty, \ p, q \in [1, \infty);$$

(4) The space  $C_b = C_b(\mathbb{R}, X)$  of bounded continuous X-valued functions with the norm

$$||x||_{\infty} = \sup_{t \in \mathbb{R}} ||x(t)||, \ x \in C_b;$$

- (5) The subspace  $C_{ub} = C_{ub}(\mathbb{R}, X) \subset C_b$  of uniformly continuous functions;
- (6) The subspace  $C_0 = C_0(\mathbb{R}, X) \subset C_{ub}$  of continuous functions vanishing at infinity:  $x \in C_0$  if  $\lim_{|t| \to \infty} ||x(t)|| = 0$ ;
- (7) The subspace  $C_{sl,\infty} = C_{sl,\infty}(\mathbb{R}, X) \subset C_{ub}$  of slowly varying at infinity functions:  $x \in C_{sl,\infty}$  if  $\lim_{|\tau| \to \infty} ||x(\tau + t) x(\tau)|| = 0$  for all  $t \in \mathbb{R}$  (see [11]);
- (8) The subspace  $C_{\omega} = C_{\omega}(\mathbb{R}, X) \subset C_{ub}$  of  $\omega$ -periodic functions,  $\omega \in \mathbb{R}$ ;
- (9) The subspace  $AP = AP(\mathbb{R}, X) \subset C_{ub}$  of (Bohr) almost periodic functions [2, 33];
- (10) The subspace  $AP_{\infty} = AP_{\infty}(\mathbb{R}, X) \subset C_{ub}$  of almost periodic at infinity functions [9] defined by

$$AP_{\infty} = \overline{\operatorname{span}}\{e^{i\lambda} x : \lambda \in \mathbb{R}, x \in C_{sl,\infty}\}.$$

(11) The spaces  $C^k = C^k(\mathbb{R}, X)$ ,  $k \in \mathbb{N}$ , of k times continuously differentiable functions with a bounded k-th derivative and the norm

$$||x||_{(k)} = ||x||_{\infty} + ||x^{(k)}||_{\infty} < \infty;$$

(12) The Hölder spaces  $C^{k,\alpha} = C^{k,\alpha}(\mathbb{R}, X), k \in \mathbb{N} \cup \{0\}, \alpha \in (0, 1]$ :

$$C^{k,\alpha} = \left\{ x \in C^k : \|x^{(k)}\|_{C^{0,\alpha}} = \sup_{t \neq s \in \mathbb{R}} \frac{|x(t) - x(s)|}{|t - s|^{\alpha}} < \infty \right\},$$

$$\|x\|_{C^{k,\alpha}} = \|x\|_{C^k} + \|x^{(k)}\|_{C^{0,\alpha}}.$$

Remark 2.1. We note that Definition 2.1 in this paper differs from Definition 2.1 in [9], which is more narrow. In particular, here we do not assume that the space  $\mathcal{F}$  is solid and this allows us to consider the spaces of periodic and almost periodic functions.

**Definition 2.2.** A homogeneous space  $\mathcal{F} = \mathcal{F}(\mathbb{R}, X)$  is called *spectrally homogeneous* if for all  $\lambda \in \mathbb{R}$  and  $x \in \mathcal{F}$  we have  $e^{i\lambda \cdot}x \in \mathcal{F}$  and  $\|e^{i\lambda \cdot}x\| = \|x\|$ .

Among all of the homogeneous spaces in Example 2.1 the only class of spaces that are not spectrally homogeneous are the spaces of  $C_{\omega}$  of  $\omega$ -periodic functions.

In a spectrally homogeneous space  $\mathcal{F}$  there is a well-defined isometric representation  $V \in \mathbb{R} \to B(\mathcal{F})$  given by

(2.3) 
$$V(\lambda)x(t) = e^{i\lambda t}x(t), \ \lambda, t \in \mathbb{R}, x \in \mathcal{F}.$$

# 2.2. Banach $L^1$ -modules and the Beurling spectrum.

Properties (4) and (5) in Definition 2.1 ensure that a homogeneous Banach space is a non-degenerate Banach  $L^1(\mathbb{R})$ -module [7, 28]. In this subsection we present the necessary definitions and results from the spectral theory of such Banach modules. The proofs omitted in the presentation and further details can be found in [7, 12, 14, 28].

**Definition 2.3.** A Banach space  $\mathcal{X}$  is a Banach  $L^1(\mathbb{R})$ -module if there is a bilinear map  $(f, x) \mapsto fx : L^1(\mathbb{R}) \times \mathcal{X} \to \mathcal{X}$  such that

- (1)  $(f * g)x = f(gx), f, g \in L^1(\mathbb{R}), x \in \mathcal{X};$
- (2)  $||fx|| \le ||f|| ||x||$ ,  $f \in L^1(\mathbb{R})$ ,  $x \in \mathcal{X}$ . The module structure is *non-degenerate* if, in addition,
- (3) fx = 0 for all  $f \in L^1(\mathbb{R})$  implies  $x = 0 \in \mathcal{X}$ .

We say that the structure of a Banach  $L^1(\mathbb{R})$ -module  $\mathcal{X}$  is associated with a representation  $U: \mathbb{R} \to B(\mathcal{X})$  if (2.4)

$$U(t)(fx) = f_t x = f(U(t)x), f \in L^1(\mathbb{R}), x \in \mathcal{X}, f_t(s) = f(s+t).$$

As we mentioned above, any homogeneous function space  $\mathcal{X} = \mathcal{F}(\mathbb{R}, X)$  is a non-degenerate Banach  $L^1(\mathbb{R})$ -module. Its structure is given by

(2.5) 
$$(fx)(t) = (f*x)(t) = \int_{\mathbb{R}} f(t-s)x(s)ds = \int_{\mathbb{R}} f(s)x(t-s)ds,$$

 $f \in L^1(\mathbb{R}), x \in \mathcal{F}$ , and is associated with the translation representation S defined by (2.2).

**Definition 2.4.** The *Beurling spectrum* of an element  $x \in \mathcal{X}$  is the subset  $\Lambda(x) \subseteq \mathbb{R}$  the complement of which is given by

$$\{\lambda \in \mathbb{R} : \text{ there is } f \in L^1(\mathbb{R}) \text{ such that } \hat{f}(\lambda) \neq 0 \text{ and } fx = 0\}.$$

Remark 2.2. In homogeneous Banach spaces the Beurling spectrum  $\Lambda(x)$  coincides with the support of the (distributional) Fourier transform of  $x \in \mathcal{F}$ .

In the next lemma we present basic properties of the Beurling spectrum that will be used throughout the paper. We refer to [7, 12, 36] and references therein for the proof.

**Lemma 2.1.** Let  $\mathcal{X}$  be a non-degenerate Banach  $L^1(\mathbb{R})$ -module. Then

- (i):  $\Lambda(x)$  is closed for every  $x \in \mathcal{X}$  and  $\Lambda(x) = \emptyset$  if and only if x = 0;
- (ii):  $\Lambda(Ax+By) \subseteq \Lambda(x) \cup \Lambda(y)$  for all  $A, B \in B(\mathcal{X})$  that commute with all operators  $x \mapsto fx$ ,  $f \in L^1(\mathbb{R})$ ;
- (iii):  $\Lambda(fx) \subseteq (\text{supp } \hat{f}) \cap \Lambda(x) \text{ for all } f \in L^1(\mathbb{R}) \text{ and } x \in \mathcal{X};$
- (iv): fx = 0 if  $(\operatorname{supp} \hat{f}) \cap \Lambda(x)$  is countable and  $\hat{f}(\lambda) = 0$  for all  $\lambda \in (\operatorname{supp} \hat{f}) \cap \Lambda(x)$ ,  $f \in L^1(\mathbb{R})$ ,  $x \in \mathcal{X}$ ;
- (v): fx = x if  $\Lambda(x)$  is a compact set, the boundary of  $\Lambda(x)$  is countable, and  $\hat{f} \equiv 1$  on  $\Lambda(x)$ ,  $f \in L^1(\mathbb{R})$ ,  $x \in \mathcal{X}$ .

Given a closed set  $\Delta \subset \mathbb{R}$  we shall denote by  $\mathcal{X}(\Delta)$  the (closed) spectral submodule of all vectors  $x \in \mathcal{X}$  such that  $\Lambda(x) \subseteq \Delta$ . The symbol  $\mathcal{X}_{Comp}$  will stand for the set of all vectors x such that  $\Lambda(x)$  is compact. If the module structure is associated with a representation U, by  $\mathcal{X}_U$  we shall denote the submodule of U-continuous vectors, i.e., the set of all vectors  $x \in \mathcal{X}$  such that the function  $t \mapsto U(t)x : \mathbb{R} \to \mathcal{X}$  is continuous.

Remark 2.3. Observe that any spectral submodule  $\mathcal{F}(\Delta)$  of a homogeneous function space  $\mathcal{F}(\mathbb{R}, X)$  is itself a homogeneous function space. It may happen that  $\mathcal{F}(\Delta) = \{0\}$  even if  $\Delta \neq \emptyset$ . For example, if  $\mathcal{F} = L^p(\mathbb{R}, X)$ ,  $1 \leq p < \infty$ , and  $\Delta$  is finite then  $\mathcal{F}(\Delta) = \{0\}$ . If  $\Delta$  is compact then  $\mathcal{F}(\Delta) \subset C_{ub}$  and each  $x \in \mathcal{F}(\Delta)$  extends to an entire function of exponential type  $\omega = \max\{|\lambda|, \lambda \in \Delta\}$  [11].

The following lemma opens up the possibility of applying our main results to non-linear equations.

**Lemma 2.2.** Let  $\mathcal{X}$  be a non-degenerate Banach  $L^1(\mathbb{R})$ -module. Assume that the module structure is associated with a representation U as in (2.4). Let  $F: \mathcal{X}^n \to \mathcal{X}$ ,  $\mathcal{X}^n = \underbrace{\mathcal{X} \times \cdots \times \mathcal{X}}_{n \text{ times}}$ , be an n-linear map

such that for any  $t \in \mathbb{R}$  we have

$$U(t)(F(x_1,\ldots,x_n))=F(U(t)x_1,\ldots,U(t)x_n).$$

Then

$$\Lambda(F(x_1,\ldots,x_n))\subseteq\overline{\Lambda(x_1)+\ldots+\Lambda(x_n)}.$$

*Proof.* In case  $\mathcal{X}$  is a Banach algebra and  $F(x_1, x_2) = x_1 x_2$ , this result appears in [12, 14]. In the general case the proof may be repeated nearly verbatim with obvious modifications.

### 2.3. Vectors with the absolutely summable spectrum.

In this subsection we define the class  $\mathcal{X}_{as}$  of vectors in a Banach  $L^1(\mathbb{R})$ -module  $\mathcal{X}$  that have absolutely summable spectrum. In the scalar case this class was introduced in [6]. In this exposition we follow [14] where the general definition appears. We use the family of functions  $(\phi_{\alpha})$ ,  $\alpha \in \mathbb{R}$ , defined via the Fourier transform by

(2.6) 
$$\hat{\phi}_a(\lambda) = \hat{\phi}(\lambda - a), \ a \in \mathbb{R},$$

where

$$\hat{\phi}(\lambda) \equiv \hat{\phi}_0(\lambda) = (1 - |\lambda|)\chi_{[-1,1]}(\lambda),$$

and  $\chi_E$  is, as usually, the characteristic function of the set E.

**Definition 2.5.** The class  $\mathcal{X}_{as}$  of vectors in a Banach  $L^1(\mathbb{R})$ -module  $\mathcal{X}$  that have absolutely summable spectrum is

(2.8) 
$$\mathcal{X}_{as} = \left\{ x \in \mathcal{X} : \|x\|_{\widetilde{as}} = \int_{\mathbb{R}^d} \|\phi_a x\| da < \infty \right\},$$

where the functions  $\phi_a$  are defined by (2.6) and (2.7).

In [14], one can find plenty of examples of the spaces with absolutely summable spectrum. Classical Wiener amalgam spaces [21, 23] are among the well-studied spaces that arise in such a way.

Remark 2.4. In the above definition, instead of the family of functions  $(\phi_a)$  one can use just about any bounded uniform partition of unity [21, 22]. One would obtain the same space as a result [14]. This is analogous to using different window functions in the short time Fourier transform [26].

In [14] we have shown that  $\mathcal{X}_{as}$  is a Banach space with the norm  $\|\cdot\|_{\tilde{as}}$ . For our purposes, however, it is often more convenient to use an equivalent norm given by

(2.9) 
$$||x||_{as} = 5 \sum_{n \in \mathbb{Z}} ||\phi_n x||.$$

In [14] we obtained the inequalities

$$(2.10) ||x||_{\tilde{as}} \le ||x||_{as} \le 20 ||x||_{\tilde{as}}, \ x \in \mathcal{X}_{as}.$$

In [14] we have also shown that if  $\mathcal{X}$  is a Banach algebra and the module structure is associated with a group of algebra automorphisms, then  $\mathcal{X}_{as}$  is also a Banach algebra and, due to the choice of the constant in (2.9),

$$(2.11) ||xy||_{as} \le ||x||_{as} ||y||_{as}, \ x, y \in \mathcal{X}_{as}.$$

Moreover, the key result of [14] states that the algebra  $\mathcal{X}_{as}$  is inverse closed, i.e. if  $x \in \mathcal{X}_{as}$  is invertible in  $\mathcal{X}$ , then  $x^{-1} \in \mathcal{X}_{as}$ . The following analog of (2.11) can be proved in exactly the same way.

**Proposition 2.3.** Let  $\mathcal{X}$  be a non-degenerate Banach  $L^1(\mathbb{R})$ -module. Assume that the module structure is associated with a representation U as in (2.4). Let  $F: \mathcal{X}^n \to \mathcal{X}$ , be an n-linear map such that for any  $t \in \mathbb{R}$  we have

$$U(t)(F(x_1,\ldots,x_n)) = F(U(t)x_1,\ldots,U(t)x_n).$$

Then

(2.12) 
$$||F(x_1, \dots, x_n)||_{as} \le ||F|| \cdot \prod_{k=1}^n ||x_k||_{as}.$$

It has been observed by many people, see, e.g., [27] and [14, Remark 3.5], that smoothness of the function x is closely related to the spectral decay of x. In particular, we have  $C^{1,\alpha} \subset \mathcal{X}_{as}$ ,  $\alpha > 0$ . Below we prove a slightly weaker sufficient condition that uses the following modulus of continuity.

**Definition 2.6.** Let  $\mathcal{X}$  be a non-degenerate Banach  $L^1(\mathbb{R})$ -module with the structure associated with an isometric representation  $U: \mathbb{R} \to B(\mathcal{X})$ . For  $x \in \mathcal{X}$ , its modulus of continuity  $\omega_x$  is defined by

$$\omega_x(t) = \sup_{|s| < t} ||U(s)x - x||, \ t \ge 0.$$

Remark 2.5. The basic properties of the modulus of continuity can be found, for example in [16, 29]. Here we mention the obvious facts that

$$\lim_{t\to 0}\omega_x(t)=0, x\in\mathcal{X}_U,$$

and  $\omega_x$  is subadditive, that is

$$\omega_x(t+s) \le \omega_x(t) + \omega_x(s), \ x \in \mathcal{X}.$$

As an immediate consequence of subbaditivity we get

(2.13) 
$$\omega_x(ks) \le k\omega_x(s)$$
 and  $\frac{\omega_x(1)}{k} \le \omega_x(\frac{1}{k}), x \in \mathcal{X}, k \in \mathbb{N}, s \ge 0.$ 

Using the monotonicity of  $\omega_x$  we also get

(2.14) 
$$\omega_x(\lambda s) \le (\lambda + 1)\omega_x(s), \ \lambda, s > 0, \ x \in \mathcal{X}.$$

**Lemma 2.4.** For  $x \in \mathcal{X}$  we have

(2.15) 
$$\|\phi_k x\| \le Const \cdot \omega_x(\frac{1}{|k|}), k \in \mathbb{Z} \setminus \{0\}.$$

*Proof.* Observe that for  $k \neq 0$ 

$$\phi_k x = \int_{\mathbb{R}} \phi(t) e^{ikt} U(-t) x dt = -\int_{\mathbb{R}} \phi(\tau + \frac{\pi}{k}) e^{ikt} U(-\tau - \frac{\pi}{k}) x d\tau,$$

where we made the change of variables  $t = \tau + \pi/k$ . Averaging the above two expressions we get

$$\phi_{k}x = \frac{1}{2} \int_{\mathbb{R}} e^{ikt} (\phi(t)U(-t) - \phi(t + \frac{\pi}{k})U(-t - \frac{\pi}{k}))xdt$$

$$= \frac{1}{2} \int_{\mathbb{R}} e^{ikt} (\phi(t) - \phi(t + \frac{\pi}{k}))U(-t)xdt$$

$$+ \frac{1}{2} \int_{\mathbb{R}} e^{ikt} \phi(t + \frac{\pi}{k})(U(-t) - U(-t - \frac{\pi}{k}))xdt.$$

Hence,

$$\|\phi_k x\| \le \frac{1}{2} \|S(\frac{\pi}{k})\phi - \phi\|_1 \|x\| + \frac{1}{2}\omega_x(\frac{\pi}{|k|}).$$

Direct computation (as well as [12, Theorem 3.7]) implies that

$$||S(\frac{\pi}{k})\phi - \phi||_1 \le Const \cdot \frac{1}{|k|}.$$

Finally, using (2.13) and (2.14) we get (2.15).

**Theorem 2.5.** Assume that  $\mathcal{X}$  is a non-degenerate Banach  $L^1(\mathbb{R})$ module with the structure associated with a representation  $U: \mathbb{R} \to B(\mathcal{X})$ . Let B be the infinitesimal generator of U and assume that  $x \in D(B)$ . Assume also that y = Bx satisfies

(2.16) 
$$\sum_{k \in \mathbb{N}} \frac{\omega_y(1/k)}{k} < \infty.$$

Then  $x \in \mathcal{X}_{as}$ .

*Proof.* For  $k \in \mathbb{N} \setminus \{1\}$ , let  $f_k \in L^1(\mathbb{R})$  be such that

$$\hat{f}_k(\lambda) = \left\{ \begin{array}{cc} \frac{1}{i\lambda}, & \lambda \ge k - 1; \\ \frac{1}{i(2k - 2 - \lambda)}, & \lambda < k - 1. \end{array} \right.$$

Similarly, for  $-k \in \mathbb{N} \setminus \{1\}$ , let  $f_k \in L^1(\mathbb{R})$  be such that

$$\hat{f}_k(\lambda) = \left\{ \begin{array}{cc} \frac{1}{i\lambda}, & \lambda \le k+1; \\ \frac{1}{i(2k+2-\lambda)}, & \lambda > k+1. \end{array} \right.$$

Observe that for  $k \in \mathbb{Z} \setminus \{-1, 0, 1\}$  we have  $||f_k||_1 = \frac{1}{|k|-1}$  and the function  $\varphi_k = f_k * \phi_k \in L^1(\mathbb{R})$  satisfies

$$\widehat{\varphi}_k(\lambda) = \frac{\widehat{\phi}_k(\lambda)}{i\lambda}, \lambda \in \mathbb{R} \setminus \{0\}.$$

Hence, for  $x \in D(B)$  integration by parts yields  $\phi_k x = \varphi_k(Bx)$ ,  $k \in \mathbb{Z}\setminus\{-1,0,1\}$ . Finally, using (2.15) for y = Bx, we get

$$\|\phi_k x\| \le \|f_k\|_1 \|\|\phi_k y\| \le \frac{Const}{|k| - 1} \cdot \omega_y(\frac{1}{|k|}).$$

and the result follows.

Observe that condition (2.16) is satisfied automatically if  $x \in D(B^{1+\alpha})$ ,  $\alpha > 0$ . We also have the following corollary.

Corollary 2.6. Let  $\mathcal{X} = C_b(\mathbb{R}, X)$ . Then for any  $\alpha > 0$  we have  $C^{1,\alpha} \subset \mathcal{X}_{as}$ .

*Proof.* In this case B = d/dt. Since,  $x \in C^{1,\alpha}$  implies

$$\omega_{x'}(t) \le ||x||_{C^{1,\alpha}} \cdot t^{\alpha}, \ t > 0,$$

the series (2.16) converges.

Remark 2.6. If  $x \in \mathcal{F} = C_0(\mathbb{R}, X)$  and  $x = \hat{y}$  for some  $y \in L^1(\mathbb{R})$ , then  $x \in \mathcal{F}_{as}$  and  $||x||_{as} \leq 5||y||_1$ .

We conclude the section with the following useful result.

**Lemma 2.7.** If  $\mathcal{F}$  is a (spectrally) homogeneous space then  $\mathcal{F}_{as}$  is also a (spectrally) homogeneous space.

*Proof.* Since convolution operators commute with translation, the verification of the properties of a (spectrally) homogeneous space is straightforward and is left to the reader.  $\Box$ 

### 3. Basic properties of the operator $\mathscr L$

In this section we collect and enhance some of the known spectral properties of abstract parabolic operators.

Recall that by  $A: D(A) \subseteq X \to X$  we denote the infinitesimal generator of a  $C_0$ -Semigroup T. In a homogeneous space  $\mathcal{F}(\mathbb{R}, X)$  we define the differential operator  $\mathcal{L} = \mathcal{L}_{\mathcal{F}} = -d/dt + A$  in (1.1) as follows [5, 9, 33].

**Definition 3.1.** A function  $x \in \mathcal{F}$  belongs to the domain  $D(\mathcal{L})$  of the operator  $\mathcal{L}$  if there is a function  $y \in \mathcal{F}$  such that for all  $s \leq t$  in  $\mathbb{R}$  we have

(3.1) 
$$x(t) = T(t-s)x(s) - \int_{s}^{t} T(t-\tau)y(\tau)d\tau.$$

For  $x \in D(\mathcal{L})$  we let  $\mathcal{L}x = y$ , if x and y satisfy (3.1).

We remark that the operator  $\mathcal{L}$  is well defined as it is not hard to see that for  $x \in D(\mathcal{L})$  there is a unique y such that (3.1) is satisfied. We also remark that  $\mathcal{F} \subset \mathcal{S}^1$  implies  $D(\mathcal{L}) \subset C_{ub}$ .

The operator  $\mathcal{L}$  is *invertible* if it is *injective*, i.e.  $\ker \mathcal{L} = \{0\}$ , and *surjective*, i.e. its range  $\operatorname{Im} \mathcal{L} = \mathcal{L}D(\mathcal{L})$  satisfies  $\operatorname{Im} \mathcal{L} = \mathcal{F}$ .

We begin studying the spectral properties of the operator  $\mathcal{L}_{\mathcal{F}}$  with the following key property of its kernel. It was originally proved in [3] for  $\mathcal{F} = C_b$ .

**Lemma 3.1.** Assume  $x \in \ker \mathcal{L}_{\mathcal{F}}$ . Then

$$(3.2) i\Lambda(x) \subseteq \sigma(A) \cap (i\mathbb{R}).$$

*Proof.* Let  $x \in \ker \mathscr{L}_{\mathcal{F}}$ . Then from (3.1) we get x(t) = T(t-s)x(s),  $s \leq t$ . Then in view of this equality, for any  $f \in L^1(\mathbb{R})$ , we have

$$T(t-s)(f*x)(s) = \int_{\mathbb{R}} f(s-\tau)T(t-s)x(\tau)d\tau$$
$$= \int_{\mathbb{R}} f(s-\tau)x(t-s+\tau)d\tau = (f*x)(t).$$

Hence, (3.1) implies  $f * x \in \ker \mathcal{L}_{\mathcal{F}}$ .

Let  $i\lambda_0 \notin \sigma(A)$  and  $f \in L^1(\mathbb{R})$  be such that  $\hat{f}(\lambda_0) \neq 0$ , supp  $\hat{f}$  is compact, and  $i \operatorname{supp} \hat{f} \subset \rho(A)$  Then, according to the above, y = f \* x satisfies y(s+t) = T(t)y(s) for all  $s \in \mathbb{R}$  and  $t \geq 0$ . Observe that since  $f \in C^{\infty}$ , i.e. f is differentiable infinitely many times, we have  $y \in C^{\infty}$ . Moreover,

$$y'(s+t) = T(t)Ay(s) = AT(t)y(s), \ s \in \mathbb{R}, t \ge 0,$$

and, therefore, plugging in t = 0, we get y' - Ay = 0.

Let  $\psi \in L^1(\mathbb{R})$  be such that  $\hat{\psi} = 1$  in a neighborhood of supp  $\hat{f}$ , supp  $\hat{\psi}$  is compact, and  $i \operatorname{supp} \hat{\psi} \subset \rho(A)$ . Consider  $F \in L^1(\mathbb{R}, B(\mathcal{F}))$  defined by

$$F(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{\psi}(\lambda) R(i\lambda, A) e^{i\lambda t} d\lambda, t \in \mathbb{R};$$

the above integral makes sense because  $i \operatorname{supp} \hat{\psi} \subset \rho(A)$ . Observe that

(3.3) 
$$\hat{F}(\lambda) = \begin{cases} \hat{\psi}(\lambda)R(i\lambda, A), & i\lambda \in \rho(A); \\ 0, & i\lambda \notin \rho(A); \end{cases}$$

is compactly supported and infinitely differentiable. Then, since the operator A is closed,

$$0 = F * (Ay - y') = A(F * y) - F' * y - F_0 * y,$$

where  $F_0 = AF - F'$  has Fourier transform

$$\widehat{F}_0(\lambda) = \widehat{\psi}(\lambda) A R(i\lambda, A) - i\lambda \widehat{\psi}(\lambda) R(i\lambda, A) = \widehat{\psi}(\lambda) I.$$

Hence, 
$$f * x = y = F_0 * y = 0$$
 and, therefore,  $\lambda_0 \notin \Lambda(x)$ .

Corollary 3.2. If the generator A satisfies (1.3) then the operator  $\mathcal{L}_{\mathcal{F}}$  is injective.

*Proof.* Assume that  $\mathscr{L}_{\mathcal{F}}x = 0$ . Immediately from Lemma 3.1 we get  $\Lambda(x) = \emptyset$ . Hence, x = 0 and, the operator  $\mathscr{L}_{\mathcal{F}}$  is injective.

Next we proceed to use the above result to obtain invertibility conditions for  $\mathscr{L}_{\mathcal{F}}$  in the case when the semigroup T is hyperbolic, i.e. it satisfies

(3.4) 
$$\sigma(T(1)) \cap \mathbb{T} = \emptyset.$$

For such semigroups we have

$$\sigma(T(1)) = \sigma_{in} \cup \sigma_{out},$$

where  $\sigma_{in}$  is the spectral component inside the unit disc and  $\sigma_{out} = \sigma(T(1)) \setminus \sigma_{in}$ . We let  $P_{in}$  and  $P_{out}$  be the corresponding spectral projections

$$P_{in} = \frac{1}{2\pi i} \int_{\mathbb{T}} (T(1) - \lambda I)^{-1} d\lambda, \quad P_{out} = I - P_{in},$$

and represent the space X as a direct sum

$$X = X_{in} \oplus X_{out}, \quad X_{in} = P_{in}X, \ X_{out} = P_{out}X.$$

From the definition of the spectral projections it follows that  $P_{in}$  and  $P_{out}$  commute with the operators T(t),  $t \ge 0$ . Therefore,  $X_{in}$  and  $X_{out}$ 

are invariant subspaces for these operators and we can consider the (restriction) semigroup  $T_{in}$  and the group  $T_{out}$  defined by

$$T_{in}: \mathbb{R}_+ \to B(X_{in}), \ T_{in}(t) = T(t)|_{X_{in}};$$

$$T_{out}: \mathbb{R} \to B(X_{out}), \ T_{out}(t) = \begin{cases} T(t)|_{X_{out}}, & t \ge 0; \\ (T(-t)|_{X_{out}})^{-1}; & t < 0. \end{cases}$$

The following theorem is one of the main results of this paper. As we mentioned in the introduction, its special cases appear in [5, 9, 15, 17].

**Theorem 3.3.** If T is a hyperbolic semigroup, the operator  $\mathscr{L}_{\mathcal{F}}$  from Definition 3.1 is invertible. The inverse  $\mathscr{L}_{\mathcal{F}}^{-1} \in B(\mathcal{F})$  is defined by

$$(3.5) \quad (\mathscr{L}_{\mathcal{F}}^{-1}y(t) = (G*y)(t) = \int_{\mathbb{R}} G(t-\tau)y(\tau)d\tau, \ t \in \mathbb{R}, \ y \in \mathcal{F},$$

where the Green function  $G \in L^1_s(\mathbb{R}, B(\mathcal{F}))$  is given by

(3.6) 
$$G(t) = \begin{cases} -T(t)P_{in}, & t \ge 0; \\ T_{out}(t)P_{out}; & t < 0. \end{cases}$$

*Proof.* It is immediate from Definition 2.1(4) that the right hand side of (3.5) defines a bound operator in  $B(\mathcal{F})$ . We need to check that this operator is, indeed, the inverse of  $\mathcal{L}_{\mathcal{F}}$ .

Since the semigroup T is hyperbolic, the spectral inclusion theorem [20, Theorem IV.3.6] implies that the operator A satisfies (1.3). Therefore, the operator  $\mathcal{L}_{\mathcal{F}}$  is injective by Corollary 3.2.

It remains to show that given  $y \in \mathcal{F}$  and x = G \* y we have  $\mathscr{L}_{\mathcal{F}} x = y$ . We get

$$x(t) - T(t - s)x(s) = (G * y)(t) - T(t - s)(G * y)(s)$$

$$= \int_{t}^{\infty} T_{out}(t - \tau)P_{out}y(\tau)d\tau - \int_{-\infty}^{t} T(t - \tau)P_{in}y(\tau)d\tau$$

$$- \int_{s}^{\infty} T(t - s)T_{out}(s - \tau)P_{out}y(\tau)d\tau + \int_{-\infty}^{s} T(t - \tau)P_{out}y(\tau)d\tau$$

$$= - \int_{s}^{t} T(t - \tau)P_{out}y(\tau)d\tau - \int_{s}^{t} T(t - \tau)P_{in}y(\tau)d\tau$$

$$= - \int_{s}^{t} T(t - \tau)y(\tau)d\tau,$$

and the result follows from (3.1).

Corollary 3.4. If T is a hyperbolic semigroup, then the generator A satisfies (1.3) and (1.4).

*Proof.* As we mentioned above (1.3) follows from the spectral mapping theorem. The inequality (1.4) follows since the Fourier transform  $\widehat{G}$  of the Green function G satisfies  $\widehat{G}(\lambda) = R(i\lambda, A)$ , see [15] for details.  $\square$ 

Corollary 3.5. The operator  $\mathscr{L}_{\mathcal{F}}$  in Definition 3.1 is closed.

Proof. Let  $\omega \in \mathbb{R}$  be such that the semigroup  $T_{\omega}$ ,  $T_{\omega}(t) = T(t)e^{-\omega t}$ , satisfies  $||T_{\omega}(t)|| \to 0$  as  $t \to \infty$ . The value  $\omega_0 = \min \omega$ , where the minimum is taken over all  $\omega$  with the above property is usually called the growth bound of the semigroup T [20, Definition I.5.6]. Theorem 3.3 applied to  $T_{\omega}$  implies that the operator  $\mathcal{L}_{\mathcal{F}} - \omega I$  is invertible and  $(\mathcal{L}_{\mathcal{F}} - \omega I)^{-1}y = G_{\omega} * y$ , where

$$G_{\omega}(t) = \begin{cases} -T_{\omega}(t), & t \ge 0; \\ 0; & t < 0. \end{cases}$$

Hence,  $\rho(\mathscr{L}_{\mathcal{F}}) \neq \emptyset$  and the operator  $\mathscr{L}_{\mathcal{F}}$  is closed.

In what follows we shall denote by  $\tilde{S}: L^1(\mathbb{R}) \to B(\mathcal{F})$  the algebra homomorphism given by

$$\tilde{S}(f)x = f * x, \ f \in L^1(\mathbb{R}), x \in \mathcal{F}.$$

The next result asserts that in any homogenous space  $\mathcal{F}$  the operator  $\mathscr{L}_{\mathcal{F}}$  commutes with the operators  $\tilde{S}(f)$ ,  $f \in L^1(\mathbb{R})$ .

**Lemma 3.6.** For all  $f \in L^1(\mathbb{R})$  and  $x \in D(\mathcal{L}_F)$  we have

$$\mathscr{L}_{\mathcal{F}}\tilde{S}(f)x = \tilde{S}(f)\mathscr{L}_{\mathcal{F}}x.$$

*Proof.* Assume  $\lambda > \omega_0$  where  $\omega_0$  is the growth bound of the semigroup T. From (3.5) we deduce that  $R(\lambda, \mathcal{L}_{\mathcal{F}})\tilde{S}(f) = \tilde{S}(f)R(\lambda, \mathcal{L}_{\mathcal{F}})$ . Let  $x \in D(\mathcal{L}_{\mathcal{F}})$  and  $y = (\mathcal{L}_{\mathcal{F}} - \lambda I)x$ . Then

$$\tilde{S}(f)x = \tilde{S}(f)R(\lambda, \mathscr{L}_{\mathcal{F}})(\mathscr{L}_{\mathcal{F}} - \lambda I)x = R(\lambda, \mathscr{L}_{\mathcal{F}})\tilde{S}(f)y$$

implies  $\tilde{S}(f)x \in D(\mathscr{L}_{\mathcal{F}})$  and

$$(\mathscr{L}_{\mathcal{F}} - \lambda I)\tilde{S}(f)x = \tilde{S}(f)y,$$

from where the result immediately follows.

Corollary 3.7. Assume  $\Delta \subset \mathbb{R}$  is closed. Then the spectral submodule  $\mathcal{F}(\Delta)$  is an invariant subspace of the operator  $\mathcal{L}_{\mathcal{F}}$  and the restriction of  $\mathcal{L}_{\mathcal{F}}$  to  $\mathcal{F}(\Delta)$  coincides with the operator  $\mathcal{L}_{\mathcal{F}(\Delta)}$  given by Definition 3.1.

*Proof.* Assume  $x \in \mathcal{F}(\Delta)$  and  $y = \mathscr{L}_{\mathcal{F}}x$ . Let  $f \in L^1(\mathbb{R})$  be such that supp  $\hat{f} \cap \Delta = \emptyset$ . Then  $0 = f * x = \mathscr{L}_{\mathcal{F}}(f * x) = f * (\mathscr{L}_{\mathcal{F}}y)$  and, therefore,  $y \in \mathcal{F}(\Delta)$ .

Next, we use the above commutativity relation to extend the result of Lemma 3.1 to the non-homogeneous case.

**Lemma 3.8.** Assume  $x \in D(\mathcal{L}_{\mathcal{F}})$  and  $y = \mathcal{L}_{\mathcal{F}}x$ . Then

$$\Lambda(x) \subseteq \Lambda(y) \cup \{\lambda \in \mathbb{R} : i\lambda \in \sigma(A)\}.$$

Proof. Assume  $\lambda \notin \Delta_0 = \Lambda(y) \cup \{\lambda \in \mathbb{R} : i\lambda \in \sigma(A)\}$  and let  $f \in L^1(\mathbb{R})$  be such that  $\hat{f}(\lambda) \neq 0$ , supp  $\hat{f}$  is compact, and supp  $\hat{f} \cap \Delta_0 = \emptyset$ . From Definition 3.1, Lemma 3.6, and Lemma 2.1(iv) we deduce that  $f * x \in D(\mathcal{L}_{\mathcal{F}})$  and  $\mathcal{L}_{\mathcal{F}}(f * x) = f * y = 0$ . Hence, Lemma 3.1 implies  $i\Lambda(f * x) \subseteq \sigma(A) \cap (i\mathbb{R})$ . On the other hand,  $\Lambda(f * x) \subseteq \text{supp } \hat{f} \cap \Lambda(x)$  from Lemma 2.1(iii). Hence,  $\Lambda(f * x) = \emptyset$ , f \* x = 0, and  $\lambda \notin \Lambda(x)$ .  $\square$ 

The following corollary is immediate in view of Lemma 2.1(i).

Corollary 3.9. Assume  $\Delta \subset \mathbb{R}$  is closed and

$$(3.7) i\Delta \cap \sigma(A) = \emptyset.$$

Then  $\ker \mathscr{L}_{\mathcal{F}(\Delta)} = \{0\}.$ 

**Theorem 3.10.** Assume  $\Delta \subset \mathbb{R}$  is compact and the generator A satisfies (3.7). Then the operator  $\mathcal{L}_{\mathcal{F}(\Delta)}$  is invertible.

*Proof.* In view of Corollary 3.9 we only need to prove that the operator  $\mathcal{L}_{\mathcal{F}(\Delta)}$  is onto. Let  $y \in \mathcal{F}(\Delta)$ , i.e.  $\Lambda(y) \subseteq \Delta$ . Let also  $f \in L^1(\mathbb{R})$  be such that  $\hat{f} = 1$  in a neighborhood of  $\Delta$ ,  $i \operatorname{supp} \hat{f} \subset \rho(A)$  is compact, and  $\hat{f} \in C^{\infty}$ , i.e.  $\hat{f}$  is differentiable infinitely many times. Consider the function  $F \in L^1(\mathbb{R}, B(\mathcal{F}))$  defined by

$$F(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\lambda) R(i\lambda, A) e^{i\lambda t} d\lambda, t \in \mathbb{R};$$

the above integral makes sense because  $i \operatorname{supp} \hat{f} \subset \rho(A)$ . Observe that

(3.8) 
$$\hat{F}(\lambda) = \begin{cases} \hat{f}(\lambda)R(i\lambda, A), & i\lambda \in \rho(A); \\ 0, & i\lambda \notin \rho(A); \end{cases}$$

is compactly supported and infinitely differentiable, and  $F * y \in \mathcal{F}(\Delta)$  by Lemma 2.1(iii). Moreover, since the operators  $\mathscr{L}$  and A are closed, we can write

$$-\frac{d}{dt}(F*y)(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \int_{\mathbb{R}} i\lambda \hat{F}(\lambda) e^{i\lambda(t-s)} y(s) d\lambda ds$$

and

$$A(F * y)(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \int_{\mathbb{R}} \hat{f}(\lambda)(I + i\lambda R(i\lambda, A))e^{i\lambda(t-s)}y(s)d\lambda ds.$$

Hence,  $F * y \in D(\mathcal{L})$  and  $\mathcal{L}(F * y) = f * y = y$ , where the last equality follows from Lemma 2.1(iv).

We conclude this paragraph with a result on the spectrum of the operator  $\mathcal{L}_F$  in a spectrally homogeneous space. For several specific homogeneous spaces this result was proved in [5].

**Theorem 3.11.** Assume  $\mathcal{F}$  is a spectrally homogeneous space. Then

$$\sigma(\mathscr{L}_{\mathcal{F}}) = \sigma(\mathscr{L}_{\mathcal{F}}) + i\mathbb{R}.$$

*Proof.* Directly from Definition 3.1 and (2.3) we have

$$V(\lambda)\mathscr{L}_{\mathcal{F}}V(-\lambda) = \mathscr{L}_{\mathcal{F}} + i\lambda I, \ \lambda \in \mathbb{R},$$

and the result follows.

## 4. Invertibility of the operator $\mathscr{L}$ in $\mathcal{F}_{as}$

The other main result of this paper is the following theorem.

**Theorem 4.1.** Let  $\mathcal{F} = \mathcal{F}(\mathbb{R}, X)$  be a homogeneous function space and  $\mathcal{F}_{as} \subset \mathcal{F}$  be the space of functions with the absolutely summable spectrum. Assume that an operator  $\mathcal{L} = \mathcal{L}_{\mathcal{F}_{as}} : D(\mathcal{L}) \subseteq \mathcal{F}_{as} \to \mathcal{F}_{as}$  from Definition 3.1 is such that the generator A of the semigroup T satisfies (1.3) and (1.4). Then the operator  $\mathcal{L}$  is invertible,  $\mathcal{L}^{-1} \in B(\mathcal{F}_{as})$ , and

$$(4.1) \|\mathscr{L}^{-1}\| \le \frac{18}{\pi} M \left( 4 + 4M + 2M^2 \right)^{1/2}, \ M = \sup_{\lambda \in i\mathbb{R}} \|R(\lambda, A)\|.$$

The proof of the above result is based on several lemmas.

**Lemma 4.2.** Assume that  $\Phi \in C^2(\mathbb{R}, B(X))$  and  $\phi \in L^1(\mathbb{R})$  is defined by (2.7), i.e.

$$\hat{\phi}(\lambda) = \hat{\phi}_0(\lambda) = (1 - |\lambda|)\chi_{[-1,1]}(\lambda), \ \lambda \in \mathbb{R}.$$

Let  $\Phi_0 = (\Phi \hat{\phi})^{\vee}$ , i.e.  $\Phi_0$  is the inverse Fourier transform of the function  $\Phi \hat{\phi}$ . Then  $\Phi_0 \in L^1(\mathbb{R}, B(X))$  and

*Proof.* Observe that the definition of  $\Phi_0$  implies that  $\Phi_0 \in C_b(\mathbb{R}, B(X))$  and

(4.3) 
$$\|\Phi_0(t)\| \le \frac{1}{2\pi} \|\Phi\|_{\infty}, \ t \in \mathbb{R}.$$

We also have

$$\Phi_0(t) = \frac{1}{2\pi} \int_{\mathbb{R}} \Phi(\lambda) \hat{\phi}(\lambda) e^{i\lambda t} d\lambda = \frac{1}{2\pi} \int_{-1}^1 \Phi(\lambda) (1 - |\lambda|) e^{i\lambda t} d\lambda$$

$$= \frac{1}{2\pi} \left( \int_{-1}^{0} \Phi(\lambda)(1+\lambda) e^{i\lambda t} d\lambda + \int_{0}^{1} \Phi(\lambda)(1-\lambda) e^{i\lambda t} d\lambda \right).$$

Applying integration by parts in the above integrals, we get

$$\Phi_0(t) = \frac{1}{2\pi i t} \Big[ \Phi(0) - \int_{-1}^0 (\Phi(\lambda) + \Phi'(\lambda)(1+\lambda)) e^{i\lambda t} d\lambda - \Phi(0) - \int_0^1 (-\Phi(\lambda) + \Phi'(\lambda)(1-\lambda)) e^{i\lambda t} d\lambda \Big]$$

$$= \frac{1}{2\pi i t} \Big[ \int_0^1 \Phi(\lambda) e^{i\lambda t} d\lambda - \int_{-1}^0 \Phi(\lambda) e^{i\lambda t} d\lambda - \int_{-1}^1 \Phi'(\lambda)(1-|\lambda|) e^{i\lambda t} d\lambda \Big].$$

Applying integration by parts once again, we get

$$\Phi_{0}(t) = \frac{1}{2\pi t^{2}} \Big[ 2\Phi(0) - \Phi(1)e^{it} - \Phi(-1)e^{-it} + 2\int_{0}^{1} \Phi'(\lambda)e^{i\lambda t}d\lambda - 2\int_{-1}^{0} \Phi'(\lambda)e^{i\lambda t}d\lambda - \int_{-1}^{1} \Phi''(\lambda)(1 - |\lambda|)e^{i\lambda t}d\lambda \Big].$$

Hence, for  $t \neq 0$ , we have

Using (4.3) and (4.4) we get  $\Phi_0 \in L^1(\mathbb{R}, B(X))$  since for any  $\alpha > 0$ 

$$\begin{split} \|\Phi_0\|_1 &= \int_{|t| \le \alpha} \|\Phi_0(t)\| dt + \int_{|t| \ge \alpha} \|\Phi_0(t)\| dt \\ &\le \frac{1}{\pi} \left( \alpha \|\Phi\|_{\infty} + (4\|\Phi\|_{\infty} + 4\|\Phi'\|_{\infty} + \|\Phi''\|_{\infty}) \int_{\alpha}^{\infty} \frac{dt}{t^2} \right) \\ &= \frac{1}{\pi} \left( \alpha \|\Phi\|_{\infty} + \frac{1}{\alpha} (4\|\Phi\|_{\infty} + 4\|\Phi'\|_{\infty} + \|\Phi''\|_{\infty}) \right). \end{split}$$

Plugging in 
$$\alpha = \left(\frac{4\|\Phi\|_{\infty} + 4\|\Phi'\|_{\infty} + \|\Phi''\|_{\infty}}{\|\Phi\|_{\infty}}\right)^{1/2}$$
 we get (4.2).

Corollary 4.3. Assume that  $\Phi \in C^2(\mathbb{R}, B(X))$  and  $\phi_n \in L^1(\mathbb{R})$  is defined by (2.6) and (2.7). Let  $\Phi_n = (\Phi \hat{\phi}_n)^{\vee}$ . Then  $\Phi_n \in L^1(\mathbb{R}, B(X))$  and

*Proof.* The result follows by applying the lemma to the function  $S(-n)\Phi$ , where S is the translation representation (2.2).

**Corollary 4.4.** Assume that the generator A of a semigroup T satisfies (1.3) and (1.4). Let  $R_n = (R(i \cdot A)\hat{\phi}_n)^{\vee}$ . Then  $R_n \in L^1(\mathbb{R}, B(X))$  and

(4.6) 
$$||R_n||_1 \le \frac{2}{\pi} M \left(4 + 4M + 2M^2\right)^{1/2}.$$

*Proof.* The result follows from (4.5) since  $\frac{d}{d\lambda}R(i\lambda, A) = iR^2(i\lambda, A)$  and  $\frac{d^2}{d\lambda^2}R(i\lambda, A) = 2R^3(i\lambda, A)$ .

We are now ready to complete the proof of Theorem 4.1.

*Proof of Theorem 4.1.* Since  $\mathcal{F}_{as}$  is itself a homogeneous space, Corollary 3.2 applies and the operator  $\mathcal{L} = \mathcal{L}_{\mathcal{F}_{as}}$  is injective.

To prove surjectivity, consider  $y \in \mathcal{F}_{as}$ . Let  $y_n = \phi_n * y$  so that  $\Lambda(y_n) \subseteq [n-1,n+1] = \Delta(n)$ . Let  $R_n$  be defined as in Corollary 4.4. Then from the proof of Theorem 3.10 and Lemma 2.1(v) we see that  $x_n = R_n * y = R_n * (y_{n-1} + y_n + y_{n+1}) \in \mathcal{F}(\Delta(n))$  satisfies  $\mathcal{L}x_n = y_n$ . Let  $x = \sum_{n \in \mathbb{Z}} x_n$ , where the series converges absolutely since  $y \in \mathcal{F}_{as}$  and  $R_n, n \in \mathbb{Z}$ , satisfy (4.6). Since the operator  $\mathcal{L}$  is closed, we conclude that  $\mathcal{L}x = y$ .

It remains to estimate  $||x||_{as}$ . Observe that

$$\phi_n * x = \phi_n * (x_{n-1} + x_n + x_{n+1}).$$

Hence, (4.6) implies

$$\|\phi_n * x\| \le \frac{2}{\pi} M \left(4 + 4M + 2M^2\right)^{1/2} \sum_{k=n-1}^{n+1} (\|y\|_{k-1} + \|y\|_k + \|y\|_{k+1}),$$

and the postulated estimate for  $\|\mathcal{L}\|^{-1}$  follows.

Remark 4.1. Observe that if  $\mathcal{F} = C_{ub}$ , and  $\mathscr{L}_{\mathcal{F}}$  satisfies the conditions of Theorem 4.1, then  $\mathscr{L}_{\mathcal{F}}\mathcal{F} \supset \mathscr{L}_{\mathcal{F}}\mathcal{F}_{as} \supset C^{1,\alpha}$ ,  $\alpha > 0$ . Moreover, for any function  $y \in C^{1,\alpha}$  the equation  $\mathscr{L}_{\mathcal{F}}x = y$  has a unique solution  $x \in \mathcal{F}_{as}$ .

## 5. Examples

We begin this section with an example of an operator A and a homogeneous space  $\mathcal{F}(\mathbb{R}, X)$  such that A satisfies (1.3) and (1.4) but the operator  $\mathcal{L} = \mathcal{L}_{\mathcal{F}}$  is not invertible. This examples appears in [20, Counterexample IV.2.7], we provide it in order to point out a feature that seems to be common for all such examples.

We let

$$X = C_0(\mathbb{R}^+) \cap L^1_{\nu}(\mathbb{R}^+), \ \nu(s) = e^s,$$

where  $L^1_{\nu}$  is the Beurling algebra of all measurable functions that are summable with the weight  $\nu$ . The norm in  $L^1_{\nu}(\mathbb{R}^+)$  is given by

$$||x||_{\nu} = \int_0^\infty |x(s)|e^s ds$$

so that  $||x||_X = ||x||_\infty + ||x||_\nu$ . We let  $A = \frac{d}{ds}$  be the generator of the semigroup  $T: \mathbb{R}_+ \to X$  given by  $T(t)x(s) = x(s+t), \ x \in X$ ,  $s,t \in \mathbb{R}^+$ . Observe that  $||T(t)|| = 1, \ t \geq 0$ , the growth bound of T is equal to 0, and the spectral bound  $s(A) = \sup\{\Re e\lambda, \lambda \in \sigma(A)\}$  satisfies  $s(A) \leq -1$ . Since

$$R(i\lambda, A)x(s) = \int_0^\infty e^{-\lambda t}x(s+t)dt,$$

the operator A indeed satisfies (1.3) and (1.4). However, if we let  $\mathcal{F}$  be, for example,  $C_0(\mathbb{R}, X)$ , Theorem 1.1 would imply that  $\mathcal{L}_{\mathcal{F}}$  is not invertible since the spectral radius of the operator T(1) is equal to 1. Observe that in this case  $\sigma(T(1)) = \mathbb{T}$ . To the best of our knowledge, the semigroups in all of the known examples of this kind have this property.

We conclude the paper with an application of our results to the following non-linear equation in  $\mathcal{F} = C_b$ :

(5.1) 
$$x'(t) = (Ax)(t) + y(t) + F(x(t)), \ y \in \mathcal{F}_{as},$$

where F is the polynomial

$$F(z) = F_1(z) + F_2(z, z) + \ldots + F_n(z, z, \ldots, z), \ z \in X,$$

and each  $F_k$ , k = 1, ..., n, is a k-linear map.

We assume that the operator A satisfies (1.3) and (1.4). Then  $x \in \mathcal{F}_{as}$  is a *mild solution* of the equation (5.1) if it satisfies

$$x = z + \Phi x$$
,

where  $z = \mathscr{L}_{\mathcal{F}_{as}}^{-1} y$  and the non-linear map  $\Phi : \mathcal{F}_{as} \to \mathcal{F}_{as}$  is given by  $\Phi = \mathscr{L}_{\mathcal{F}_{as}}^{-1} \circ F$ . Observe that the map  $\Phi$  is Lipschitz in any ball  $B_{\beta}(0)$  of radius  $\beta$  centered at  $0 \in \mathcal{F}_{as}$ , that is

$$\|\Phi(x) - \Phi(y)\|_{as} \le L_{F,A}(\beta) \|x - y\|_{as}, \ x, y \in B_{\beta}(0).$$

Moreover, because of (4.1), the Lipschitz constant satisfies

(5.2) 
$$L_{F,A}(\beta) \le \frac{18}{\pi} M \left( 4 + 4M + 2M^2 \right)^{1/2} \sum_{k=1}^{n} k \|F_k\| \beta^{k-1}.$$

**Theorem 5.1.** Assume that  $\beta > 0$ ,  $y \in \mathcal{F}_{as}$ , and F are such that  $L_{F,A}(\beta) < 1$  and  $\|\Phi(z)\|_{\mathcal{F}_{as}} < \beta(1 - L_{A,F}(\beta))$ . Then the non-linear equation (5.1) has a unique mild solution  $x \in \mathcal{F}_{as}$  and  $\|x - z\|_{as} \leq \beta$ .

*Proof.* The solution is obtained by the method of simple iterations as in [19, Theorem 10.1.2].  $\Box$ 

Remark 5.1. Results of this paper can be extended in a straightforward way to the case of differential inclusions

$$\frac{dx}{dt} \in \mathscr{A}x + y, \ x, y \in \mathcal{F}_{as}(\mathbb{R}, X),$$

where  $\mathscr{A}$  is a linear relation on X [18, 8, 10, 13].

#### References

- [1] W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace transforms and Cauchy problems*, vol. 96 of Monographs in Mathematics, Birkhäuser/Springer Basel AG, Basel, second ed., 2011.
- [2] R. Balan and I. Krishtal, An almost periodic noncommutative Wiener's lemma, J. Math. Anal. Appl., 370 (2010), pp. 339–349.
- [3] A. G. Baskakov, Spectral tests for the almost periodicity of the solutions of functional equations, Mat. Zametki, 24 (1978), pp. 195–206, 301. English translation: Math. Notes 24 (1978), no. 1–2, pp. 606–612 (1979).
- [4] ——, Spectral analysis of linear differential operators, and semigroups of difference operators, Dokl. Akad. Nauk, 343 (1995), pp. 295–298.
- [5] —, Semigroups of difference operators in the spectral analysis of linear differential operators, Funktsional. Anal. i Prilozhen., 30 (1996), pp. 1–11, 95. English translation: Funct. Anal. Appl. 30 (1996), no. 3, pp. 149–157 (1997).
- [6] —, Asymptotic estimates for elements of matrices of inverse operators, and harmonic analysis, Sibirsk. Mat. Zh., 38 (1997), pp. 14–28, i. English translation: Siberian Math. J. 38 (1997), no. 1, pp. 10–22.
- [7] —, Theory of representations of Banach algebras, and abelian groups and semigroups in the spectral analysis of linear operators, Sovrem. Mat. Fundam. Napravl., 9 (2004), pp. 3–151 (electronic). English translation: J. Math. Sci. (N. Y.) 137 (2006), no. 4, pp. 4885–5036.
- [8] ——, Linear relations as generators of semigroups of operators, Mat. Zametki, 84 (2008), pp. 175–192.
- [9] —, Analysis of linear differential equations by methods of the spectral theory of difference operators and linear relations, Uspekhi Mat. Nauk, 68 (2013), pp. 77–128. English translation: Russian Math. Surveys 68 (2013), no. 1, pp. 69–116.
- [10] A. G. BASKAKOV AND K. I. CHERNYSHOV, Spectral analysis of linear relations, and degenerate semigroups of operators, Mat. Sb., 193 (2002), pp. 3–42. English translation: Sb. Math. 193 (2002), no. 11–12, pp. 1573–1610.
- [11] A. G. BASKAKOV AND N. S. KALUZHINA, Beurling's theorem for functions with essential spectrum from homogeneous spaces and stabilization of solutions of parabolic equations, Mathematical Notes, 92 (2012), pp. 587–605.
- [12] A. G. BASKAKOV AND I. A. KRISHTAL, Harmonic analysis of causal operators and their spectral properties, Izv. Ross. Akad. Nauk Ser. Mat., 69 (2005), pp. 3–54. English translation: Izv. Math. 69 (2005), no. 3, pp. 439–486.
- [13] —, On completeness of spectral subspaces of linear relations and ordered pairs of linear operators, J. Math. Anal. Appl., 407 (2013), pp. 157–178.

- [14] ——, Memory estimation of inverse operators, J. Funct. Anal., 267 (2014), pp. 2551–2605.
- [15] A. G. BASKAKOV AND Y. N. SINTYAEV, Difference operators in the investigation of differential operators: estimates for solutions, Differ. Uravn., 46 (2010), pp. 210–219. English translation: Differ. Equ. 46 (2010), no. 2, 214–223.
- [16] P. L. Butzer and H. Berens, Semi-groups of operators and approximation, Die Grundlehren der mathematischen Wissenschaften, Band 145, Springer-Verlag New York Inc., New York, 1967.
- [17] C. CHICONE AND Y. LATUSHKIN, Evolution semigroups in dynamical systems and differential equations, vol. 70 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 1999.
- [18] R. Cross, *Multivalued linear operators*, vol. 213 of Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, Inc., New York, 1998.
- [19] J. DIEUDONNÉ, Foundations of modern analysis, Pure and Applied Mathematics, Vol. X, Academic Press, New York, 1960.
- [20] K.-J. ENGEL AND R. NAGEL, One-parameter semigroups for linear evolution equations, vol. 194 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.
- [21] H. G. FEICHTINGER, Banach convolution algebras of Wiener type, in Proc. Conf. "Functions, Series, Operators", Budapest, Aug. 1980, vol. 35 of Coll. Math. Soc. Bolyai Janos, 1980, pp. 509–524.
- [22] H. G. FEICHTINGER AND P. GRÖBNER, Banach spaces of distributions defined by decomposition methods. I, Math. Nachr., 123 (1985), pp. 97–120.
- [23] J. J. F. FOURNIER AND J. STEWART, Amalgams of  $L^p$  and  $l^q$ , Bull. Amer. Math. Soc. (N.S.), 13 (1985), pp. 1–21.
- [24] L. Gearhart, Spectral theory for contraction semigroups on Hilbert space, Trans. Amer. Math. Soc., 236 (1978), pp. 385–394.
- [25] G. Greiner, J. Voigt, and M. Wolff, On the spectral bound of the generator of semigroups of positive operators, J. Operator Theory, 5 (1981), pp. 245–256.
- [26] K. GRÖCHENIG, Foundations of time-frequency analysis, Applied and Numerical Harmonic Analysis, Birkhäuser Boston Inc., Boston, MA, 2001.
- [27] K. GRÖCHENIG AND A. KLOTZ, Noncommutative approximation: inverseclosed subalgebras and off-diagonal decay of matrices, Constr. Approx., 32 (2010), pp. 429–466.
- [28] E. HEWITT AND K. A. ROSS, Abstract harmonic analysis. Vol. I, vol. 115 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, second ed., 1979. Structure of topological groups, integration theory, group representations.
- [29] N. P. Korneĭčuk, Ekstremalnye zadachi teorii priblizheniya, Izdat. "Nauka", Moscow, 1976.
- [30] Y. LATUSHKIN AND S. MONTGOMERY-SMITH, Lyapunov theorems for Banach spaces, Bull. Amer. Math. Soc. (N.S.), 31 (1994), pp. 44–49.
- [31] Y. Latushkin and S. Montgomery-Smith, Evolutionary semigroups and Lyapunov theorems in Banach spaces, J. Funct. Anal., 127 (1995), pp. 173–197.

- [32] Y. LATUSHKIN AND R. SHVYDKOY, Hyperbolicity of semigroups and Fourier multipliers, in Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000), vol. 129 of Oper. Theory Adv. Appl., Birkhäuser, Basel, 2001, pp. 341–363.
- [33] B. M. LEVITAN AND V. V. ZHIKOV, Almost periodic functions and differential equations, Cambridge University Press, Cambridge, 1982. Translated from the Russian by L. W. Longdon.
- [34] V. Q. Phong, On stability of  $C_0$ -semigroups, Proc. Amer. Math. Soc., 129 (2001), pp. 2871–2879.
- [35] J. PRÜSS, On the spectrum of  $C_0$ -semigroups, Trans. Amer. Math. Soc., 284 (1984), pp. 847–857.
- [36] H. Reiter, Classical harmonic analysis and locally compact groups, Clarendon Press, Oxford, 1968.

DEPARTMENT OF APPLIED MATHEMATICS AND MECHANICS, VORONEZH STATE UNIVERSITY, RUSSIA 394693, EMAIL: MMIO@AMM.VSU.RU

DEPARTMENT OF MATHEMATICAL SCIENCES, NORTHERN ILLINOIS UNIVERSITY, DEKALB, IL 60115, EMAIL: KRISHTAL@MATH.NIU.EDU