

GROUPS OF COMPONENTS AND WEIL RESTRICTION

ALESSANDRA BERTAPELLE AND CRISTIAN D. GONZÁLEZ-AVILÉS

ABSTRACT. We determine the behavior under Weil restriction of the group of connected components of the special fiber of an arbitrary smooth group scheme (whose Weil restriction exists) over an arbitrary (commutative and unital) local ring. Applications to Néron models are given.

1. INTRODUCTION

Let R be a discrete valuation ring with fraction field K and residue field k and let k^s be a fixed separable algebraic closure of k . Let A_K be an abelian variety over K with Néron model A over R and let $\pi_0(A_s)$ denote the k -group scheme of connected components of the special fiber A_s of A . Let A_K^\vee be the dual (i.e., Picard) variety of A_K and write A^\vee and $\pi_0(A_s^\vee)$ for the corresponding objects associated to the abelian variety A_K^\vee . When k is perfect, Grothendieck's pairing $\pi_0(A_s)(k^s) \times \pi_0(A_s^\vee)(k^s) \rightarrow \mathbb{Q}/\mathbb{Z}$ [13, IX, (1.2.2)] has been verified to be perfect in all cases except when K has positive characteristic and k is infinite; see [2, §1] for further comments. More recently a general proof of the perfectness of Grothendieck's pairing has been announced in [17]. However, when k is *imperfect*, there exist examples that show that the above pairing is no longer perfect. The first such examples were constructed in [2, comment after Corollary 2.5] using the Weil restriction functor. It is for this reason, at least, that a full understanding of the behavior of the groups of connected components of abelian (or more general smooth group) varieties under Weil restriction is desirable. Previously, this has been achieved only in the case of abelian varieties under various restrictions. See Remark 4.3. In this paper we completely determine the behavior under Weil restriction of the group of connected components of the special fiber of an arbitrary (possibly non commutative) smooth group scheme over an arbitrary (possibly non noetherian) local ring. Our main result is the following

Theorem 1.1. *Let R'/R be a finite flat extension of local rings with associated residue field extension k'/k . Let G' be a smooth R' -group scheme such that the Weil restriction $\mathfrak{R}_{R'/R}(G')$ exists. Then $\mathfrak{R}_{k'/k}(\pi_0(G'_s))$ exists and there exists a canonical isomorphism of étale k -group schemes*

$$\pi_0(\mathfrak{R}_{R'/R}(G')_s) \simeq \mathfrak{R}_{k'/k}(\pi_0(G'_s)).$$

Date: March 1, 2024.

2010 Mathematics Subject Classification. 14L15, 11G99.

Key words and phrases. group of connected components, Weil restriction, Grothendieck's pairing.

A. B. was partially supported by PRAT 2013 "Arithmetic of Varieties over Number Fields" CPDA 135371/13.

C. G.-A. was partially supported by Fondecyt grant 1120003.

The theorem applies, in particular, to smooth and quasi-projective R' -group schemes since $\mathfrak{R}_{R'/R}(G')$ is well-known to exist in this case. More general conditions that guarantee the existence of $\mathfrak{R}_{R'/R}(G')$ can be found in [4, §7.6, Theorem 4, p. 194] and [3, Theorem 2.15].

The proof of the theorem, which is simpler than the proofs of the particular cases alluded to above, essentially relies only on basic properties of the Weil restriction functor and the following key fact: if k is a field, R' is a finite local k -algebra and G' is a smooth R' -group scheme with geometrically connected special fiber, then $\mathfrak{R}_{R'/k}(G')$ is geometrically connected.

Some applications of Theorem 1.1 are discussed at the end of the paper.

We thank Brian Conrad for helping us shorten our original proof of the main theorem. We also thank Bas Edixhoven and Dino Lorenzini for helpful comments.

2. PRELIMINARIES

All rings considered in this paper are commutative and unital. If X is a scheme, the topological space underlying X will be denoted by $|X|$. The identity morphism of X will be denoted by 1_X .

Let S be a scheme, let X be an S -scheme and let $S'' \rightarrow S' \rightarrow S$ be morphisms of schemes. We will make the identification

$$(2.1) \quad (X \times_S S') \times_{S'} S'' = X \times_S S''.$$

If $A \rightarrow B$ is a ring homomorphism and X is an A -scheme, X_B will denote $X \times_{\mathrm{Spec} A} \mathrm{Spec} B$.

Let $W \rightarrow S' \rightarrow S$ be morphisms of schemes. If Z is an S -scheme and $\mathrm{pr}_1: Z \times_S S' \rightarrow Z$ is the first projection then, by the universal property of the fiber product, the map

$$(2.2) \quad \mathrm{Hom}_{S'}(W, Z \times_S S') \xrightarrow{\sim} \mathrm{Hom}_S(W, Z), \quad g \mapsto \mathrm{pr}_1 \circ g,$$

is a bijection.

Let A be an artinian local ring with residue field k and let G be an A -group scheme *locally of finite type*. We will identify G with the functor on (Sch/A) represented by G . Let $G_s := G \times_{\mathrm{Spec} A} \mathrm{Spec} k$ be the special fiber of G ¹. Now let $|G|^0 \subset |G|$ denote the connected component of the identity $1 \in G$ and define a subfunctor G^0 of G by

$$(2.3) \quad G^0(T) = \{u \in G(T) : u(|T|) \subseteq |G|^0\},$$

where T is any A -scheme. Since G is locally of finite type over A , (2.3) is represented by an open and normal subgroup scheme G^0 of G [7, VI_A, Proposition 2.3.1]. Note also that, since every subgroup scheme of G is closed [7, VI_A, Corollary 0.5.2], G^0 is open and closed in G . Assume now that G is locally of finite type and *flat* over A . Then, by [7, VI_A, Proposition 5.5.1(i) and Theorem 3.2(vi)], the fppf quotient sheaf of groups

$$\pi_0(G) := G^0 \backslash G$$

¹ Here the subscript “s” indicates “special fiber” rather than the closed point of $\mathrm{Spec} A$. We use this notation because it remains unchanged by a change of rings, i.e., if A' is another artinian local ring with residue field k' and G' is an A' -group scheme, then it should be clear that G'_s means $G' \times_{\mathrm{Spec} A'} \mathrm{Spec} k'$.

is represented by an étale A -group scheme and the canonical morphism

$$(2.4) \quad p_G: G \rightarrow \pi_0(G)$$

is faithfully flat and locally of finite presentation. Further, by [7, VI_B, Proposition 3.3], there exists a canonical isomorphism of k -group schemes $(G_s)^0 \simeq (G^0)_s$ and therefore $\pi_0(G_s) \simeq \pi_0(G)_s$.

Note that, if G is *smooth* over A , then G^0 is smooth as well and therefore (2.4) is a smooth morphism by [11, IV₄, Proposition 17.5.1] and [7, VI_B, Proposition 1.3].

Let $f: S' \rightarrow S$ be a morphism of schemes and let X' be an S' -scheme. We will say that *the Weil restriction of X' along f exists* (or, more concisely, that $\mathfrak{R}_{S'/S}(X')$ *exists*) if the contravariant functor $(\text{Sch}/S) \rightarrow (\text{Sets}), T \mapsto \text{Hom}_{S'}(T \times_S S', X')$, is representable, i.e., if there exists a pair $(\mathfrak{R}_{S'/S}(X'), q_{X', S'/S})$, where $\mathfrak{R}_{S'/S}(X')$ is an S -scheme and $q_{X', S'/S}: \mathfrak{R}_{S'/S}(X')_{S'} \rightarrow X'$ is an S' -morphism of schemes, such that the map

$$(2.5) \quad \text{Hom}_S(T, \mathfrak{R}_{S'/S}(X')) \xrightarrow{\sim} \text{Hom}_{S'}(T \times_S S', X'), \quad g \mapsto q_{X', S'/S} \circ g_{S'}$$

is a bijection [10, (1.1.8), p. 22]. The pair $(\mathfrak{R}_{S'/S}(X'), q_{X', S'/S})$ (or, more concisely, the scheme $\mathfrak{R}_{S'/S}(X')$) is called the *Weil restriction of X' along f* . If $S' = \text{Spec } A'$ and $S = \text{Spec } A$ are affine, we will write $(\mathfrak{R}_{A'/A}(X'), q_{X', A'/A})$ for $(\mathfrak{R}_{S'/S}(X'), q_{X', S'/S})$ and refer to $\mathfrak{R}_{A'/A}(X')$ as the *Weil restriction of X' along A'/A* .

If X' is an S' -scheme such that $\mathfrak{R}_{S'/S}(X')$ exists and $T \rightarrow S$ is any morphism of schemes, then $\mathfrak{R}_{S'_T/T}(X' \times_{S'} S'_T)$ exists as well and (2.1), (2.2) and (2.5) yield a canonical isomorphism of T -schemes

$$(2.6) \quad \mathfrak{R}_{S'/S}(X') \times_S T \xrightarrow{\sim} \mathfrak{R}_{S'_T/T}(X' \times_{S'} S'_T).$$

Further, if $S'' \rightarrow S' \rightarrow S$ are morphisms of schemes and X'' is an S'' -scheme such that $\mathfrak{R}_{S''/S'}(X'')$ exists, then (2.5) implies that $\mathfrak{R}_{S'/S}(\mathfrak{R}_{S''/S'}(X''))$ exists if, and only if, $\mathfrak{R}_{S''/S}(X'')$ exists. If this is the case, then there exists a canonical isomorphism of S -schemes

$$(2.7) \quad \mathfrak{R}_{S'/S}(\mathfrak{R}_{S''/S'}(X'')) \xrightarrow{\sim} \mathfrak{R}_{S''/S}(X'').$$

Let $f: S' \rightarrow S$ be a finite and locally free (i.e., flat and of finite presentation) morphism of schemes and let X' be an S' -scheme. By [4, §7.6, Theorem 4, p. 194], $\mathfrak{R}_{S'/S}(X')$ exists if, for each point $s \in S$, every finite set of points of $X' \times_S \text{Spec } k(s)$ is contained in an open affine subscheme of X' . This condition is satisfied if X' is quasi-projective over S' [11, II, Definition 5.3.1 and Corollary 4.5.4]. A weaker condition for the existence of $\mathfrak{R}_{S'/S}(X')$ is given in [3, Theorem 2.15].

Proposition 2.8. *Let k be a field and let A be a finite local k -algebra with residue field k' . Let X be an étale A -scheme. Then the following hold.*

- (i) $\mathfrak{R}_{k'/A}(X_s)$ exists and is canonically isomorphic to X .
- (ii) $\mathfrak{R}_{A/k}(X)$ and $\mathfrak{R}_{k'/k}(X_s)$ exist and are canonically isomorphic.

Proof. Let T be any A -scheme. Since $T_s \rightarrow T$ is a nilpotent immersion and X is formally étale over A , the canonical map $\text{Hom}_A(T, X) \rightarrow \text{Hom}_{k'}(T_s, X_s)$, $g \mapsto g_s$, is a bijection [11, IV₄, Remark 17.1.2(iv)]. Thus $(\mathfrak{R}_{k'/A}(X_s), q_{X_s, k'/A}) = (X, 1_{X_s})$ is the Weil restriction of X_s along k'/A . Assertion (i) is now clear. Now, by (i), $\mathfrak{R}_{A/k}(X)$ exists if, and only if, $\mathfrak{R}_{A/k}(\mathfrak{R}_{k'/A}(X_s)) \simeq \mathfrak{R}_{k'/k}(X_s)$ (2.7) exists and, if

this is the case, then $\mathfrak{R}_{A/k}(X)$ and $\mathfrak{R}_{k'/k}(X_s)$ are canonically isomorphic étale k -schemes. Thus it remains to show that $\mathfrak{R}_{k'/k}(X_s)$ exists. By [4, §7.6, Theorem 4, p. 194], it suffices to check that any finite set of points of X_s is contained in an affine open subscheme of X_s . Since X_s is an étale k' -scheme, X_s is isomorphic to a sum $\coprod_{i \in I} \text{Spec } k'_i$, where each k'_i is a finite and separable extension of k' [11, IV₄, Corollary 17.4.2(d')]. It is now clear that the stated condition holds. \square

3. PROOF OF THE MAIN THEOREM

An *extension of local rings* is a flat homomorphism $R \rightarrow R'$ of local rings. Then $R \rightarrow R'$ is faithfully flat and therefore injective [16, Chapter 2, (4.A), Corollary, p. 27, and (4.C)(i), p. 28]. Let \mathfrak{m} and \mathfrak{m}' denote the maximal ideals of R and R' , respectively. Then $R \rightarrow R'$ induces an extension of residue fields $k = R/\mathfrak{m} \hookrightarrow k' = R'/\mathfrak{m}'$.

Proposition 3.1. *Let k be a field and let A be a finite local k -algebra with residue field k' . Let G be a smooth A -group scheme such that $\mathfrak{R}_{A/k}(G)$ exists. Then $\mathfrak{R}_{k'/k}(\pi_0(G_s))$ exists and there exists a canonical isomorphism of étale k -group schemes $\pi_0(\mathfrak{R}_{A/k}(G)) \simeq \mathfrak{R}_{k'/k}(\pi_0(G_s))$.*

Proof. The existence assertion follows from Proposition 2.8(ii). Now, since G^0 is an open and closed subgroup scheme of G , $\mathfrak{R}_{A/k}(G^0)$ is an open and closed subgroup scheme of $\mathfrak{R}_{A/k}(G)$ by [4, §7.6, Proposition 2, p. 192]. Further, since $(G^0)_s$ is geometrically connected [7, VI_A, Proposition 2.4(i)], $\mathfrak{R}_{A/k}(G^0)$ is geometrically connected as well by [6, Proposition A.5.9]². It follows that $\mathfrak{R}_{A/k}(G^0) = \mathfrak{R}_{A/k}(G^0)^0$. Now, since $p_G: G \rightarrow \pi_0(G)$ is smooth and surjective, the induced smooth morphism $\mathfrak{R}_{A/k}(G) \rightarrow \mathfrak{R}_{A/k}(\pi_0(G))$ is surjective by [6, Corollary A.5.4(1)]. Thus there exists an exact and commutative diagram of sheaves of groups on $(\text{Sch}/k)_{\text{fppf}}$

$$\begin{array}{ccccccc} 1 & \longrightarrow & \mathfrak{R}_{A/k}(G)^0 & \longrightarrow & \mathfrak{R}_{A/k}(G) & \longrightarrow & \pi_0(\mathfrak{R}_{A/k}(G)) \longrightarrow 1 \\ & & \parallel & & \parallel & & \downarrow \wr \\ 1 & \longrightarrow & \mathfrak{R}_{A/k}(G^0) & \longrightarrow & \mathfrak{R}_{A/k}(G) & \longrightarrow & \mathfrak{R}_{A/k}(\pi_0(G)) \longrightarrow 1, \end{array}$$

which yields a canonical isomorphism of étale k -group schemes $\pi_0(\mathfrak{R}_{A/k}(G)) \simeq \mathfrak{R}_{A/k}(\pi_0(G))$. Now Proposition 2.8(ii) completes the proof. \square

We may now prove Theorem 1.1.

The existence assertion follows from Proposition 2.8(ii). Let $A := R' \otimes_R k = R'/\mathfrak{m}R'$, which is a finite local k -algebra with residue field k' and let $S' = \text{Spec } R'$. Then $G'_A := G' \times_{S'} \text{Spec } A$ is a smooth A -group scheme. Now, since $\mathfrak{R}_{R'/R}(G')$ exists by hypothesis, $\mathfrak{R}_{A/k}(G'_A)$ exists as well and the map (2.6) for $(X', S'/S, T) = (G', R'/R, k)$ is an isomorphism of smooth k -group schemes $\mathfrak{R}_{R'/R}(G')_s \simeq \mathfrak{R}_{A/k}(G'_A)$. On the other hand, Proposition 3.1 yields a canonical isomorphism of étale k -group schemes $\pi_0(\mathfrak{R}_{A/k}(G'_A)) \simeq \mathfrak{R}_{k'/k}(\pi_0(G'_s))$, whence the theorem follows.

²The proof in [loc.cit.] remains valid if the quasi-projectivity assumption made there is replaced by the condition that the Weil restrictions involved exist.

4. SOME APPLICATIONS

We present the following consequences of Theorem 1.1.

Corollary 4.1. *Let R'/R be a finite extension of local rings such that the associated extension of residue fields k'/k is purely inseparable. Let G' be a smooth R' -group scheme such that $\mathfrak{R}_{R'/R}(G')$ exists. Then there exists a canonical isomorphism of étale k' -group schemes $\pi_0(\mathfrak{R}_{R'/R}(G')_s)_{k'} \simeq \pi_0(G'_s)$.*

Proof. If $\chi: \pi_0(\mathfrak{R}_{R'/R}(G')_s) \xrightarrow{\sim} \mathfrak{R}_{k'/k}(\pi_0(G'_s))$ is the isomorphism of Theorem 1.1, then the diagram of étale k' -group schemes

$$\begin{array}{ccc} \pi_0(\mathfrak{R}_{R'/R}(G')_s)_{k'} & \xrightarrow{\chi_{k'}} & \mathfrak{R}_{k'/k}(\pi_0(G'_s))_{k'} \\ \downarrow \pi_0((q_{G', R'/R})_{k'}) & & \downarrow q_{\pi_0(G'_s), k'/k} \\ \pi_0(G'_s) & \xlongequal{\quad} & \pi_0(G'_s) \end{array}$$

commutes. Further, since $\mathrm{Spec} k' \rightarrow \mathrm{Spec} k$ is a finite and locally free universal homeomorphism, the right-hand vertical map in the above diagram is an isomorphism³. Thus the left-hand vertical map is an isomorphism as well, which completes the proof. \square

Corollary 4.2. *Let R'/R be a finite extension of discrete valuation rings with associated residue and fraction field extensions k'/k and K'/K . Let G' be a smooth, separated and commutative R' -group scheme which is a Néron model⁴ of its generic fiber G'_η . Then $\mathfrak{R}_{R'/R}(G')$ exists, is a Néron model of $\mathfrak{R}_{K'/K}(G'_\eta)$ and there exists a canonical isomorphism of étale k -group schemes $\pi_0(\mathfrak{R}_{R'/R}(G')_s) \simeq \mathfrak{R}_{k'/k}(\pi_0(G'_s))$.*

Proof. The existence of $\mathfrak{R}_{R'/R}(G')$, as well as the fact that it is a Néron model of $\mathfrak{R}_{K'/K}(G'_\eta)$, follow from [4, §10.1, proof of Proposition 4, p. 291]. The corollary is now immediate from the theorem. \square

Remark 4.3. In the case where R'/R is a finite extension of discrete valuation rings, the isomorphism of Corollary 4.2 was discussed in [15, Proposition 3.19] when the associated extension of fraction fields K'/K is Galois, k is perfect and G' is the Néron model of an abelian variety over K' . Corollary 4.1 is proved in [2, Proposition 1.1] (see also [9, proof of Theorem 1]) when K'/K is separable, G' is as above and R is henselian. The proofs of both results make use of a filtration introduced by B. Edixhoven.

Let R'/R be a finite extension of discrete valuation rings with associated fraction field extension K'/K and residue field extension k'/k . Let \bar{k} be a fixed algebraic closure of k containing k' and let k^s and $(k')^s$ denote, respectively, the separable closures of k and k' inside \bar{k} . Now let $A_{K'}$ be an abelian variety over K' with Néron model A over R' and recall from [13, IX, (1.2.2)] Grothendieck's pairing

$$(4.4) \quad \pi_0(A_s)((k')^s) \times \pi_0(A_s^\vee)((k')^s) \rightarrow \mathbb{Q}/\mathbb{Z},$$

where A^\vee is the Néron model of the abelian variety $A_{K'}^\vee$, dual to $A_{K'}$. When the preceding pairing is perfect, $\pi_0(A_s)((k')^s)$ and $\pi_0(A_s^\vee)((k')^s)$ are isomorphic as abelian groups. As noted by D. Lorenzini, it is not known, in general, whether

³Proofs of these facts, currently available at <http://arxiv.org/abs/1501.05621v3>, will appear in [3]

⁴Only assumed to be locally of finite type [4, §10.1. Definition 1, p. 289].

$\pi_0(A_s)((k')^s)$ and $\pi_0(A_s^\vee)((k')^s)$ are always isomorphic as abelian groups, i.e., even when (4.4) is not perfect. See [14, p. 1, footnote] and [15, p. 2032]. Regarding this question, we observe the following consequence of the previous corollary.

Corollary 4.5. *Let R'/R be a finite extension of discrete valuation rings with associated fraction field extension K'/K . Let $A_{K'}$ be an abelian variety over K' with dual abelian variety $A_{K'}^\vee$, and let A and A^\vee denote, respectively, the Néron models of $A_{K'}$ and $A_{K'}^\vee$ over R' . Assume that Grothendieck's pairing (4.4) is perfect. Then there exists an isomorphism of abelian groups*

$$\pi_0(\mathfrak{R}_{R'/R}(A)_s)(k^s) \simeq \pi_0(\mathfrak{R}_{R'/R}(A^\vee)_s)(k^s).$$

Proof. By Corollary 4.2, we have

$$\pi_0(\mathfrak{R}_{R'/R}(A)_s)(k^s) \simeq \mathfrak{R}_{k'/k}(\pi_0(A_s))(k^s) = \pi_0(A_s)(k' \otimes_k k^s).$$

Now, by [5, V, §6, no. 7, Theorem 4, p. A.V.34] and an inductive limit argument, $k' \otimes_k k^s$ is a finite reduced k^s -algebra. Consequently (see, e.g., [1, Theorem 8.7, p. 90]), $k' \otimes_k k^s \simeq \prod_{i \in I} k_i$ for some finite set I , where each k_i is a finite field extension of k^s . Thus $\pi_0(A_s)(k' \otimes_k k^s) \simeq \prod_{i \in I} \pi_0(A_s)(k_i)$. Now, since each k_i/k^s is purely inseparable, the associated morphism $\text{Spec } k_i \rightarrow \text{Spec } k^s$ is a universal homeomorphism. Thus, by [12, VIII, Corollary 1.2], the canonical homomorphism $\pi_0(A_s)(k^s) \rightarrow \pi_0(A_s)(k_i)$ is an isomorphism for each $i \in I$. Further, since $(k')^s/k^s$ is also purely inseparable, the canonical homomorphism $\pi_0(A_s)(k^s) \rightarrow \pi_0(A_s)((k')^s)$ is an isomorphism as well. We conclude that $\pi_0(A_s)(k' \otimes_k k^s) \simeq \pi_0(A_s)((k')^s)^I$. Similarly, $\pi_0(A_s^\vee)(k' \otimes_k k^s) \simeq \pi_0(A_s^\vee)((k')^s)^I$. Since $\pi_0(A_s)((k')^s)$ and $\pi_0(A_s^\vee)((k')^s)$ are isomorphic by the perfectness of (4.4), the corollary follows. \square

The interest of the above corollary is that, when K'/K is separable so that $\mathfrak{R}_{K'/K}(A_{K'})$ is an abelian variety (see [8, Proposition 4.1]), Grothendieck's pairing

$$(4.6) \quad \pi_0(\mathfrak{R}_{R'/R}(A)_s)(k^s) \times \pi_0(\mathfrak{R}_{R'/R}(A^\vee)_s)(k^s) \rightarrow \mathbb{Q}/\mathbb{Z}$$

may not be perfect [2, comment after Corollary 2.2], although, by the corollary, the groups involved are isomorphic when (4.4) is perfect. In other words, Corollary 4.5 suggests that the answer to Lorenzini's question may well be positive.

REFERENCES

- [1] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Co., Reading, MA, 1969.
- [2] A. Bertapelle and S. Bosch, *Weil restriction and Grothendieck's duality conjecture*, J. Algebraic Geom. **9** (2000), 155–164.
- [3] A. Bertapelle and C. D. González-Avilés, *The Greenberg functor revisited*. In preparation. Preliminary version available at <http://arxiv.org/abs/1311.0051v3>.
- [4] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Math. **21**, Springer-Verlag, Berlin, 1990.
- [5] N. Bourbaki, *Algebra II*, Chapters 4–7, Masson, Paris, 1990. ISBN 0-387-19375-8.
- [6] B. Conrad, O. Gabber and G. Prasad, *Pseudo-reductive Groups*, New Math. Monogr. **17**, Cambridge University Press, Cambridge, 2010.
- [7] M. Demazure and A. Grothendieck (Eds.), *Schémas en groupes. Séminaire de Géométrie Algébrique du Bois Marie 1962–64 (SGA 3)*, augmented and corrected 2008–2011 re-edition of the original by P. Gille and P. Polo. Available at <http://www.math.jussieu.fr/~polo/SGA3>.
- [8] B. Edixhoven, *Néron models and tame ramification*, Compos. Math. **81** (1992), 291–306.
- [9] B. Edixhoven, Q. Liu and D. Lorenzini, *The p -part of the group of components of a Néron model*, J. Algebraic Geom. **5** (1996), 801–813.

- [10] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique I. Le langage des schémas*, Grundlehren Math. Wiss. **166**, Springer-Verlag, Berlin, 1971.
- [11] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique*, Publ. Math. Inst. Hautes Études Sci. **8** (= EGA II) (1961), **24** (= EGA IV₂) (1965), **32** (= EGA IV₄) (1967).
- [12] A. Grothendieck et al., *Théorie des topos et cohomologie étale des schémas (SGA 4-2)*. *Séminaire de Géométrie Algébrique du Bois Marie 1963-64*, Lecture Notes in Math. **270**, Springer-Verlag, Berlin, 1972.
- [13] A. Grothendieck et al., *Groupes de monodromie en géométrie algébrique (SGA 7-1)*. *Séminaire de Géométrie Algébrique du Bois Marie 1967-69*, Lecture Notes in Math. **288**, Springer-Verlag, Berlin, 1972.
- [14] D. Lorenzini, *Grothendieck's pairing for Jacobians and base change*, J. Number Theory **128** (2008), 1448–1457.
- [15] D. Lorenzini, *Torsion and Tamagawa numbers*, Ann. Inst. Fourier **61**, no. 5 (2011), 1995–2037.
- [16] H. Matsumura, *Commutative Algebra*, Second edition. Math. Lecture Note Ser. **56**. The Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980. ISBN 0-8053-7024-2.
- [17] T. Suzuki, *Grothendieck's pairing on Neron component groups: Galois descent from the semistable case*, arXiv:1410.3046v2.

UNIVERSITÀ DEGLI STUDI DI PADOVA, DIPARTIMENTO DI MATEMATICA, VIA TRIESTE 63, I-35121 PADOVA

E-mail address: `alessandra.bertapelle@unipd.it`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LA SERENA, CISTERNAS 1200, LA SERENA 1700000, CHILE

E-mail address: `cgonzalez@userena.cl`