

# FAMILIES OF CALABI-YAU HYPERSURFACES IN $\mathbb{Q}$ -FANO TORIC VARIETIES

MICHELA ARTEBANI, PAOLA COMPARIN, AND ROBIN GUILBOT

**ABSTRACT.** We provide a sufficient condition for a general hypersurface in a  $\mathbb{Q}$ -Fano toric variety to be a Calabi-Yau variety in terms of its Newton polytope. Moreover, we define a generalization of the Berglund-Hübsch-Krawitz construction in case the ambient is a  $\mathbb{Q}$ -Fano toric variety with torsion free class group and the defining polynomial is not necessarily of Delsarte type. Finally, we introduce a duality between families of Calabi-Yau hypersurfaces which includes both Batyrev and Berglund-Hübsch-Krawitz mirror constructions. This is given in terms of a polar duality between pairs of polytopes  $\Delta_1 \subseteq \Delta_2$ , where  $\Delta_1$  and  $\Delta_2^*$  are canonical.

## INTRODUCTION

A wide class of Calabi-Yau varieties is given by anticanonical hypersurfaces, or more generally complete intersections, in Fano toric varieties. A special interest for such families of Calabi-Yau's arised after the work of Batyrev [5], who defined a duality between the anticanonical linear series of toric Fano varieties which satisfies the requirement of topological mirror symmetry [5, 8]:

$$(1) \quad h_{st}^{p,q}(X) = h_{st}^{n-p,q}(X^*), \quad 0 \leq p, q \leq n,$$

where  $X, X^*$  are general elements in the dual linear series,  $n = \dim(X) = \dim(X^*)$  and  $h_{st}^{p,q}$  denote the string-theoretic Hodge numbers. In the case of hypersurfaces, Batyrev mirror construction relies on the polar duality between reflexive polytopes. A different class of Calabi-Yau varieties can be constructed by considering quotients of quasismooth (or transverse) hypersurfaces in weighted projective spaces, i.e. defined by homogeneous polynomials whose affine cone is singular only at the vertex. For such Calabi-Yau varieties, in case they are defined by Delsarte type equations, the physicists Berglund and Hübsch [10] defined a transposition rule for the defining polynomial. The construction has been later refined by Krawitz [33], who introduced the action of finite diagonal symplectic groups. More precisely, the Berglund-Hübsch-Krawitz (BHK for short) transposition rule is a correspondence

$$\{W = 0\} \subset \mathbb{P}(w)/\tilde{G} \longleftrightarrow \{W^* = 0\} \subset \mathbb{P}(w^*)/\tilde{G}^*,$$

---

2010 *Mathematics Subject Classification.* 14M25, 14J32, 14J33.

*Key words and phrases.*  $\mathbb{Q}$ -Fano toric varieties, Calabi-Yau hypersurfaces, canonical polytope, reflexive polytope, Berglund-Hübsch-Krawitz construction.

The first author has been partially supported by Proyecto Fondecyt Regular N. 1130572 and Proyecto Anillo ACT 1415 PIA Conicyt. The second author has been partially supported by Proyecto Fondecyt Postdoctorado N. 3150015 and Proyecto Anillo ACT 1415 PIA Conicyt. The third author was supported by the CNPq, Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brasil, and by the NCN project 2013/08/A/ST1/00804.

where  $W^*$ ,  $w^*$  and  $\tilde{G}^*$  are suitably defined transposed versions of the polynomial  $W$ , the set of weights  $w$  and the group  $\tilde{G}$  (see Section 4). Recently Chiodo and Ruan [17] proved that the BHK transposition rule actually satisfies (1) in terms of the Chen-Ruan orbifold cohomology.

The motivation of this work is first to relate the two constructions of Calabi-Yau varieties described above and secondly to define a duality generalizing both Batyrev and BHK duality. A way of unifying the BHK duality with Batyrev-Borisov duality of reflexive Gorenstein cones has been proposed in [12], where the author develops the vertex algebras approach to mirror symmetry for the BHK construction and suggest a common framework for the latter and Batyrev-Borisov duality (see [12, §7]). The present paper makes the previous framework more explicit in the hypersurface case in terms of polytopes and identifies the needed regularity condition to obtain a duality between families of Calabi-Yau varieties.

More precisely, given a  $\mathbb{Q}$ -Fano toric variety  $X$  with canonical singularities, we consider families of anticanonical hypersurfaces of  $X$  with fixed Newton polytope. Called  $\Theta$  the anticanonical polytope of  $X$  and  $\Delta$  the given Newton polytope, one such family will be denoted by  $\mathcal{F}_{\Delta, \Theta^*}$ . We prove the following result, where we recall that a canonical polytope is a lattice polytope whose unique lattice interior point is the origin.

**Theorem 1.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities and let  $\Delta \subset M_{\mathbb{Q}}$  be a canonical polytope contained in the anticanonical polytope  $\Theta$  of  $X$ . Then a general hypersurface in  $\mathcal{F}_{\Delta, \Theta^*}$  is a Calabi-Yau variety.*

This result gives examples of families of Calabi-Yau varieties in dimension  $\geq 5$  whose general element is not quasismooth and it is not birational to a hypersurface in a toric Fano variety (see Table 2 for some of them).

Theorem 1 also suggests the definition of a duality between families of Calabi-Yau varieties with fixed Newton polytope in  $\mathbb{Q}$ -Fano toric varieties. We will say that a pair  $(\Delta_1, \Delta_2)$  of polytopes is a *good pair* if  $\Delta_1 \subseteq \Delta_2$  and both  $\Delta_1$  and  $\Delta_2^*$  are canonical. Clearly the polar  $(\Delta_2^*, \Delta_1^*)$  of a good pair is still a good pair. This involution on good pairs produces the duality

$$\mathcal{F}_{\Delta_1, \Delta_2^*} \subseteq |-K_{X_{\Delta_2}}| \longleftrightarrow \mathcal{F}_{\Delta_2^*, \Delta_1} \subseteq |-K_{X_{\Delta_1^*}}|.$$

This coincides with Batyrev duality when  $\Delta_1 = \Delta_2$ , since canonical polytopes whose polar is canonical are exactly reflexive polytopes.

Moreover, Theorem 1 allows to define a *generalized BHK transposition rule* where the weighted projective space  $\mathbb{P}(w)$  is replaced by a  $\mathbb{Q}$ -Fano toric variety with torsion free class group and canonical singularities. More precisely, a generalized BHK family  $\mathcal{F}(A, \tilde{G})$  in  $X/\tilde{G}$  is a family of hypersurfaces of anticanonical degree defined by a matrix  $A$  of exponents for the equations and a symplectic group  $\tilde{G}$ . We can associate to it a pair of polytopes  $\Delta_1 \subseteq \Delta_2$ , where  $\Delta_1$  is the Newton polytope of a general element in  $\mathcal{F}(A, \tilde{G})$  and  $\Delta_2$  is the anticanonical polytope of  $X/\tilde{G}$ . The following shows that the generalized BHK transposition rule can be seen as a duality between good pairs.

**Theorem 2.** *Let  $(\Delta_1, \Delta_2)$  be the pair associated to a generalized Berglund-Hübsch-Krawitz family  $\mathcal{F}(A, \tilde{G})$ . Then  $(\Delta_1, \Delta_2)$  is a good pair and the pair associated to  $\mathcal{F}(A^T, G^*)$  is  $(\Delta_2^*, \Delta_1^*)$ .*

The theorem relies on a toric description of the generalized BHK construction, which had been given in the classical case in [42, §2] and generalizes [42, Proposition 2.9].

The paper is organized as follows. In Section 1 we recall some definitions and basic results about toric varieties and polytopes. In Section 2 we study hypersurfaces in toric varieties and we describe their regularity properties according to the Newton polytope. In Section 3 we prove Theorem 1 and we define the duality between good pairs. Section 4 is devoted to the definition of the generalized BHK mirror construction and to the proof of Theorem 2. Finally Section 5.1 contains some remarks about the stringy Hodge numbers of our families.

## 1. TORIC BACKGROUND

**1.1. Toric varieties.** We start recalling some standard facts in toric geometry, see for example [20]. Let  $N$  denote a lattice and let  $M = \text{Hom}(N, \mathbb{Z})$  be its dual. Let  $\Delta$  be a polytope in  $M_{\mathbb{Q}}$ , i.e. the convex hull of a finite subset of  $M_{\mathbb{Q}}$ . The *polar* of  $\Delta$  is the polyhedron

$$\Delta^* = \{y \in N_{\mathbb{Q}} : \langle x, y \rangle \geq -1, \forall x \in \Delta\},$$

which clearly contains the origin in its interior and whose facets are contained in the affine hyperplanes of equation  $\langle y, v_i \rangle = -1$ , where  $v_i$  is a vertex of  $\Delta$ .

It is well known that to any polytope  $\Delta$  as above one can associate a toric variety  $X = X_{\Delta}$  together with a  $\mathbb{Q}$ -divisor  $D$ . The variety  $X$  is the toric variety associated to the normal fan  $\Sigma_{\Delta}$  to  $\Delta$ . If  $n_1, \dots, n_r$  are the primitive generators of the one-dimensional cones of  $\Sigma_{\Delta}$  and  $D_1, \dots, D_r$  are the corresponding integral torus-invariant divisors, then  $D = -\sum_i h_{\Delta}(n_i)D_i$ , where  $h_{\Delta}$  is the strictly upper convex function

$$h_{\Delta} : N_{\mathbb{Q}} \rightarrow \mathbb{Q}, \quad h_{\Delta}(y) = \min_{x \in \Delta} \{\langle x, y \rangle\}.$$

Now let  $P : \mathbb{Z}^r \rightarrow N$  be the homomorphism defined by  $P(e_i) = n_i$ , which will be called *P-morphism* of the toric variety. We denote by  $P^T$  its transpose and by  $Q$  the homomorphism defined by the following exact sequence:

$$0 \longrightarrow M \xrightarrow{P^T} \mathbb{Z}^r \xrightarrow{Q} K \longrightarrow 0,$$

where  $K$  is isomorphic to the Class group of  $X$ .

The *Cox ring*  $\mathcal{R}(X)$  is the polynomial ring  $\mathbb{C}[T_1, \dots, T_r]$ , where  $T_i$  is the defining element of the divisor  $D_i$ , graded by  $K$ :  $\deg(T_i) = Q(e_i)$ . Let  $\bar{X} = \text{Spec } \mathcal{R}(X) \cong \mathbb{C}^r$ . By Cox's construction [20, Theorem 5.1.11] a toric variety  $X$  associated to a fan  $\Sigma$  can be described as a GIT-quotient of a quasitorus

$$X \cong (\bar{X} \setminus V(I)) // G_{\Sigma},$$

where  $V(I) = V(x^{\hat{\sigma}} \mid \sigma \in \Sigma)$  is the irrelevant locus, that is the subvariety of  $\bar{X}$  defined by the vanishing of every monomial of the form  $x^{\hat{\sigma}} = \prod_{\rho \notin \sigma(1)} x_{\rho}$  for  $\sigma \in \Sigma$ , and

$$G_{\Sigma} = \text{Hom}(\text{Cl}(X), \mathbb{C}^*) \simeq \text{Spec}(\mathbb{C}[\text{Cl}(X)]).$$

The big torus  $\hat{T} = (\mathbb{C}^*)^r$  naturally acts on the characteristic space  $\hat{X} = \bar{X} \setminus V(I)$  by coordinatewise multiplication. Observe that  $G_{\Sigma}$  is the kernel of the natural

homomorphism

$$\Psi : \hat{T} \rightarrow T, \quad (\lambda_1, \dots, \lambda_r) \mapsto (u \mapsto \prod_{j=1}^r \lambda_j^{\langle u, n_j \rangle}),$$

where we identify  $T$  with  $\text{Hom}(M, \mathbb{C}^*)$ .

We finally recall how quotients of toric varieties by finite subgroups of the torus can be described. Let  $X = X_{\Sigma, N}$  be a toric variety associated to a fan  $\Sigma \subset N_{\mathbb{Q}}$  and let  $N \rightarrow N'$  be a lattice monomorphism with finite cokernel  $G$ . The fan  $\Sigma$  in  $N'_{\mathbb{Q}} = N_{\mathbb{Q}}$  defines a toric variety  $X'$  and the inclusion of lattices induces a morphism of toric varieties  $X \rightarrow X'$  which is a good geometric quotient by the action of the group  $G$  [20, Proposition 3.3.7].

**Lemma 1.1.** *Let  $X = X_{\Sigma, N}$  be a toric variety associated to a fan  $\Sigma \subset N_{\mathbb{Q}}$  with torsion free class group, let  $\iota : N \rightarrow N'$  be a lattice monomorphism with finite cokernel  $G$  and  $\pi : X \rightarrow X'$  be the associated finite quotient. If the primitive generators  $n_1, \dots, n_r \in N$  of the rays of the fan of  $X$  are primitive in  $N'$ , then the homomorphism  $\pi^* : \mathcal{R}(X') \rightarrow \mathcal{R}(X)$  can be taken to be the identity and  $\text{Cl}(X') \cong \text{Cl}(X) \oplus G$ .*

*Proof.* Let  $P, P'$  be the  $P$ -morphisms of  $X, X'$  respectively. By the primitivity assumption on the  $n_i$ 's in  $N'$ , we have  $P' = \iota \circ P$ . Thus we have the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \xrightarrow{P'^T} & \mathbb{Z}^r & \longrightarrow & K' \longrightarrow 0 \\ & & \downarrow \iota^\vee & & \downarrow id & & \downarrow pr_1 \\ 0 & \longrightarrow & M & \xrightarrow{P^T} & \mathbb{Z}^r & \longrightarrow & K \longrightarrow 0 \end{array}$$

Observe that the central vertical arrow describes the map  $\pi^*$  in Cox coordinates. Since  $K$  is free, then  $K'$  is isomorphic to  $K \oplus \ker(pr_1)$ . Chasing in the diagram one finds that  $\ker(pr_1)$  is isomorphic to  $M/M' \cong G$ .  $\square$

**1.2. Polytopes.** We recall that a lattice polytope  $\Delta$  in  $M_{\mathbb{Q}}$  containing the origin in its interior is

- *reflexive* if  $\Delta^*$  is a lattice polytope;
- *canonical* if the origin is its unique interior lattice point;
- $\mathbb{Q}$ -*Fano* if its vertices are primitive in  $M$ .

A reflexive polytope can be defined equivalently as a lattice polytope with the origin in its interior such that the integral distance between any of its facets and the origin is equal to one ([5, Theorem 4.1.6]). Clearly with our definition  $\Delta$  is reflexive if and only if  $\Delta^*$  is reflexive. A canonical polytope is clearly  $\mathbb{Q}$ -Fano since, given a non-primitive vertex  $mv$ ,  $m \in \mathbb{Z}_{>0}$ , the vector  $v$  would be a non zero interior lattice point. Thus we have the following implications:

$$\text{reflexive} \Rightarrow \text{canonical} \Rightarrow \mathbb{Q}\text{-Fano}.$$

**Remark 1.2.** In [27] Kasprzyk provided a classification of three dimensional canonical polytopes, available in the Graded Ring Database <http://www.grdb.co.uk/>, and described an approach to the classification in higher dimension. Polytopes with the origin in their interior, sometimes called polytopes with the IP property in the literature, had a key role in the classification of reflexive polytopes of dimension  $\leq 4$  by Kreuzer and Skarke [35–37].

We recall that a projective normal variety  $X$  is  $\mathbb{Q}$ -Fano if  $-K_X$  is  $\mathbb{Q}$ -Cartier and ample (in particular it is  $\mathbb{Q}$ -Gorenstein) and Fano if moreover  $-K_X$  is Cartier (in particular it is Gorenstein). Moreover, the following holds (see [20, Theorem 6.2.1, Proposition 11.4.12, Theorem 8.3.4]).

**Theorem 1.3.** *Let  $\Delta \subset N_{\mathbb{Q}}$  be a lattice polytope containing the origin in its interior. Then  $X_{\Delta^*}$  is*

- $\mathbb{Q}$ -Fano if and only if  $\Delta$  is  $\mathbb{Q}$ -Fano;
- $\mathbb{Q}$ -Fano with canonical singularities if and only if  $\Delta$  is canonical;
- Fano if and only if  $\Delta$  (or  $\Delta^*$ ) is reflexive.

We finally introduce the concept of good pair, a key word in the paper.

**Definition 1.4.** Let  $\Delta_1, \Delta_2$  be two polytopes in  $M_{\mathbb{Q}}$ . We will say that  $(\Delta_1, \Delta_2)$  is a *good pair* if  $\Delta_1 \subseteq \Delta_2$ , and  $\Delta_1, \Delta_2^*$  are canonical (in particular  $\Delta_1$  and  $\Delta_2^*$  are both lattice polytopes).

The following shows that a lattice polytope containing the origin in its interior is canonical as soon as its dual is big enough.

**Lemma 1.5.** *Let  $\Delta \subset M_{\mathbb{Q}}$  be a lattice polytope containing the origin as an interior point. Then the origin is the only lattice interior point of  $\Delta^*$ .*

*Proof.* By definition of the polar polytope it is clear that it contains the origin in its interior. Suppose now that  $\Delta^*$  contains a lattice point  $n \neq 0$  in its interior. In particular for  $\varepsilon > 0$  small enough  $(1 + \varepsilon)n$  is contained in  $\Delta^*$ . By definition of the polar polytope, this implies that  $\langle u, (1 + \varepsilon)n \rangle \geq -1$  for all  $u \in \Delta$ . Thus every lattice point  $u$  in  $\Delta$  satisfies

$$\langle u, n \rangle \geq \frac{-1}{(1 + \varepsilon)} > -1$$

and hence  $\langle u, n \rangle \geq 0$  since  $\langle u, n \rangle$  is an integer. It follows that the lattice polytope  $\Delta$  is contained in the half space  $H_n = \{u \in M_{\mathbb{R}} \mid \langle u, n \rangle \geq 0\}$ , contradicting the fact that  $\Delta$  contains the origin as an interior point.  $\square$

As an immediate consequence we have:

**Corollary 1.6.** *Let  $\Delta_1$  and  $\Delta_2$  two polytopes in  $M_{\mathbb{Q}}$  with  $\Delta_1 \subseteq \Delta_2$ . Then  $(\Delta_1, \Delta_2)$  is a good pair if and only if  $\Delta_1$  and  $\Delta_2^*$  are lattice polytopes containing the origin as an interior point.*

This also implies the following result, explaining how good pairs behave when changing the underlying lattice up to finite index.

**Lemma 1.7.** *Let  $(\Delta_1, \Delta_2)$  be a good pair in  $M_{\mathbb{Q}}$  and let  $M' \subset M$  be an inclusion of lattices with finite index. If the vertices of  $\Delta_1$  belong to  $M'$ , then  $(\Delta_1, \Delta_2)$  is a good pair in  $M'_{\mathbb{Q}}$ .*

## 2. ANTICANONICAL HYPERSURFACES

Let  $X$  be a projective toric variety defined by a fan  $\Sigma \subset N_{\mathbb{Q}}$  and let  $n_1, \dots, n_r \in N$  be the primitive generators of the one dimensional cones of  $\Sigma$ . The *anticanonical polytope* of  $X$  is the polytope

$$\Theta = \{m \in M_{\mathbb{Q}} : \langle m, n_i \rangle \geq -1, \forall i\}.$$

The lattice points of  $\Theta$  naturally give a basis for the Riemann-Roch space of the divisor  $-K_X = \sum_i D_i$ . In fact, given  $u \in \Theta \cap M$ , the vector  $P^T(u) + \mathbf{1} \in \mathbb{Z}^r$ , where  $\mathbf{1}$  is the vector with all entries equal to 1, is the vector of exponents of a monomial  $m_u$  in the Cox ring  $\mathcal{R}(X)$  of degree  $[-K_X]$ . Conversely, any such monomial can be obtained in the same way.

Given a hypersurface  $D$  of  $X$  of degree  $[-K_X]$  defined by  $f = 0$  in Cox coordinates, we define the *support*  $\text{supp}(f)$  to be the set of  $u \in M$  such that  $m_u$  is a monomial of  $f$ . Moreover we define the *Newton polytope* of  $D$  as the convex hull of the points in its support. Given a lattice polytope  $\Delta$  contained in  $\Theta$  we will denote by  $\mathcal{F}_{\Delta, \Theta^*}$  the family of anticanonical hypersurfaces of  $X$  whose Newton polytope is equal to  $\Delta$ .

**2.1. Regularity of hypersurfaces.** In this section we will translate some basic regularity properties of hypersurfaces in  $\mathcal{F}_{\Delta, \Theta^*}$  in terms of geometric properties of  $\Delta$ . We recall that a hypersurface  $D$  of a projective toric variety  $X$  is called *well-formed* if

$$\text{codim}_D(D \cap \text{Sing}(X)) \geq 2,$$

where  $\text{Sing}(X)$  is the singular locus of  $X$ .

**Example 2.1.** In case  $X$  is a normalized weighted projective space, i.e.  $X = \mathbb{P}(w_1, \dots, w_n)$  with  $\gcd(w_1, \dots, w_n) = 1$ , it is known [26] that the general anticanonical hypersurface is well-formed if and only if

$$\gcd(w_1, \dots, \hat{w}_i, \dots, \hat{w}_j, \dots, w_n) \mid \sum_k w_k.$$

We will need the following result, where  $D_I := \cap_{i \in I} D_i$  for  $I \subseteq \{1, \dots, r\}$ .

**Lemma 2.2.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities and let  $\Delta$  be a lattice polytope contained in its anticanonical polytope. Then  $D_I$  is not empty if and only if  $\{n_i : i \in I\}$  is contained in a facet of  $\Theta^*$  and a hypersurface in  $\mathcal{F}_{\Delta, \Theta^*}$  contains  $D_I$  if and only if  $\{n_i : i \in I\}$  is not contained in a facet of  $\Delta^*$ .*

*Proof.* Let  $\Theta$  be the anticanonical polytope of  $X$ . By the assumption on  $X$ , a fan  $\Sigma$  for  $X$  is given by the cones over the facets of  $\Theta^*$  and the  $n_i$  are the vertices of  $\Theta^*$ . The stratum  $D_I$  is not empty if and only if the set  $\{n_i : i \in I\}$  is contained in a cone of  $\Sigma$ , or equivalently if the  $n_i$  are contained in a facet of  $\Theta^*$ . This gives the first statement.

Let  $u \in \Delta \cap M$ , and  $m_u$  be the corresponding monomial in homogeneous coordinates. The zero set of the monomial  $m_u$  is given by

$$\text{div}(m_u) = \sum_{i=1}^r (\langle u, n_i \rangle + 1) D_i.$$

A hypersurface  $f = 0$  in  $\mathcal{F}_{\Delta, \Theta^*}$  contains  $D_I$  if and only if any monomial  $m_u$  in  $f$  vanishes along some of the  $D_i$ 's with  $i \in I$ . This means that for all  $u \in \text{supp}(f)$  there exists some  $i \in I$  with  $\langle u, n_i \rangle > -1$ . In particular this holds for the vertices of  $\Delta$ , which means that the  $n_i$ 's do not all belong to a single facet of  $\Delta^*$ .  $\square$

**Remark 2.3.** In the following results we will usually ask  $D$  to be a *general element* in  $\mathcal{F}_{\Delta, \Theta^*}$ . This means that  $D = \{f = 0\}$ , where the Newton polytope of  $D$  is  $\Delta$  and the coefficients of  $f$  are general. Observe that the support of  $f$  is not necessarily equal to  $\Delta \cap M$ .

**Proposition 2.4.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities and let  $\Delta \subset M_{\mathbb{Q}}$  be a lattice polytope contained in the anticanonical polytope  $\Theta$  of  $X$ . A general hypersurface in  $\mathcal{F}_{\Delta, \Theta^*}$  is:*

- i) *irreducible if and only if  $n_i$  belongs to the boundary of  $\Delta^*$  for any  $i$ ;*
- ii) *well-formed if and only if, anytime  $n_i, n_j$  belong to a facet of  $\Theta^*$  and not to a facet of  $\Delta^*$ , the segment joining them doesn't contain any lattice point;*
- iii) *normal if, anytime  $n_i, n_j$  belong to a facet of  $\Theta^*$  and not to a facet of  $\Delta^*$ ,  $n_i + n_j$  is not in the interior of  $\Delta^*$ .*

*Proof.* Since  $X$  is  $\mathbb{Q}$ -Fano with canonical singularities, the  $n_i$ 's are the vertices of  $\Theta^*$  and the origin is the only interior lattice point of both  $\Theta$  and  $\Theta^*$ . In what follows  $D$  denotes a general element in  $\mathcal{F}_{\Delta, \Theta^*}$ .

By the second Bertini's theorem  $D$  is reducible if and only if it contains one of the integral invariant divisors  $D_i$  for the torus action as a component. Thus i) follows from Lemma 2.2.

By the same Lemma,  $D$  contains the stratum  $D_{ij}$  if and only if  $n_i, n_j$  are contained in a facet of  $\Theta^*$  and not in a facet of  $\Delta^*$ . Moreover,  $X$  is singular along  $D_{ij}$  if and only if the triangle  $0, n_i, n_j$  contains a lattice point  $n$  outside its vertices. Since the only interior lattice point of  $\Theta^*$  is the origin, this means that  $n$  belongs to the segment between  $n_i, n_j$ . This gives ii).

Let  $p : \hat{X} \rightarrow X$  be the characteristic space of  $X$ , let  $\hat{D} = p^{-1}(D)$  and let  $\hat{D}_i = p^{-1}(D_i)$ . By Serre's criterion [25, Proposition 8.23, Ch.II]  $\hat{D}$  is normal if and only if it is smooth in codimension one. By the first Bertini's theorem this happens if and only if  $\hat{D}_{ij} := \hat{D}_i \cap \hat{D}_j$  is not contained in the singular locus of  $\hat{D}$ , whenever it is not empty. By Lemma 2.2  $\hat{D}_{ij}$  is not empty and it is contained in  $\hat{D}$  when  $n_i, n_j$  belong to the same facet of  $\Theta^*$  but not to a facet of  $\Delta^*$ . Under these conditions,  $\hat{D}$  is singular along  $\hat{D}_{ij}$  if and only if, for any  $u \in \text{supp}(f)$ ,  $(u, n_j) > -1$  whenever  $(u, n_i) = 0$ , and similarly changing the role of  $i$  and  $j$  (this is equivalent to ask that the partial derivatives of the equation of  $\hat{D}$  in homogeneous coordinates vanish along  $\hat{D}_{ij}$ ). Since there exists no  $u \in \Delta \cap M$  such that  $(u, n_i) = (u, n_j) = -1$ , this is equivalent to ask that  $(u, n_i + n_j) > -1$  for all  $u \in \Delta \cap M$ , i.e. that  $n_i + n_j$  belongs to the interior of  $\Delta^*$ .

We recall that  $p$  is a GIT quotient for the action of a quasi-torus  $T$ . The divisor  $\hat{D}$  is  $T$ -invariant, being defined by a homogeneous polynomial in  $\mathcal{R}(X)$ . This implies that  $p|_{\hat{D}} : \hat{D} \rightarrow D$  is still a GIT quotient for the action of the group  $T/T_0$ , where  $T_0$  is the subgroup of  $T$  acting trivially on  $\hat{D}$ . Since  $\hat{D}$  is normal, it follows that  $D$  is normal (see for example [20, Lemma 5.0.4]). This proves iii).  $\square$

**2.2. Hypersurfaces with canonical singularities.** Let  $X$  be a  $\mathbb{Q}$ -Gorenstein normal variety over  $\mathbb{C}$  of dimension  $\geq 2$  and let  $D$  be a  $\mathbb{Q}$ -Cartier divisor on  $X$ . Given a resolution  $f : \tilde{X} \rightarrow X$ , that is a proper birational morphism such that  $\tilde{X}$  is smooth, one can write

$$(K_{\tilde{X}} + f_*^{-1}D) - f^*(K_X + D) \equiv \sum_{i=1}^r a(X, D, E_i)E_i,$$

where  $E_1, \dots, E_r$  are the distinct irreducible components of the exceptional divisor of  $f$  and  $f_*^{-1}D$  denotes the proper birational transform of  $D$  (see [32, Remark 6.6] for the precise meaning of this equation). The *discrepancy* of the pair  $(X, D)$ ,



denoted by  $\text{discrep}(X, D)$ , is the infimum of the values  $a(X, D, E)$ , as  $E$  varies over all exceptional divisors of the resolutions of  $X$ . The pair  $(X, D)$  is *canonical* if  $\text{discrep}(X, D) \geq 0$  and  $X$  has *canonical singularities* if  $(X, 0)$  is canonical.

In order to compute the discrepancy of a pair  $(X, D)$  it is enough to consider the minimum over the values  $a(X, D, E)$  as  $E$  varies among the exceptional divisors of a given *log resolution* of  $(X, D)$ , i.e. a resolution such that  $\text{Exc}(f) + f_*^{-1}(D)$  has pure codimension 1 and is a divisor with simple normal crossings [32, Definition 6.21]. Such a resolution always exists by a theorem of Hironaka [30, Theorem 10.45]. We recall the following result, which relates the discrepancy of a pair  $(X, D)$  to the discrepancy of  $D$ .

**Theorem 2.5** ([28]). *Let  $X$  be a normal variety over  $\mathbb{C}$  and let  $D$  be a normal divisor on  $X$  such that  $K_X + D$  is  $\mathbb{Q}$ -Cartier and  $\text{codim}_D(\text{Sing}(X) \cap D) \geq 2$ . Then*

$$(2) \quad \text{discrep}(D) \geq \text{discrep}(\text{center} \subset D, X, D),$$

where the right hand side is the infimum of the values  $a(X, D, E)$ , where  $f(E) \subset Z$ . In particular  $D$  has canonical singularities if the pair  $(X, D)$  is canonical.

*Proof.* The inequality follows from [31, Proposition 5.46] taking  $Z = S$  and  $B = 0$  or [28, §17.2]. The last statement immediately follows since  $\text{discrep}(\text{center} \subset D, X, D) \geq \text{discrep}(X, D)$ .  $\square$

**Proposition 2.6.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities and let  $\Delta \subset M_{\mathbb{Q}}$  be a lattice polytope contained in the anticanonical polytope  $\Theta$  of  $X$ . If  $\Delta$  is a canonical polytope then the general element  $D$  of  $\mathcal{F}_{\Delta, \Theta^*}$  is well-formed, normal and has canonical singularities.*

*Proof.* If  $\Delta$  is canonical, then  $\Delta^*$  is a polytope and its only interior lattice point is the origin by Lemma 1.5. By Proposition 2.4 we have that  $D$  is well-formed since, if  $n_i, n_j$  belong to a facet of  $\Theta^*$  and not to a facet of  $\Delta^*$ , then the segment joining them intersects the boundary of  $\Delta^*$  only at  $n_i, n_j$ . Moreover, by the same proposition,  $D$  is normal.

Since  $D$  is general, there exists a toric resolution of singularities  $f : \tilde{X} \rightarrow X$  which is a log resolution for  $D$ , obtained by means of a refinement  $\tilde{\Sigma}$  of the fan  $\Sigma$  of  $X$ . This can be obtained taking first a toric resolution of the singularities of  $X$  and then successive toric blow-ups along the base locus of  $\mathcal{F}_{\Delta, \Theta^*}$  until its proper transform is base point free. By the first Bertini's theorem the general element  $\tilde{D}$  of such proper transform is smooth. Moreover, the same theorem implies that  $\tilde{D}$  intersects transversally each component of the exceptional locus, since its restriction to any such component is base point free.

Let  $E$  be an exceptional divisor of  $f$  and let  $n \in N$  be the primitive generator of the corresponding ray of  $\tilde{\Sigma}$ . Observe that

$$\text{mult}_E(K_{\tilde{X}} - f^*(K_X)) = -1 - \text{mult}_E f^*(K_X)$$

since  $E$  is one of the integral torus invariant divisors of  $\tilde{X}$ . We can write  $D = \text{div}(\chi) - K_X$ , where  $\chi$  is a general linear combination of  $\chi^u$ , where  $u$  belongs to the support  $\text{supp}(\phi)$  of a defining equation  $\phi$  of  $D$ . Then

$$\begin{aligned} \text{mult}_E(f^*(D) - f_*^{-1}(D)) &= \text{mult}_E(f^*(D)) = \text{mult}_E(f^*\text{div}(\chi)) - \text{mult}_E f^*(K_X) \\ &= \min_{u \in \text{supp}(\phi)} \{\text{mult}_E(f^*\chi^u)\} - \text{mult}_E f^*(K_X) = \min_{u \in \text{supp}(\phi)} \{(u, n)\} - \text{mult}_E f^*(K_X), \end{aligned}$$



where the third equality is due to the generality assumption on  $D$  and the last equality to the fact that

$$\text{mult}_E f^* \text{div}(\chi^u) = \text{mult}_E \text{div}(f^* \chi^u) = (u, n).$$

This gives

$$(3) \quad a(X, D, E) = -1 - \min_{u \in \text{supp}(\phi)} \{(u, n)\} = -1 - \min_{u \in \Delta \cap M} \{(u, n)\},$$

where the second equality is due to the fact that  $\text{supp}(\phi)$  contains the vertices of  $\Delta$ . Such discrepancy is non-negative since  $\Delta^*$  has no non-zero interior lattice point. Theorem 2.5 thus implies that  $D$  has canonical singularities.  $\square$

**Corollary 2.7.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities,  $\Delta$  be a canonical polytope contained in its anticanonical polytope  $\Theta$  and  $D$  be a general element in  $\mathcal{F}_{\Delta, \Theta^*}$ . A birational toric morphism  $f : \tilde{X} \rightarrow X$  induces a crepant morphism  $f^*(D) \rightarrow D$  if and only if the rays of the fan of  $\tilde{X}$  are generated by nonzero lattice points in  $\Delta^*$ .*

*Proof.* It follows from (3) that  $a(X, D, E) = 0$  if and only if  $n \in \Delta^*$ .  $\square$

**Remark 2.8.** By the proof of Proposition 2.6 the discrepancies of a general anticanonical hypersurface  $D$  only depend on its Newton polytope  $\Delta$ .

**Remark 2.9.** As a consequence of the adjunction Conjecture [30, Theorem 4.9] formulated by Shokurov and Kollár, the inequality (3) is actually an equality. We now show that, under such conjecture, the condition on the polytope  $\Delta$  in Proposition 2.6 is also a necessary condition for  $D$  to be normal with canonical singularities. Assume that  $n \in N$  is a non-zero primitive vector in the interior of  $\Delta^*$ . Let  $\tilde{\Sigma}$  be a smooth fan refining the star subdivision of the fan  $\Sigma$  of  $X$  induced by  $n$ . This gives a resolution  $f$  of  $X$  and  $n$  corresponds to an exceptional divisor  $E$  of  $f$ . Let  $\sigma$  be the cone of  $\Sigma$  containing  $n$  in its interior. The primitive generators of the rays of  $\sigma$  are not contained in a facet of  $\Delta^*$ , since otherwise this would also be a facet of  $\Theta^*$  and  $n$  would be an interior point of  $\Theta^*$ , contradicting the fact that  $X$  has canonical singularities (see Theorem 1.3). Thus  $f(E) \subseteq D$  by Lemma 2.2. Since  $n$  is in the interior of  $\Delta^*$ , the computation in the proof of Proposition 2.6 gives that  $a(X, D, E) < 0$ , thus by [30, Theorem 4.9]  $D$  has a non-canonical singularity.

**Corollary 2.10.** *Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety with canonical singularities. If the convex hull of the lattice points of the anticanonical polytope of  $X$  is a canonical polytope, then a general anticanonical hypersurface of  $X$  is well-formed, normal and has canonical singularities.*

*Proof.* It follows from Proposition 2.6 taking  $\Delta$  to be the the convex hull of  $\Theta \cap M$ .  $\square$

**2.3. Quasismooth hypersurfaces.** A hypersurface  $D$  of a projective toric variety  $X$  is called *quasismooth* (or *transverse*) if  $p^{-1}(D)$  is smooth, where  $p : \hat{X} \rightarrow X$  is the quotient map in the Cox construction of  $X$ .

If  $X$  is a weighted projective space, a quasismooth hypersurface of  $X$  of dimension  $\geq 3$  is known to be well-formed, unless it is isomorphic to a toric stratum [21, Proposition 6]. This result can be generalized as follows.

**Proposition 2.11.** *Let  $X$  be a projective toric variety whose irrelevant locus has codimension  $> 4$  in  $\hat{X}$ . A quasismooth hypersurface  $D$  of  $X$  is either well-formed or it is isomorphic to a toric stratum of  $X$ .*

*Proof.* Let  $f$  be a defining element for  $D$  in the Cox ring  $R(X) = \mathbb{C}[x_1, \dots, x_r]$ . Assume that  $D$  is not well-formed, in particular it contains a codimension two toric stratum of  $X$ . Thus we can assume  $f$  to be of the form

$$f(x_1, \dots, x_r) = x_1 f_1 + x_2 f_2.$$

Computing the partial derivatives of  $f$  one can see that they all vanish along the subset  $S$  of  $\hat{X}$  defined by  $\{x_1 = x_2 = f_1 = f_2 = 0\}$ . If neither  $f_1$  or  $f_2$  is constant, we have that

$$\dim(S) \geq \dim(\hat{X}) - 4 > \dim(\bar{X} - \hat{X}),$$

contradicting the fact that  $D$  is quasismooth. Thus we can assume that  $f_1$  is constant, so that  $f(x_1, \dots, x_r) = \alpha x_1 + x_2 f_2$  is isomorphic to the stratum  $x_1 = 0$  by the isomorphism  $(x_2, \dots, x_r) \mapsto (-x_2 \alpha^{-1} f_2, x_2, \dots, x_r)$ .  $\square$

Quasismooth and well-formed anticanonical hypersurfaces give a class of hypersurfaces with canonical singularities. However, as we will observe later, such class is quite small in dimension bigger than three.

**Proposition 2.12.** *Let  $D$  be an anticanonical hypersurface of a projective toric variety  $X$ . If  $D$  is quasismooth and well-formed, then  $D$  has canonical singularities.*

*Proof.* Since  $D$  is quasismooth, then  $D$  is normal by the proof of Proposition 2.4. Moreover, by [30, Proposition 4.5, (1) and (5)] the adjunction formula holds for  $D$  and gives that  $K_D \sim \mathcal{O}_D$ . In particular  $D$  has Gorenstein singularities. Moreover, since  $\hat{D}$  is smooth, the singularities of  $D$  are rational [15, Corollaire]. By [29, Corollary 11.13] Gorenstein rational singularities are canonical.  $\square$

**Remark 2.13.** In [5] Batyrev considered a different notion of regularity: an anticanonical hypersurface  $Y$  of a toric Fano variety  $X$  is regular if the intersection of  $Y$  with any toric stratum of  $X$  is either empty or smooth of codimension one. Any regular hypersurface is quasismooth, see [9, Proposition 4.15] or [19, Proposition 5.3].

**2.4. Examples.** Given a  $\mathbb{Q}$ -Fano toric variety with canonical singularities and with anticanonical polytope  $\Theta$ , we can consider three different properties for the convex hull  $\bar{\Theta}$  of the lattice points of  $\Theta$ : canonical, reflexive and quasismooth. Here we say that  $\bar{\Theta}$  is *quasismooth* if such property holds for the general anticanonical hypersurface. In the case of weighted projective spaces it is known that the three concepts are equivalent in dimension 2 and 3. In dimension 4, canonical implies reflexive [43, Theorem, §3]. Moreover, for weighted projective spaces of any dimension, quasismooth implies canonical [43, Lemma 2]. In higher dimension the concepts of reflexive and quasismooth are unrelated and there are examples of canonical polytopes which are neither quasismooth nor reflexive. In Table 1 we show the number of weight systems  $w = (w_1, \dots, w_6)$  with  $w_i \leq 10$  such that the anticanonical polytope  $\Theta$  of  $\mathbb{P}(w)$  is reflexive ( $F$ ),  $\bar{\Theta}$  is reflexive ( $R$ ),  $\bar{\Theta}$  is canonical ( $C$ ) and not reflexive and we distinguish whether the general anticanonical hypersurface of  $\mathbb{P}(w)$  is quasismooth ( $Q$ ) or not. The properties of the anticanonical polytope can be checked by means of Magma [14] and the programs available here:

<http://goo.gl/A7W17Z>.

See also the Calabi-Yau data webpage by Kreuzer and Skarke

<http://hep.itp.tuwien.ac.at/~kreuzer/CY/>.

weights up to	$F$	$R$	$Q$ and not $R$	$C$ not $R$ and not $Q$
2	3	4	1	0
3	6	13	5	2
4	10	39	11	3
5	15	83	30	30
6	28	164	45	63
7	31	300	89	193
8	44	524	133	358
9	52	833	190	747
10	71	1278	269	1221

TABLE 1. Counting weight systems in dimension 5

**Remark 2.14.** Let  $X$  be a  $\mathbb{Q}$ -Fano toric variety and assume that there exists a toric Fano variety  $X'$  and a birational toric map  $X \rightarrow X'$  which induces a bijection between the anticanonical linear series  $|-K_X|$  and  $|-K_{X'}|$ . By standard facts in toric geometry, this map is induced by an isomorphism  $\varphi : M \rightarrow M'$  which gives a bijection between the lattice points of the anticanonical polytope  $\Theta$  of  $X$  and those of the anticanonical polytope  $\Theta'$  of  $X'$ . In particular  $\varphi_{\mathbb{Q}}$  induces an isomorphism between the convex hulls of the lattice points of  $\Theta$  and  $\Theta'$ . Since  $X'$  is Fano, the polytope  $\Theta'$  is reflexive by [5, Theorem 4.1.9], thus the convex hull of the lattice points of  $\Theta$  is reflexive. This shows that, if  $\bar{\Theta}$  is not reflexive, then the general anticanonical hypersurface of  $X$  is not (torically) birational to a hypersurface in a toric Fano variety.

**Example 2.15** ( $R$ ,  $C$  and  $Q$ ). In dimension 3 there are 104 weighted projective spaces with canonical singularities; for 95 of them the convex hull of lattice points of the anticanonical polytope is a canonical, reflexive and quasismooth polytope. Moreover, for 14 of these weight systems the anticanonical polytope is reflexive, i.e. the weighted projective space is Fano.

**Example 2.16** ( $R$  and not  $Q$ ).  $X = \mathbb{P}(1, 1, 1, 3, 4)$  is a toric  $\mathbb{Q}$ -Fano variety such that  $\bar{\Theta}$  is reflexive. Observe that an anticanonical hypersurface of  $X$  is defined by an equation of the form:

$$f(x_1, \dots, x_5) = x_1 f_1 + x_2 f_2 + x_3 f_3 + x_4 f_4,$$

since it has degree 10 and there is no power of such degree in the variable  $x_5$ . All partial derivatives vanish at the point  $(0 : 0 : 0 : 0 : 1)$  since  $f_1, f_2, f_3, f_4$  do not contain a power of  $x_5$ . Thus the general anticanonical hypersurface of  $X$  is not quasismooth.

**Example 2.17** ( $Q$  and not  $R$ ).  $X = \mathbb{P}(1, 1, 1, 1, 1, 2)$  is a toric  $\mathbb{Q}$ -Fano variety such that  $\bar{\Theta}$  is canonical and not reflexive. A general anticanonical hypersurface of  $X$  is defined by an equation of the form

$$f(x_1, \dots, x_6) = x_1 f_1 + x_2 f_2 + x_3 f_3 + x_4 f_4 + x_5 f_5.$$

By Bertini's theorem, the singular locus of  $f$  in  $\mathbb{C}^6$  is contained in the base locus of the corresponding linear system, which only contains the point  $(0 : 0 : 0 : 0 : 0 : 1)$ . The partial derivatives of  $f$  do not vanish at such point since we can assume that  $f_1$  (for example) contains the monomial  $x_6^3$ . Thus  $f = 0$  is quasismooth. More examples are given in Table 2.

**Example 2.18** ( $C$  not  $R$  and not  $Q$ ).  $X = \mathbb{P}(1, 1, 2, 3, 3, 3)$  is a toric  $\mathbb{Q}$ -Fano variety such that  $\Theta$  is canonical and not reflexive and such that the general anticanonical hypersurface is not quasismooth. More examples are given in Table 2.

$C$ , not $R$ , $Q$	$C$ , not $R$ , not $Q$
( 1, 1, 1, 1, 1, 2 )	( 1, 1, 2, 3, 3, 3 )
( 1, 2, 2, 2, 3, 4 )	( 1, 1, 2, 3, 3, 4 )
( 1, 1, 1, 1, 2, 3 )	( 1, 1, 1, 2, 3, 3 )
( 1, 1, 2, 2, 2, 3 )	
( 1, 2, 3, 3, 3, 3 )	
( 1, 1, 1, 3, 3, 4 )	
( 1, 2, 2, 3, 3, 4 )	
( 1, 1, 2, 2, 3, 4 )	
( 1, 1, 3, 3, 3, 4 )	
( 1, 2, 2, 3, 4, 4 )	
( 1, 1, 1, 2, 2, 3 )	

TABLE 2. Weights  $\leq 4$  having quasismooth and non-quasismooth general anticanonical hypersurface

### 3. A DUALITY BETWEEN FAMILIES OF CALABI-YAU HYPERSURFACES

We recall that an  $n$ -dimensional normal projective variety  $Y$  is a *Calabi-Yau variety* if it has canonical singularities,  $K_Y \cong \mathcal{O}_Y$  and  $h^i(Y, \mathcal{O}_Y) = 0$  for  $0 < i < n$ .

In [5, Theorem 4.1.9] Batyrev proved that a projective toric variety  $X_\Delta$  is Fano, or equivalently its anticanonical polytope  $\Delta$  is reflexive, if and only if regular anticanonical hypersurfaces  $Y$  of  $X$  are Calabi-Yau varieties. Moreover he defines a duality between families of anticanonical hypersurfaces of Fano toric varieties:

$$\mathcal{F}_{\Delta, \Delta^*} \subseteq |-K_{X_\Delta}| \longleftrightarrow \mathcal{F}_{\Delta^*, \Delta} \subseteq |-K_{X_{\Delta^*}}|.$$

In this section we will introduce a generalization of this duality in case  $X$  is  $\mathbb{Q}$ -Fano and the family of hypersurfaces is not necessarily the full anticanonical linear system. Such generalization is based on the result given in Theorem 1; using the characterization of Proposition 2.6 we can now prove it. Observe that by Remark 2.9, if the equality holds in (3), this would provide a characterization of  $\mathbb{Q}$ -Fano toric varieties whose general anticanonical hypersurfaces are Calabi-Yau.

*Proof of Theorem 1.* In what follows  $D$  will denote a general anticanonical hypersurface of  $\mathcal{F}_{\Delta, \Theta^*}$ . By Proposition 2.6  $D$  is well-formed, normal and has canonical singularities. By [30, Proposition 4.5, (1) and (5)] the adjunction formula holds for  $D$ , giving that  $K_D$  is trivial. Moreover we have the exact sequence

$$0 \longrightarrow \mathcal{O}_X(K_X) \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_D \longrightarrow 0,$$

which induces the exact sequence

$$\dots \longrightarrow H^i(X, \mathcal{O}_X) \longrightarrow H^i(D, \mathcal{O}_D) \longrightarrow H^{i+1}(X, \mathcal{O}_X(K_X)) \longrightarrow \dots$$

Since  $h^i(X, \mathcal{O}_X) = 0$  for  $i > 0$  and  $h^{i+1}(X, \mathcal{O}_X(K_X)) = h^{\dim(X)-i-1}(X, \mathcal{O}_X)$  for  $i < \dim(X)-1$  by Serre-Grothendieck duality, we obtain the vanishing of  $h^i(D, \mathcal{O}_D)$  for  $0 < i < \dim(D)$ . Thus  $D$  is a Calabi-Yau variety.  $\square$

**Remark 3.1.** In [39, Theorem 2.25] the author states that a general anticanonical hypersurface of a projective toric variety is a Calabi-Yau variety if and only if the convex hull of the lattice points of its anticanonical polytope is reflexive. This is not true in general, see Example 2.17.

Observe that a good pair of polytopes  $\Delta_1 \subset \Delta_2$  naturally produces a family of Calabi-Yau varieties in a  $\mathbb{Q}$ -Fano projective toric variety. In fact, the toric variety  $X := X_{\Delta_2}$  defined by the normal fan to  $\Delta_2$  is  $\mathbb{Q}$ -Fano with canonical singularities by Theorem 1.3 and  $\Delta_2$  is its anticanonical polytope. The subpolytope  $\Delta_1 \subseteq \Delta_2$  identifies the family  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  of anticanonical hypersurfaces of  $X$ , whose general element is a Calabi-Yau variety by Proposition 2.6. By our definition of good pair we immediately have that if  $(\Delta_1, \Delta_2)$  is a good pair in  $M_{\mathbb{Q}}$ , then its polar  $(\Delta_2^*, \Delta_1^*)$  is a good pair in  $N_{\mathbb{Q}}$ . This provides a duality between families of Calabi-Yau hypersurfaces of  $\mathbb{Q}$ -Fano toric varieties:

$$\mathcal{F}_{\Delta_1, \Delta_2^*} \subseteq |-K_{X_{\Delta_2}}| \longleftrightarrow \mathcal{F}_{\Delta_2^*, \Delta_1^*} \subseteq |-K_{X_{\Delta_1^*}}|.$$

**Proposition 3.2.** *If  $\Delta_1 = \Delta_2$ , then the duality between good pairs is Batyrev duality.*

*Proof.* If  $\Delta_1 = \Delta_2$ , then  $\Delta_2$  and  $\Delta_2^*$  are lattice polytopes, thus  $\Delta_2$  is reflexive. By Theorem 1.3 this means that  $X_{\Delta_2}$  is a Fano variety. Moreover  $\mathcal{F}_{\Delta_2, \Delta_2^*}$  is the family of all anticanonical hypersurfaces of  $X_{\Delta_2}$ .  $\square$

**Example 3.3.** Let  $X = \mathbb{P}^2$  and let  $\Delta_2$  be its anticanonical polytope. Let  $\Delta_1$  be the lattice polytope whose vertices are  $(1, 0), (0, 1), (-1, 0), (0, -1)$ . The pair  $(\Delta_1, \Delta_2)$  is a good pair and the toric variety  $Y := X_{\Delta_1^*}$  is  $\mathbb{P}^1 \times \mathbb{P}^1$ . The family  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  in  $X$  is given by  $\alpha_1 x_1^2 x_2 + \alpha_2 x_2^2 x_1 + \alpha_3 x_1 x_3^2 + \alpha_4 x_2 x_3^2 + \alpha_5 x_1 x_2 x_3$  where  $\alpha_i \in \mathbb{C}^*$ , whereas the dual family  $\mathcal{F}_{\Delta_2^*, \Delta_1^*}$  is defined by the equation  $\beta_1 y_1^2 y_3 y_4 + \beta_2 y_1 y_2 y_3^2 + \beta_3 y_2^2 y_4^2 + \beta_4 y_1 y_2 y_3 y_4$ ,  $\beta_i \in \mathbb{C}^*$  in homogeneous coordinates of  $Y$  (in fact up to rescaling the variables one can put all coefficients  $\beta_i$  to be equal to one).




FIGURE 1. The good pairs  $(\Delta_1, \Delta_2)$  and  $(\Delta_2^*, \Delta_1^*)$

**Remark 3.4.** Families of Calabi-Yau varieties associated to pairs of nested reflexive polytopes already appeared in the literature in relation with the phenomenon of extremal transition [23, 38, 41]. An extremal transition between two families  $\mathcal{F}$  and  $\mathcal{G}$  of Calabi-Yau manifolds occurs when there is a degeneration of  $\mathcal{F}$  to a singular

family  $\mathcal{F}_0$  whose resolution is isomorphic to  $\mathcal{G}$ . An inclusion of reflexive polytopes  $\Delta_1 \subset \Delta_2$  can produce such a transition since the family  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  is a degeneration of  $\mathcal{F}_{\Delta_2, \Delta_2^*}$  (obtained putting some coefficients to zero) and is birational to the family  $\mathcal{F}_{\Delta_1, \Delta_1^*}$ . Such extremal transitions have been used to prove connectedness of moduli spaces of Calabi-Yau manifolds [3, 4, 11, 16].

**Proposition 3.5.** *Let  $(\Delta_1, \Delta_2)$  and  $(\Delta'_1, \Delta_2)$  be two good pairs. Then the dual families of  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta'_1, \Delta_2^*}$  are birational.*

*Proof.* This follows from the fact that the toric varieties  $X_{(\Delta'_1)^*}$  and  $X_{(\Delta_1)^*}$  are compactifications of the same torus  $T_N = \text{Spec } \mathbb{C}[N]$  and that the dual families in  $T_N$  are both defined by linear combinations of the monomials corresponding to the points of the polytope  $\Delta_2^*$ .  $\square$

In the next section we will show that the duality between good pairs also includes Berglund-Hübsch-Krawitz duality. This implies that Proposition 3.5 can be seen as a generalization of [42, Theorem 3.1]. See also [22] for a recent generalization of the proposition in a more general setting and in terms of derived equivalences.

#### 4. BERGLUND-HÜBSCH-KRAWITZ (BHK) DUALITY

**4.1. The BHK construction.** We will recall a mirror construction due to the physicists Berglund and Hübsch [10] and later refined by Krawitz in [33]. Let  $\mathbb{P}(w) = \mathbb{P}(w_1, \dots, w_n)$  be a normalized weighted projective space and let  $W$  be a homogeneous polynomial of Delsarte type, i.e. having the same number of monomials and variables. Up to rescaling the variables, we can assume that

$$(4) \quad W(x_1, \dots, x_n) = \sum_{i=1}^n \prod_{j=1}^n x_j^{a_{ij}},$$

so that  $W$  is uniquely determined by its matrix of exponents  $A = (a_{ij})$ . We will denote by  $Y_W$  the hypersurface defined by  $W$  in  $\mathbb{P}(w)$  and we will assume that

- (i)  $A$  is invertible over  $\mathbb{Q}$ ,
- (ii)  $Y_W$  is quasismooth,
- (iii)  $\deg(W) = \sum_{i=1}^n w_i$  (*Calabi-Yau condition*).

The assumptions (ii) and (iii) imply that  $Y_W$  is a Calabi-Yau variety by Proposition 2.12 and [18, Lemma 1.11].

**Remark 4.1.** By the proof of [43, Lemma 2] the condition of quasismoothness implies that the matrix  $A$  is invertible over  $\mathbb{Q}$ . Thus condition (i) in the above construction is redundant. Moreover, it can be easily proved that asking  $A$  to be invertible over  $\mathbb{Q}$  is equivalent to say that the convex hull of the elements  $u_1, \dots, u_n \in M$  corresponding to the monomials of  $W$  is a simplex in  $M_{\mathbb{R}}$ .

If we now consider the transposed matrix of  $A$ , this defines in the same way a Delsarte type polynomial  $W^*$ . A set of weights  $w^* = (w_1^*, \dots, w_n^*)$  which makes  $W^*$  homogeneous is given by the smallest integer multiple of the vector

$$q^* = (A^T)^{-1} \cdot \mathbf{1},$$

where  $\mathbf{1}$  denotes the column vector with all entries equal to 1. By the quasismoothness assumption it follows that  $w^*$  can be chosen with all positive entries (see Remark 4.2). Thus  $W^*$  defines a hypersurface  $Y_{W^*}$  in  $\mathbb{P}(w^*)$ . By [34, Theorem 1]  $W^*$  is still quasismooth and an easy computation shows that it satisfies the

Calabi-Yau condition in  $\mathbb{P}(w^*)$ . Thus  $Y_{W^*}$  is a Calabi-Yau variety. The Berglund-Hübsch-Krawitz construction gives a duality

$$Y_W/\tilde{G} \longleftrightarrow Y_{W^*}/\tilde{G}^*,$$

where  $\tilde{G}$  denotes a quotient group  $G/J$ , with  $J \subseteq \mathrm{SL}_n(\mathbb{C})$  the subgroup of diagonal automorphisms inducing the identity on  $\mathbb{P}(w)$  and  $G$  a subgroup of diagonal automorphisms in  $\mathrm{SL}_n(\mathbb{C})$  containing  $J$  and acting trivially on  $W$ , i.e.

$$W(g(x)) = W(x), \quad \forall g \in G.$$

The transposed group  $\tilde{G}^*$  is defined as  $G^*/J^*$ , where  $J^*$  is the analogous of  $J$  for  $\mathbb{P}(w^*)$  and  $G^*$  is defined by

$$(5) \quad G^* := \left\{ \prod_{j=1}^n (\rho_j^*)^{\alpha_j} \mid \prod_{j=1}^n x_j^{\alpha_j} \text{ is } G\text{-invariant} \right\},$$

where  $\rho_j^* := \mathrm{diag}(\exp(2\pi i a^{j1}), \dots, \exp(2\pi i a^{jn}))$  and  $a^{ji}$  are the entries of  $A^{-1}$ . Several equivalent definitions for the transposed group can be found in [1, §3]. The groups  $\tilde{G}$  and  $\tilde{G}^*$  both act simplyctically [1, Proposition 2.3], thus  $Y_W/\tilde{G}$  and  $Y_{W^*}/\tilde{G}^*$  are both Calabi-Yau varieties. In fact we will prove in Proposition 4.7 that they are anticanonical hypersurfaces of the  $\mathbb{Q}$ -Fano toric varieties  $\mathbb{P}(w)/\tilde{G}$  and  $\mathbb{P}(w^*)/\tilde{G}^*$ .

In [17, Theorem 2] Chiodo and Ruan proved that  $Y_W/\tilde{G}$  and  $Y_{W^*}/\tilde{G}^*$  have symmetric Hodge diamonds for the Chen-Ruan orbifold cohomology.

**Remark 4.2.** Observe that, by the above definition, we have that

$$A^T q^* = \mathbf{1} \iff \sum_{i=1}^n q_i^* P^T(u_i) = 0 \iff \sum_{i=1}^n q_i^* u_i = 0,$$

where  $u_1, \dots, u_n \in M$  are the points corresponding to the monomials of  $W$ , i.e.  $P^T(u_i) + \mathbf{1}$  is the  $i$ -th row of  $A$ . Moreover

$$\sum_{i=1}^n q_i^* = \mathbf{1}^T (A^T)^{-1} \mathbf{1} = \mathbf{1} A^{-1} \mathbf{1} = 1.$$

Thus the entries of the vector  $q^*$  are the barycentric coordinates of the origin in the simplex with vertices  $u_1, \dots, u_n$ . In particular all the entries of  $q^*$  are positive if and only if the origin lies in the interior of the simplex. Since  $X_W$  is quasismooth, by [43, Lemma 2] the simplex contains the origin in its interior.

**4.2. The generalized BHK construction.** We now describe a natural generalization of the BHK construction where weighted projective spaces are replaced by their higher class group rank analogs:  $\mathbb{Q}$ -Fano toric varieties with torsion-free class group.

**Construction 4.3.** Let  $X = X_\Sigma$  be a  $\mathbb{Q}$ -Fano toric variety with torsion free class group. Denote by  $n_1, \dots, n_r$  the minimal generators of the rays of  $\Sigma$  and by  $\Delta_2$  the anticanonical polytope of  $X$ . Let  $\Delta_1 \subseteq \Delta_2$  be a lattice polytope containing the origin in its interior and let us denote by  $u_1, \dots, u_s$  its vertices.

Note that by Corollary 1.6  $(\Delta_1, \Delta_2)$  is a good pair and hence  $u_1, \dots, u_s$  are primitive lattice vectors. Moreover  $n_1, \dots, n_r$  generate the whole lattice  $N$  since  $\mathrm{Cl}(X)$  is torsion-free. Instead of considering a single polynomial  $W$  (made unique



by a rescaling of the variables) as in the original construction, we consider a family of polynomials of the form

$$W(x_1, \dots, x_r) = \sum_{i=1}^s \alpha_i \prod_{j=1}^r x_j^{a_{ij}}, \quad \alpha_i \in \mathbb{C}^*,$$

uniquely determined by the matrix

$$A = (a_{ij})_{\substack{i=1, \dots, s \\ j=1, \dots, r}} \quad \text{with } a_{ij} = \langle u_i, n_j \rangle + 1.$$

By construction  $W$  is homogeneous of degree  $[-K_X]$  and the equation  $W = 0$  defines a hypersurface  $Y_W$  of  $X$ . Observe that, by the proof of [43, Lemma 2], the quasismoothness condition for  $W$  implies that its Newton polytope contains the origin in its interior.

An important fact to notice is that by the hypotheses made on  $u_1, \dots, u_s$ , the matrix  $A$  determines the variety  $X$ .

**Lemma 4.4.** *Since  $\Delta_1$  is a lattice polytope containing the origin in its interior, it is possible to recover the toric variety  $X$  from the matrix  $A$ .*

*Proof.* Since the polytopes  $\Delta_1$  and  $\Delta_2^*$  are full dimensional, we know that the matrix  $A - \mathbf{1} = (\langle u_i, n_j \rangle)_{i=1, \dots, s, j=1, \dots, r}$  has rank  $n$ , so that we may choose  $n$  linearly independent rows  $l_{i_1}, \dots, l_{i_n}$  of it. These define a submatrix  $P'$ , equal to the product  $UP$  where the rows of  $U$  are the coefficients of  $u_{i_1}, \dots, u_{i_n}$  and  $P$  is the matrix of the  $P$ -morphism of  $X$  (see §1).

The lattice  $N' = UN$  is a sublattice of  $N$  of index  $|N/N'| = |\det(U)|$  and by [20, Prop. 3.3.7], the group  $K'$  in the short exact sequence

$$(6) \quad 0 \longrightarrow (N')^\vee = M' \xrightarrow{(P')^T} \mathbb{Z}^r \xrightarrow{Q'} K' \longrightarrow 0,$$

is isomorphic to the class group of the geometric quotient  $X/H$  where  $H = N/N'$ . Since  $P' = UP$  we have  $K' \simeq K \times H$  and we can factorize the sequence as

$$(7) \quad 0 \longrightarrow M \xrightarrow{P^T} M' \xrightarrow{U^T} \mathbb{Z}^r \xrightarrow{Q'} K' \xrightarrow{F} K \longrightarrow 0,$$

where  $M$  is a sublattice of  $M'$  of index  $|\det(U)|$  and the map  $F : K' \simeq K \times H \rightarrow K$  is the first projection.

It follows that knowing just  $A$ , we can form a short exact sequence (6) by finding a suitable submatrix of  $A - \mathbf{1}$  of rank  $n$ . Then by computing the torsion part of  $K'$  we can deduce the map  $F$  of (7) and hence the sublattice  $M = \ker(F \circ Q')$  of  $M'$ . Up to a unimodular basis change, this gives  $P^T$  and hence the minimal generators  $n_1, \dots, n_r$  of  $\Sigma$ . This is enough to recover  $X$  since it is  $\mathbb{Q}$ -Fano. Indeed by [40, Prop. 4.3] the cones of the fan  $\Sigma$  are the cones over the faces of  $\Delta_2^* = \text{Conv}(n_1, \dots, n_r)$ .  $\square$

Let us now turn to the groups of diagonal automorphisms of  $X$  that stabilize  $W$ . The stabilizer of the monomials of  $W$  under the induced action of the big torus  $\hat{T} \cong (\mathbb{C}^*)^r$  on the ring  $\mathbb{C}[x_1, \dots, x_r]$  is

$$\text{Aut}(W) = \left\{ (\lambda_1, \dots, \lambda_r) \in \hat{T} \mid \prod_{j=1}^r \lambda_j^{a_{ij}} = 1, \text{ for } i = 1, \dots, s \right\}.$$

Let  $\Psi$  and  $G_\Sigma = \ker(\Psi)$  as defined in section 1. The group  $\text{Aut}(W)$  does not contain  $G_\Sigma$  in general, but if we consider the special linear parts  $J_\Sigma = G_\Sigma \cap \text{SL}_r(\mathbb{C})$  and

$\mathrm{SL}(W) = \mathrm{Aut}(W) \cap \mathrm{SL}_r(\mathbb{C})$ , we have  $J_\Sigma \subset \mathrm{SL}(W)$ . Indeed, for all  $(\lambda_1, \dots, \lambda_r) \in \hat{T} \cap \mathrm{SL}_r(\mathbb{C})$  we have  $\prod_{j=1}^r \lambda_j = 1$  and since  $a_{ij} = \langle u_i, n_j \rangle + 1$  we have

$$\mathrm{SL}(W) = \{\lambda \in \hat{T} \cap \mathrm{SL}_r(\mathbb{C}) : \Psi(\lambda)(u) = 1, \forall u \in M_W\},$$

where  $M_W$  is the sublattice of  $M$  generated by  $u_1, \dots, u_s$ . Moreover

$$J_\Sigma = \{\lambda \in \hat{T} \cap \mathrm{SL}_r(\mathbb{C}) : \Psi(\lambda)(u) = 1, \forall u \in M\}.$$

Note that contrary to the original BHK construction, the group  $\mathrm{SL}(W)$  here can be infinite. But when this is the case, then  $J_\Sigma$  is infinite as well and the quotient is finite.

**Lemma 4.5.** *The homomorphism  $\Psi$  defines a perfect pairing*

$$\bar{\Psi} : \mathrm{SL}(W)/J_\Sigma \times M/M_W \rightarrow \mathbb{C}^*, ([\lambda], [u]) \mapsto \Psi(\lambda)(u).$$

*In particular  $\mathrm{SL}(W)/J_\Sigma$  is isomorphic to  $M/M_W$ .*

*Proof.* The homomorphism  $\bar{\Psi}$  is clearly well-defined and it induces an injective homomorphism from  $\mathrm{SL}(W)/J_\Sigma$  to  $\mathrm{Hom}(M/M_W, \mathbb{C}^*)$  by definition of  $J_\Sigma$ . Let  $(u_1, \dots, u_n)$  be a basis of  $M$  such that there exist positive integers  $k_1, \dots, k_n \in \mathbb{N}$  with  $(k_1 u_1, \dots, k_n u_n)$  a basis of  $M_W$ . We have  $M/M_W \simeq \bigoplus_{i=1}^n \mathbb{Z}/k_i \mathbb{Z}$ . To prove the surjectivity we will prove that there exists an element of  $\mathrm{SL}(W)$  whose image is the homomorphism sending  $u_1$  to  $e^{2\pi i/k_1}$  and  $u_i$  to 1 for  $i > 1$ . Consider the dual basis of  $(k_1 u_1, \dots, k_n u_n)$ . This is a  $\mathbb{Z}$ -basis of  $M_W^\vee$ , let us denote it by  $(m_1, \dots, m_n)$ .

Now pick a  $r$ -uple of rational numbers  $w_1, \dots, w_r \in \mathbb{Q}$  such that  $\sum_{j=1}^r w_j n_j = m_1$  and write  $e^{i2\pi w} = (e^{i2\pi w_1}, \dots, e^{i2\pi w_r})$ . Since  $m_1$  is an element of  $M_W^\vee$ , we have

$$\sum_{j=1}^r w_j \langle n_j, u \rangle = \langle m_1, u \rangle \in \mathbb{Z} \quad \text{for all } u \in M_W,$$

which implies that  $e^{2\pi i w} \in \mathrm{SL}(W)$ . It remains to observe that

$$e^{2\pi i \sum_{j=1}^r w_j \langle u_1, n_j \rangle} = e^{2\pi i/k_1}, \quad e^{2\pi i \sum_{j=1}^r w_j \langle u_i, n_j \rangle} = 1 \text{ for } i > 1.$$

A similar argument holds for any  $u_i$ , proving the surjectivity.  $\square$

We now choose a subgroup  $G$  of  $\mathrm{SL}(W)$  containing  $J_\Sigma$ , or equivalently a subgroup  $\tilde{G} = G/J_\Sigma$  of  $\mathrm{SL}(W)/J_\Sigma$ . Observe that by Lemma 4.5 there exists a unique lattice  $M_G$  satisfying  $M_W \subseteq M_G \subseteq M$  such that

$$(8) \quad G = \left\{ \lambda \in \hat{T} \cap \mathrm{SL}_r(\mathbb{C}) \mid \Psi(\lambda)(u) = 1, \forall u \in M_G \right\}.$$

We thus define an orbifold  $Y_{W,G} = Y_W/\tilde{G}$ , the family of which we denote by  $\mathcal{F}(A, \tilde{G})$ .

The following generalizes [42, Lemma 2.3].

**Proposition 4.6.** *Let  $X, G$  be as in Construction 4.3. Then the quotient map  $X \rightarrow X/G$  induces the identity  $\mathcal{R}(X/G) \rightarrow \mathcal{R}(X)$  and  $\mathrm{Cl}(X/G) \cong \mathrm{Cl}(X) \oplus G$ .*

*Proof.* By Lemma 1.1 it is enough to prove that the quotient map is induced by a lattice monomorphism  $N \rightarrow N_G$  such that the primitive generators  $n_1, \dots, n_r \in N$  of the rays of the fan of  $X$  remain primitive in  $N_G$ . In fact, assume that  $n_i = kn$  for some  $n \in N_G$  and some integer  $k$ . Since  $N_G \subset M_W^\vee$ , then  $a_{ij} - 1 = \langle n_i, u_j \rangle = k \langle n, u_j \rangle \in k\mathbb{Z}$  for all  $j$ . Since a monomial in  $W$  can not contain all the variables (otherwise it would correspond to  $0 \in M$ ), this gives  $k = 1$ .  $\square$

**Proposition 4.7.** *The orbifolds  $Y_{W,G}$  are Calabi-Yau varieties for general  $\alpha_i \in \mathbb{C}^*$ .*

*Proof.* By construction  $Y_W$  is an anticanonical hypersurface in  $X$ . The quotient  $Y_W \rightarrow Y_W/G$  is clearly induced by the quotient  $X \rightarrow X/G$  induced by the inclusion  $N \subseteq N_G = M_G^\vee$ . Let  $\Delta_2 \subset M_\mathbb{Q}$  be the anticanonical polytope of  $X$  and  $\Delta_1$  be the Newton polytope of  $Y_W$ . The anticanonical polytope of  $X/G$  is equal to  $\Delta_2 \subset (M_G)_\mathbb{Q} = M_\mathbb{Q}$  by Proposition 4.6. The monomials in the defining equation of  $Y_W/G$  are the lattice points  $u_i \in M_W \subset M_G$ . Thus, the Newton polytope of  $Y_W/G$  is clearly  $\Delta_1$ . Since  $(\Delta_1, \Delta_2)$  is a good pair in  $M_\mathbb{Q}$  by Corollary 1.6, then the same holds in  $(M_G)_\mathbb{Q}$  by Lemma 1.7. By Theorem 1 and Remark 2.3 the orbifold  $Y_{W,G}$ , whose monomials in Cox coordinates correspond to the vertices of  $\Delta_1$ , are Calabi-Yau varieties.  $\square$

Let us now define the dual family  $\mathcal{F}(A^T, \tilde{G}^*)$ . Let  $X^*$  be the toric variety whose fan is the collection of cones over the faces of  $\Delta_1$ , considered with respect to the lattice  $M_W$  generated by  $u_1, \dots, u_s$ . The transposed matrix  $A^T$  of  $A$  gives the family

$$W^*(y_1, \dots, y_s) = \sum_{i=1}^r \beta_i \prod_{j=1}^s y_j^{a_{ji}}, \quad \beta_i \in \mathbb{C}^*,$$

where  $y_1, \dots, y_s$  are the Cox coordinates of  $X^*$ . Observe that by Lemma 4.4 the toric variety  $X^*$  can be recovered uniquely by the matrix  $A^T$ .

We now define the transposed of the group  $G$ . If  $M^\vee \subseteq M_G^\vee \subseteq M_W^\vee$  denote the dual lattices and  $\hat{T}^\vee = (\mathbb{C}^*)^s$  is an algebraic torus of dimension  $s$ , we define

$$(9) \quad G^* := \left\{ \nu \in \hat{T}^\vee \cap \mathrm{SL}_s(\mathbb{C}) \mid \Psi^*(\nu)(n) = 1, \text{ for all } n \in M_G^\vee \right\},$$

where

$$\Psi^* : \hat{T}^\vee \rightarrow \mathrm{Hom}(N, \mathbb{C}), \quad \nu \mapsto (n \mapsto \prod_{i=1}^s \nu_i^{\langle u_i, n \rangle}).$$

Let us show that this generalizes the original definition of the transposed group given by Krawitz (5) (see also [12, Proposition 2.3.1]).

**Lemma 4.8.** *When the good pair  $(\Delta_1, \Delta_2)$  is formed by simplices then the definitions (5) and (9) coincide.*

*Proof.* The proof relies on the fact that when  $r = s = n + 1$  the diagonal groups preserving the polynomials  $W$  and  $W^*$  are finite, hence formed of  $(n + 1)$ -uples of roots of unity. Let us write  $e^{2\pi i v} = (e^{2\pi i v_0}, \dots, e^{2\pi i v_n})$  for all  $v \in \mathbb{Q}^{n+1}$ . Given a subgroup  $G$  with  $J_\Sigma \subset G \subset \mathrm{SL}(W)$  we will denote by  $G_1^*$  its transposed as defined in (5) and  $G_2^*$  its transposed as in (9). It is proven in [1, §3.5] that

$$G_1^* := \{ e^{2\pi i v} \in \hat{T}^\vee \cap \mathrm{SL}_{n+1}(\mathbb{C}) \mid v^T A w \in \mathbb{Z} \text{ for all } e^{2\pi i w} \in G \}.$$

We first remark that for all  $(v, w) \in \mathbb{Q}^{n+1} \times \mathbb{Q}^{n+1}$  we have

$$e^{2\pi i v}, e^{2\pi i w} \in \mathrm{SL}_{n+1}(\mathbb{C}) \Rightarrow \sum_{i,j=0}^n v_i w_j \in \mathbb{Z},$$

and that by (8), for all  $w \in \mathbb{Q}^{n+1}$  such that  $e^{2\pi i w} \in \mathrm{SL}_{n+1}(\mathbb{C})$  we have

$$e^{2\pi i w} \in G \Leftrightarrow \forall u \in M_G, \sum_{j=0}^n \langle u, n_j \rangle w_j \in \mathbb{Z} \Leftrightarrow \sum_{j=0}^n w_j n_j \in M_G^\vee.$$

It follows that for all  $v \in \mathbb{Q}^{n+1}$  such that  $e^{2\pi i v} \in \mathrm{SL}_{n+1}(\mathbb{C})$  we have

$$\begin{aligned} e^{2\pi i v} \in G_1^* &\Leftrightarrow \sum_{i,j=0}^n v_i (\langle u_i, n_j \rangle + 1) w_j \in \mathbb{Z} \text{ for all } e^{2\pi i w} \in G \\ &\Leftrightarrow \langle \sum_{i=0}^n v_i u_i, \sum_{j=0}^n w_j n_j \rangle \in \mathbb{Z} \text{ for all } e^{2\pi i w} \in G \\ &\Leftrightarrow \sum_{i=0}^n v_i u_i \in M_G. \end{aligned}$$

On the other hand, writing  $\nu = e^{2\pi i v}$ , definition (9) is equivalent to

$$G_2^* := \{e^{2\pi i v} \in \hat{T}^\vee \cap \mathrm{SL}_{n+1}(\mathbb{C}) \mid \sum_{i=0}^n v_i \langle u_i, n \rangle \in \mathbb{Z} \text{ for all } n \in M_G^\vee\}.$$

Since for all  $v \in \mathbb{Q}^{n+1}$  we have

$$\sum_{i=0}^n v_i \langle u_i, n \rangle \in \mathbb{Z} \text{ for all } n \in M_G^\vee \Leftrightarrow \sum_{i=0}^n v_i u_i \in M_G,$$

we get  $G_1^* = G_2^*$  so that the two definitions are indeed equivalent.  $\square$

**Remark 4.9.** In [1] it is proved that  $G^*$  is the orthogonal complement of  $G$  with respect to the bilinear form

$$\mathrm{SL}(W)/J_W \times \mathrm{SL}(W^*)/J_{W^*} \rightarrow \mathbb{C}^*, (e^{2\pi i v}, e^{2\pi i w}) \mapsto e^{2\pi i v^T A w}.$$

It can be easily proved that the same description can be given for the generalized Berglund-Hübsch-Krawitz construction.

**Definition 4.10.** The *generalized Berglund-Hübsch-Krawitz dual family* of the family  $\mathcal{F}(A, \tilde{G})$  is the family  $\mathcal{F}(A^T, \tilde{G}^*)$  formed by the orbifolds

$$Y_{W^*, \tilde{G}^*} = \{W^* = 0\}/\tilde{G}^*.$$

We want to emphasize that the above construction is a direct generalization of the original BHK construction, simply replacing canonical simplices (with the quasismoothness condition) by arbitrary canonical polytopes. It is quite striking that the transposition rule extends naturally to the case of a non square matrix, thanks to Cox construction of homogeneous coordinates on the ambient toric varieties.

We conclude with the proof of Theorem 2, which shows that the generalized BHK construction can be described as a duality of good pairs.

*Proof of Theorem 2.* Let  $\Delta_2 \subset M_{\mathbb{Q}}$  be the anticanonical polytope of  $X$  and  $\Delta_1 \subset \Delta_2$  be the Newton polytope of  $Y_W$ . We recall that the anticanonical polytope of  $X/\tilde{G}$  is the same polytope  $\Delta_2 \subset (M_G)_{\mathbb{Q}}$  and the Newton polytope of  $Y_{W,G}$  is  $\Delta_1 \subset (M_G)_{\mathbb{Q}}$ . Thus  $(\Delta_1, \Delta_2)$  is a good pair in  $(M_G)_{\mathbb{Q}}$  as already explained in the proof of Proposition 4.7. We now consider the polar pair  $(\Delta_2^*, \Delta_1^*)$  as a good pair in  $(M_W^\vee)_{\mathbb{Q}}$ . By construction  $X^*$  is defined by the fan over the faces of  $\Delta_1$  and the vertices of  $\Delta_2^*$  correspond to the monomials of  $W^*$ . Thus  $\Delta_1^*$  is the anticanonical polytope of  $X^*$  and  $\Delta_2^*$  is the Newton polytope of  $Y_{W^*}$ . Looking at the definition of the transposed group  $G^*$  one immediately sees that the pair  $(\Delta_2^*, \Delta_1^*)$  seen in  $(M_G^\vee)_{\mathbb{Q}}$  defines the toric variety  $X^*/\tilde{G}^*$  and the hypersurface  $Y_{W^*, G^*}$ .  $\square$

**Example 4.11.** Let  $X = \mathbb{P}(5, 5, 4, 4, 2)$  and let  $\Delta_1 \subset M_{\mathbb{Q}}$  be the convex hull of the lattice points  $u_1, \dots, u_5 \in M$  corresponding to the monomials

$$x_1^3 x_2, x_2^4, x_3^5, x_4^5, x_5^{10}.$$

The polytope  $\Delta_1$  is a simplex. We denote by  $\Delta_2 \subset M_{\mathbb{Q}}$  the anticanonical polytope of  $X$ , thus  $(\Delta_1, \Delta_2)$  is a good pair and they both are reflexive. Since  $\Delta_1$  and  $\Delta_2$  are simplices, then the diagonal group of automorphisms preserving

$$W = x_1^3 x_2 + x_2^4 + x_3^5 + x_4^5 + x_5^{10}$$

is finite and  $\mathrm{SL}(W)/J_W \cong \mathbb{Z}/5\mathbb{Z}$ . The polynomial  $W$  is quasismooth since it satisfies the conditions of [34, Theorem 1]. The BHK construction gives a dual Calabi-Yau hypersurface in the toric variety  $Y$  which is the quotient of  $\mathbb{P}(10, 6, 6, 5, 3)$  by the group of order five generated by  $g = (1, e^{2\pi i \frac{1}{5}}, e^{2\pi i \frac{4}{5}}, 1, 1)$ . The hypersurface is defined by the following polynomial in the Cox coordinates of  $Y$ :

$$W^* = y_1^3 + y_1 y_4^4 + y_2^5 + y_3^5 + y_5^{10}.$$

The threefold defined by  $W$  in  $X$  is a birational projective model for the Borcea-Voisin Calabi-Yau threefold obtained from the elliptic curve  $E := \{x_0^2 + x_1^3 x_2 + x_2^4 = 0\} \subset \mathbb{P}(2, 1, 1)$  and the K3 surface  $S := \{y_0^2 + y_1^5 + y_2^5 + y_3^{10} = 0\} \subset \mathbb{P}(5, 2, 2, 1)$  (see [2, Proposition 4.4]).

**Example 4.12.** Let  $X = \mathbb{P}(5, 5, 4, 4, 2)$  as in Example 4.11 and let  $\Delta'_1 \subset \Delta_2$  be the convex hull of the lattice points  $u_1, \dots, u_8 \in M$  corresponding to the monomials

$$x_1^3 x_2, x_2^4, x_3^5, x_4^5, x_5^{10}, x_1^2 x_5^5, x_1^2 x_4^2 x_5, x_1^2 x_3^2 x_5.$$

Observe that  $\Delta_1 \subset \Delta'_1$  and  $\Delta'_1$  is not a simplex. We have that  $(\Delta'_1, \Delta_2)$  is a good pair of reflexive polytopes and one can prove that the general element of  $\mathcal{F}_{\Delta'_1, \Delta_2^*}$  is quasismooth. The lattice generated by  $u_1, \dots, u_8$  is equal to  $M$  so that the only possible choice for  $\tilde{G}$  is the trivial group. Thus starting from  $\Delta'_1$  one obtains the family of Calabi-Yau hypersurfaces  $\mathcal{F}(A', G)$

$$\alpha_1 x_1^3 x_2 + \alpha_2 x_2^4 + \alpha_3 x_3^5 + \alpha_4 x_4^5 + \alpha_5 x_5^{10} + \alpha_6 x_1^2 x_5^5 + \alpha_7 x_1^2 x_4^2 x_5 + \alpha_8 x_1^2 x_3^2 x_5 = 0.$$

The generalized BHK construction gives the dual family  $\mathcal{F}((A')^T, G^*)$

$$\beta_1 y_1^3 y_3^2 y_4^2 y_5^2 + \beta_2 y_1 y_2^4 + \beta_3 y_3^5 y_4 y_5 y_6^{10} + \beta_4 y_4^2 y_7^5 + \beta_5 y_5^2 y_8^5 = 0$$

in the toric variety whose Cox ring is  $\mathbb{C}[y_1, \dots, y_8]$  with Class group isomorphic to  $\mathbb{Z}^4$  and grading matrix

$$\begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 5 & 0 & 0 & 10 & 1 & 4 & 0 \\ 0 & 5 & 0 & 5 & 5 & 1 & 2 & 2 \\ 0 & 7 & 1 & 4 & 9 & 1 & 4 & 2 \end{pmatrix}.$$

## 5. ABOUT THE STRINGY HODGE NUMBERS OF GOOD PAIRS

The polar duality between good pairs provides a duality between families of Calabi-Yau hypersurfaces of  $\mathbb{Q}$ -Fano toric varieties. Of course a natural question arises:

*Question.* Do the families  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta_2^*, \Delta_1}$  associated to a good pair  $(\Delta_1, \Delta_2)$  satisfy the topological mirror test, i.e.

$$h_{st}^{p,q}(X) = h_{st}^{n-p,q}(X^*), \quad 0 \leq p, q \leq n$$

for general  $X \in \mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $X^* \in \mathcal{F}_{\Delta_2^*, \Delta_1}$ ?

This is known to be true for anticanonical hypersurfaces of toric Fano varieties (i.e. when  $\Delta_1 = \Delta_2$ ) by Batyrev and Borisov [8] and in the classical Berglund-Hübsch case by Chiodo and Ruan [17] (i.e. when  $\Delta_1, \Delta_2$  are simplices and quasismoothness holds) in terms of Chen-Ruan orbifold cohomology. Observe that in the latter case orbifold Hodge numbers are known to be the same as the stringy Hodge numbers (see [44] and [13]). Unfortunately we are unable to give a complete answer at the moment. Example 5.3 shows that the answer is negative in general and Example 5.4 leads us to think that the answer could be positive if quasismoothness holds.

We recall that in case the polytope is reflexive of dimension four Hodge numbers can be computed in terms of combinatorial properties of the polytope [5, Corollary 4.5.1].

The following result allows to compute the stringy Hodge numbers for good pairs of four dimensional reflexive polytopes. We recall that, given a reflexive polytope  $\Delta$ , an *MPCP resolution* of the toric variety  $X_\Delta$  is defined by a complete, simplicial projective fan which is a subdivision of the normal fan  $\Sigma_\Delta$  to  $\Delta$  and whose rays are generated by nonzero lattice points of  $\Delta^*$ . In [24] the author introduces the more general notion of  $\Delta$ -*maximal fan*, which has the same definition of an MPCP resolution except for the fact that it is not necessarily projective and it does not need to refine  $\Sigma_\Delta$ . By [5, Corollary 4.2.3] an MPCP resolution induces a crepant resolution of the general anticanonical hypersurface of  $X_\Delta$  in case  $\Delta$  is four dimensional. By [24, Theorem 4.9], given any  $\Delta$ -maximal fan  $\Sigma$  for a four dimensional reflexive polytope  $\Delta$ , the general member of the anticanonical linear series of the toric variety  $X_\Sigma$  is smooth.

**Proposition 5.1.** *Let  $\Delta_1 \subseteq \Delta_2$  be four dimensional reflexive polytopes. The general elements of the families  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta_1, \Delta_1^*}$  have the same Hodge numbers.*

*Proof.* Let  $\Sigma_2$  be the normal fan to  $\Delta_2$  and  $\Sigma'_2$  be the fan of an MPCP resolution of  $X_{\Delta_2}$ . By [24, Lemma 6.1] there exists a projective  $\Delta_1$ -maximal fan  $\tilde{\Sigma}_2$  which refines  $\Sigma'_2$ , and thus  $\Sigma_2$ . Let  $\phi : X_{\tilde{\Sigma}_2} \rightarrow X_{\Sigma_2}$  be the corresponding toric morphism. By [24, Theorem 4.9] the general member  $\tilde{Y}$  of the anticanonical linear series of  $X_{\tilde{\Sigma}_2}$  is smooth. Moreover, by Corollary 2.7,  $\tilde{Y}$  is a crepant resolution of the general member  $Y$  of  $\mathcal{F}(\Delta_1)$ . Now let  $\tilde{Z}$  be a crepant resolution of the general anticanonical hypersurface  $Z$  of  $X_{\Delta_1}$ , induced by an MPCP resolution. By [6, Theorem 3.12]  $\tilde{Y}$  and  $Y$  have the same stringy Hodge numbers, and the same holds for  $\tilde{Z}$  and  $Z$ . Since  $\tilde{Y}$  and  $\tilde{Z}$  are birational smooth Calabi-Yau threefolds (since they are given by the same Newton polytope), then they have the same Hodge numbers [7, Theorem 1.1], which achieves the proof.  $\square$

**Corollary 5.2.** *Let  $(\Delta_1, \Delta_2)$  be a good pair of reflexive, four dimensional polytopes. The associated families  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta_2^*, \Delta_1}$  satisfy the topological mirror test if and only if the general elements of the families  $\mathcal{F}_{\Delta_1, \Delta_1^*}$  and  $\mathcal{F}_{\Delta_2, \Delta_2^*}$  have the same Hodge numbers.*

*Proof.* It follows from Proposition 5.1 and [5, Corollary 4.5.1], which gives that  $h^{1,1}(\Delta_2) = h^{2,1}(\Delta_2^*)$  and viceversa.  $\square$

**Example 5.3.** Consider the following four dimensional reflexive polytopes:

$$\begin{aligned}\Delta_1 &= \text{Conv}((1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1), (-1, -1, -1, -1)), \\ \Delta_2 &= \text{Conv}((1, 0, 0, 0), (0, 1, 0, 0), (0, 0, 1, 0), (0, 0, 0, 1), \\ &\quad (-1, -1, -1, -1), (1, 1, 1, 1), (0, 0, 0, -1)).\end{aligned}$$

By Proposition 5.1 and [5, Corollary 4.5.1] the Hodge numbers of  $(\Delta_1, \Delta_2)$  are  $(h^{1,1}, h^{2,1}) = (101, 1)$  (observe that the variety associated to the normal fan to  $\Delta_1$  is  $\mathbb{P}^4$ ), whereas the Hodge numbers of the dual pair are  $(3, 79)$ . Thus the dual families  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta_2^*, \Delta_1}$  do not satisfy the topological mirror test. A direct computation shows the general member of  $\mathcal{F}_{\Delta_1, \Delta_2^*}$  is not quasismooth. This example appears in [23, §3].

**Example 5.4.** Let  $\mathcal{F}_{\Delta'_1, \Delta_2^*}$  be the family of hypersurfaces of  $\mathbb{P}(5, 5, 4, 4, 2)$  associated to the reflexive polytope  $\Delta'_1 \subset \Delta_2$  defined in Example 4.12. The Hodge numbers of  $\Delta'_1$  and  $\Delta_2$  are equal to  $(h^{1,1}, h^{2,1}) = (15, 39)$  by Batyrev formulas. Thus by Corollary 5.2 the dual families  $\mathcal{F}_{\Delta'_1, \Delta_2^*}$  and  $\mathcal{F}_{\Delta_2^*, \Delta'_1}$  satisfy the topological mirror test.

Combinatorially, the duality of good pairs can be seen as included in the unified setting for mirror symmetry sketched in [12, §7], except for the regularity condition described in [12, Proposition 7.1.3]. We intend to explore such condition, in terms of combinatorial properties of polytopes, in a further work.

*Acknowledgments.* It is a pleasure to thank Gilberto Bini, Samuel Boissière, Antonio Laface and Alessandra Sarti for several useful discussions.

## REFERENCES

- [1] Michela Artebani, Samuel Boissière, and Alessandra Sarti, *The Berglund-Hübsch-Chiodo-Ruan mirror symmetry for K3 surfaces*, J. Math. Pures Appl. (9) **102** (2014), no. 4, 758–781. MR3258129 ↑15, 18, 19
- [2] ———, *Borcea-Voisin Calabi-Yau threefolds and invertible potentials (with an appendix in collaboration with Sara A. Filippini)*, Math. Nachr. 1–11 (2015). ↑20
- [3] A. C. Avram, P. Candelas, D. Jančić, and M. Mandelberg, *On the connectedness of the moduli space of Calabi-Yau manifolds*, Nuclear Phys. B **465** (1996), no. 3, 458–472. MR1392861 (97f:14038) ↑14
- [4] A. C. Avram, M. Kreuzer, M. Mandelberg, and H. Skarke, *The web of Calabi-Yau hypersurfaces in toric varieties*, Nuclear Phys. B **505** (1997), no. 3, 625–640. MR1490782 (99a:14056) ↑14
- [5] Victor V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, J. Algebraic Geom. **3** (1994), no. 3, 493–535. ↑1, 4, 10, 11, 12, 21, 22
- [6] ———, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, Integrable systems and algebraic geometry (Kobe/Kyoto, 1997), 1998, pp. 1–32. MR1672108 (2001a:14039) ↑21
- [7] ———, *Birational Calabi-Yau n-folds have equal Betti numbers*, New trends in algebraic geometry (Warwick, 1996), 1999, pp. 1–11. MR1714818 (2000i:14059) ↑21
- [8] Victor V. Batyrev and Lev A. Borisov, *Mirror duality and string-theoretic Hodge numbers*, Invent. Math. **126** (1996), no. 1, 183–203. MR1408560 (97k:14039) ↑1, 21
- [9] Victor V. Batyrev and David A. Cox, *On the Hodge structure of projective hypersurfaces in toric varieties*, Duke Math. J. **75** (1994), no. 2, 293–338. MR1290195 (95j:14072) ↑10
- [10] Per Berglund and Tristan Hübsch, *A generalized construction of mirror manifolds*, Nuclear Phys. B **393** (1993), no. 1–2, 377–391. MR1214325 (94k:14031) ↑1, 14
- [11] Per Berglund, Sheldon Katz, and Albrecht Klemm, *Mirror symmetry and the moduli space for generic hypersurfaces in toric varieties*, Nuclear Phys. B **456** (1995), no. 1–2, 153–204. MR1372213 (97a:32016) ↑14



- [12] Lev A. Borisov, *Berglund-Hübsch mirror symmetry via vertex algebras*, Comm. Math. Phys. **320** (2013), no. 1, 73–99. MR3046990 ↑2, 18, 22
- [13] Lev A. Borisov and Anvar R. Mavlyutov, *String cohomology of Calabi-Yau hypersurfaces via mirror symmetry*, Adv. Math. **180** (2003), no. 1, 355–390. MR2019228 (2005b:32056) ↑21
- [14] Wieb Bosma, John Cannon, and Catherine Playoust, *The Magma algebra system. I. The user language, computational algebra and number theory* (London, 1993), J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. ↑10
- [15] Jean-François Boutot, *Singularités rationnelles et quotients par les groupes réductifs*, Invent. Math. **88** (1987), no. 1, 65–68. MR877006 (88a:14005) ↑10
- [16] Ti-Ming Chiang, Brian R. Greene, Mark Gross, and Yakov Kanter, *Black hole condensation and the web of Calabi-Yau manifolds*, Nuclear Phys. B Proc. Suppl. **46** (1996), 82–95. *S-duality and mirror symmetry* (Trieste, 1995). MR1411461 (97g:32022) ↑14
- [17] Alessandro Chiodo and Yongbin Ruan, *LG/CY correspondence: the state space isomorphism*, Adv. Math. **227** (2011), no. 6, 2157–2188. MR2807086 (2012g:14069) ↑2, 15, 21
- [18] Alessio Corti and Vasily Golyshev, *Hypergeometric equations and weighted projective spaces*, Sci. China Math. **54** (2011), no. 8, 1577–1590. MR2824960 ↑14
- [19] David A. Cox, *Toric residues*, Ark. Mat. **34** (1996), no. 1, 73–96. ↑10
- [20] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR2810322 (2012g:14094) ↑3, 4, 5, 7, 16
- [21] Alexandru Dimca, *Singularities and coverings of weighted complete intersections*, J. Reine Angew. Math. **366** (1986), 184–193. MR833017 (87g:14056) ↑9
- [22] Charles F. Doran, David Favero, and Tyler L. Kelly, *Equivalences of families of stacky toric calabi-yau hypersurfaces*, available at <http://arxiv.org/abs/1503.04888>. ↑14
- [23] Karl Fredrickson, *Extremal transitions from nested reflexive polytopes*, Comm. Math. Phys. **335** (2015), no. 3, 1381–1395. MR3320316 ↑13, 22
- [24] ———, *Generalized compactifications of batyrev hypersurface families* (2015), available at <http://arxiv.org/abs/1410.8287>. ↑21
- [25] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52. MR0463157 (57 #3116) ↑7
- [26] Anthony R. Iano-Fletcher, *Working with weighted complete intersections*, Explicit birational geometry of 3-folds, 2000, pp. 101–173. MR1798982 (2001k:14089) ↑6
- [27] Alexander M. Kasprzyk, *Canonical toric Fano threefolds*, Canad. J. Math. **62** (2010), no. 6, 1293–1309. MR2760660 (2012c:14087) ↑4
- [28] János Kollár, *Flips and abundance for algebraic threefolds*, Société Mathématique de France, Paris, 1992. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991, Astérisque No. 211 (1992). MR1225842 (94f:14013) ↑8
- [29] ———, *Singularities of pairs*, Algebraic geometry-Santa Cruz 1995, 1997, pp. 221–287. MR1492525 (99m:14033) ↑10
- [30] ———, *Singularities of the minimal model program*, Cambridge Tracts in Mathematics, vol. 200, Cambridge University Press, Cambridge, 2013. With a collaboration of Sándor Kovács. MR3057950 ↑8, 9, 10, 12
- [31] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. MR1658959 (2000b:14018) ↑8
- [32] János Kollár, Karen E. Smith, and Alessio Corti, *Rational and nearly rational varieties*, Cambridge Studies in Advanced Mathematics, vol. 92, Cambridge University Press, Cambridge, 2004. MR2062787 (2005i:14063) ↑7, 8
- [33] Marc Krawitz, *FJRW rings and Landau-Ginzburg mirror symmetry*, ProQuest LLC, Ann Arbor, MI, 2010. Thesis (Ph.D.)—University of Michigan. MR2801653 ↑1, 14
- [34] Maximilian Kreuzer and Harald Skarke, *On the classification of quasihomogeneous functions*, Comm. Math. Phys. **150** (1992), no. 1, 137–147. MR1188500 (93k:32075) ↑14, 20
- [35] ———, *Classification of reflexive polyhedra in three dimensions*, Adv. Theor. Math. Phys. **2** (1998), no. 4, 853–871. MR1663339 (99m:14100) ↑4
- [36] ———, *Complete classification of reflexive polyhedra in four dimensions*, Adv. Theor. Math. Phys. **4** (2000), no. 6, 1209–1230. MR1894855 (2003e:52020) ↑4

- [37] ———, *Reflexive polyhedra, weights and toric Calabi-Yau fibrations*, Rev. Math. Phys. **14** (2002), no. 4, 343–374. MR1901220 (2003b:14062) ↑4
- [38] David R. Morrison, *Through the looking glass*, Mirror symmetry, III (Montreal, PQ, 1995), 1999, pp. 263–277. MR1673108 (2000d:14049) ↑13
- [39] Max Pumperla, *Global aspects of moduli spaces and mirror symmetry*, PhD thesis, 2008. ↑13
- [40] M. Reid, *Decomposition of toric morphisms*, Arithmetic and geometry, 1983, pp. 395–418. ↑16
- [41] Michele Rossi, *Geometric transitions*, J. Geom. Phys. **56** (2006), no. 9, 1940–1983. MR2240431 (2007e:14061) ↑13
- [42] Mark Shoemaker, *Birationality of Berglund-Hübsch-Krawitz mirrors*, Comm. Math. Phys. **331** (2014), no. 2, 417–429. MR3238520 ↑3, 14, 17
- [43] Harald Skarke, *Weight systems for toric Calabi-Yau varieties and reflexivity of Newton polyhedra*, Modern Phys. Lett. A **11** (1996), no. 20, 1637–1652. MR1397227 (97e:14054) ↑10, 14, 15, 16
- [44] Takehiko Yasuda, *Twisted jets, motivic measures and orbifold cohomology*, Compos. Math. **140** (2004), no. 2, 396–422. MR2027195 (2004m:14037) ↑21

DEPARTAMENTO DE MATEMÁTICA,  
 UNIVERSIDAD DE CONCEPCIÓN,  
 CASILLA 160-C, CONCEPCIÓN, CHILE  
*E-mail address:* `martebani@udec.cl`

DEPARTAMENTO DE MATEMÁTICA,  
 UNIVERSIDAD DE CONCEPCIÓN,  
 CASILLA 160-C, CONCEPCIÓN, CHILE  
*E-mail address:* `pcomparin@udec.cl`

INSTITUTO NACIONAL DE MATEMÁTICA PURA E APLICADA,  
 ESTRADA DONA CASTORINA, 110  
 RIO DE JANEIRO, BRASIL  
*E-mail address:* `rguilbot@math.cnrs.fr`