

Extraction of Neutrino Flux from the Inclusive Muon Cross Section

Tomoya Murata*

Osaka University

E-mail: murata@kern.phys.sci.osaka-u.ac.jp

Toru Sato

Osaka University

E-mail: tsato@phys.sci.osaka-u.ac.jp

We have studied a method to extract neutrino flux from the data of neutrino-nucleus reaction by using maximum entropy method. We demonstrate a promising example to extract neutrino flux from the inclusive cross section of muon production without selecting a particular reaction process such as quasi-elastic nucleon knockout.

16th International Workshop on Neutrino Factories and Future Neutrino Beam Facilities

25 -30 August, 2014

University of Glasgow, United Kingdom

*Speaker.

1. Introduction

A precise description of the neutrino-nucleus interaction is crucial for the study the neutrino properties such as CP violating phase and mass hierarchy[1, 2, 3, 4] when one extracts the neutrino flux from the neutrino data. The quasi-elastic (QE) reaction is the key mechanism to reconstruct the neutrino flux from the data of neutrino-nucleus reaction. Neutrino energy(E_ν) can be determined from the muon scattering angle(θ) and energy(E_μ) by using the kinematics of the reaction of free nucleon at rest.

$$E_\nu = \frac{E_\mu M_N - m_\mu^2/2}{M_N - E_\mu + p_\mu \cos \theta} \quad (1.1)$$

Since the reaction takes place inside nuclear many body system, various initial and final state interactions can spoil the useful relation of Eq. (1.1). The observed quasi-elastic like events are mixture of the various reaction mechanism, which contribute the reaction amplitudes and, in principle, interference among various mechanism exists. The central issue of the current theoretical research of neutrino-nucleus reaction is to improve the accuracy of the estimation of various nuclear effects[5, 6] by testing the model from the neutrino data.

Because of the rather wide energy band of the neutrino flux in the current long baseline experiments, the observed cross section of neutrino reaction ($\langle \sigma \rangle$) is a result of neutrino flux(Φ_ν) average of the cross section (σ).

$$\langle \sigma \rangle = \int dE_\nu \sigma(E_\nu) \Phi_\nu(E_\nu) \quad (1.2)$$

One can regard the extraction of neutrino flux Φ_ν from the data $\langle \sigma \rangle$ as an inverse problem to obtain Φ_ν from $\langle \sigma \rangle$ provided the cross section σ is well under control. In this report we will examine a method to solve the inverse problem, where the neutrino flux can be extracted without using the formula (1.1) nor the separation of 'true' QE events. For this purpose, we use the maximum entropy method (MEM) which are widely used in the field of condensed matter physics and Lattice QCD[7, 8, 9].

We briefly outline the application of MEM to extract neutrino flux from data in section 2. To examine the usefulness of the approach, we generate pseudo data of neutrino reaction. The pseudo data are generated by using a model of neutrino reaction and neutrino flux, which are explained in section 3. Here, as an example of the pseudo data, we use double differential cross section of muon for the inclusive neutrino nucleus reaction. Results on how well the neutrino flux can be extracted from the pseudo 'data' without assuming QE mechanism is shown in section 4.

2. Maximum Entropy Method

In this section we briefly explain the maximum entropy method. The most plausible neutrino flux (Φ_ν^{MEM}) from the experimental data based on the Bayes' theorem is given by the functional integral of the neutrino flux Φ

$$\Phi_\nu^{MEM} = \int [d\Phi] \Phi P[\Phi | \bar{D}, I], \quad (2.1)$$

where $P[\Phi|\bar{D}, I]$ is a conditional probability of neutrino flux Φ for given data of neutrino reaction $\bar{D} = \{< \sigma >_{exp}\}$ and the prior information on the neutrino flux I . I is called as default model of neutrino flux. Introducing auxiliary variable α , we can write the above formula as

$$\Phi_v^{MEM} = \int d\alpha \Phi_v^\alpha P[\alpha|\bar{D}, I], \quad (2.2)$$

where Φ_v^α is neutrino flux for given α , data $\bar{D}$ and prior information I given as,

$$\Phi_v^\alpha = \int [d\Phi] \Phi P[\Phi|\alpha, \bar{D}, I] \propto \int [d\Phi] \Phi P[\bar{D}|\Phi, \alpha, I] P[\Phi|\alpha, I]. \quad (2.3)$$

Here $P[\bar{D}|\Phi, \alpha, I]$ and $P[\Phi|\alpha, I]$ are called as likefood function and prior probability, respectively. The likefood function is written as

$$P[\bar{D}|\Phi, I] = \frac{1}{Z_\chi} \exp\left(-\frac{1}{2}\chi^2\right), \quad (2.4)$$

with

$$\begin{aligned} \chi^2 &= \sum_{l=1} \frac{(\bar{D}_l - D_l)^2}{\sigma_l^2} \\ Z_\chi &= \prod_{l=1} \sqrt{2\pi\sigma_l^2}. \end{aligned} \quad (2.5)$$

Here $\bar{D}_l$ and D_l are cross sections $< \sigma >_{exp}$ and $< \sigma >$.

The prior probability can be expressed as

$$P[\Phi|\alpha, I] = \frac{1}{Z_S} \exp(\alpha S), \quad (2.6)$$

where S is the Shannon-Jaynes entropy,

$$S = \sum_i \left(\Phi_i - m_i - \Phi_i \ln \frac{\Phi_i}{m_i} \right). \quad (2.7)$$

m_i is default model of neutrino flux at neutrino energy E_i . Combining the likefood function and the prior probability, neutrino flux Φ_v^α is the maximum probability of $P[\Phi|\alpha, \bar{D}, I] \propto e^{Q(\Phi)}$ with $Q(\Phi) = \alpha S - \frac{1}{2}\chi^2$.

Finally the α dependent flux is integrated with the probability $P[\alpha|\bar{D}, I]$, which has a sharp peak as a function of α written as

$$P[\alpha|\bar{D}, I] = \int [d\Phi] P[\Phi|\alpha, \bar{D}, I] \propto P[\alpha|I] \int [d\Phi] \frac{1}{Z_S Z_\chi} e^{Q(\Phi)} \quad (2.8)$$

Assuming constant $P[\alpha|I]$, we obtain the neutrino flux Φ_v^{MEM} by integrating α around the sharp peak of $P[\alpha|\bar{D}, I]$ in Eq. 2.2.

One can also estimate the errors of the extracted neutrino flux Φ_v^{MEM} within MEM. For given α , the average flux $< \Phi_\alpha >_S$ in energy region S of neutrino is defined as

$$\langle \Phi_v^\alpha \rangle_S = \frac{\int_S dE_v \Phi_v^\alpha(E_v)}{\int_S dE_v}. \quad (2.9)$$

The covariance of $\langle \Phi_V^\alpha \rangle_S$ can be written as

$$\begin{aligned} \langle (\delta \Phi_V^\alpha)^2 \rangle_S &= \int_{S^2} dE_V dE'_V \delta \Phi(E_V) \Phi(E'_V) P[\Phi | \bar{D}, \alpha, I] / \int_{S^2} dE_V dE'_V \\ &\simeq - \int_{S^2} dE_V dE'_V \left(\frac{\delta^2 Q}{\delta \Phi(E_V) \delta \Phi(E'_V)} \right)_{\Phi=\Phi_\alpha}^{-1} / \int_{S^2} dE_V dE'_V. \end{aligned} \quad (2.10)$$

Then the error of the extracted flux is obtained by integrating over α as

$$\langle (\delta \Phi_V^{MEM})^2 \rangle_S = \int d\alpha \langle (\delta \Phi_V^\alpha)^2 \rangle_S P[\alpha | \bar{D}, I]. \quad (2.11)$$

3. Model of inclusive neutrino-nucleus cross section and neutrino flux

To examine the method outlined in the previous section, we construct pseudo data of neutrino-nucleus reaction by using a model of neutrino-nucleus reaction and neutrino flux. The neutrino flux is obtained by applying the formula of two-flavor oscillation as

$$\Phi(E_V) = \left(1 - \sin^2 2\theta_{23} \sin^2 \left(\frac{\Delta m_{23}^2 L}{4E_V} \right) \right) \Phi_V^0(E_V) \quad (3.1)$$

Here we used $\Delta m_{23}^2 = 2.5 \times 10^{-3} \text{eV}^2$, $\sin^2 2\theta_{23} = 1.0$ and $L = 295 \text{km}$. The initial flux $\Phi_V^0(E_V)$ is taken from near detector neutrino flux of T2K experiment Ref. [10]. The solid(black) curve in Fig.




Figure 1: Neutrino flux. The solid(black) and the dashed(red) lines show the neutrino flux Φ_V and Φ_V^0 .

1 is the neutrino flux we use to calculate the pseudo data, while the original flux Φ_V^0 is shown in dashed(red) curve for comparison.

As a observable to test the method of flux extraction, we use double differential cross section of muon of neutrino induced inclusive reaction on ^{12}C , $\nu_\mu + ^{12}\text{C} \rightarrow \mu^- + X$

$$\frac{d^2 \sigma}{dE_\mu d\Omega_\mu}. \quad (3.2)$$

In the energy region of neutrino flux in Fig. 1, the main reaction mechanism is quasi elastic scattering and pion production through the $\Delta(1232)$ excitation. The quasi elastic nucleon knockout

process is calculated using modified Fermi-gass model which incorporates the spectral function [12, 13]. For the delta excitation region we use the neutrino pion production amplitudes calculated in a formalism in which the resonance contributions and the background amplitudes are treated on the same footing[14]. Detail on the reaction model is described in Ref. [11]. The model of nuclear reaction is rather simple without taking into account the final state interaction, however the reliability of our formalism is tested against the electron nucleus scattering cross sections in the same theoretical framework; the calculated cross sections agree reasonably well with the existing data. Therefore, we believe it is good enough for the purpose of testing the method of neutrino flux extraction.

We now are able to calculate the flux averaged double differential cross section $d^2\sigma/dE_\mu d\Omega_\mu$ from the model of neutrino-nucleus reaction and neutrino flux Φ_ν in Eq. (3.1).

$$\left\langle \frac{d^2\sigma}{dE_\mu d\Omega_\mu} \right\rangle = \int dE_\nu \frac{d^2\sigma}{dE_\mu d\Omega_\mu} \Phi(E_\nu), \quad (3.3)$$

The pseudo data of double differential cross section is obtained as following. We assume muon energy resolution is 100MeV and evaluate the average of cross section as,

$$\left\langle \frac{d^2\sigma}{dE_\mu d\Omega_\mu} \right\rangle = \int_{E_1}^{E_2} dE_\mu \frac{1}{E_2 - E_1} \left\langle \frac{d^2\sigma}{dE_\mu d\Omega_\mu} \right\rangle \quad (3.4)$$

We assume Gaussian distribution of the data with the mean given by Eq. (3.4) and the standard deviation by the 10% of the value of flux averaged double differential cross section at $E_\mu=1\text{GeV}$. Fig. 2 shows the obtained pseudo data of the muon energy distribution at the muon scattering angle $\theta_\mu = 10^\circ$.




Figure 2: Pseudo data of neutrino induced inclusive muon production cross section on ^{12}C . The scattering angle of the muon is $\theta_\mu = 10^\circ$.

4. Results and discussion

We extract neutrino flux from the 19 points of pseudo data of the muon energy distribution shown in Fig. 2. They give $\bar{D}_l$ and σ_l of Eq. (2.5). D_l is calculated from the theoretical cross section of Eq. (3.2).

At first, we study the effects of the default model I on the extracted neutrino flux. We have examined two default models. The one is constant flux and the other is initial flux Φ_V^0 shown in the dashed(red) and solid(black) curve in the left panel of Fig. 3. The extracted flux using each default




Figure 3: Default model dependence of extracted neutrino flux. The solid(black) and dashed(red) curves in the left panel show Φ_V^0 and constant flux as default model. The extracted flux Φ_V^{MEM} corresponding to each default model is shown in the right panel.

model is shown in the right panel of Fig. 3. One can see that, in this case, the extracted flux is very stable against the choice of the default model.

We now compare the extracted flux Φ_V^{MEM} (solid curve) with the expected 'exact' neutrino flux Φ_V (dashed curve) in Fig. 4. Extracted flux agrees well with the 'exact' flux over the wide muon energy region. The extraction of the flux is only valid on the energy above the reaction threshold and the flux extraction did not work well near the threshold energy. We noticed that the extracted flux has structures in the $0.7 < E_\nu < 1$ GeV region. One can judge the structure is real




Figure 4: Comparison of extracted flux Φ_V^{MEM} (solid curve) and 'true' flux Φ_V (dashed curve).

or not by estimating the error of extracted flux. In Table 1, the neutrino flux averaged in the peak region ($0.28 < E_\nu < 0.42$ GeV), dip region ($0.52 < E_\nu < 0.70$ GeV) and higher energy regions ($0.72 < E_\nu < 0.82$ GeV), ($0.86 < E_\nu < 0.94$ GeV) and ($1.00 < E_\nu < 1.08$ GeV) are given. The second row shows the calculated 'exact' value of the average flux and the third row shows the flux obtained from MEM together with the error calculated from Eq. (2.11). The central values of the

Φ_{ν}^{MEM} agree well with the 'exact' ones within the errors in all the energy regions. The size of the oscillation of Φ_{ν}^{MEM} seen in $0.7 < E_{\nu} < 1$ GeV is smaller than the estimated errors in that region. Therefore one can conclude that we could not take the oscillation of the obtained flux seriously from the error estimation even without knowing the 'exact' flux.

$E_{\nu}(\text{GeV})$	0.28 – 0.42	0.52 – 0.70	0.72-0.82	0.86-0.94	1.00-1.08
Exact	53.5	3.9	10.5	10.9	8.7
MEM	56.5 ± 6.5	5.1 ± 1.7	14.0 ± 3.7	11.6 ± 3.7	10.7 ± 1.9

Table 1: Energy averaged neutrino flux.

In summary, we have examined MEM to extract the neutrino flux from the inclusive muon production cross section. The results show the proposed method can be useful alternative tool to extract the neutrino flux provided accurate theoretical cross section is known. Further work to find the optimal observables for the efficient extraction of flux with the refined model of neutrino-nucleus reaction should be done in near future.

The authors would like thank to Drs. M. Kitazawa and S. Nakamura for illuminating discussions. This work was supported by JSPS KAKENHI Grant Nos. 24540273 and 25105010.

References

- [1] K. Abe *et al.* (Hyper-Kamiokande working group), arXiv 1109.3262[hep-ex] (2011).
- [2] C. Adams *et al.* (LBNE collaboration) arXiv:1307.7335[hep-ex] (2013).
- [3] S. K. Agarwalla *et al.* (LAGUNA-LBNO collaboration), JHEP05, 094 (2014).
- [4] S. Bhadra *et al.* (The nuPRISM Collaboration) arXiv:1412.3086 [physics.ins-det] (2014).
- [5] L. Alvarez-Ruso, Y. Hayato and J. Nieves, New Jour. Phys. **16**, 075015 (2014).
- [6] G. T. Garvey, D. A. Harris, H. A. Tanaka, R. Tayloe and G. P. Zeller, arXiv:1412.4294[hep-ex] (2014).
- [7] R. N. Silver, D. S. Sivia and J. E. Gubernatis, Phys. Rev. **B41** 2380 (1990).
- [8] J. E. Gubernatis, M. Jarrell, R. N. Silver and D. S. Sivia, Phys. Rev. **B44**, 6011 (1991).
- [9] M. Asakawa, T. Hatsuda and Y. Nakahara, Prog. Part. Nucl. Phys. **46** 459 (2001).
- [10] K. Abe *et al.* (T2K Collaboration) Phys. Rev. **D87** 012001 (2013).
- [11] B. Szczerbinska, T. Sato, K. Kubodera and T. -S. H. Lee, Phys. Lett. **B649**, 132 (2007).
- [12] O. Benhar, A. Fabrocini, S. Fantoni and I. Sick, Nucl. Phys. **A579** 493 (1994).
- [13] O. Benhar, N. Farina, H. Nakamura, M. Sakuda and R. Seki, Phys. Rev. **D72** 053005 (2005).
- [14] T. Sato, D. Uno and T. -S. H. Lee, Phys. Rev. **C67** 065201 (2003).