ON BIRATIONAL GEOMETRY OF MINIMAL THREEFOLDS WITH NUMERICALLY TRIVIAL CANONICAL DIVISORS

CHEN JIANG

ABSTRACT. For a minimal 3-fold X with $K_X \equiv 0$ and a nef and big Weil divisor L on X, we investigate the birational geometry inspired by L. We prove that |mL| and $|K_X + mL|$ give birational maps for all $m \geq 17$. The result remains true under weaker assumption that L is big and has no stable base components.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Rational map defined by a Weil divisor	4
2.2. Reid's Riemann–Roch formula	5
2.3. Some facts about minimal 3-folds with $K \equiv 0$	7
3. Gorenstein case	8
4. Birationality criterion	9
4.1. Main reduction	9
4.2. Key theorem	10
4.3. Criterion	13
5. Birationality on polarized triples	16
5.1. Estimation of ρ_0	17
5.2. Estimation of m_1	19
5.3. Proof of Theorems 1.5 and 1.6	19
References	24

1. INTRODUCTION

A normal projective variety X is said to be minimal if X has at worst \mathbb{Q} -factorial terminal singularities and the canonical divisor K_X is nef. According to Minimal Model Program, minimal varieties form a fundamental class in birational geometry.

Given an *n*-dimensional normal projective variety X with mild singularities and a big Weil divisor L on X, we are interested in the geometry of the rational map $\Phi_{|mL|}$ defined by the linear system |mL|. By definition,

Date: December 19, 2021.

²⁰⁰⁰ Mathematics Subject Classification. 14E05, 14J30.

Key words and phrases. minimal varieties, birationality, boundedness.

The author is supported by Grant-in-Aid for JSPS Fellows (KAKENHI No. 25-6549) and Program for Leading Graduate Schools, MEXT, Japan.

 $\Phi_{|mL|}$ is birational onto its image when m is sufficiently large. Therefore it is interesting to find such a practical number m(n), depending only on dim X, which stably guarantees the birationality of $\Phi_{|mL|}$. In fact, the following three special cases are the most interesting:

Case 1. K_X is nef and big, $L = K_X$;

Case 2. $K_X \equiv 0$, L is an arbitrary nef and big Weil divisor;

Case 3. $-K_X$ is nef and big, $L = -K_X$.

It is an interesting exercise to deal the case X being a smooth curve or surface. We recall some known results on surfaces.

Theorem 1.1 (cf. Bombieri [2], Reider [25]). Let S be a smooth surface.

- (1) If K_S is nef and big, then $|mK_S|$ gives a birational map for $m \ge 5$;
- (2) If $K_S \equiv 0$, then |mL| gives a birational map for $m \geq 3$ and L an arbitrary nef and big divisor;
- (3) If $-K_S$ is nef and big, then $|-mK_S|$ gives a birational map for $m \ge 3$.

Smooth threefolds were studied by Matsuki [19], M. Chen [7], Ando [1], Fukuda [12], Oguiso [22], and many others, and we have the following known results.

Theorem 1.2 (Chen [7], Fukuda [12]). Let X be a smooth 3-fold.

- (1) If K_X is nef and big, then $|mK_X|$ gives a birational map for $m \ge 6$;
- (2) If $K_X \equiv 0$, then |mL| gives a birational map for $m \ge 6$ and L an arbitrary nef and big divisor;
- (3) If $-K_X$ is nef and big, then $|-mK_X|$ gives a birational map for $m \ge 4$.

When X is a 3-fold with \mathbb{Q} -factorial terminal singularities, Case 1 was systematically treated by J. A. Chen and M. Chen [4]-[6] and Case 3 is systematically treated by M. Chen and the author [10].

Theorem 1.3 (Chen–Chen [6], Chen–Jiang [10]). Let X be a 3-fold with \mathbb{Q} -factorial terminal singularities.

- (1) If K_X is nef and big, then $|mK_X|$ gives a birational map for $m \ge 61$;
- (2) If $-K_X$ is nef and big, then $|-mK_X|$ gives a birational map for $m \ge 97$;
- (3) If $-K_X$ is ample with $\rho(X) = 1$, then $|-mK_X|$ gives a birational map for $m \ge 39$;

Now we consider Case 2, when X is a minimal 3-fold with $K_X \equiv 0$ and L is an arbitrary nef and big Weil divisor on X. If X is smooth, then |mL| gives a birational map for $m \ge 6$ by Fukuda [12]. If X is with Gorenstein terminal singularities and $q(X) := h^1(\mathcal{O}_X) = 0$, then |mL| gives a birational map for $m \ge 5$ by Oguiso–Peternell [23].

The motivation of this paper is to study the birational geometry of minimal 3-fold with $K \equiv 0$. For an arbitrary nef and big Weil divisor L on X, we investigate the birationality of the linear system |mL| and the adjoint linear system $|K_X + mL|$. We remark that the behavior of these two linear systems are a litte different even though they are numerically equivalent. For example, if the local index i(X) > 1, then i(X)L is always a Cartier divisor while $K_X + i(X)L$ can never be (see Section 2).

 $\mathbf{2}$

The difficulty arises from the singularities of X, and the assumption that L is only a Weil divisor. If we assume that L is Cartier, then the problem becomes relatively easy and can be treated by the method of Fukuda [12] using Reider's theorem [25]. On the other hand, fortunately, the singularities of minimal 3-folds with $K \equiv 0$ is not so complicated due to Kawamata [14] and Morrison [21], and this makes it possible to deal with the birationality problem.

As the main result, we prove the following theorem.

Theorem 1.4. Let X be a minimal 3-fold with $K_X \equiv 0$ and a nef and big Weil divisor L. Then |mL| and $|K_X + mL|$ give birational maps for all $m \geq 17$.

In fact, we prove a more general theorem.

Theorem 1.5. Let X be a minimal 3-fold with $K_X \equiv 0$, a nef and big Weil divisor L, and a Weil divisor $T \equiv 0$. Then $|K_X + mL + T|$ gives a birational map for all $m \geq 17$.

Moreover, by Log Minimal Model Program, the assumption that L is nef can be weaken. We say that a divisor D has no stable base components if |mD| has no base components for sufficiently divisible m.

Theorem 1.6. Let X be a minimal 3-fold with $K_X \equiv 0$, a big Weil divisor L without stable base components, and a Weil divisor $T \equiv 0$. Then $|K_X + mL + T|$ gives a birational map for all $m \ge 17$. In particular, |mL| and $|K_X + mL|$ give birational maps for all $m \ge 17$.

As a by-product, we prove a direct generalization of Fukuda [12] and Oguiso–Peternell [23] which is optimal by the general weighted hypersurface $X_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$.

Theorem 1.7 (=Theorem 3.2). Let X be a minimal Gorenstein 3-fold with $K_X \equiv 0$, a nef and big Weil divisor L, and a Weil divisor $T \equiv 0$. Then $|K_X + mL + T|$ gives a birational map for all $m \geq 5$.

For the convenience, we introduce the following definition.

Definition 1.8. (X, L, T) is called a *polarized triple* if X is a minimal 3-fold with q(X) = 0 and $K_X \equiv 0$, L is a nef and big Weil divisor, and T is a numerically trivial Weil divisor on X.

Note that we assume q(X) = 0 in the definition. The case that q(X) > 0 is relatively easy and we treat it in Section 3 (see Theorem 3.2).

This paper is organized as follows. In Section 2, we recall some basic knowledge and facts. In Section 3, we treat Gorenstein case. We study the birationality of polarized triples in Section 4 and give an effective criterion for the birationality of linear systems. In the last section, to apply the birationality criterion, we estimate several quantities of polarized triples. We prove Theorems 1.5 and 1.6 in the last part.

Acknowledgment. The author would like to express his gratitude to his supervisor Professor Yujiro Kawamata for suggestions and encouragement. He appreciates the very effective discussion with Professors Meng Chen and Keiji Oguiso during the preparation of this paper. Part of this paper was

written during the author's visit to Fudan University and he would like to thank for the hospitality and support. The author would like to thank the anonymous reviewer for his valuable comments and suggestions to improve the explanation of the paper.

2. Preliminaries

Throughout we work over an algebraically closed field k of characteristic 0 (for instance, $k = \mathbb{C}$). We adopt the standard notation in Kollár–Mori [18], and will freely use them.

Let X be a minimal 3-fold with $K_X \equiv 0$. Denote by i(X) the local index of X, i.e. the Cartier index of K_X . By Kawamata [15, Corollary 5.2], for arbitrary Weil divisor D on X, i(X)D is a Cartier divisor. By Kawamata [13, Theorem 8.2], $K_X \sim_{\mathbb{Q}} 0$ and we define the global index

$$I(X) = \min\{m \in \mathbb{N} \mid mK_X \sim 0\}.$$

Note that i(X)|I(X).

For two linear systems |A| and |B|, we write $|A| \leq |B|$ if there exists an effective divisor F such that

$$|B| \supset |A| + F.$$

In particular, if $A \leq B$ as divisors, then $|A| \leq |B|$.


2.1. Rational map defined by a Weil divisor. Consider a Q-Cartier Weil divisor D on X with $h^0(X, D) \ge 2$. We study the rational map defined by |D|, say

$$X \xrightarrow{\Phi_D} \mathbb{P}^{h^0(D)-1}$$

which is not necessarily well-defined everywhere. By Hironaka's big theorem, we can take successive blow-ups $\pi: Y \to X$ such that:

- (1) Y is smooth projective;
- (2) the movable part |M| of the linear system $|\lfloor \pi^*(D) \rfloor|$ is base point free and, consequently, the rational map $\gamma := \Phi_D \circ \pi$ is a morphism;
- (3) the support of the union of $\pi_*^{-1}(D)$ and the exceptional divisors of π is of simple normal crossings.

Let $Y \xrightarrow{f} \Gamma \xrightarrow{s} Z$ be the Stein factorization of γ with $Z := \gamma(Y) \subset \mathbb{P}^{h^0(D)-1}$. We have the following commutative diagram.



Case (f_{np}) . If dim $(\Gamma) \geq 2$, a general member S of |M| is a smooth projective surface by Bertini's theorem. We say that |D| is not composed with a pencil of surfaces.

Case (f_p) . If dim $(\Gamma) = 1$, i.e. dim $\overline{\Phi_D(X)} = 1$, a general fiber S of f is an irreducible smooth projective surface by Bertini's theorem. We may write

$$M = \sum_{i=1}^{a} S_i \equiv aS$$

where S_i is a smooth fiber of f for all i. We say that |D| is composed with a pencil of surfaces. It is clear that $a \ge h^0(D) - 1$. Furthermore, $a = h^0(D) - 1$ if and only if $\Gamma \cong \mathbb{P}^1$, and then we say that |D| is composed with a rational pencil of surfaces. In particular, if q(X) = 0, then $\Gamma \cong \mathbb{P}^1$ since $g(\Gamma) \le q(Y) = q(X) = 0$.

For another Weil divisor D' satisfying $h^0(X, D') > 1$, we say that |D| and |D'| are composed with the same pencil if |D| and |D'| are composed with pencils and they define the same fibration structure $Y \to \Gamma$ on some smooth model Y. In particular, |D| and |D'| are not composed with the same pencil if one of them is not composed with a pencil.

Define

$$\iota = \iota(D) := \begin{cases} 1, & \text{Case } (f_{np}); \\ a, & \text{Case } (f_{p}). \end{cases}$$

Clearly, in both cases, $M \equiv \iota S$ with $\iota \geq 1$.

Definition 2.1. For both Case (f_{np}) and Case (f_p) , we call S a general irreducible element of |M|.

We may also define "a general irreducible element" of a moving linear system on any surface in the similar way.

2.2. Reid's Riemann–Roch formula. A basket B is a collection of pairs of integers (permitting weights), say $\{(b_i, r_i) \mid i = 1, \dots, s; b_i \text{ is coprime to } r_i\}$. For simplicity, we will alternatively write a basket as follows, say

$$B = \{(1,2), (1,2), (2,5)\} = \{2 \times (1,2), (2,5)\}.$$

Let X be a 3-fold with \mathbb{Q} -factorial terminal singularities. According to Reid [24], for a Weil divisor D on X,

$$\chi(D) = \chi(\mathcal{O}_X) + \frac{1}{12}D(D - K_X)(2D - K_X) + \frac{1}{12}(D \cdot c_2) + \sum_Q c_Q(D),$$

where the last sum runs over Reid's basket of orbifold points. If the orbifold point Q is of type $\frac{1}{r_Q}(1, -1, b_Q)$ and $i_Q = i_Q(D)$ is the local index of divisor D at Q (i.e. $D \sim i_Q K_X$ around Q, $0 \leq i_Q < r$), then

$$c_Q(D) = -\frac{i_Q(r_Q^2 - 1)}{12r_Q} + \sum_{j=0}^{i_Q - 1} \frac{\overline{jb_Q}(r_Q - \overline{jb_Q})}{2r_Q}.$$

Here the symbol $\overline{\cdot}$ means the smallest residue mod r and $\sum_{j=0}^{-1} := 0$. We can write Reid's basket as $B_X = \{(b_Q, r_Q)\}_Q$. Note that we may assume $0 < b_Q \leq \frac{r_Q}{2}$. Recall that $i(X) = \text{l.c.m.}\{r_Q \in B_X\}$.

Let X be a minimal 3-fold with $K_X \equiv 0$. Note that for arbitrary nef and big Weil divisor H, Kawamata–Viehweg vanishing theorem [17, Theorem 1-2-5] implies

$$h^{i}(H) = h^{i}(K_{X} + (H - K_{X})) = 0$$

for all i > 0. For a nef and big Weil divisor L and a Weil divisor $T \equiv 0$, Reid's formula gives

$$h^{0}(mL+T) = \chi(\mathcal{O}_{X}) + \frac{m^{3}}{6}L^{3} + \frac{m}{12}(L \cdot c_{2}) + \sum_{Q} c_{Q}(mL+T).$$

We make some remarks on estimating this formula. Recall that by Miyaoka [20], c_2 is pseudo-effective and hence $(L \cdot c_2) \ge 0$ holds. Also Reid's formula or Kawamata [14, Theorem 2.4] gives

$$\chi(\mathcal{O}_X) = \sum_Q \frac{r_Q^2 - 1}{24r_Q}.$$
 (2.1)

We define

$$\lambda(L) := \frac{1}{6}L^3 + \frac{1}{12}(L \cdot c_2).$$

Note that $\lambda(L)$ is a numerical invariant of L. We can rewrite Reid's formula as following:

$$h^{0}(mL+T) = \chi(\mathcal{O}_{X}) + \frac{m^{3}-m}{6}L^{3} + m\lambda(L) + \sum_{Q} c_{Q}(mL+T).$$

We have the following lemma.

Lemma 2.2. $i(X)\lambda(L) \in \mathbb{Z}_{>0}$. In particular, $\lambda(L) \geq \frac{1}{i(X)}$.

Proof. For a singular point Q of type (b, r), note that if i runs over $\{0, 1, \dots, r-1\}$ then so does the local index of $L + iK_X$ at Q. Hence we have

$$\sum_{i=0}^{r-1} c_Q(L+iK_X)$$

$$= \sum_{i=0}^{r-1} \left(-\frac{i(r^2-1)}{12r} + \sum_{j=0}^{i-1} \frac{\overline{jb}(r-\overline{jb})}{2r} \right)$$

$$= -\frac{(r-1)(r^2-1)}{24} + \sum_{i=1}^{r-1} \sum_{j=0}^{i-1} \frac{\overline{jb}(r-\overline{jb})}{2r}$$

$$= -\frac{(r-1)(r^2-1)}{24} + \sum_{j=0}^{r-2} \sum_{i=j+1}^{r-1} \frac{\overline{jb}(r-\overline{jb})}{2r}$$

$$= -\frac{(r-1)(r^2-1)}{24} + \sum_{j=1}^{r-2} (r-1-j) \frac{\overline{jb}(r-\overline{jb})}{2r}$$

$$= -\frac{(r-1)(r^2-1)}{24} + \sum_{k=2}^{r-1} (k-1) \frac{\overline{kb}(r-\overline{kb})}{2r} \qquad (k=r-j)$$

 $\mathbf{6}$

On birational geometry of 3-folds with $K\equiv 0$

$$\begin{aligned} &= -\frac{(r-1)(r^2-1)}{24} + \frac{1}{2}\sum_{k=1}^{r-1}((r-1-k)+(k-1))\frac{\overline{kb}(r-\overline{kb})}{2r} \\ &= -\frac{(r-1)(r^2-1)}{24} + \frac{r-2}{2}\sum_{k=0}^{r-1}\frac{\overline{kb}(r-\overline{kb})}{2r} \\ &= -\frac{(r-1)(r^2-1)}{24} + \frac{r-2}{2}\sum_{j=0}^{r-1}\frac{j(r-j)}{2r} \\ &= -\frac{r^2-1}{24}. \end{aligned}$$

Hence by Reid's formula,

$$\sum_{i=0}^{i(X)-1} h^{0}(L+iK_{X})$$

$$= \sum_{i=0}^{i(X)-1} \left(\chi(\mathcal{O}_{X}) + \lambda(L) + \sum_{Q} c_{Q}(L+iK_{X}) \right)$$

$$= i(X)\chi(\mathcal{O}_{X}) + i(X)\lambda(L) + \sum_{Q} \left(-\frac{r_{Q}^{2}-1}{24} \cdot \frac{i(X)}{r_{Q}} \right)$$

$$= i(X)\lambda(L).$$

Hence $i(X)\lambda(L) \in \mathbb{Z}$. On the other hand, $\lambda(L) > 0$ since L is nef and big. \Box

We have the following lemma for intersection numbers.

Lemma 2.3. Let X be a normal projective 3-fold with Q-factorial terminal singularities. Recall that i(X) is the local index of X, i.e. the Cartier index of K_X . Then for Weil divisors D_1 , D_2 , and D_3 on X, $(i(X)D_1 \cdot D_2 \cdot D_3) \in \mathbb{Z}$. In particular, if L is a nef and big Weil divisor on X, then $L^3 \geq \frac{1}{i(X)}$.

Proof. Recall that by Kawamata [15, Corollary 5.2], $i(X)D_1$ is Cartier. Take a resolution of isolated singularities $\phi : W \to X$. We may write $K_W = \phi^*(K_X) + \Delta$ where Δ is an exceptional effective \mathbb{Q} -divisor over those isolated terminal singularities on X. Denote by D'_i the strict transform of D_i on W for i = 1, 2, 3. By intersection theory, we have

$$(i(X)D_1 \cdot D_2 \cdot D_3)_X = (\phi^*(i(X)D_1) \cdot \phi^*(D_2) \cdot D'_3)_W = (\phi^*(i(X)D_1) \cdot D'_2 \cdot D'_3)_W$$

is an integer.

2.3. Some facts about minimal 3-folds with $K \equiv 0$. We collect some facts about minimal 3-folds with $K \equiv 0$ proved by Kawamata [14] and Morrison [21].

Theorem 2.4 ([14, 21]). Let X be a minimal 3-fold with $K_X \equiv 0$. The following facts hold:

(1) $0 \leq \chi(\mathcal{O}_X) \leq 4;$

- (2) $\chi(\mathcal{O}_X) = 0$ if and only if X has Gorenstein singularities;
- (3) If q(X) > 0, then X is smooth;
- (4) If q(X) = 0 and $\chi(\mathcal{O}_X) \ge 2$, then $I(X) \in \{2, 3, 4, 6\}$;
- (5) If q(X) = 0 and $\chi(\mathcal{O}_X) = 1$, then

 $I(X) \in \{2, 3, 4, 5, 6, 8, 10, 12\};$

(6) If $I(X) \in \{5, 8, 10, 12\}$, then $\chi(\mathcal{O}_X) = 1$, $q(X) = h^2(\mathcal{O}_X) = 0$, i(X) = I(X), and the singular points can be described explicitly by Morrison [21, Proposition 3].

Proof. (1) is proved by Kawamata [14, Theorem 3.1]. (2) is a direct consequence of equality (2.1). (3) is proved by Kawamata [13] and [14] (see [21, Section 1]). (4) is proved by Morrison [21, Proposition 1, Proof of Theorem 1] and (5) is proved by Morrison [21, Proposition 3, Proof of Theorem 2]. (6) is a direct consequence of (2)-(5) and Morrison [21, Proposition 3]. \Box

3. Gorenstein case

Throughout this section, we assume that X is a minimal Gorenstein 3fold with $K_X \equiv 0$ and L is a nef and big Weil divisor on X. Note that L is a Cartier divisor since i(X) = 1. Recall that we have a *canonical model* $\mu: (X, L) \to (Z, H)$ such that Z is a 3-fold with canonical singularities and $\mu^*K_Z = K_X$, H is an ample Catier divisor with $L = \mu^*H$ (cf. [23, Lemma 0.2]).

Lemma 3.1 (cf. [23, Lemma 1.1]). Let D be a divisor on X. Then

(1) $(D \cdot L^2)^2 \ge (D^2 \cdot L)(L^3);$

- (2) $D \cdot L^2 \equiv \overline{D^2} \cdot L \mod 2;$
- (3) If $D \cdot L^2 = 1$ and $D^2 \cdot L \ge 0$, then $D^2 \cdot L = L^3 = 1$.

Proof. See the proof of [23, Lemma 1.1]. Note that $K_X \equiv 0$ is sufficient in the proof.

We prove Theorem 1.5 for the Gorenstein case. It is a direct generalization of Fukuda [12] and Oguiso–Peternell [23], and we follow their ideas.

Theorem 3.2. Let X be a minimal Gorenstein 3-fold with $K_X \equiv 0$, a nef and big Weil divisor L, and a Weil divisor $T \equiv 0$. Then $|K_X + mL + T|$ gives a birational map for all $m \geq 5$.

Proof. Note that L and T are Cartier divisors since i(X) = 1.

Case 1. dim $\Phi_{|L|}(X) \ge 1$.

Take a resolution $\pi: Y \to X$. Consider the linear system $|K_Y + m\pi^*L + \pi^*T|$. Note that

$$\dim \Phi_{|\pi^*L|}(Y) = \dim \Phi_{|L|}(X) \ge 1.$$

By [12, Key Lemma] with $R = \pi^* L$, $r_0 = 4$, and $r_1 = 1$, $|K_Y + m\pi^* L + \pi^* T|$ gives a birational map for all $m \ge 5$. So $|K_X + mL + T|$ gives a birational map for all $m \ge 5$.

Case 2. dim $\Phi_{|L|}(X) \le 0$.

In this case, since $h^0(L) > 0$ by Riemann–Roch formula, we have $h^0(L) = 1$. By Riemann–Roch formula again,

$$h^{0}(2L) = \frac{1}{6}(2^{3} - 2)L^{3} + 2h^{0}(L) = L^{3} + 2.$$

First, we assume that |2L| is composed with a pencil of surfaces. Set D := 2L and keep the same notation as in Subsection 2.1. Then we have

$$2\pi^*(L) \ge M \equiv aS \ge (h^0(2L) - 1)S = (L^3 + 1)S.$$

Thus we have $2L^3 \geq (L^3 + 1)(\pi^*(L)^2 \cdot S)$. This implies that $L^2 \cdot \pi_*S = \pi^*(L)^2 \cdot S = 1$ since $\pi^*(L)^2 \cdot S > 0$. On the other hand, $L \cdot (\pi_*S)^2 = \pi^*(L) \cdot \pi^*\pi_*S \cdot S \geq 0$. Hence by Lemma 3.1(3), $L \cdot (\pi_*S)^2 = L^3 = 1$. Hence $M \equiv (L^3 + 1)S = 2S$, in particular, |2L| is composed with a rational pencil (see Case (f_p) in subsection 2.1). Consider the canonical model (Z, H). Since $h^0(H) = h^0(L) = 1$ and $H^3 = L^3 = 1$, there exists an irreducible surface G such that $|H| = \{G\}$. Denote by G' the strict transform of G. Then we may write $2L \sim 2G' + 2E$ for some μ -exceptional divisor E. Note that $Mov|2L| = |2\pi_*S|$, hence $2\pi_*S \sim 2G' + F$ for some effective μ -exceptional divisor F. Note that $|2\pi_*S|$ is a rational pencil by construction, which means that, every element in $|2\pi_*S|$ can be written as the form $\pi_*S_1 + \pi_*S_2$ with some $S_1 \sim S_2 \sim S$. Hence $\pi_*S_1 + \pi_*S_2 = 2G' + F$ (not only linear equivalence but equality). Hence $\pi_*S_1 = G' + E'$ for some effective μ -exceptional divisor E' by the irreduciblity of G'. But this implies dim $|\pi_*S| = 0$, a contradiction.

Hence |2L| is not composed with a pencil of surface. Set D := 2L and keep the same notation as in Section 2.1. Then we have $2\pi^*(L) = |M| + F$ such that |M| is base point free. Consider a smooth element N in |M|. Note that $|K_X + mL + T|$ gives a birational map if so does the restriction $|K_Y + N + \pi^*((m-2)L + T)||_N$ by Lemma 4.2 and birationality principle (cf. [23, Lemma 1.3]). On the other hand, Kawamata–Viehweg vanishing theorem and adjunction formula give

$$|K_Y + N + \pi^*((m-2)L + T)||_N = |K_N + \pi^*((m-2)L + T)|_N|.$$

Reider's theorem (cf. [25]) implies that $|K_N + \pi^*((m-2)L+T)|_N|$ gives a birational map for $m \ge 5$ if $(\pi^*L)^2 \cdot N \ge 2$. Now we assume that $(\pi^*L)^2 \cdot N = 1$, then Lemma 3.1(3) implies that $L^3 = L^2 \cdot \pi_* N = L \cdot (\pi_*N)^2 = 1$. Consider the canonical model (Z, H). Since $h^0(H) = H^3 = 1$, and $L^2 \cdot \pi_* N = 1$, a similar argument implies dim $|\pi_*N| = 0$, a contradiction.

We completed the proof.

By Theorems 3.2 and 2.4(2)(3), to prove Theorem 1.5, we only need to consider polarized triples (X, L, T) with $\chi(\mathcal{O}_X) > 0$.

4. BIRATIONALITY CRITERION

In this section, we give a criterion for the birationality of polarized triples. The methods using in this section are mainly developed in Chen–Chen [5] and Chen [9], and latter modified in Chen–Jiang [10].

4.1. Main reduction. Firstly, we reduce the birationality problem on X to that on its smooth model Y.

Lemma 4.1 (cf. [9, Lemma 2.5]). Let W be a normal projective variety on which there is an integral Weil \mathbb{Q} -Cartier divisor D. Let $h: V \longrightarrow W$ be any resolution of singularities. Assume that E is an effective exceptional \mathbb{Q} -divisor on V with $h^*(D) + E$ a Cartier divisor on V. Then

$$h_*\mathcal{O}_V(h^*(D) + E) = \mathcal{O}_W(D)$$

where $\mathcal{O}_W(D)$ is the reflexive sheaf corresponding to the Weil divisor D.

Lemma 4.2. Let X be a normal projective variety with \mathbb{Q} -factorial terminal singularities, D be a Weil divisor on X and $\pi : Y \longrightarrow X$ be a resolution. Then $\Phi_{|K_X+D|}$ is birational if and only if so is $\Phi_{|K_Y+\lceil \pi^*(D)\rceil|}$.

Proof. Recall that

$$K_Y = \pi^*(K_X) + E_\pi$$

where E_{π} is an effective Q-Cartier Q-divisor since X has at worst terminal singularities. We have

$$K_Y + \lceil \pi^*(D) \rceil$$

= $\pi^*(K_X) + E_\pi + \pi^*(D) + E$
= $\pi^*(K_X + D) + E_\pi + E$

where $E_{\pi} + E$ is an effective Q-divisor on Y exceptional over X. Lemma 4.1 implies

$$\pi_*\mathcal{O}_Y(K_Y + \lceil \pi^*(D) \rceil) = \mathcal{O}_X(K_X + D).$$

Hence $\Phi_{|K_X+D|}$ is birational if and only if so is $\Phi_{|K_Y+\lceil \pi^*(D)\rceil\rceil}$.

4.2. Key theorem. Let (X, L, T) be a polarized triple. Take a Weil divisor L_0 such that $L_0 \equiv L$. Suppose that $h^0(m_0L_0) \ge 2$ for some integer $m_0 > 0$. Suppose that $m_1 \ge m_0$ is an integer with $h^0(m_1L_0) \ge 2$ and that $|m_1L_0|$ and $|m_0L_0|$ are not composed with the same pencil. Note that in application, if $|m_0L_0|$ is not composed with a pencil, we can just take $m_1 = m_0$.

Set $D := m_0 L_0$ and keep the same notation as in Subsection 2.1. We may modify the resolution π in Subsection 2.1 such that the movable part $|M_m|$ of $|\lfloor \pi^*(mL_0) \rfloor|$ is base point free for all $m_0 \leq m \leq m_1$. Set $\iota_m := \iota(mL_0)$ defined in Subsection 2.1. Recall that, for any integer m with $h^0(mL_0) > 1$,

$$\iota_m = \begin{cases} 1, & \text{if } |mL_0| \text{ is not composed with a pencil;} \\ h^0(mL_0) - 1, & \text{if } |mL_0| \text{ is composed with a pencil.} \end{cases}$$

Pick a general irreducible element S of $|M_{m_0}|$. We have

$$m_0 \pi^*(L_0) = \iota_{m_0} S + F_{m_0}$$

for some effective \mathbb{Q} -divisor F_{m_0} . In particular, we see that

$$\frac{m_0}{\iota_{m_0}}\pi^*(L_0) - S \sim_{\mathbb{Q}} \text{effective } \mathbb{Q}\text{-divisor.}$$

Define the real number

$$\mu_0 = \mu_0(|S|) := \inf\{t \in \mathbb{Q}^+ \mid t\pi^*(L_0) - S \sim_{\mathbb{Q}} \text{ effective } \mathbb{Q}\text{-divisor}\}.$$

Remark 4.3. Clearly, we have $0 < \mu_0 \leq \frac{m_0}{\iota_{m_0}} \leq m_0$. For all k such that $|kL_0|$ and $|m_0L_0|$ are composed with the same pencil, we have

$$k\pi^*(L_0) = \iota_k S + F_k$$

for some effective \mathbb{Q} -divisor F_k , and hence $\mu_0 \leq \frac{k}{\iota_k}$.

By the assumption on $|m_1L_0|$, we know that $|G| = |M_{m_1}|_S|$ is a base point free linear system on S and $h^0(S, G) \ge 2$. Denote by C a general irreducible element of |G|. Note that since $K_X \equiv 0$, K_Y is pseudo-effective and hence $g(C) \ge 1$. Since $m_1\pi^*(L_0) \ge M_{m_1}$, we have

$$m_1 \pi^*(L_0)|_S \equiv C + H$$

where H is an effective \mathbb{Q} -divisor on S.

We define two numbers which will be the key invariants accounting for the birationality of $\Phi_{|K_X+mL+T|}$. They are

$$\zeta := (\pi^*(L) \cdot C)_Y = (\pi^*(L_0) \cdot C)_Y = (\pi^*(L_0)|_S \cdot C)_S \text{ and}$$

$$\epsilon(m) := (m - \mu_0 - m_1)\zeta.$$

Note that ζ and $\epsilon(m)$ are birational invariants by projection formula. Hence we can modify π if necessary. Also note that $\zeta > 0$ since L is nef and big and C is free.

While studying the birationality of $\Phi_{|K_X+mL+T|}$, we will always check that the linear system $\Lambda_m := |K_Y + \lceil \pi^*(mL+T) \rceil|$ satisfies the following assumption for some integer m > 0.

Assumption 4.4. Keep the notation as above.

- (1) The linear system Λ_m distinguishes different general irreducible elements of $|M_{m_0}|$ (namely, $\Phi_{\Lambda_m}(S') \neq \Phi_{\Lambda_m}(S'')$ for two different general irreducible elements S', S'' of $|M_{m_0}|$).
- (2) The linear system $\Lambda_m|_S$ distinguishes different general irreducible elements of the linear system $|G| = |M_{m_1}|_S|$ on S.

The following is the key theorem in this section.

Theorem 4.5. Let (X, L, T) be a polarized triple. Keep the notation as above. Let m > 0 be an integer. If Assumption 4.4 is satisfied and $\epsilon(m) > 2$, then $\Phi_{|K_X+mL+T|}$ is birational onto its image.

Proof. By Lemma 4.2, we only need to prove the birationality of Φ_{Λ_m} . Since Assumption 4.4(1) is satisfied, the usual birationality principle reduces the birationality of Φ_{Λ_m} to that of $\Phi_{\Lambda_m}|_S$ for a general irreducible element Sof $|M_{m_0}|$. Similarly, due to Assumption 4.4(2), we only need to prove the birationality of $\Phi_{\Lambda_m}|_C$ for a general irreducible element C of |G|. Now we show how to restrict the linear system Λ_m to C.

Now assume $\epsilon(m) > 0$. We can find a sufficiently large integer n so that there exists a number $\mu_n \in \mathbb{Q}^+$ with $0 \le \mu_n - \mu_0 \le \frac{1}{n}$, $\lceil \epsilon(m,n) \rceil = \lceil \epsilon(m) \rceil$ where $\epsilon(m,n) := (m - \mu_n - m_1)\zeta$ and

$$\iota_n \pi^*(L_0) \sim_{\mathbb{Q}} S + E_n$$

for an effective Q-divisor E_n . In particular, $\epsilon(m,n) > 0$, and $\epsilon(m,n) > 2$ if $\epsilon(m) > 2$. Re-modify the resolution π in Subsection 2.1 so that E_n has simple normal crossing support.

For the given integer m > 0, we have

l

$$|K_Y + \lceil \pi^*(mL + T) - E_n \rceil| \leq |K_Y + \lceil \pi^*(mL + T) \rceil|.$$
(4.1)

Since $\epsilon(m, n) > 0$, the Q-divisor

 $\pi^*(mL+T) - E_n - S \equiv (m - \mu_n)\pi^*(L)$

is nef and big and thus

$$H^{1}(Y, K_{Y} + \lceil \pi^{*}(mL + T) - E_{n} \rceil - S) = 0$$

by Kawamata–Viehweg vanishing theorem. Hence we have surjective map

$$H^{0}(Y, K_{Y} + \lceil \pi^{*}(mL + T) - E_{n} \rceil) \longrightarrow H^{0}(S, K_{S} + L_{m,n})$$

$$(4.2)$$

where

$$L_{m,n} := \left(\left\lceil \pi^*(mL+T) - E_n \right\rceil - S \right) |_S \ge \left\lceil \mathcal{L}_{m,n} \right\rceil$$
(4.3)

and $\mathcal{L}_{m,n} := (\pi^*(mL+T) - E_n - S)|_S$. Moreover, we have

$$m_1 \pi^*(L_0)|_S \equiv C + H$$

for an effective \mathbb{Q} -divisor H on S by the setting. Thus the \mathbb{Q} -divisor

$$\mathcal{L}_{m,n} - H - C \equiv (m - \mu_n - m_1)\pi^*(L)|_S$$

is nef and big by $\epsilon(m,n)>0.~$ By Kawamata–Viehweg vanishing theorem again,

$$H^1(S, K_S + \lceil \mathcal{L}_{m,n} - H \rceil - C) = 0.$$

Therefore, we have surjective map

$$H^{0}(S, K_{S} + \lceil \mathcal{L}_{m,n} - H \rceil) \longrightarrow H^{0}(C, K_{C} + D_{m,n})$$
(4.4)

where

$$D_{m,n} := \left\lceil \mathcal{L}_{m,n} - H - C \right\rceil |_C \ge \left\lceil \mathcal{D}_{m,n} \right\rceil$$
(4.5)

and $\mathcal{D}_{m,n} := (\mathcal{L}_{m,n} - H - C)|_C$ with deg $\mathcal{D}_{m,n} = \epsilon(m, n)$.

Now by inequalities (4.1), (4.3), (4.5), and surjective maps (4.2), (4.4), to prove the birationality of $\Phi_{\Lambda_m}|_C$, it is sufficient to prove that $|K_C + \lceil \mathcal{D}_{m,n}\rceil|$ gives a birational map. Clearly this is the case whenever $\epsilon(m) > 2$, which in fact implies deg($\lceil \mathcal{D}_{m,n}\rceil$) $\geq \lceil \epsilon(m,n)\rceil \geq 3$ and $K_C + \lceil \mathcal{D}_{m,n}\rceil$ is very ample. We complete the proof.

Corollary 4.6. Keep the same notation as above. For any integer m > 0, set

$$\epsilon(m,0) := (m - \frac{m_0}{\iota_{m_0}} - m_1)\zeta.$$

If $\epsilon(m,0) > 0$, then

 $\Lambda_m|_S \succeq |K_S + L_m|$ where $L_m := (\lceil \pi^*(mL + T) - \frac{1}{\iota_{m_0}} F_{m_0} \rceil - S)|_S.$

Proof. Recall that

$$m_0 \pi^*(L_0) = \iota_{m_0} S + F_{m_0}.$$

First of all,

$$|K_Y + \lceil \pi^*(mL + T) - \frac{1}{\iota_{m_0}} F_{m_0} \rceil| \leq |K_Y + \lceil \pi^*(mL + T) \rceil|.$$

In fact, as long as $\epsilon(m,0) > 0$, the proof of Theorem 4.5 is valid except for the last paragraph. In explicit, sujective map (4.2) reads the following surjective map

$$H^{0}(Y, K_{Y} + \lceil \pi^{*}(mL + T) - \frac{1}{\iota_{m_{0}}}F_{m_{0}}\rceil) \longrightarrow H^{0}(S, K_{S} + L_{m})$$

where

$$L_m = (\lceil \pi^*(mL + T) - \frac{1}{\iota_{m_0}} F_{m_0} \rceil - S)|_S$$

Hence we have proved the statement.

4.3. Criterion. In order to apply Theorem 4.5, we need to verify Assumption 4.4 and $\epsilon(m) > 2$ in advance, for which one of the crucial steps is to estimate the lower bound of ζ .

Lemma 4.7. (1) If $\epsilon(m) > 1$, then $\zeta \geq \frac{2g(C) - 2 + \lceil \epsilon(m) \rceil}{m}$; (2) Moreover,

$$\zeta \ge \frac{2g(C) - 1}{\mu_0 + m_1 + 1};$$

- (3) If g(C) = 1, then $\zeta \ge 1$;
- (4) $i(X)\zeta \in \mathbb{Z}_{>0}$.

Proof. (1). Recall that since $K_X \equiv 0$, K_Y is pseudo-effective and hence $g(C) \geq 1$. In the proof of Theorem 4.5, if $\epsilon(m) > 1$ then $|K_C + \lceil \mathcal{D}_{m,n} \rceil|$ is base point free with

$$\deg(K_C + \lceil \mathcal{D}_{m,n} \rceil) \ge 2g(C) - 2 + \lceil \epsilon(m,n) \rceil = 2g(C) - 2 + \lceil \epsilon(m) \rceil.$$

Denote by \mathcal{N}_m the movable part of $|K_S + [\mathcal{L}_{m,n} - H]|$. Note that

$$H^{0}(\mathcal{O}_{Y}(\lfloor \pi^{*}(K_{X}+mL+T) \rfloor))$$

$$\cong H^{0}(\mathcal{O}_{X}(K_{X}+mL+T))$$

$$\cong H^{0}(\mathcal{O}_{Y}(K_{Y}+\lceil \pi^{*}(mL+T) \rceil)).$$

Denote by \mathcal{M}_m the movable part of $|[\pi^*(K_X + mL + T)]|$. Noting the relations (4.1-4.5) while applying [8, Lemma 2.7], we get

$$\pi^*(K_X + mL + T)|_S \ge \mathcal{M}_m|_S \ge \mathcal{N}_m$$

and $\mathcal{N}_m|_C \geq K_C + \lceil \mathcal{D}_{m,n} \rceil$ since the latter one is base point free. So we have

$$m\zeta = \pi^*(K_X + mL + T)|_S \cdot C \ge \mathcal{N}_m \cdot C \ge \deg(K_C + \lceil \mathcal{D}_{m,n} \rceil).$$

Hence

$$m\zeta \ge 2g(C) - 2 + \lceil \epsilon(m) \rceil$$

(2). Take $m' = \min\{m \mid \epsilon(m) > 1\}$, then (1) implies $\zeta \geq \frac{2g(C)}{m'}$. We may assume that $m' > \mu_0 + m_1 + 1$ otherwise $\zeta \geq \frac{2g(C)}{\mu_0 + m_1 + 1}$. Hence

$$\epsilon(m'-1) = (m'-1-\mu_0-m_1)\zeta$$

$$\geq (m'-1-\mu_0-m_1)\frac{2g(C)}{m'}$$

By the minimality of m', it follows that $\epsilon(m'-1) \leq 1$. Hence $m' \leq \frac{2g(C)}{2g(C)-1}(\mu_0 + m_1 + 1)$. Then

$$\zeta \geq \frac{2g(C)}{m'} \geq \frac{2g(C) - 1}{\mu_0 + m_1 + 1}.$$

(3). Recall that

$$K_Y = \pi^* K_X + E_\pi \equiv E_\pi,$$

13

where E_{π} is an effective Q-Cartier Q-divisor whose support contains all π exceptional divisors since X has at worst terminal singularities. If g(C) = 1,
then

$$0 = ((K_S + C) \cdot C)_S$$

= $(K_Y \cdot C)_Y + (S \cdot C)_Y + (C^2)_S$
= $(E_\pi \cdot C)_Y + (S \cdot C)_Y + (C^2)_S$.

Since C is free on a free surface S, $(C^2)_S$, $(S \cdot C)_Y$, and $(E_{\pi} \cdot C)_Y$ are nonnegative. Hence $(E_{\pi} \cdot C)_Y = 0$, which implies that $(E \cdot C)_Y = 0$ for any π -exceptional divisor E on Y since X has at worst terminal singularities. Hence $\zeta = (\pi^* L \cdot C)_Y$ is an integer. On the other hand, $\zeta > 0$. Hence $\zeta \ge 1$. (4). It follows from the fact that i(X)L is Cartier.

Proposition 4.8. Let m > 0 be an integer. Keep the same notation as in Subsection 4.2. Then

$$\zeta \ge \left\lceil i(X) \min\left\{1, \frac{3}{\mu_0 + m_1 + 1}\right\}\right\rceil / i(X).$$

Proof. If g(C) = 1, by Lemma 4.7(3), $\zeta \ge 1$; if $g(C) \ge 2$, by Lemma 4.7(2), $\zeta \ge \frac{3}{\mu_0 + m_1 + 1}$. Then by Lemma 4.7(4),

$$i(X)\zeta \ge \left\lceil i(X)\min\left\{1,\frac{3}{\mu_0+m_1+1}\right\}\right\rceil.$$

We complete the proof.

Define

 $\rho_0 := \min\{k \in \mathbb{Z}_{>0} \mid h^0(mL + T') > 0 \text{ for all } m \ge k \text{ and for all } T' \equiv 0\},$ where T' is assumed to be a Weil divisor.

To verify Assumption 4.4(1), we have the following proposition.

Proposition 4.9. Let (X, L, T) be a polarized triple. Keep the same notation as Subsection 4.2. Then Assumption 4.4(1) is satisfied for all $m \ge m_0 + \rho_0$.

Proof. We have

$$K_{Y} + \lceil \pi^{*}(mL + T) \rceil$$

$$\geq K_{Y} + \lceil \pi^{*}(mL + T - m_{0}L_{0}) + M_{m_{0}} \rceil$$

$$= (K_{Y} + \lceil \pi^{*}(mL + T - m_{0}L_{0}) \rceil) + M_{m_{0}}$$

$$\geq M_{m_{0}}.$$

The last inequality is due to

$$h^{0}(K_{Y} + \lceil \pi^{*}(mL + T - m_{0}L_{0}) \rceil)$$

= $h^{0}(K_{X} + mL + T - m_{0}L_{0}) > 0$

when $m \ge m_0 + \rho_0$ by Lemma 4.1 and the definition of ρ_0 .

When $f: Y \to \Gamma$ is of type (f_{np}) , [26, Lemma 2] implies that Λ_m can distinguish different general irreducible elements of $|M_{m_0}|$. When f is of type (f_p) , since the rational pencil $|M_{m_0}|$ (recall that q(X) = 0) can already separate different fibers of f, Λ_m can automatically distinguish different general irreducible elements of $|M_{m_0}|$.

14

It is slightly more complicated to verify Assumption 4.4(2).

Lemma 4.10 (cf. [9, Lemma 3.7]). Let R be a smooth projective surface with a base point free linear system |G|. Let Q be an arbitrary \mathbb{Q} -divisor on R. Denote by C a general irreducible element of |G|. Then the linear system $|K_R + \lceil Q \rceil + G|$ can distinguish different general irreducible elements of |G|under one of the following conditions:

- (1) |G| is not composed with an irrational pencil of curves and $K_R + \lceil Q \rceil$ is effective;
- (2) |G| is composed with an irrational pencil of curves, g(C) > 0, and Q is nef and big;

Proof. The statement corresponding to (1) follows from [26, Lemma 2] and the fact that a rational pencil can automatically distinguish its different general irreducible elements.

For situation (2), we pick a general irreducible element C of |G|. Then, since $h^0(R,G) \ge 2$, $G \equiv sC$ for some integer $s \ge 2$ and $C^2 = 0$. Denote by C_1 and C_2 two general irreducible elements of |G| such that $C_1 + C_2 \le |G|$. Then Kawamata–Viehweg vanishing theorem gives the surjective map

$$H^{0}(R, K_{R} + \lceil Q \rceil + G) \longrightarrow H^{0}(C, K_{C_{1}} + D_{1}) \oplus H^{0}(C_{2}, K_{C_{2}} + D_{2})$$

where $D_i := (\lceil Q \rceil + G - C_i)|_{C_i}$ with $\deg(D_i) \ge Q \cdot C_i > 0$ for i = 1, 2. If g(C) > 0, Riemann-Roch formula gives $h^0(C_i, K_{C_i} + D_i) > 0$ for i = 1, 2.

1,2. Thus $|K_R + \lceil Q \rceil + G|$ can distinguish C_1 and C_2 .

Proposition 4.11. Let (X, L, T) be a polarized triple. Keep the same notation as in Subsection 4.2. Then Assumption 4.4(2) is satisfied for all $m \ge m_0 + m_1 + \rho_0$.

Proof. Assuming $m \ge m_0 + m_1 + 1$, we have $\epsilon(m, 0) > 0$, and Corollary 4.6 implies that

$$\Lambda_m|_S \succeq |K_S + L_m|.$$

It suffices to prove that $|K_S + L_m|$ can distinguish different general irreducible elements of |G|.

For a suitable integer m > 0, we have

$$K_{S} + L_{m}$$

= $K_{Y}|_{S} + \left[\pi^{*}(mL + T) - \frac{1}{\iota_{m_{0}}}F_{m_{0}}\right]|_{S}$
 $\geq (K_{Y} + \left[\pi^{*}(mL + T - (m_{0} + m_{1})L_{0})\right])|_{S} + M_{m_{1}}|_{S}$

Thus, if |G| is not composed with an irrational pencil of curves, $|K_S + L_m|$ can distinguish different irreducible elements provided that

$$K_Y + \left[\pi^*(mL + T - (m_0 + m_1)L_0)\right]$$

is effective, which holds for $m - m_0 - m_1 \ge \rho_0$.

Assume |G| is composed with an irrational pencil of curves, we have

$$K_S + L_m$$

$$\geq K_S + \left\lceil (\pi^*(mL+T) - \frac{1}{\iota_{m_0}} F_{m_0} - S) |_S \right\rceil$$

$$\geq K_{S} + \left[(\pi^{*}(mL + T - m_{1}L_{0}) - \frac{1}{\iota_{m_{0}}}F_{m_{0}} - S)|_{S} \right] + M_{m_{1}}|_{S}.$$

We can take $Q = (\pi^*(mL + T - m_1L_0) - \frac{1}{\iota_{m_0}}F_{m_0} - S)|_S$ in Lemma 4.10 since $\epsilon(m, 0) > 0$. Since g(C) > 0, Lemma 4.10(2) implies that Assumption 4.4(2) is satisfied for $m \ge m_0 + m_1 + 1$.

We complete the proof.

In summary, we have a criterion for birationality.

Theorem 4.12. Let (X, L, T) be a polarized triple. Keep the same notation as in Subsection 4.2. Then $|K_X + mL + T|$ gives a birational map if

$$m > \max\left\{m_0 + m_1 + \rho_0 - 1, \mu_0 + m_1 + \frac{2}{\zeta}\right\}.$$

This theorem is optimal in some sense by the following examples.

Example 4.13 ([11, 14.3 Theorem]). Consider the general weighted hypersurface $X_{10} \subset \mathbb{P}(1, 1, 1, 2, 5)$ which is a smooth Calabi–Yau 3-fold. Take $L = \mathcal{O}_X(1)$ and $T = K_X \sim 0$. Then |5L| gives a birational map but |4L| does not.

On the other hand, we may take $m_0 = m_1 = \mu_0 = \rho_0 = 1$ and $\zeta \ge 1$. Hence Theorem 4.12 implies that |mL| gives a birational map for all $m \ge 5$.

Example 4.14 ([11, 14.3 Theorem]). Consider the general weighted hypersurface $X_8 \subset \mathbb{P}(1, 1, 1, 1, 4)$ which is a smooth Calabi–Yau 3-fold. Take $L = \mathcal{O}_X(1)$ and $T = K_X \sim 0$. Then |4L| gives a birational map but |3L| does not.

On the other hand, we may take $m_0 = m_1 = \mu_0 = \rho_0 = 1$. Note that $S \in |L|$ and $C \in |L|_S|$. Hence $\zeta = L^3 = 2$. Then Theorem 4.12 implies that |mL| gives a birational map for all $m \ge 4$.

Example 4.15 ([3, Theorem 4.5]). Consider the general weighted complete intersection $X_{2,6} \subset \mathbb{P}(1,1,1,1,1,3)$ which is a terminal Calabi–Yau 3-fold. Take $L = \mathcal{O}_X(1)$ and $T = K_X \sim 0$. Then |3L| gives a birational map but |2L| does not.

On the other hand, we may take $m_0 = m_1 = \mu_0 = \rho_0 = 1$. Note that $S \in |L|$ and $C \in |L|_S|$. Hence $\zeta = L^3 = 4$. Then Theorem 4.12 implies that |mL| gives a birational map for all $m \geq 3$.

5. BIRATIONALITY ON POLARIZED TRIPLES

In this section, we consider the birationality problem on polarized triples. By Theorem 4.12, we need to estimate m_0 , m_1 , ρ_0 , μ_0 , and ζ . First we will give estimation for ρ_0 by Reid's formula. Then we reduce the estimation of m_1 to the estimation of Hilbert function of L so that we can estimate both m_0 and m_1 by Reid's formula. Note that μ_0 can be estimated by Remark 4.3 and ζ can be estimated by Proposition 4.8 once we have estimation of m_0 and m_1 .

We always assume that (X, L, T) is a polarized triple with $\chi(\mathcal{O}_X) > 0$ in this section.

5.1. Estimation of ρ_0 . In this subsection, we estimate ρ_0 . Note that by Theorem 2.4(4)(5) and the fact that i(X)|I(X), we have

$$i(X) \in \{2, 3, 4, 5, 6, 8, 10, 12\}.$$

Since we need to estimate the Hilbert function of some divisor D, we need to estimate the singular part $c_Q(D)$ in Reid's formula. We list all the possible values for $c_Q(D)$ with all the possible singularities in Table A. The first row corresponds to the local index $i_Q(D)$ of D and the first column corresponds to the possible singularities of Q. In the estimation, we will always replace $c_Q(D)$ by the minimal value in the list corresponding to Q.

$i_Q(D)$	0	1	2	3	4	5		
(1, 2)	0	-1/8						
(1, 3)	0	-2/9	-1/9					
(1, 4)	0	-5/16	-1/4	-1/16				
(1,5)	0	-2/5	-2/5	-1/5	0			
(2,5)	0	-2/5	-1/5	-1/5	-1/5			
(1, 6)	0	-35/72	-5/9	-3/8	-1/9	5/72		
(1, 8)	0	-21/32	-7/8	-25/32	-1/2	-5/32		
(3,8)	0	-21/32	-3/8	-9/32	-1/2	-5/32		
(1, 10)	0	-33/40	-6/5	-49/40	-1	-5/8		
(3, 10)	0	-33/40	-3/5	-9/40	-3/5	-5/8		
(1, 12)	0	-143/144	-55/36	-27/16	-14/9	-175/144		
(5, 12)	0	-143/144	-19/36	-11/16	-5/9	-31/144		
$i_Q(D)$	6	7	8	9	10	11		
(1,8)	1/8	7/32						
(3,8)	-3/8	-9/32						
(1, 10)	-1/5	7/40	2/5	3/8				
(3, 10)	-1/5	-9/40	-3/5	-9/40				
(1, 12)	-3/4	-35/144	2/9	9/16	25/36	77/144		
(5, 12)	-3/4	-35/144	-7/9	-7/16	-11/36	-67/144		
Table A: table of $c_0(D)$								

Table A: table of $c_Q(D)$

Note that for a singular point Q of index $r \in \{2, 3, 4, 5\}$ and for any Weil divisor D,

$$c_Q(D) \geq -\frac{r^2-1}{12r}.$$

To estimate ρ_0 , we discuss on the value of i(X). Fix a Weil divisor $T' \equiv 0$. Recall that $L^3 \geq \frac{1}{i(X)}$ and $\lambda(L) \geq \frac{1}{i(X)}$.

If $i(X) \in \{2, 3, 4, 5\}$, by Reid's formula and equality (2.1),

$$h^{0}(mL + T') \geq \chi(\mathcal{O}_{X}) + \frac{m^{3} - m}{6}L^{3} + m\lambda(L) - \sum_{Q} \frac{r_{Q}^{2} - 1}{12r_{Q}}$$
$$\geq \frac{m^{3} + 5m}{6i(X)} - \chi(\mathcal{O}_{X}).$$

Recall that $\chi(\mathcal{O}_X) \leq 4$ (or $\chi(\mathcal{O}_X) = 1$ if i(X)=5), hence

$$\rho_0 \leq \begin{cases}
3, & \text{if } i(X) = 5; \\
4, & \text{if } i(X) \in \{2, 3\}; \\
5, & \text{if } i(X) = 4.
\end{cases}$$

If i(X) = 6, then we write $B_X = \{a \times (1,2), b \times (1,3), c \times (1,6)\}$. By equality (2.1),

$$24\chi(\mathcal{O}_X) = \frac{3}{2}a + \frac{8}{3}b + \frac{35}{6}c.$$

Hence $c < \frac{144}{35}\chi(\mathcal{O}_X)$. By Reid's formula and equality (2.1),

$$h^{0}(mL + T') \geq \chi(\mathcal{O}_{X}) + \frac{m^{3} - m}{6}L^{3} + m\lambda(L) - \frac{1}{8}a - \frac{2}{9}b - \frac{5}{9}c$$
$$= \frac{m^{3} - m}{6}L^{3} + m\lambda(L) - \chi(\mathcal{O}_{X}) - \frac{5}{72}c$$
$$> \frac{m^{3} + 5m}{36} - \frac{9}{7}\chi(\mathcal{O}_{X}).$$

Recall that $\chi(\mathcal{O}_X) \leq 4$, hence $\rho_0 \leq 6$.

If i(X) = 8, by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{3 \times (1,2), (1,4), (b_1,8), (b_2,8)\}$ for $b_1, b_2 = 1$ or 3. By Reid's formula,

$$h^{0}(mL+T') \ge 1 + \frac{m^{3}-m}{6}L^{3} + m\lambda(L) - 3 \times \frac{1}{8} - \frac{5}{16} - 2 \times \frac{7}{8}$$
$$= \frac{m^{3}+5m}{48} - \frac{23}{16}.$$

Hence $\rho_0 \leq 4$.

If i(X) = 10, by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{3 \times (1,2), (b_1,5), (b_2,5), (c,10)\}$ for $b_1, b_2 = 1$ or 2, c = 1 or 3. By Reid's formula,

$$h^{0}(mL+T') \ge 1 + \frac{m^{3}-m}{6}L^{3} + m\lambda(L) - 3 \times \frac{1}{8} - 2 \times \frac{2}{5} - \frac{49}{40}$$
$$= \frac{m^{3}+5m}{60} - \frac{7}{5}.$$

Hence $\rho_0 \leq 5$.

If i(X) = 12, recall that by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{2 \times (1,2), 2 \times (1,3), (1,4), (b,12)\}$ for b = 1 or 5. By Reid's formula,

$$h^{0}(mL+T') \ge 1 + \frac{m^{3}-m}{6}L^{3} + m\lambda(L) - 2 \times \frac{1}{8} - 2 \times \frac{2}{9} - \frac{5}{16} - \frac{27}{16}$$
$$= \frac{m^{3}+5m}{72} - \frac{61}{36}.$$

Hence $\rho_0 \leq 5$.

In summary, we proved the following proposition.

Proposition 5.1. We have the following estimation for ρ_0 :

$$\rho_0 \leq \begin{cases}
3, & \text{if } i(X) = 5; \\
4, & \text{if } i(X) \in \{2, 3, 8\}; \\
5, & \text{if } i(X) \in \{4, 10, 12\}; \\
6, & \text{if } i(X) = 6.
\end{cases}$$

5.2. Estimation of m_1 . We give a criterion for a linear system not composing with a pencil of surfaces by looking at its Hilbert function.

Proposition 5.2. Let L_0 be a nef and big Weil divisor. If

$$h^0(mL_0) > i(X)L_0^3m + 1$$

for some integer m, then $|mL_0|$ is not composed with a pencil of surfaces.

Proof. Assume that $|mL_0|$ is composed with a pencil of surfaces. Set $D := mL_0$ and keep the same notation as in Subsection 2.1. Then we have $m\pi^*(L_0) \ge M \equiv (h^0(mL_0) - 1)S$. Note that by Lemma 2.3, $i(X)\pi^*(L_0)^2 \cdot S$ is an integer. On the other hand, $\pi^*(L_0)^2 \cdot S$ is positive since $\pi^*(L_0)|_S$ is nef and big on S. Hence $\pi^*(L_0)^2 \cdot S \ge \frac{1}{i(X)}$. Thus we have $mL_0^3 \ge (h^0(mL_0) - 1)(\pi^*(L_0)^2 \cdot S) \ge \frac{1}{i(X)}(h^0(mL_0) - 1)$, a contradiction. \Box

5.3. Proof of Theorems 1.5 and 1.6. In this subsection, we prove Theorems 1.5 and 1.6 by estimating m_0 and m_1 .

Proof of Theorem 1.5. To prove Theorem 1.5, by Section 3, we only need to consider polarized triples (X, L, T) with $\chi(\mathcal{O}_X) > 0$. We discuss on the value of i(X). Recall that

$$i(X) \in \{2, 3, 4, 5, 6, 8, 10, 12\}.$$

Recall again that $L^3 \geq \frac{1}{i(X)}$ and $\lambda(L) \geq \frac{1}{i(X)}$. The main problem is to estimate $\sum_Q c_Q$. In the proof, we often use the fact that if Q is a cyclic singular point and D is a Weil divisor with local index $i_Q(D) = 0$, then $c_Q(D) = 0$.

Case 1. i(X) = 2 or 3.

In this case, by Reid's formula,

$$h^{0}(i(X)L) \geq \chi(\mathcal{O}_{X}) + \frac{i(X)^{3}}{6}L^{3} > 1,$$

$$h^{0}(2i(X)L) \geq \chi(\mathcal{O}_{X}) + \frac{8i(X)^{3}}{6}L^{3} > 2i(X)^{2}L^{3} + 1$$

Hence we can take $L_0 = L$, $m_0 = i(X)$, and $m_1 = 2i(X)$. Then we have $\mu_0 \leq i(X)$ by Remark 4.3. By Proposition 4.8, $\zeta \geq \frac{1}{i(X)}$. By Proposition 5.1, $\rho_0 \leq 4$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 5i(X) + 1$.

Case 2. i(X) = 4.

In this case, by the proof of Lemma 2.2,

$$\sum_{i=0}^{3} h^0(5L + iK_X) = 4\lambda(5L).$$

Hence there exists i_0 such that

$$h^0(5L + i_0 K_X) \ge \lambda(5L).$$

Take $L_0 = L + i_0 K_X$. Then

$$h^{0}(5L_{0}) = h^{0}(5L + i_{0}K_{X} + 4i_{0}K_{X})$$

= $h^{0}(5L + i_{0}K_{X})$
 $\geq \lambda(5L)$
= $20L_{0}^{3} + 5\lambda(L)$
 $> 5i(X)L_{0}^{3} + 1.$

On the other hand,

$$h^{0}(4L_{0}) = \chi(\mathcal{O}_{X}) + \frac{4^{3} - 4}{6}L_{0}^{3} + 4\lambda(L) > 4.$$

Hence $h^0(4L_0) \ge 5$ and $|5L_0|$ is not composed with a pencil. Take $m_0 = 4$. By Proposition 5.1, $\rho_0 \le 5$.

If $|4L_0|$ is composed with a pencil, then we have $\mu_0 \leq 1$ by Remark 4.3 and we can take $m_1 = 5$. By Proposition 4.8, $\zeta \geq \frac{1}{2}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 14$.

If $|4L_0|$ is not composed with a pencil, then we have $\mu_0 \leq 4$ and we can take $m_1 = 4$. By Proposition 4.8, $\zeta \geq \frac{1}{2}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 13$.

Case 3. i(X) = 6.

In this case, recall that $1 \leq \chi(\mathcal{O}_X) \leq 4$ and we write $B_X = \{a \times (1,2), b \times (1,3), c \times (1,6)\}$. By equality (2.1),

$$24\chi(\mathcal{O}_X) = \frac{3}{2}a + \frac{8}{3}b + \frac{35}{6}c.$$

If $\chi(\mathcal{O}_X) = 1$, there is only one solution satisfying i(X) = 6, which is $B_X = \{5 \times (1,2), 4 \times (1,3), (1,6)\}$. We can take $L_0 = L + i_0 K_X$ for some i_0 such that the local index of L_0 at the point (1,6) is 0. Note that

$$\sum_{Q} c_Q(kL_0) \ge \begin{cases} -5 \times \frac{1}{8}, & \text{if } k = 3; \\ -4 \times \frac{2}{9}, & \text{if } k = 4; \\ -5 \times \frac{1}{8} - 4 \times \frac{2}{9}, & \text{if } k = 7. \end{cases}$$

By Reid's formula,

$$h^{0}(3L_{0}) = 1 + \frac{3^{3} - 3}{6}L_{0}^{3} + 3\lambda(L) + \sum_{Q} c_{Q}(3L_{0}) > 1,$$

$$h^{0}(4L_{0}) = 1 + \frac{4^{3} - 4}{6}L_{0}^{3} + 4\lambda(L) + \sum_{Q} c_{Q}(4L_{0}) > 2,$$

$$h^{0}(7L_{0}) = 1 + \frac{7^{3} - 7}{6}L_{0}^{3} + 7\lambda(L) + \sum_{Q} c_{Q}(7L_{0}) > 7i(X)L_{0}^{3} + 1.$$

Hence $h^0(3L_0) \ge 2$ and $|7L_0|$ is not composed with a pencil. Take $m_0 = 3$. By Proposition 5.1, $\rho_0 \le 6$.

If $|4L_0|$ and $|3L_0|$ are composed with the same pencil, then we have $\mu_0 \leq 2$ by Remark 4.3. Take $m_1 = 7$. By Proposition 4.8, $\zeta \geq \frac{1}{3}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 16$.

If $|4L_0|$ and $|3L_0|$ are not composed with the same pencil, then we can take $m_1 = 4$ and we have $\mu_0 \leq 3$. By Proposition 4.8, $\zeta \geq \frac{1}{2}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 13$.

Now we assume that $\chi(\mathcal{O}_X) \geq 2$. Note that for any Weil divisor D and singular point Q of index r,

$$c_Q(3D) + c_Q(3D + 3K_X) = \begin{cases} -\frac{1}{8}, & \text{if } r = 2; \\ 0, & \text{if } r = 3; \\ -\frac{3}{8}, & \text{if } r = 6. \end{cases}$$

Hence

$$h^{0}(3L) + h^{0}(3L + 3K_{X})$$

= $2\chi(\mathcal{O}_{X}) + 2\lambda(3L) + \sum_{Q}(c_{Q}(3L) + c_{Q}(3L + 3K_{X}))$
= $2\chi(\mathcal{O}_{X}) + 2\lambda(3L) - \frac{1}{8}a - \frac{3}{8}c$
 $\geq 2\lambda(3L).$

Therefore there exists a Weil divisor $L_0 \equiv L$ such that

$$h^0(3L_0) \ge \lambda(3L) = 4L^3 + 3\lambda(L) > 1.$$

On the other hand,

$$h^{0}(6L_{0}) \ge \chi(\mathcal{O}_{X}) + \frac{6^{3}}{6}L_{0}^{3} > 6i(X)L_{0}^{3} + 1.$$

Hence $|6L_0|$ is not composed with a pencil.

Hence we can take $m_0 = 3$ and $m_1 = 6$. Then we have $\mu_0 \leq 3$ by Remark 4.3. By Proposition 4.8, $\zeta \geq \frac{1}{3}$. By Proposition 5.1, $\rho_0 \leq 6$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 16$.

Case 4. i(X) = 5.

In this case, recall that by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{(b_1, 5), (b_2, 5), (b_3, 5), (b_4, 5), (b_5, 5)\}$ for $b_i = 1$ or 2 for $1 \leq i \leq 5$. We can take $L_0 = L + i_0 K_X$ for some i_0 such that the local index of L_0 at the point $(b_1, 5)$ is 0. Note that $\sum_Q c_Q(kL_0) \geq -4 \times \frac{2}{5}$ for all k. By Reid's formula,

$$h^{0}(4L_{0}) = 1 + \frac{4^{3} - 4}{6}L_{0}^{3} + 4\lambda(L) + \sum_{Q} c_{Q}(4L_{0}) > 2,$$

$$h^{0}(5L_{0}) = 1 + \frac{5^{3} - 5}{6}L_{0}^{3} + 5\lambda(L) = 6,$$

$$h^{0}(6L_{0}) = 1 + \frac{6^{3} - 6}{6}L_{0}^{3} + 6\lambda(L) + \sum_{Q} c_{Q}(6L_{0}) > 6i(X)L_{0}^{3} + 1$$

Hence $h^0(4L_0) \ge 3$ and $|6L_0|$ is not composed with a pencil. Take $m_0 = 4$. By Proposition 5.1, $\rho_0 \le 3$.

If $|5L_0|$ and $|4L_0|$ are composed with the same pencil, then we have $\mu_0 \leq 1$ by Remark 4.3. Take $m_1 = 6$. By Proposition 4.8, $\zeta \geq \frac{2}{5}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 13$.

If $|4L_0|$ is composed with a pencil, and $|5L_0|$ and $|4L_0|$ are not composed with the same pencil, then we can take $m_1 = 5$ and we have $\mu_0 \leq 2$. By Proposition 4.8, $\zeta \geq \frac{2}{5}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 13$.

If $|4L_0|$ is not composed with a pencil, then we can take $m_1 = 4$ and we have $\mu_0 \leq 4$. By Proposition 4.8, $\zeta \geq \frac{2}{5}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 14$.

Case 5. i(X) = 8.

In this case, recall that by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{3 \times (1,2), (1,4), (b_1,8), (b_2,8)\}$ for $b_1, b_2 = 1$ or 3. We can take $L_0 = L + i_0 K_X$ for some i_0 such that the local index of L_0 at the point $(b_1, 8)$ is 0. Note that

$$\sum_{Q} c_Q(kL_0) \ge \begin{cases} -\frac{7}{8}, & \text{if } k = 4; \\ -\frac{5}{16} - \frac{7}{8}, & \text{if } k = 6. \end{cases}$$

By Reid's formula,

$$h^{0}(4L_{0}) = 1 + \frac{4^{3} - 4}{6}L_{0}^{3} + 4\lambda(L) + \sum_{Q} c_{Q}(4L_{0}) > 1,$$

$$h^{0}(6L_{0}) = 1 + \frac{6^{3} - 6}{6}L_{0}^{3} + 6\lambda(L) + \sum_{Q} c_{Q}(6L_{0}) > 4,$$

$$h^{0}(8L_{0}) = 1 + \frac{8^{3} - 8}{6}L_{0}^{3} + 8\lambda(L) > 8i(X)L_{0}^{3} + 1.$$

Hence $h^0(4L_0) \ge 2$ and $|8L_0|$ is not composed with a pencil. Take $m_0 = 4$. By Proposition 5.1, $\rho_0 \le 4$.

If $|6L_0|$ and $|4L_0|$ are composed with the same pencil, then we have $\mu_0 \leq \frac{6}{4}$ by Remark 4.3. Take $m_1 = 8$. By Proposition 4.8, $\zeta \geq \frac{3}{8}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 16$.

If $|6L_0|$ and $|4L_0|$ are not composed with the same pencil, then we can take $m_1 = 6$ and we have $\mu_0 \leq 4$. By Proposition 4.8, $\zeta \geq \frac{3}{8}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 16$.

Case 6. i(X) = 10.

In this case, recall that by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{3 \times (1,2), (b_1,5), (b_2,5), (c,10)\}$ for $b_1, b_2 = 1$ or 2, c = 1 or 3. We can take $L_0 = L + i_0 K_X$ for some i_0 such that the local index of L_0 at the point (c, 10) is 0. Note that for an even integer k, $\sum_Q c_Q(kL_0) \geq -2 \times \frac{2}{5}$. By Reid's formula,

$$h^{0}(4L_{0}) = 1 + \frac{4^{3} - 4}{6}L_{0}^{3} + 4\lambda(L) + \sum_{Q} c_{Q}(4L_{0}) > 1;$$

$$h^{0}(6L_{0}) = 1 + \frac{6^{3} - 6}{6}L_{0}^{3} + 6\lambda(L) + \sum_{Q} c_{Q}(6L_{0}) > 4;$$

On birational geometry of 3-folds with $K \equiv 0$

$$h^{0}(8L_{0}) = 1 + \frac{8^{3} - 8}{6}L_{0}^{3} + 8\lambda(L) + \sum_{Q} c_{Q}(8L_{0}) > 8i(X)L_{0}^{3} + 1.$$

Hence $h^0(4L_0) \ge 2$ and $|8L_0|$ is not composed with a pencil. Take $m_0 = 4$. By Proposition 5.1, $\rho_0 \le 5$.

If $|6L_0|$ and $|4L_0|$ are composed with the same pencil, then we have $\mu_0 \leq \frac{3}{2}$ by Remark 4.3. Take $m_1 = 8$. By Proposition 4.8, $\zeta \geq \frac{3}{10}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 17$.

If $|6L_0|$ and $|4L_0|$ are not composed with the same pencil, then we can take $m_1 = 6$ and we have $\mu_0 \leq 4$. By Proposition 4.8, $\zeta \geq \frac{3}{10}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 17$.

Case 7. i(X) = 12.

In this case, recall that by Morrison [21, Proposition 3], we have i(X) = I(X), $\chi(\mathcal{O}_X) = 1$, and $B_X = \{2 \times (1,2), 2 \times (1,3), (1,4), (b,12)\}$ for b = 1 or 5. We can take $L_0 = L + i_0 K_X$ for some i_0 such that the local index of L_0 at the point (b, 12) is 0. Note that

$$\sum_{Q} c_Q(kL_0) \ge \begin{cases} -2 \times \frac{1}{8} - \frac{5}{16}, & \text{if } k = 3, 9; \\ -\frac{5}{16}, & \text{if } k = 6. \end{cases}$$

By Reid's formula,

$$h^{0}(3L_{0}) = 1 + \frac{3^{3} - 3}{6}L_{0}^{3} + 3\lambda(L) + \sum_{Q} c_{Q}(3L_{0}) > 1,$$

$$h^{0}(6L_{0}) = 1 + \frac{6^{3} - 6}{6}L_{0}^{3} + 6\lambda(L) + \sum_{Q} c_{Q}(6L_{0}) > 4,$$

$$h^{0}(9L_{0}) = 1 + \frac{9^{3} - 9}{6}L_{0}^{3} + 9\lambda(L) + \sum_{Q} c_{Q}(9L_{0}) > 9i(X)L_{0}^{3} + 1$$

Hence $h^0(3L_0) \ge 2$ and $|9L_0|$ is not composed with a pencil. Take $m_0 = 3$. By Proposition 5.1, $\rho_0 \le 5$.

If $|6L_0|$ and $|3L_0|$ are composed with the same pencil, then we have $\mu_0 \leq \frac{3}{2}$ by Remark 4.3. Take $m_1 = 9$. By Proposition 4.8, $\zeta \geq \frac{1}{3}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 17$.

If $|6L_0|$ and $|3L_0|$ are not composed with the same pencil, then we have $\mu_0 \leq 3$ and we can take $m_1 = 6$. By Proposition 4.8, $\zeta \geq \frac{1}{3}$. By Theorem 4.12, $|K_X + mL + T|$ gives a birational map for $m \geq 16$.

Proof of Theorem 1.6. Since L has no stable components, take a sufficient divisible k such that $kL \sim M$ is movable and effective and take a sufficient small rational number $\delta > 0$ such that $(X, \delta M)$ is terminal. Run a $(K_X + \delta M)$ -MMP with scaling of an ample divisor, it terminates on X' by Kawamata [16]. Since $i(X)l(K_X + \delta M) \sim i(X)l\delta M$ is movable for l sufficient divisible, this MMP $\psi : X \to X'$ does not contract any divisors, hence isomorphic in codimension one. Hence $(X', \delta \psi_* M)$ is terminal and so is X'. Hence X' is a minimal 3-fold with $K_{X'} = \psi_* K_X \equiv 0$ and $\psi_* L$ is a nef and big Weil divisor by MMP. Note that $|K_X + mL + T|$ gives a birational

map if and only if so does $|K_{X'} + m\psi_*L + \psi_*T|$, hence Theorem 1.6 follows from Theorem 1.5.

References

- Ando, T.: Pluricanonical systems of algebraic varieties of general type of dimension ≤ 5. In: Algebraic geometry (Sendai, 1985), Adv. Stud. Pure Math., 10, pp. 1–10. North-Holland, Amsterdam (1987)
- [2] Bombieri, E.: Canonical models of surfaces of general type. Publ. Math. IHES 42, 171–219 (1973)
- Chen, J-J., Chen, J.A., Chen, M.: On quasismooth weighted complete intersections. J. Algebraic Geom. 20, 239–262 (2011)
- [4] Chen, J.A., Chen, M.: Explicit birational geometry of threefolds of general type, I. Ann. Sci. Éc. Norm. Supér (4) 43, 365–394 (2010)
- [5] Chen, J.A., Chen, M.: Explicit birational geometry of threefolds of general type, II. J. Differential Geom. 86, 237–271 (2010)
- [6] Chen, J.A., Chen, M.: Explicit birational geometry of threefolds of general type, III. Compositio Math., 151, 1041–1082 (2015)
- [7] Chen, M.: On pluricanonical maps for threefolds of general type. J. Math. Soc. Japan, 50, 615–621 (1998)
- [8] Chen, M.: Canonical stability in terms of singularity index for algebraic threefolds. Math. Proc. Cambridge Philos. Soc. 13, 241–264 (2001)
- [9] Chen, M.: On anti-pluricanonical systems of Q-Fano 3-folds. Sci. China Math. 54, 1547–1560 (2011)
- [10] Chen, M., Jiang, C.: On the anti-canonical geometry of Q-Fano threefolds. Preprint, arXiv:1408.6349, to appear in J. Differential Geom.
- [11] Iano-Fletcher, A.R.: Working with weighted complete intersections. In: Explicit birational geometry of 3-folds, London Math. Soc. Lecture Note Ser., 281, pp. 101–173. Cambridge Univ. Press, Cambridge (2000)
- [12] Fukuda, S.: A note on Ando's paper "Pluricanonical systems of algebraic varieties of general type of dimension ≤ 5 ". Tokyo J. of Math. 14, 479–487 (1991)
- [13] Kawamata, Y.: Minimal models and the Kodaira dimension of algebraic fiber spaces.
 J. Reine Angew. Math. 363, 1–46 (1985)
- [14] Kawamata, Y.: On the plurigenera of minimal algebraic 3-folds with $K \equiv 0$. Math. Ann. 275, 539–546 (1986)
- [15] Kawamata, Y.: Crepant blowing-up of 3-dimensional canonical singularities and its application to degeneration of surfaces. Ann. of Math. 127, 93–163 (1988)
- [16] Kawamata, Y.: Termination of log-flips for algebraic 3-folds. Internat. J. Math. 3, 653–659 (1992)
- [17] Kawamata, Y., Matsuda, K., Matsuki, K.: Introduction to the minimal model problem. In: Algebraic geometry (Sendai, 1985), Adv. Stud. Pure Math., 10, pp. 283–360. North-Holland, Amsterdam (1987)
- [18] Kollár, J., Mori, S.: Birational geometry of algebraic varieties, Cambridge tracts in mathematics, vol. 134. Cambridge University Press (1998)
- [19] Matsuki, K.: On the value n which makes the n-ple canonical map birational for a 3-fold of general type. J. Math. Soc. Japan 38, 339–359 (1986)
- [20] Miyaoka, Y.: The Chern classes and Kodaira dimension of a minimal variety. In: Algebraic geometry (Sendai, 1985), Adv. Stud. Pure Math., 10, pp. 449–476. North-Holland, Amsterdam (1987)
- [21] Morrison, D.: A remark on Kawamata's paper "On the plurigenera of minimal algebraic 3-folds with $K \equiv 0$ ". Math. Ann. 275, 547–553 (1986)
- [22] Oguiso, K.: On polarized Calabi–Yau 3-folds. J. Fac. Sci. Univ. Tokyo 38, 395–429 (1991)
- [23] Oguiso, K., Peternell, T.: On polarized canonical Calabi–Yau threefolds. Math. Ann. 301, 237–248 (1995)

- [24] Reid, M.: Young person's guide to canonical singularities. In: Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Sympos. Pure Math., 46, Part 1, pp. 345–414. Amer. Math. Soc., Providence, RI (1987)
- [25] Reider, I.: Vector bundles of rank 2 and linear systems on algebraic surfaces. Ann. of Math. 127, 309–316 (1988)
- [26] Tankeev, S.G.: On n-dimensional canonically polarized varieties and varieties of fundamental type. Izv. A. N. SSSR, Ser. Math. 35, 31–44 (1971)

Graduate School of Mathematical Sciences, the University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8914, Japan.

E-mail address: cjiang@ms.u-tokyo.ac.jp