RANK TWO NILPOTENT CO-HIGGS SHEAVES ON COMPLEX SURFACES

M. CORRÊA

Dedicated to Jose Seade, for his 60th birthday.

ABSTRACT. Let (\mathcal{E}, ϕ) be a rank two co-Higgs vector bundles on a Kähler compact surface X with $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ nilpotent. If (\mathcal{E}, ϕ) is semistable, then one of the following holds up to finite étale cover:

- i) X is uniruled.
- *ii*) X is a torus and (\mathcal{E}, ϕ) is strictly semi-stable.
- *iii*) X is a properly elliptic surface and (\mathcal{E}, ϕ) is strictly semi-stable.

1. INTRODUCTION.

A generalised complex structure on a real manifold X of dimension 2n, as defined by Hitchin [10], is a rank-2n isotropic subbundle $E^{0,1} \subset (T_X \oplus T_X^*)^{\mathbb{C}}$ such that

- i) $E^{0,1} \oplus \overline{E^{0,1}} = (T_X \oplus T_X^*)^{\mathbb{C}}$
- ii) $C^{\infty}(E^{0,1})$ is closed under the Courant bracket.

On a manifold with a generalized complex structure M. Gualtieri in [7] defined the notion of a generalized holomorphic bundle. More precisely, a generalized holomorphic bundle on a generalized complex manifold, is a vector bundle \mathcal{E} with a differential operator $\overline{D}: C^{\infty}(\mathcal{E}) \longrightarrow C^{\infty}(\mathcal{E} \otimes E^{0,1})$ such that for all smooth function f and all section $s \in C^{\infty}(\mathcal{E})$ the following holds

- i) $\overline{D}(fs) = \overline{\partial}(fs) + f\overline{D}(s)$
- ii) $\overline{D}^2 = 0.$

In the case of an ordinary complex structure and $\overline{D} = \overline{\partial} + \phi$, for operators

$$\overline{\partial}: C^{\infty}(\mathcal{E}) \longrightarrow C^{\infty}(\mathcal{E} \otimes \overline{T_X^*})$$

and

$$\phi: C^{\infty}(\mathcal{E}) \longrightarrow C^{\infty}(\mathcal{E} \otimes T_X),$$

the vanishing $\overline{D}^2 = 0$ means that $\overline{\partial}^2 = 0$, $\overline{\partial}\phi = 0$ and $\phi \wedge \phi = 0$. By a classical result of Malgrange the condition $\overline{\partial}^2 = 0$ implies that \mathcal{E} is a holomorphic vector bundle. On the other hand, $\overline{\partial}\phi = 0$ implies that ϕ is a holomorphic global section

$$\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$$

which satisfies an integrability condition $\phi \wedge \phi = 0$. A co-Higgs sheaf on a complex manifold X is a sheaf \mathcal{E} together with a section $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ (called a *Higgs fields*) for which $\phi \wedge \phi = 0$. General properties of co-Higgs bundles were studied in [9, 20]. There is a motivation in physics for studying co-Higgs bundles, see [8], [12] and [26].

There are no stable co-Higgs bundles with nonzero Higgs field on curves C of genus g > 1. (When g = 1, a co-Higgs bundle is the same thing as a Higgs bundle

Date: January 27, 2022.

²⁰¹⁰ Mathematics Subject Classification. Primary 14D20- 53D18, 14D06.

Key words and phrases. Higgs bundles-stable vector bundles.

M. CORRÊA

in the usual sense.) In fact, contracting with a holomorphic differential gives a non-trivial endomorphism of \mathcal{E} commuting with ϕ which is impossible in the stable case [9, 21]. S. Rayan showed in [19] the non-existence of stable co-Higgs bundles with non trivial Higgs field on K3 and general-type surfaces. In this note we prove the following result.

Theorem 1.1. Let (\mathcal{E}, ϕ) be a rank two co-Higgs vector bundles on a Kähler compact surface X with $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ nilpotent. If (\mathcal{E}, ϕ) is semi-stable, then one of the following holds up to finite étale cover:

- i) X is uniruled.
- ii) X is a torus and (\mathcal{E}, ϕ) is strictly semi-stable.
- iii) X is a properly elliptic surface and (\mathcal{E}, ϕ) is strictly semi-stable.

It follows direct of proof of Theorem 1.1 that the we can consider a more general classes of singular projective surfaces .

Theorem 1.2. Let (\mathcal{E}, ϕ) be a rank two co-Higgs torsion-free sheaf on a normal projective surface X with $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ nilpotent. If (\mathcal{E}, ϕ) is stable, then X is uniruled.

Finally, in this work we consider a relation between co-Higgs bundles and Poisson geometry on \mathbb{P}^1 -bundles. In [23] Polishchuk associated to each rank-2 co-Higgs bundle (\mathcal{E}, ϕ) a Poisson structure on its projectivized bundle $\mathbb{P}(\mathcal{E})$. This relation was explained by Rayan in [20] as follows:

let $Y := \mathbb{P}(\mathcal{E})$ and consider the natural projection $\pi : Y \to X$. The exact sequence

$$0 \longrightarrow T_{X|Y} \longrightarrow T_Y \longrightarrow \pi^* T_X \longrightarrow 0$$

implies that $T_{X|Y} \otimes \pi^* T_X \subset \bigwedge^2 T_Y$. Since $T_{X|Y} = \operatorname{Aut}(\mathbb{P}(\mathcal{E})) = \operatorname{Aut}(\mathcal{E})/\mathbb{C}^*$ we get that

$$\pi_*(T_{X|Y} \otimes \pi^* T_X) = \pi_* T_{X|Y} \otimes T_X = End_0(\mathcal{E}) \otimes T_X,$$

where $End_0(\mathcal{E})$ denotes the trace-free endomorphisms of \mathcal{E} . Therefore, we can associate a trace-free co-Higgs fields $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ a bi-vector $\pi^* \phi \in H^0(X, T_X|_Y \otimes \pi^*T_X) \subset H^0(X, \bigwedge^2 T_Y)$ on $\mathbb{P}(\mathcal{E})$. The co-Higgs condition $\phi \wedge \phi = 0$ implies that bi-vector $\pi^* \phi$ is integrable, see the introduction of [20]. The codimension one foliation on $\mathbb{P}(\mathcal{E})$ is the called *foliation by symplectic leaves* induced by Poisson struture .

We get an interisting consequence of the proof Theorem 1.2.

Corollary 1.1. If (\mathcal{E}, ϕ) is locally free, stable and nilpotent, then the closure of the all leaves of the foliation by symplectic leaves on $\mathbb{P}(\mathcal{E})$ are rational surfaces.

2. Semi-stable co-Higgs sheaves

Definition 2.1. A co-Higgs sheaf on a complex manifold X is a sheaf \mathcal{E} together with a section $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ (called a Higgs fields) for which $\phi \wedge \phi = 0$.

Denote by $End_0(\mathcal{E}) := ker(tr : End(\mathcal{E}) \longrightarrow \mathcal{O}_X)$ the trace-free part of the endomorphism bundle of \mathcal{E} . Since

$$End(\mathcal{E}) = End_0(\mathcal{E}) \oplus \mathcal{O}_X$$

we have that $End(\mathcal{E}) \otimes T_X = (End_0(\mathcal{E}) \otimes T_X) \oplus T_X$. Thus, the Higgs field $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ can be decomposed as $\phi = \phi_1 + \phi_2$, where ϕ_1 is the trace-free part and ϕ_2 is a global vector field on X. In particular, if the surface X has no global holomorphic vector fields, then every Higgs field is trace-free.

Definition 2.2. Let (X, ω) be a polarized Kähler compact manifold. We say that (\mathcal{E}, ϕ) is semi-stable if

$$\frac{c_1(\mathcal{F}) \cdot [\omega]}{rank(\mathcal{F})} \le \frac{c_1(\mathcal{E}) \cdot [\omega]}{rank(\mathcal{E})}$$

for all coherent subsheaves $0 \neq \mathcal{F} \subsetneq \mathcal{E}$ satisfying $\Phi(\mathcal{F}) \subseteq \mathcal{F} \otimes T_X$, and stable if the inequality is strict for all such \mathcal{F} . We say that (\mathcal{E}, ϕ) is strictly semi-stable if (\mathcal{E}, ϕ) is semi-stable but non-stable.

3. Holomorphic foliations

Definition 3.1. Let X be a connected complex manifold. A one-dimensional holomorphic foliation is given by the following data:

- i) an open covering $\mathcal{U} = \{U_{\alpha}\}$ of X;
- ii) for each U_{α} an holomorphic vector field ζ_{α} ;
- iii) for every non-empty intersection, $U_{\alpha} \cap U_{\beta} \neq \emptyset$, a holomorphic function

 $f_{\alpha\beta} \in \mathcal{O}_X^*(U_\alpha \cap U_\beta);$

such that $\zeta_{\alpha} = f_{\alpha\beta}\zeta_{\beta}$ in $U_{\alpha} \cap U_{\beta}$ and $f_{\alpha\beta}f_{\beta\gamma} = f_{\alpha\gamma}$ in $U_{\alpha} \cap U_{\beta} \cap U_{\gamma}$.

We denote by $K_{\mathcal{F}}$ the line bundle defined by the cocycle $\{f_{\alpha\beta}\} \in \mathrm{H}^1(X, \mathcal{O}^*)$. Thus, a one-dimensional holomorphic foliation \mathcal{F} on X induces a global holomorphic section $\zeta_{\mathcal{F}} \in \mathrm{H}^0(X, T_X \otimes K_{\mathcal{F}})$. The line bundle $K_{\mathcal{F}}$ is called the *canonical bundle* of \mathcal{F} . Two sections $\zeta_{\mathcal{F}}$ and $\eta_{\mathcal{F}}$ of $\mathrm{H}^0(X, T_X \otimes K_{\mathcal{F}})$ are equivalent, if there exists a never vanishing holomorphic function $\varphi \in \mathrm{H}^0(X, \mathcal{O}^*)$, such that $\zeta_{\mathcal{F}} = \varphi \cdot \eta_{\mathcal{F}}$. It is clear that $\zeta_{\mathcal{F}}$ and $\eta_{\mathcal{F}}$ define the same foliation. Thus, a holomorphic foliation \mathcal{F} on X is an equivalence of sections of $H^0(X, T_X \otimes K_{\mathcal{F}})$.

4. Examples

4.1. Canonical example of split co-Higgs bundles. Here we will give an example which naturally generalizes the canonical example given by Rayan on [20, Chapter 6]. Let (X, ω) be a polarized Kähler compact manifold. Suppose that there exists a global section $\zeta \in H^0(X, Hom(N, TX \otimes L)) \simeq H^0(X, TX \otimes L \otimes N^*)$. Now, consider the vector bundle

$$\mathcal{E} = L \oplus N.$$

Define the following co-Higgs fields

$$\phi: L \oplus N \longrightarrow (T_X \otimes L) \oplus (T_X \otimes N) \in H^0(X, End(\mathcal{E}) \otimes T_X)$$

by $\phi(s,t) = (\zeta(t),0)$. Since $\phi \circ \phi = 0 \in H^0(End(\mathcal{E}) \otimes T_X \otimes T_X)$ we get that $\phi \wedge \phi = 0$. Moreover, observe that the kernel of ϕ is the ϕ -invariant line bundle L. On the other hand, the line bundle L is destabilising only when

$$[2c_1(L) - c_1(\mathcal{E})] \cdot [\omega] = [c_1(L) - c_1(N)] \cdot [\omega] > 0.$$

That is, if

$$[c_1(L)] \cdot [\omega] > [c_1(N)] \cdot [\omega].$$

4.2. Co-Higgs bundles on ruled surfaces. Let C be a curve of genus g > 1. Now, consider the ruled surface $X := \mathbb{P}(K_C \oplus \mathcal{O}_C)$ and

$$\pi: \mathbb{P}(K_C \oplus \mathcal{O}_C) \longrightarrow C$$

the natural projection. Consider a Poisson structure on X given by a bivector $\sigma \in H^0(X, \wedge^2 T_X)$. Let (\mathcal{E}, ϕ) be a nilpotent Higgs bundle on C. S. Rayan showed in [20] that $(\pi^* \mathcal{E}, \sigma(\pi^* \phi))$ is a stable co-Higgs bundle on X.

4.3. Co-Higgs orbibundles on weighted projective spaces. Let w_0, w_1, w_2 be positive integers, set $|w| := w_0 + w_1 + w_2$. Assume that w_0, w_1, w_2 are relatively prime. Define an action of \mathbb{C}^* in $\mathbb{C}^3 \setminus \{0\}$ by

$$\begin{array}{lll}
\mathbb{C}^* \times (\mathbb{C}^3 \setminus \{0\}) &\longrightarrow & (\mathbb{C}^3 \setminus \{0\}) \\
\lambda.(z_0, z_1, z_2) &\longmapsto & (\lambda^{w_0} z_0, \lambda^{w_1} z_1, \lambda^{w_2} z_2)
\end{array}$$
(4.1)

and consider the weighted projective plane

$$\mathbb{P}(w_0, w_1, w_2) := (\mathbb{C}^3 \setminus \{0\}) / \sim$$

induced by the action above. We will denote this space by $\mathbb{P}(\omega)$. On $\mathbb{P}(\omega)$ we have an Euler sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}(\omega)} \xrightarrow{\varsigma} \bigoplus_{i=0}^{2} \mathcal{O}_{\mathbb{P}(\omega)}(\omega_{i}) \longrightarrow T\mathbb{P}(\omega) \longrightarrow 0,$$

where $\mathcal{O}_{\mathbb{P}(\omega)}$ is the trivial line orbibundle and $T\mathbb{P}(\omega) = \operatorname{Hom}(\Omega^{1}_{\mathbb{P}(\omega)}, \mathcal{O}_{\mathbb{P}(\omega)})$ is the tangent orbibundle of $\mathbb{P}(\omega)$. The map ς is given explicitly by $\varsigma(1) = (\omega_{0}z_{0}, \omega_{1}z_{1}, \omega_{2}z_{2})$. Now, let (\mathcal{E}, ϕ) be a co-Higgs orbibundle on $\mathbb{P}(\omega)$. Tensoring the Euler sequence by $End(\mathcal{E})$, we obtain

$$0 \longrightarrow End(\mathcal{E}) \longrightarrow \bigoplus_{i=0}^{2} End(\mathcal{E})(\omega_{i}) \longrightarrow End(\mathcal{E}) \otimes T\mathbb{P}(\omega) \longrightarrow 0.$$

Thus, the co-Higgs fields ϕ can be represented, in homogeneous coordinates, by

$$\phi = \phi_0 \otimes \frac{\partial}{\partial z_0} + \phi_1 \otimes \frac{\partial}{\partial z_1} + \phi_2 \otimes \frac{\partial}{\partial z_2},$$

where $\phi_i \in H^0(\mathbb{P}(\omega), End(\mathcal{E})(w_i))$, for all i = 0, 1, 2, and $\phi + \theta \otimes R_{\omega}$ define the same co-Higgs field as ϕ , where R_{ω} is the adapted radial vector field

$$R_{\omega} = \omega_0 z_0 \frac{\partial}{\partial z_0} + \omega_1 z_1 \frac{\partial}{\partial z_1} + \omega_2 z_2 \frac{\partial}{\partial z_2},$$

with θ a endomorphism of \mathcal{E} . Suppose that

$$\mathcal{E} = \mathcal{O}(m_1) \oplus \mathcal{O}(m_2)$$

and that there exists a stable ϕ for \mathcal{E} such that $m_1 \geq m_2$. Then

$$|m_1 - m_2| \le \max_{0 \le i \ne j \le 2} \{\omega_i + \omega_j\}.$$

In fact, this is a consequence of Bott's Formulae for weighted projective spaces. It follows from (see [6]) that

$$\mathrm{H}^{0}(\mathbb{P}(\omega), T\mathbb{P}(\omega) \otimes \mathcal{O}_{\omega}(k)) \simeq \mathrm{H}^{0}(\mathbb{P}(\omega), \Omega^{1}_{\mathbb{P}(\omega)}(\sum_{i=0}^{2} \omega_{i} + k)) \neq \emptyset$$

if and only if $k > -\max_{0 \le i \ne j \le 2} \{\omega_i + \omega_j\}$. This generalize the example given by S. Rayan in [19].

4.4. Co-Higgs bundles on two dimensional complex tori. Let X be a two dimensional complex torus and a co-Higgs bundle $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$. Then ϕ is equivalente to a pair of commutative endomorphism of \mathcal{E} . In fact, since the tangent bundle T_X is holomorphically trivial, we can take a trivialization by choosing two linearly independent global vector fields $v_1, v_2 \in H^0(X, T_X)$. Then, we can write

$$\phi = \phi_1 \otimes v_1 + \phi_2 \otimes v_2.$$

The condition $\phi \wedge \phi = 0$ implies that

We have a canonical nilpotent co-Higgs bundle (\mathcal{E}, ϕ) , where $\mathcal{E} = T_X = \mathcal{O} \oplus \mathcal{O}$ and

$$\left(\begin{array}{cc} 0 & v \\ 0 & 0 \end{array}\right),$$

where v is a global vector field on X.

5. Proof of Theorem

By using that the Higgs field $\phi \in H^0(X, End(\mathcal{E}) \otimes T_X)$ is nilpotent we have that $Ker(\phi) =: L$ is a well defined line bundle on X. Thus, we have a exact sequence

$$0 \to L \longrightarrow \mathcal{E} \longrightarrow \mathcal{I}_Z \otimes N \longrightarrow 0,$$

where the nilpotent Higgs field ϕ factors as

The morphism $\mathcal{I}_Z \otimes N \to L \otimes T_X$ induces a holomorphic foliation on X which induces a global section $\zeta_{\phi} \in H^0(X, T_X \otimes L \otimes N^*)$. Since $\det(\mathcal{E}) = L \otimes N$ we conclude that $L \otimes N^* = L^2 \otimes \det(\mathcal{E}^*)$. Then

$$\zeta_{\phi} \in H^0(X, T_X \otimes L^2 \otimes \det(\mathcal{E}^*)).$$

Let $K := L^2 \otimes \det(\mathcal{E}^*)$ the canonical bundle of the foliation \mathcal{F} associated to the co-Higgs fields ϕ . If \mathcal{E} is semi-stable then

$$[c_1(K)] \cdot [\omega] = [2c_1(L) - c_1(\mathcal{E})] \cdot [\omega] \le 0$$

for some Kähler class ω . If $K \cdot [\omega] < 0$, it follows from [14] that K is not pseudoeffective [5]. It follows from Brunella's theorem [1] that X is uniruled.

Now, suppose X is a normal projective surface and that $K \cdot H = 0$, for some H ample By Hodge index theorem we have that $K^2 \cdot H^2 \leq (K \cdot H)^2 = 0$, then $K^2 \leq 0$. Suppose that $K^2 < 0$. We have that $D = H + \epsilon K$ is a \mathbb{Q} -divisor ample for $0 < \epsilon << 1$, see [15, proposition 1.3.6]. Thus, we have that

$$K \cdot D = K \cdot H + \epsilon^2 K^2 = \epsilon^2 K^2 < 0.$$

By Bogomolov-McQuillan-Miyaoka's theorem [3] we conclude that X is uniruled. If $K^2 = 0$, then K is numerically trivial. This fact is well known, but for convenience of the reader we give a proof. Suppose that there exists $C \subset X$ such that $K \cdot C > 0$. Now, Consider the divisor $B = (H^2)C - (H \cdot C)H$. Then $B \cdot H = 0$ and $K \cdot B = (H^2)K \cdot C < 0$. Define F = mK + B for 0 < m << 1. Therefore $F \cdot H = 0$ and $F^2 > 0$. This is a contradiction by the Hodge index Theorem. In this case \mathcal{E} is strictly semi-stable.

Now, we apply the classification, up to finite étale cover, of holomorphic foliations on projective surfaces with canonical bundle numerically trivial [22], [17], [2]. Therefore, up to finite étale cover, either:

- i) X is uniruled;
- ii) X is a torus;
- *iii*) k(X) = 1 and $X = B \times C$ with $g(B) \ge 2$, C is elliptic. That is, X is a sesquielliptic surface.

If X is Kähler and non-algebraic it follows from [2] that, up to finite étale cover, either X has a unique elliptic fibration or X is a torus.

M. CORRÊA

6. Proof of Corollary 1.1

Since (\mathcal{E}, ϕ) is nilpotent and stable the co-Higgs fields induces a foliation \mathcal{F} by rational curves on X. Now, consider the projective bundle $\pi : \mathbb{P}(\mathcal{E}) \to X$. Then the foliation by symplectic leaves \mathcal{G} on $\mathbb{P}(\mathcal{E})$ is the pull-back of \mathcal{F} by π . In particular, a closure of the a leaf of the foliation by symplectic leaves \mathcal{G} is of type $\pi^{-1}(f(\mathbb{P}^1))$, where $f : \mathbb{P}^1 \to X$ is the uniformization of a rational leaf of \mathcal{F} . Clearly $\pi^{-1}(f(\mathbb{P}^1))$ is a rational surface.

ACKNOWLEGMENTS. We are grateful to Arturo Fernandez-Perez, Renato Martins and Marcos Jardim for pointing out corrections. We are grateful to Henrique Bursztyn for interesting conversations about Poisson Geometry.

References

- M. Brunella, A positivity property for foliations on compact Kähler manifolds, Internat. J. Math. 17 (2006), 35-43.
- M. Brunella, Foliations on complex projective surfaces, in Dynamical Systems, Part II, Scuola Norm.Sup., Pisa 2003, 49-77.
- [3] F. Bogomolov, M. McQuillan, Rational curves on foliated varieties, preprint IHES M/01/07 (2001)
- [4] K. Corlette, Flat G-bundles with canonical metrics. J. Differential Geom. 28, 3 (1988), 361-382.
- [5] J.-P. Demailly, L2-vanishing theorems for positive line bundles and adjunction theory, Transcendental Methods in Algebraic Geometry, Springer Lecture Notes 1646 (1996), 1-97
- [6] I. Dolgachev, Weighted projective varieties, Group actions and vector fields, pp. 34-71, Lecture Notes in Mathematics 956, Springer-Verlag, Berlin-Heidelberg, New York (1982).
- M Gualtieri , Generalized complex geometry, Ann. of Math.(2), 2011 arXiv:math/0401221v1. DPhil thesis, University of Oxford.
- [8] S. Gukov, E. Witten, Branes and quantization. Adv. Theor. Math. Phys. 13, 5 (2009), 1445-1518.
- [9] N. Hitchin, Generalized holomorphic bundles and the B-field action, J. Geom. and Phys. 61 (2011), no. 1, 352-362.
- [10] N. Hitchin, Generalized Calabi-Yau manifolds. Q.J.Math. 54,3(2003),281-308.
- [11] N. Hitchin, The self-duality equations on a Riemann surface. Proc. London Math. Soc. (3) 55, 1 (1987), 59-126.
- [12] A. Kapustin, Y. Li, Open-string BRST cohomology for generalized complex branes. Adv. Theor. Math. Phys. 9, 4 (2005), 559-574.
- [13] S. Kebekus, L.Sola-Conde, M. Toma. Rationally connected foliations after Bogomolov and McQuillan (mit Luis Sol und Matei Toma) Journal Algebraic Geometry16 (2007), 65-81.
- [14] A. Lamari, Le cone Kählerien d'une surface, J. Math. Pures Appl. 78 (1999), 249-263
- [15] R. K. Lazarsfeld, *Positivity in algebraic geometry*, Vol. I, Springer, 2000, A Series of Modern Surveys in Mathematics, No. 48.
- [16] M. McQuillan, Noncommutative Mori theory, preprint IHES M/00/15 (2000) (re- vised: M/01/42 (2001))
- [17] M. McQuillan, Diophantine approximations and foliations, Publ. Math. IHES 87 (1998), 121-174
- [18] Y. Miyaoka. Deformation of a morphism along a foliation. Algebraic Geometry Bowdoin 1985. Proc. Symp. Pure Math. 46 (1987), 245-268.
- [19] Constructing co-Higgs bundles on \mathbb{CP}^2 . Q. J. Math. 65 (2014), no. 4, 1437-1460
- [20] S. Rayan, Geometry of co-Higgs bundles. D. Phil. thesis, Oxford, 2011. http://people.maths.ox.ac.uk/hitchin/hitchinstudents/rayan.pdf.
- [21] S. Rayan, Co-Higgs bundles on \mathbb{P}^1 , New York J. Math. 19 (2013), 925-945;
- [22] T. Peternell, Generically nef vector bundles and geometric applications, in: Complex and differential geometry, 345-368, Springer Proc. in Math. 8, Springer-Verlag, Berlin (2011).
- [23] A. Polishchuk, Algebraic geometry of Poisson brackets. J. Math. Sci. 84 (1997), 1413-1444.
- [24] C. Simpson, Moduli of representations of the fundamental group of a smooth projective variety I; Inst. Hautes Etudes Sci. Publ. Math., 79 (1994), 47-129.
- [25] C. Simpson, Moduli of representations of the fundamental group of a smooth projective variety II; Inst. Hautes Etudes Sci. Publ. Math., 80 (1994), 5-79.

[26] R. Zucchini, Generalized complex geometry, generalized branes and the Hitchin sigma model.
 J. High Energy Phys., 3 (2005), 22-54

DEPARTAMENTO DE MATEMÁTICA - ICEX, UNIVERSIDADE FEDERAL DE MINAS GERAIS, UFMG Current address: Av. Antônio Carlos 6627, 31270-901, Belo Horizonte-MG, Brasil. E-mail address: mauricio@mat.ufmg.br