

A DIRECT APPROACH TO PLATEAU'S PROBLEM IN ANY CODIMENSION

G. DE PHILIPPIS, A. DE ROSA, AND F. GHIRALDIN

ABSTRACT. This paper aims to propose a direct approach to solve the Plateau's problem in codimension higher than one. The problem is formulated as the minimization of the Hausdorff measure among a family of d -rectifiable closed subsets of \mathbb{R}^n : following the previous work [DGM14] the existence result is obtained by a compactness principle valid under fairly general assumptions on the class of competitors. Such class is then specified to give meaning to boundary conditions. We also show that the obtained minimizers are regular up to a set of dimension less than $(d - 1)$.

1. INTRODUCTION

Plateau's problem consists in looking for a surface of minimal area among those surfaces spanning a given boundary. A considerable amount of the development of Geometric Measure Theory in the last fifty years has been devoted to provide generalized concepts of surface, area and of "spanning a given boundary", in order to apply the direct methods of the calculus of variations to the Plateau's problem. In particular we recall the notions of sets of finite perimeter [De54, De55], of currents [FF60] and of varifolds [All72, All75, Alm68], introduced respectively by De Giorgi, Federer, Fleming, Almgren and Allard. A more "geometric" approach was proposed by Reifenberg in [Rei60], where Plateau's problem was set as the minimization of Hausdorff d -dimensional measure among compact sets and the notion of spanning a given boundary was given in term of inclusions of homology groups.

Any of these approach has some drawbacks: in particular, not all the "reasonable" boundaries can be obtained by the above notions and not always the solutions are allowed to have the type of singularities observed by soap bubble (the so called Plateau's laws). Recently in [HP13] Harrison and Pugh, see also [Har14], proposed a new notion of spanning a boundary, which seems to include all reasonable physical boundaries and they have been able to show, in the co-dimension one case, existence of least area surfaces spanning a given boundary.

In the recent paper [DGM14], De Lellis, Maggi and the third author have proposed a direct approach to the Plateau's problem, based on the "elementary" theory of Radon measures and on a deep result of Preiss concerning rectifiable measures. Roughly speaking they showed, in the co-dimension one case, that every time one has a class which contains "enough" competitors (namely the cone and the cup competitors, see [DGM14, Definition 1]) it is always possible to show that the infimum of the Plateau's problem is achieved by the area of a rectifiable set. They then applied this result to provide a new proof of Harrison and Pugh theorem as well as to show the existence of sliding minimizers, a new notion of minimal sets proposed by David in [Dav14, Dav13] and inspired by Almgren's $(\mathbf{M}, 0, \infty)$, [Alm76].

In this note, we extend the result [DGM14] to any co-dimension. More precisely, we prove that every time the class of competitors for the Plateau's Problem consists of rectifiable sets and it is closed by Lipschitz deformations, it is always possible to show that the infimum is achieved by a compact set K which is, away from the "boundary", an analytic manifold outside a closed set of Hausdorff dimension at most $(d - 1)$, see Theorem 1.3 below for the precise statement. We then apply this result to provide existence of sets spanning a given boundary according to

the natural generalization of the notion introduced by Harrison and Pugh, Theorem 1.3, and to show the existence of sliding minimizers in any co-dimension, Theorem 1.8.

Although the general strategy of the proof is the same of [DGM14], some non-trivial modifications have to be done in order to deal with sets of any co-dimension. In particular, with respect to [DGM14], we use a different notion of “good class”, the main reason being the following: one of the key step of the proof of our main result consists in showing a precise density lower bound for the measure obtained as limit of the sequence of Radon measures naturally associated to a minimizing sequence (K_j) , see Steps 1 and 4 in the proof of Theorem 1.3. In order to obtain such a lower bound, instead of relying on relative isoperimetric inequalities on the sphere as in [DGM14] (which are peculiar of the co-dimension one case) we use the deformation theorem of David and Semmes in [DS00] to obtain suitable competitors, following a strategy already introduced by Federer and Fleming for rectifiable currents, see [FF60] and [Alm76]. Moreover since our class is essentially closed by Lipschitz deformations, we are actually able to prove that any set achieving the infimum is a stationary varifold and that, in addition, it is smooth outside a closed set of relative co-dimension one (this does not directly follows by Allard’s regularity theorem, see Step 7 in the proof of Theorem 1.3). Simple examples show that this regularity is actually optimal.

In order to precise state our main results, let us introduce some notations and definitions, referring to Section 2 for more details. We will always work in \mathbb{R}^n and $1 \leq d \leq n$ will always be an integer number, we recall that a set K is said to be d -rectifiable if it can be covered up to an \mathcal{H}^d negligible set by countably many C^1 manifolds, see [Sim83, Chapter 3], where \mathcal{H}^d is the d -dimensional Hausdorff measure. We also let $\text{Lip}(\mathbb{R}^n)$ be the space of Lipschitz maps in \mathbb{R}^n .

Definition 1.1 (Lipschitz deformations). Given a ball $B_{x,r}$, we let $\mathfrak{D}(x, r)$ be the set of functions $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\varphi(z) = z$ in $\mathbb{R}^n \setminus B_{x,r}$ and which are smoothly isotopic to the identity inside $B_{x,r}$, namely those for which there exists a smooth isotopy $\lambda : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that

$$\lambda(0, \cdot) = \text{Id}, \quad \lambda(1, \cdot) = \varphi, \quad \lambda(t, h) = h \quad \forall (t, h) \in [0, 1] \times (\mathbb{R}^n \setminus B_{x,r}) \quad \text{and} \\ \lambda(t, \cdot) \text{ is a diffeomorphism of } \mathbb{R}^n \quad \forall t \in [0, 1].$$

We finally set $\mathsf{D}(x, r) := \overline{\mathfrak{D}(x, r)}^{C^0} \cap \text{Lip}(\mathbb{R}^n)$, the intersection of the Lipschitz maps with the closure of $\mathfrak{D}(x, r)$ with respect to the uniform topology.

The following definition describes the properties we require to the comparison sets: the key property we ask for K' to be a competitor of K is that K' must be close in energy to sets obtained from K via deformation maps in Definition 1.1. This allows a larger flexibility on the choice of the admissible sets, since a priori K' might not belong to the competition class.

Definition 1.2 (Deformed competitors and good class). Let $H \subset \mathbb{R}^n$ be closed. Given $K \subset \mathbb{R}^n \setminus H$ relatively closed countably \mathcal{H}^d -rectifiable and $B_{x,r} \subset \mathbb{R}^n \setminus H$, a *deformed competitor* for K in $B_{x,r}$ is any set of the form

$$\varphi(K) \quad \text{where} \quad \varphi \in \mathsf{D}(x, r).$$

Given a family $\mathcal{P}(H)$ of relatively closed d -rectifiable subsets $K \subset \mathbb{R}^n \setminus H$, we say that $\mathcal{P}(H)$ is a *good class* if for every $K \in \mathcal{P}(H)$, for every $x \in K$ and for a.e. $r \in (0, \text{dist}(x, H))$

$$\inf \{ \mathcal{H}^d(J) : J \in \mathcal{P}(H), J \setminus \overline{B_{x,r}} = K \setminus \overline{B_{x,r}} \} \leq \mathcal{H}^d(L) \quad (1.1)$$

whenever L is any deformed competitor for K in $B_{x,r}$.

Once we fix a closed set H , we can formulate Plateau’s problem in the class $\mathcal{P}(H)$:

$$m_0 := \inf \{ \mathcal{H}^d(K) : K \in \mathcal{P}(H) \}. \quad (1.2)$$

We will say that a sequence $(K_j) \subset \mathcal{P}(H)$ is a *minimizing sequence* if $\mathcal{H}^d(K_j) \downarrow m_0$. The following theorem is our main result and establishes the behavior of minimizing sequences.

Theorem 1.3. *Let $H \subset \mathbb{R}^n$ be closed and $\mathcal{P}(H)$ be a good class. Assume the infimum in Plateau's problem (1.2) is finite and let $(K_j) \subset \mathcal{P}(H)$ be a minimizing sequence. Then, up to subsequences, the measures $\mu_j := \mathcal{H}^d \llcorner K_j$ converge weakly* in $\mathbb{R}^n \setminus H$ to a measure $\mu = \mathcal{H}^d \llcorner K$, where $K = \text{spt } \mu \setminus H$ is a countably \mathcal{H}^d -rectifiable set. Furthermore:*

- (a) *the integral varifold naturally associated to μ is stationary in $\mathbb{R}^n \setminus H$;*
- (b) *K is a real analytic submanifold outside a relatively closed set $\Sigma \subset K$ with $\dim_{\mathcal{H}}(\Sigma) \leq d - 1$.*

In particular, $\liminf_j \mathcal{H}^d(K_j) \geq \mathcal{H}^d(K)$ and if $K \in \mathcal{P}(H)$, then K is a minimum for (1.2).

We wish to apply Theorem 1.3 to two definitions of boundary conditions. The first one is the natural generalization of the one considered in [HP13]:

Definition 1.4. Let H be a closed set in \mathbb{R}^n .

Let us consider the family

$$\mathcal{C}_H = \{ \gamma : S^{n-d} \rightarrow \mathbb{R}^n \setminus H : \gamma \text{ is a smooth embedding of } S^{n-d} \text{ into } \mathbb{R}^n \}.$$

We say that $\mathcal{C} \subset \mathcal{C}_H$ is closed by isotopy (with respect to H) if \mathcal{C} contains all elements $\gamma' \in \mathcal{C}_H$ belonging to the same smooth isotopy class $[\gamma]$ in $\mathbb{R}^n \setminus H$ of any $\gamma \in \mathcal{C}$, see [Hir94, Ch. 8]. Given $\mathcal{C} \subset \mathcal{C}_H$ closed by isotopy, we say that a relatively closed subset K of $\mathbb{R}^n \setminus H$ is a \mathcal{C} -spanning set of H if

$$K \cap \gamma \neq \emptyset \text{ for every } \gamma \in \mathcal{C}.$$

We denote by $\mathcal{F}(H, \mathcal{C})$ the family of countably \mathcal{H}^d -rectifiable sets which are \mathcal{C} -spanning sets of H .

We can prove the following closure property for the class $\mathcal{F}(H, \mathcal{C})$:

Theorem 1.5. *Let H be closed in \mathbb{R}^n and \mathcal{C} be closed by isotopy with respect to H , then:*

- (a) *$\mathcal{F}(H, \mathcal{C})$ is a good class in the sense of Definition 1.2.*
- (b) *Assume the infimum (1.2) corresponding to $\mathcal{P}(H) = \mathcal{F}(H, \mathcal{C})$ is finite, then the set K provided by Theorem 1.3 belongs to $\mathcal{F}(H, \mathcal{C})$. In particular the Plateau's problem in the class $\mathcal{F}(H, \mathcal{C})$ has a solution.*

The second type of boundary condition we want to consider is the one related to the notion of "sliding minimizers" introduced by David in [Dav14, Dav13].

Definition 1.6 (Sliding minimizers). Let $H \subset \mathbb{R}^n$ be closed and $K_0 \subset \mathbb{R}^n \setminus H$ be relatively closed. We denote by $\Sigma(H)$ the family of Lipschitz maps $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that there exists a continuous map $\Phi : [0, 1] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\Phi(1, \cdot) = \varphi$, $\Phi(0, \cdot) = \text{Id}$ and $\Phi(t, H) \subset H$ for every $t \in [0, 1]$. We then define

$$\mathcal{A}(H, K_0) = \{ K : K = \varphi(K_0) \text{ for some } \varphi \in \Sigma(H) \}$$

and say that K_0 is a sliding minimizer if $\mathcal{H}^d(K_0) = \inf \{ \mathcal{H}^d(J) : J \in \mathcal{A}(H, K_0) \}$.

Remark 1.7. For every $K_0 \subset \mathbb{R}^n \setminus H$ relatively closed and d -rectifiable, $\mathcal{A}(H, K_0)$ is a good class in the sense of Definition 1.2, since $D(x, r) \subset \Sigma(H)$ for every $B_{x,r} \subset \mathbb{R}^n \setminus H$.

Applying Theorem 1.3 to the contest of sliding minimizers we obtain the following result which is the analogous of [DGM14, Theorem 7] in any codimension. Here and in the following $U_\delta(E)$ denotes the δ -neighborhood of a set $E \subset \mathbb{R}^n$.

Theorem 1.8. *Assume that*

- (i) *K_0 is bounded d -rectifiable with $\mathcal{H}^d(K_0) < \infty$;*
- (ii) *$\mathcal{H}^d(H) = 0$ and for every $\eta > 0$ there exist $\delta > 0$ and $\Pi \in \Sigma(H)$ such that*

$$\text{Lip } \Pi \leq 1 + \eta, \quad \Pi(U_\delta(H)) \subset H.$$

Then, given any minimizing sequence (K_j) in the Plateau's problem corresponding to $\mathcal{P}(H) = \mathcal{A}(H, K_0)$ and any set K as in Theorem 1.3, we have

$$\inf \{ \mathcal{H}^d(J) : J \in \mathcal{A}(H, K_0) \} = \mathcal{H}^d(K) = \inf \{ \mathcal{H}^d(J) : J \in \mathcal{A}(H, K) \}.$$

In particular K is a sliding minimizer.

The paper is structured as follows, in Section 2 we will recall some basic definitions and recall some known theorems we are going to use, in particular Preiss rectifiability criterion and a version of the deformation theorem due to David and Semmes. In Section 3 we prove Theorem 1.3 and in Section 4 we prove Theorems 1.5 and 1.8.

Acknowledgements. The authors are grateful to Camillo De Lellis, Francesco Maggi and Emanuele Spadaro for many interesting comments and suggestions. This work has been supported by ERC 306247 *Regularity of area-minimizing currents* and by SNF 146349 *Calculus of variations and fluid dynamics*.

2. NOTATION AND PRELIMINARIES

We are going to use the following notations: $Q_{x,l}$ denotes the closed cube centered in x , with edge length l ; moreover we set

$$R_{x,a,b} := x + \left[-\frac{a}{2}, \frac{a}{2} \right]^d \times \left[-\frac{b}{2}, \frac{b}{2} \right]^{n-d} \quad \text{and} \quad B_{x,r} := \{y \in \mathbb{R}^n : |y - x| < r\}.$$

When cubes, rectangles and balls are centered in the origin, we will simply write Q_l , $R_{a,b}$ and B_r . Cubes and balls in the subspace $\mathbb{R}^d \times \{0\}^{n-d}$ are denoted with $Q_{x,l}^d$ and $B_{x,r}^d$ respectively. We also let ω_d be the Lebesgue measure of the unit ball in \mathbb{R}^d .

Let us recall the following deep structure result for Radon measures due to Preiss [Pre87, DeL08] which will play a key role in the proof of Theorem 1.3.

Theorem 2.1. *Let d be an integer and μ a locally finite measure on \mathbb{R}^n such that the d -density*

$$\theta(x) := \lim_{r \rightarrow 0} \frac{\mu(B_{x,r})}{\omega_d r^d}$$

exists and satisfies $0 < \theta(x) < +\infty$ for μ -a.e. x . Then $\mu = \theta \mathcal{H}^d \llcorner K$, where K is a countably \mathcal{H}^d -rectifiable set.

In order to apply Preiss' Theorem we will rely on the monotonicity formula for minimal surfaces, which roughly speaking can be obtained by comparing the given minimizer with a cone. To this aim let us introduce the following definition:

Definition 2.2 (Cone competitors). In the setting of Definition 1.2, the cone competitor for K in $B_{x,r}$ is the following set

$$\mathbf{C}_{x,r}(K) = (K \setminus B_{x,r}) \cup \{ \lambda x + (1 - \lambda)z : z \in K \cap \partial B_{x,r}, \lambda \in [0, 1] \}. \quad (2.1)$$

Let us note that in general a cone competitor in $B_{x,r}$ is not a deformed competitor in $B_{x,r}$. On the other hand as in [DGM14] we can show that:

Lemma 2.3. *Given a good class $\mathcal{P}(H)$ in the sense of Definition 1.2, for any $K \in \mathcal{P}(H)$ countably \mathcal{H}^d -rectifiable and for every $x \in K$, the set K verifies for a.e. $r \in (0, \text{dist}(x, H))$:*

$$\inf \{ \mathcal{H}^d(J) : J \in \mathcal{P}(H), J \setminus \overline{B_{x,r}} = K \setminus \overline{B_{x,r}} \} \leq \mathcal{H}^d(\mathbf{C}_{x,r}(K)).$$

Proof. Without loss of generality let us consider balls B_r centered at 0 with $B_r \subset \subset \mathbb{R}^n \setminus H$. We assume in addition that $K \cap \partial B_r$ is \mathcal{H}^{d-1} -rectifiable with $\mathcal{H}^{d-1}(K \cap \partial B_r) < \infty$ and that r is a

Lebesgue point of $t \in (0, \infty) \mapsto \mathcal{H}^{d-1}(K \cap \partial B_t)$. All these conditions are fulfilled for a.e. r and, again by scaling, we can assume that $r = 1$ and use B instead of B_1 . For $s \in (0, 1)$ let us set

$$\varphi_s(r) = \begin{cases} 0, & r \in [0, 1-s), \\ \frac{r-(1-s)}{s}, & r \in [1-s, 1], \\ r, & r \geq 1, \end{cases}$$

and $\phi_s(x) = \varphi_s(|x|) \frac{x}{|x|}$ for $x \in \mathbb{R}^n$. In this way, one easily checks that $\phi_s : \mathbb{R}^n \rightarrow \mathbb{R}^n \in \mathcal{D}(0, 1)$.

Since $\phi_s(K \cap \overline{B_{1-s}}) = \{0\}$, we need to show that

$$\limsup_{s \rightarrow 0^+} \mathcal{H}^d(\phi_s(K \cap (B \setminus B_{1-s}))) \leq \frac{\mathcal{H}^{d-1}(K \cap \partial B)}{d} = \mathcal{H}^d(\mathbf{C}_{x,r}(K)).$$

Let $x_0 \in K \cap \partial B_t$ and let us fix an orthonormal base ν_1, \dots, ν_d of the approximate tangent space $T_{x_0}K$ such that $\nu_i \in T_{x_0}K \cap T_{x_0}\partial B_t$ for $i \leq d-1$. Let

$$J_d^K \phi_s = \left| \left(\bigwedge^d D\phi_s \right) (T_{x_0}K) \right| = |D\phi_s(\nu_1) \wedge \dots \wedge D\phi_s(\nu_d)|$$

be the d -dimensional tangential Jacobian of ϕ_s with respect to K . Letting I be the (at most countable) set of those $t \in (0, 1)$ such that $\mathcal{H}^{d-1}(K \cap \partial B_t) > 0$, we find with the aid of the area and co-area formulas,

$$\begin{aligned} \mathcal{H}^d(\phi_s(K \cap (B \setminus B_{1-s}))) &= \int_{K \cap (B \setminus B_{1-s})} J_d^K \phi_s \, d\mathcal{H}^d \\ &= \int_{1-s}^1 dt \int_{K \cap \partial B_t} \frac{J_d^K \phi_s}{|\nu_d \cdot \hat{x}|} \, d\mathcal{H}^{d-1} + \sum_{t \in I \cap (1-s, 1)} \left(\frac{t-(1-s)}{1-s} \right)^d \mathcal{H}^d(K \cap \partial B_t), \end{aligned} \quad (2.2)$$

where $\hat{x} = x/|x|$ and we have used that $|\nu_d \cdot \hat{x}|$ is the tangential co-area factor of the map $f(x) = |x|$. We first notice that, for $t \in (1-s, 1)$, $t - (1-s) \leq s$. Moreover

$$\lim_{s \rightarrow 0} \sum_{t \in I \cap (1-s, 1)} \mathcal{H}^d(K \cap \partial B_t) = 0,$$

and thus the second term in (2.2) can be ignored. At the same time, for a constant C ,

$$J_d^K \phi_s(x) \leq C + |\hat{x} \cdot \nu_d| \varphi'_s(|x|) \left(\frac{\varphi_s(|x|)}{|x|} \right)^{d-1}, \quad \text{for } \mathcal{H}^d\text{-a.e. } x \in K.$$

The constant C gives a negligible contribution in the integral as $s \downarrow 0$; as for the second term, having $\varphi'_s = 1/s$ on $(1-s, 1)$, we find

$$\int_{1-s}^1 \mathcal{H}^{d-1}(K \cap \partial B_t) \varphi'_s(t) \left(\frac{\varphi_s(t)}{t} \right)^{d-1} dt = \frac{1}{s} \int_{1-s}^1 \mathcal{H}^{d-1}(K \cap \partial B_t) \left(\frac{\varphi_s(t)}{t} \right)^{d-1} dt.$$

Since $t = 1$ is a Lebesgue point of $t \in (0, \infty) \mapsto \mathcal{H}^{d-1}(K \cap \partial B_t)$, we have

$$\lim_{s \rightarrow 0} \frac{1}{s} \int_{1-s}^1 |\mathcal{H}^{d-1}(K \cap \partial B_t) - \mathcal{H}^{d-1}(K \cap \partial B)| dt = 0,$$

so that, combining the above remarks we find

$$\limsup_{s \rightarrow 0^+} \mathcal{H}^d(\phi_s(K \cap B)) \leq \mathcal{H}^{d-1}(K \cap \partial B) \limsup_{s \rightarrow 0^+} \frac{1}{s} \int_{1-s}^1 \left(\frac{\varphi_s(t)}{t} \right)^{d-1} dt = \frac{\mathcal{H}^{d-1}(K \cap \partial B)}{d},$$

as required. \square

The second key result we are going to use is a deformation theorem for closed sets due to David and Semmes [DS00], analogous to the one for rectifiable currents [Sim83, Fed69]. We provide a slightly extended statement for the sake of forthcoming proofs.

Before stating the theorem, let us introduce some further notation. Given a closed cube Q of edges length l in \mathbb{R}^n and $\varepsilon > 0$, we cover Q with closed smaller cubes with edges length $\varepsilon \ll l$, non empty intersection with $\text{Int}(Q)$ and such that the decomposition is centered in x (i.e. one of the subcubes is centered in x). The family of this smaller cubes is denoted $\Lambda_\varepsilon(Q)$. We set

$$\begin{aligned} C_1 &:= \bigcup \{T \cap Q : T \in \Lambda_\varepsilon(Q), T \cap \partial Q \neq \emptyset\}, \\ C_2 &:= \bigcup \{T \in \Lambda_\varepsilon(Q) : T \not\subset C_1, T \cap \partial C_1 \neq \emptyset\}, \\ Q_1 &:= \overline{Q \setminus (C_1 \cup C_2)} \end{aligned} \tag{2.3}$$

and consequently

$$\Lambda_\varepsilon(Q_1 \cup C_2) := \{T \in \Lambda_\varepsilon(Q) : T \subset (Q_1 \cup C_2)\}$$

For each nonnegative integer $m \leq n$, let $\Lambda_{\varepsilon,m}(Q_1 \cup C_2)$ denote the collection of all m -dimensional faces of cubes in $\Lambda_\varepsilon(Q_1 \cup C_2)$ and $\Lambda_{\varepsilon,m}^*(Q_1 \cup C_2)$ will be the set of the elements of $\Lambda_{\varepsilon,m}(Q_1 \cup C_2)$ which are not contained in $\partial(Q_1 \cup C_2)$. We also let $S_{\varepsilon,m}(Q_1 \cup C_2) := \bigcup \Lambda_{\varepsilon,m}(Q_1 \cup C_2)$ be the the m -skeleton of order ε in $Q_1 \cup C_2$.

Theorem 2.4. *Let $r > 0$ and E be a compact subset of Q such that $\mathcal{H}^d(E) < +\infty$ and $Q \subset B_{x,r}$. There exists a map $\Phi_{\varepsilon,E} \in \mathcal{D}(x,r)$ satisfying the following properties:*

- (1) $\Phi_{\varepsilon,E}(x) = x$ for $x \in \mathbb{R}^n \setminus (Q_1 \cup C_2)$;
- (2) $\Phi_{\varepsilon,E}(x) = x$ for $x \in S_{\varepsilon,d}(Q_1 \cup C_2)$;
- (3) $\Phi_{\varepsilon,E}(E) \subset S_{\varepsilon,d}(Q_1 \cup C_2) \cup \partial(Q_1 \cup C_2)$;
- (4) $\Phi_{\varepsilon,E}(T) \subset T$ for every $T \in \Lambda_{\varepsilon,m}(Q_1 \cup C_2)$, with $m = d, \dots, n$;
- (5) either $\mathcal{H}^d(\Phi_{\varepsilon,E}(E \cap T)) = 0$ or $\mathcal{H}^d(\Phi_{\varepsilon,E}(E \cap T)) = \mathcal{H}^d(T)$, for every $T \in \Lambda_{\varepsilon,d}^*(Q_1)$;
- (6) $\mathcal{H}^d(\Phi_{\varepsilon,E}(E \cap T)) \leq k_1 \mathcal{H}^d(E \cap T)$ for every $T \in \Lambda_\varepsilon(Q_1 \cup C_2)$;

where k_1 depends only on n and d (but not on ε).

Proof. Proposition 3.1 in [DS00] provides a map $\widetilde{\Phi}_{\varepsilon,E} \in \mathcal{D}(x,r)$ satisfying properties (1)-(4) and (6). We want to set

$$\Phi_{\varepsilon,E} := \Psi \circ \widetilde{\Phi}_{\varepsilon,E},$$

where Ψ will be defined below. We first define Ψ on every $T \in \Lambda_{\varepsilon,d}(Q_1 \cup C_2)$ distinguishing two cases

- (a) if either $\mathcal{H}^d(\widetilde{\Phi}_{\varepsilon,E}(E) \cap T) = 0$ or $\mathcal{H}^d(\widetilde{\Phi}_{\varepsilon,E}(E) \cap T) = \mathcal{H}^d(T)$ or $T \notin \Lambda_{\varepsilon,d}^*(Q_1)$, then we set $\Psi|_T = \text{Id}$;
- (b) otherwise since $\widetilde{\Phi}_{\varepsilon,E}(E)$ is compact, there exists $y_T \in T$ and $\delta_T > 0$ such that $B_{\delta_T}(y_T) \cap \widetilde{\Phi}_{\varepsilon,E}(E) = \emptyset$; we define

$$\Psi|_T(x) = x + \alpha(x - y_T) \min \left\{ 1, \frac{|x - y_T|}{\delta_T} \right\},$$

where $\alpha > 0$ such that the point $x + \alpha(x - y_T) \in (\partial T) \times \{0\}^{n-d}$.

The second step is to define Ψ on every $T' \in \Lambda_{\varepsilon,d+1}(Q_1 \cup C_2)$. Without loss of generality we can assume T' centered in 0. We divide T' in pyramids $P_{T,T'}$ with base $T \in \Lambda_{\varepsilon,d}(Q_1 \cup C_2)$ and vertex 0. Assuming $T \subset \{x_{d+1} = -\frac{\varepsilon}{2}, x_{d+2}, \dots, x_n = 0\}$ and $T' \subset \{x_{d+2}, \dots, x_n = 0\}$, we set

$$\Psi|_{P_{T,T'}}(x) = -\frac{2x_{d+1}}{\varepsilon} \Psi|_T \left(-\frac{x}{x_{d+1}} \frac{\varepsilon}{2} \right).$$

We iterate this procedure on all the dimensions till to n , defining it well in $Q_1 \cup C_2$. Since $\Psi|_{\partial(Q_1 \cup C_2)} = \text{Id}$ we can extend the map as the identity outside $Q_1 \cup C_2$. In addition one can easily check that $\Psi \in \mathbf{D}(x, r)$ and thus, since $\widetilde{\Phi}_{\varepsilon, E} \in \mathbf{D}(x, r)$ and the class $\mathbf{D}(x, r)$ is closed by composition, this concludes the proof. \square

Later we will need to implement the above deformation of a set E on a rectangle rather than a cube: the deformation theorem can be proved for very general cubical complexes, [Alm86]; however, for the sake of exposition, we limit ourselves to note the following simple observation: given a closed rectangle $R := R_{x, a, b}$, using a linear map, and covering this time with rectangles homothetic to R , one can easily deduce the same thesis as in Theorem 2.4. The only key point is the area estimate (6), which holds with a constant k_1 depending on the ratio a/b . We will apply this construction to rectangles where the ratio is approximately 2.

3. PROOF OF THEOREM 1.3

Proof of Theorem 1.3. Up to extracting subsequences we can assume the existence of a Radon measure μ on $\mathbb{R}^n \setminus H$ such that

$$\mu_j \xrightarrow{*} \mu, \quad \text{as Radon measures on } \mathbb{R}^n \setminus H, \quad (3.1)$$

where $\mu_j = \mathcal{H}^d \llcorner K_j$. We set $K = \text{spt } \mu \setminus H$ and divide the argument in several steps.

Step one: We show the existence of $\theta_0 = \theta_0(n, d) > 0$ such that

$$\mu(B_{x, r}) \geq \theta_0 \omega_d r^d, \quad x \in \text{spt } \mu \text{ and } r < d_x := \text{dist}(x, H) \quad (3.2)$$

To this end it is sufficient to prove the existence of $\beta = \beta(n, d) > 0$ such that

$$\mu(Q_{x, l}) \geq \beta l^d, \quad x \in \text{spt } \mu \text{ and } l < 2d_x / \sqrt{n}.$$

Let us assume by contradiction that there exist $x \in \text{spt } \mu$ and $l < 2d_x / \sqrt{n}$ such that

$$\frac{\mu(Q_{x, l})^{\frac{1}{d}}}{l} < \beta.$$

We claim that this assumption, for β chosen sufficiently small depending only on d and n , implies that for some $l_\infty \in (0, l)$

$$\mu(Q_{x, l_\infty}) = 0, \quad (3.3)$$

which is a contradiction with the property of x to be a point of $\text{spt } \mu$. In order to prove (3.3), we assume that $\mu(\partial Q_{x, l}) = 0$, which is true for a.e. l .

To prove (3.3) we construct a sequence of nested cubes $Q_i = Q_{x, l_i}$ such that, if β is sufficiently small it holds:

- (i) $Q_0 = Q_{x, l}$;
- (ii) $\mu(\partial Q_{x, l_i}) = 0$;
- (iii) setting $m_i := \mu(Q_i)$ it holds:

$$\frac{m_i^{\frac{1}{d}}}{l_i} < \beta;$$

- (iv) $m_{i+1} \leq (1 - \frac{1}{k_1})m_i$, where k_1 is the constant in Theorem 2.4 (6);
- (v) $(1 - 4\varepsilon_i)l_i \geq l_{i+1} \geq (1 - 6\varepsilon_i)l_i$, where

$$\varepsilon_i := \frac{1}{k\beta} \frac{m_i^{\frac{1}{d}}}{l_i} \quad (3.4)$$

and $k = \max\{6, 6/(1 - (\frac{k_1-1}{k_1})^{\frac{1}{d}})\}$ is a universal constant.

- (vi) $\lim_i m_i = 0$ and $\lim_i l_i > 0$.

Following [DS00], we are going to construct the sequence of cubes by induction: the cube Q_0 satisfies by construction hypotheses (i)-(iii). Suppose that cubes until step i are already defined.

Setting $m_i^j := \mathcal{H}^d(K_j \cap Q_i)$ we cover Q_i with the family $\Lambda_{\varepsilon_i l_i}(Q_i)$ of closed cubes with edge length $\varepsilon_i l_i$ as described in Section 2 and we set C_1^i and C_2^i for the corresponding sets defined in (2.3). We define Q_{i+1} to be the internal cube given by the construction, and we note that C_2^i and Q_{i+1} are non-empty if, for instance,

$$\varepsilon_i = \frac{1}{k\beta} \frac{m_i^{\frac{1}{d}}}{l_i} < \frac{1}{k} \leq \frac{1}{6},$$

which is guaranteed by our choice of k . Observe moreover that $C_1^i \cup C_2^i$ is a strip of width at most $2\varepsilon_i l_i$ around ∂Q_i , hence the side l_{i+1} of Q_{i+1} satisfies $(1 - 4\varepsilon_i)l_i \leq l_{i+1} < (1 - 2\varepsilon_i)l_i$.

Now we apply Theorem 2.4 to Q_i with $E = K_j$ and $\varepsilon = \varepsilon_i l_i$, obtaining the map $\Phi_{i,j} = \Phi_{K_j, \varepsilon_i l_i}$. We claim that, for every j sufficiently large,

$$m_i^j \leq k_1(m_i^j - m_{i+1}^j) + o_j(1). \quad (3.5)$$

Indeed, since (K_j) is a minimizing sequence, by the definition of good class we have that

$$\begin{aligned} m_i^j &\leq m_i + o_j(1) \leq \mathcal{H}^d(\Phi_{i,j}(K_j \cap Q_0)) + o_j(1) \\ &= \mathcal{H}^d(\Phi_{i,j}(K_j \cap Q_{i+1})) + \mathcal{H}^d(\Phi_{i,j}(K_j \cap (C_1^i \cup C_2^i))) + o_j(1) \\ &\leq k_1 \mathcal{H}^d(K_j \cap (C_1^i \cup C_2^i)) + o_j(1) = k_1(m_i^j - m_{i+1}^j) + o_j(1). \end{aligned}$$

The last inequality holds because $\mathcal{H}^d(\Phi_{i,j}(K_j \cap Q_{i+1})) = 0$, otherwise by property (5) of Theorem 2.4 there would exist $T \in \Lambda_{\varepsilon_i l_i, d}^*(Q_{i+1})$ such that $\mathcal{H}^d(\Phi_{i,j}(K_j \cap T)) = \mathcal{H}^d(T)$. Together with property (ii) this would imply

$$l_i^d \varepsilon_i^d = \mathcal{H}^d(T) \leq \mathcal{H}^d(\Phi_{i,j}(K_j) \cap Q_i) \leq k_1 \mathcal{H}^d(K_j \cap Q_i) \leq k_1 m_i^j \rightarrow k_1 m_i$$

and therefore substituting (3.4)

$$\frac{m_i}{k^d \beta^d} \leq k_1 m_i,$$

which is false if β is sufficiently small ($m_i > 0$ because $x \in \text{spt}(\mu)$). Passing to the limit in j in (3.5) we obtain (iv):

$$m_{i+1} \leq \frac{k_1 - 1}{k_1} m_i. \quad (3.6)$$

Since $l_{i+1} \geq (1 - 4\varepsilon_i)l_i$ we can slightly shrink the cube Q_{i+1} to a concentric cube Q'_{i+1} with $l'_{i+1} \geq (1 - 6\varepsilon_i)l_i > 0$, $\mu(\partial Q'_{i+1}) = 0$ and for which (iv) still holds, since just m_{i+1} decreased. With a slight abuse of notation we rename this last cube Q'_{i+1} as Q_{i+1} .

We now show (iii). Using (3.6) and condition (iii) for Q_i we obtain

$$\frac{m_{i+1}^{\frac{1}{d}}}{l_{i+1}} \leq \left(\frac{k_1 - 1}{k_1} \right)^{\frac{1}{d}} \frac{m_i^{\frac{1}{d}}}{(1 - 6\varepsilon_i)l_i} < \left(\frac{k_1 - 1}{k_1} \right)^{\frac{1}{d}} \frac{\beta}{1 - 6\varepsilon_i}$$

The last quantity will be less than β if

$$\left(\frac{k_1 - 1}{k_1} \right)^{\frac{1}{d}} \leq 1 - 6\varepsilon_i = 1 - \frac{6}{k\beta} \frac{m_i^{\frac{1}{d}}}{l_i}. \quad (3.7)$$

In turn inequality (3.7) is true because (iii) holds for Q_i , provided we choose $k \geq 6/(1 - (1 - 1/k_1)^{\frac{1}{d}})$. Furthermore, estimating $\varepsilon_0 < 1/k$ by (iii) and (v) we also have $\varepsilon_{i+1} \leq \varepsilon_i$.

We are left to prove (vi): $\lim_i m_i = 0$ follows directly from (iv); regarding the non degeneracy of the cubes, note that

$$\begin{aligned} \frac{l_\infty}{l_0} &:= \liminf_i \frac{l_i}{l_0} \geq \prod_{i=0}^{\infty} (1 - 6\varepsilon_i) = \prod_{i=0}^{\infty} \left(1 - \frac{6}{k\beta} \frac{m_i^{\frac{1}{d}}}{l_i} \right) \\ &\geq \prod_{i=0}^{\infty} \left(1 - \frac{6m_0^{\frac{1}{d}}}{k\beta l_0 \prod_{h=0}^{i-1} (1 - 6\varepsilon_h)} \left(\frac{k_1 - 1}{k_1} \right)^{\frac{i}{d}} \right) \\ &\geq \prod_{i=0}^{\infty} \left(1 - \frac{6}{k(1 - 6\varepsilon_0)^i} \left(\frac{k_1 - 1}{k_1} \right)^{\frac{i}{d}} \right), \end{aligned}$$

where we used $\varepsilon_h \leq \varepsilon_0$ in the last inequality. Since $\varepsilon_0 < 1/k$ the last product is strictly positive provided

$$k > \frac{6}{1 - \left(\frac{k_1 - 1}{k_1} \right)^{\frac{1}{d}}}$$

which is guaranteed by our choice of k . We conclude that $l_\infty > 0$ which ensures claim (3.3).

Step two: We fix $x \in \text{spt } \mu \setminus H$, and prove that

$$r \mapsto \frac{\mu(B_{x,r})}{r^d} \quad \text{is increasing on } (0, d_x). \quad (3.8)$$

The proof is a straightforward adaptation of the corresponding one in [DGM14, Theorem 2], and amounts to prove a differential inequality for the function $f(r) := \mu(B_{x,r})$. In turn, this inequality is obtained in a two step approximation: first one exploits the rectifiability of the minimizing sequence (K_j) and property (1.1) to compare K_j with the cone competitor $\mathbf{C}_{x,r}(K_j)$, see (2.1). The comparison a priori is only allowed with elements of $\mathcal{P}(H)$, so for almost every $r < d_x$ it holds:

$$\begin{aligned} f_j(r) &= \mathcal{H}^d(K_j) - \mathcal{H}^d(K_j \setminus \overline{B_{x,r}}) \leq m_0 + o_j(1) - \mathcal{H}^d(K_j \setminus \overline{B_{x,r}}) \\ &\leq o_j(1) + \inf_{K' \in \mathcal{P}(H)} \mathcal{H}^d(K') - \mathcal{H}^d(K_j \setminus \overline{B_{x,r}}) \leq o_j(1) + \inf_{\substack{K' \in \mathcal{P}(H) \\ K' \setminus \overline{B_{x,r}} = K_j \setminus \overline{B_{x,r}}}} \mathcal{H}^d(K' \cap \overline{B_{x,r}}), \end{aligned}$$

where $f_j(r) := \mathcal{H}^d(K_j \cap B_{x,r})$ and η_j is infinitesimal. Nevertheless K_j can be compared with its cone competitor, up to an error infinitesimal in j , thanks to Lemma 2.3. We recover

$$\begin{aligned} \inf_{\substack{K' \in \mathcal{P}(H) \\ K' \setminus \overline{B_{x,r}} = K_j \setminus \overline{B_{x,r}}}} \mathcal{H}^d(K' \cap \overline{B_{x,r}}) &\leq o_j(1) + \mathcal{H}^d(\mathbf{C}_{x,r}(K_j) \cap \overline{B_{x,r}}) \\ &\leq o_j(1) + \frac{r}{d} \mathcal{H}^{d-1}(K_j \cap \partial B_{x,r}) = o_j(1) + \frac{r}{d} f'_j(r). \end{aligned}$$

One then passes to the limit in j and obtains the desired monotonicity formula. We refer to [DGM14, Theorem 2] for the conclusion of the proof of (3.8).

Step three: By (3.2) and (3.8) the d -dimensional density of the measure μ , namely:

$$\theta(x) = \lim_{r \rightarrow 0^+} \frac{f(r)}{\omega_d r^d} \geq \theta_0,$$

exists, is finite and positive μ -almost everywhere. Preiss' Theorem 2.1 implies that $\mu = \theta \mathcal{H}^d \llcorner \tilde{K}$ for some countably \mathcal{H}^d -rectifiable set K and some positive Borel function θ . Since K is the support of μ , $\mathcal{H}^d(\tilde{K} \setminus K) = 0$. On the other hand, by differentiation of Hausdorff measures, (3.2) yields $\mathcal{H}^d(K \setminus \tilde{K}) = 0$. Hence K is d -rectifiable and $\mu = \theta \mathcal{H}^d \llcorner K$.

Step four: We prove that $\theta(x) \geq 1$ for every $x \in K$ such that the approximate tangent space to K exists (thus, \mathcal{H}^d -a.e. on K). Fix any such $x \in K \setminus H$ without loss of generality we can suppose that $\pi = \{x_{d+1} = \dots = x_n = 0\}$ is the approximate tangent space to K at x , and that $x = 0$: in particular,

$$\mathcal{H}^d \llcorner \frac{K}{r} \xrightarrow{*} \mathcal{H}^d \llcorner \pi, \quad \text{as } r \rightarrow 0^+.$$

The above convergence, together with the lower density estimates (3.2) imply that, for every $\varepsilon > 0$ there is $\rho > 0$ such that

$$K \cap B_r \subset \left\{ |y_{d+1}|, \dots, |y_n| < \frac{\varepsilon}{2} r \right\} \quad \forall r < \rho. \quad (3.9)$$

Let us now assume, by contradiction, that $\theta(0) < 1$. Thanks to (3.8) and (3.9), there exist $r \in (0, d_x)$ and $\alpha < 1$ such that

$$\frac{\mu(Q_\rho)}{\rho^d} \leq \alpha < 1, \quad \mu(\partial Q_r) = 0, \quad K \cap (Q_\rho \setminus R_{\rho, \varepsilon \rho}) = \emptyset \quad \forall \rho \leq r. \quad (3.10)$$

In particular, since μ_j are weakly converging to μ we get that for j large

$$\frac{\mu_j(Q_r)}{r^d} \leq \alpha < 1 \quad \text{and} \quad \mu_j(Q_r \setminus R_{r, \varepsilon r}) = o_j(1), \quad (3.11)$$

We now wish to clear the small amount of mass appearing in the complement of $R_{r, \varepsilon r}$: we achieve this by repeatedly applying Theorem 2.4. We set $Q_r \cap \{x_{d+1} \geq \frac{\varepsilon}{2} r\} =: R$, and we apply Theorem 2.4 to this rectangle with parameter εr and $E = K_j^0 := K_j$, obtaining the map $\varphi_{1,j}$. We recall that the obtained constant k_1 for the area bound is universal, since it depends on the side ratio of R , which is bounded from below by 1 and from above by 4, provided ε small enough. We set $K_j^1 := \varphi_{1,j}(K_j^0)$ and repeat the argument with $Q_r \cap \{x_{d+1} \leq -\frac{\varepsilon}{2} r\} =: R$ and $E := K_j^1$, obtaining the map $\varphi_{2,j}$. We again set $K_j^2 := \varphi_{2,j}(K_j^1)$ and iterate this procedure to the rectangles $Q_r \cap \{x_{d+2} \geq \frac{\varepsilon}{2} r\}, \dots, Q_r \cap \{x_n \leq -\frac{\varepsilon}{2} r\}$. After $2(n-d)$ iteration, we set

$$K_j^{2(n-d)} := \varphi_{2(n-d),j} \circ \dots \circ \varphi_{1,j}(K_j).$$

We are going to use the cube $Q_{r(1-\sqrt{\varepsilon})}$ because, taking ε small enough, then $\sqrt{\varepsilon} > 4C\varepsilon$, where $C > 1$ is the side ratio considered before. This allows us to claim that

$$\mathcal{H}^d(K_j^{2(n-d)} \cap (Q_{r(1-\sqrt{\varepsilon})} \setminus R_{r(1-\sqrt{\varepsilon}), 6\varepsilon r})) = 0. \quad (3.12)$$

Otherwise there would exist a d -face of a smaller rectangle $T \subset (Q_r \setminus R_{r, \varepsilon r})$ such that

$$\mathcal{H}^d(K_j^{2(n-d)} \cap T) = \mathcal{H}^d(T) \geq \varepsilon^d r^d,$$

which would lead to the following contradiction for j large:

$$\varepsilon^d r^d \leq \mathcal{H}^d(T) \leq \mathcal{H}^d(K_j^{2(n-d)} \cap (Q_r \setminus R_{r, \varepsilon r})) \leq k_1^{2(n-d)} \mathcal{H}^d(K_j \cap (Q_r \setminus R_{r, \varepsilon r})) = o_j(1)$$

In particular, we cleared any measure on every slab

$$\bigcup_{i=d+1}^n \left\{ 3\varepsilon r < |x_i| < (1-\sqrt{\varepsilon}) \frac{r}{2} \right\} \cap Q_{r(1-\sqrt{\varepsilon})}.$$

We want now to construct a map $P \in \mathbf{D}(0, r)$, collapsing $R_{r(1-\sqrt{\varepsilon}), 6\varepsilon r}$ onto the tangent plane. To this end, for $x \in \mathbb{R}^n$, $x = (x', x'')$ with $x' \in \mathbb{R}^d$ and $x'' \in \mathbb{R}^{n-d}$ we set

$$\|x'\| := \max\{|x_i| : i = 1, \dots, d\} \quad \|x''\| := \max\{|x_i| : i = d+1, \dots, n\} \quad (3.13)$$

and we define P as follows:

$$P(x) = \begin{cases} (x', g(\|x'\|) \frac{(\|x''\| - 3\varepsilon)_+}{1 - 6\varepsilon} \frac{x''}{\|x''\|} + (1 - g(\|x'\|))x'') & \text{if } \max\{\|x'\|, \|x''\|\} \leq r/2 \\ \text{Id} & \text{otherwise,} \end{cases} \quad (3.14)$$

where $g : [0, r/2] \rightarrow [0, 1]$ is a compactly supported cut off function such that

$$g \equiv 1 \quad \text{on } [0, r(1 - \sqrt{\varepsilon})/2] \quad \text{and} \quad |g'| \leq 10/r\sqrt{\varepsilon}.$$

It is not difficult to check that $P \in D(0, r)$ and that $\text{Lip } P \leq 1 + C\sqrt{\varepsilon}$ for some dimensional constant C .

We now set $\widetilde{K}_j := P(K_j^{2(n-d)})$, which verifies, thanks to (3.12),

$$\mathcal{H}^d(\widetilde{K}_j \cap (Q_{(1-\sqrt{\varepsilon})r} \setminus Q_{(1-\sqrt{\varepsilon})r}^d)) = 0 \quad (3.15)$$

and

$$\begin{aligned} \mathcal{H}^d(\widetilde{K}_j \cap (Q_r \setminus Q_{r(1-\sqrt{\varepsilon})})) &\leq (1 + C\sqrt{\varepsilon})^d \mathcal{H}^d(K_j^{2(n-d)} \cap (Q_r \setminus Q_{r(1-\sqrt{\varepsilon})})) \\ &\leq (1 + C\sqrt{\varepsilon}) k_1^{2(n-d)} \mathcal{H}^d(K_j \cap (Q_r \setminus (Q_{r(1-\sqrt{\varepsilon})} \cup R_{r,\varepsilon}))) \\ &\quad + (1 + C\sqrt{\varepsilon}) \mathcal{H}^d(K_j \cap (R_{r,\varepsilon} \setminus Q_{r(1-\sqrt{\varepsilon})})) \\ &\leq o_j(1) + (1 + C\sqrt{\varepsilon}) \mathcal{H}^d(K_j \cap (R_{r,\varepsilon} \setminus Q_{r(1-\sqrt{\varepsilon})})), \end{aligned} \quad (3.16)$$

where in the last inequality we have used (3.11). Moreover, by using (3.10), (3.11) and (3.15) we also have that, for ε small and j large:

$$\begin{aligned} \frac{\mathcal{H}^d(\widetilde{K}_j \cap Q_{r(1-\sqrt{\varepsilon})}^d)}{r^d(1 - \sqrt{\varepsilon})^d} &= \frac{\mathcal{H}^d(\widetilde{K}_j \cap Q_{r(1-\sqrt{\varepsilon})})}{r^d(1 - \sqrt{\varepsilon})^d} \leq (1 + C\sqrt{\varepsilon}) \frac{\mathcal{H}^d(K_j^{2(n-d)} \cap Q_r)}{r^d} \\ &\leq (1 + C\sqrt{\varepsilon}) \frac{\mathcal{H}^d(K_j \cap Q_r) + o_j(1)}{r^d} \\ &\leq \alpha + o_j(1) < 1. \end{aligned} \quad (3.17)$$

As a consequence of (3.17) and the compactness of \widetilde{K}_j , there exist $y'_j \in Q_{(1-\sqrt{\varepsilon})r}^d$ and $\delta_j > 0$ such that, if we set $y_j := (y'_j, 0)$, then

$$\widetilde{K}_j \cap B_{y_j, \delta_j}^d = \emptyset \quad \text{and} \quad B_{y_j, \delta_j}^d \subset Q_{(1-\sqrt{\varepsilon})r}^d. \quad (3.18)$$

After the last deformation, our set $\widetilde{K}_j \cap Q_{r(1-\sqrt{\varepsilon})}$ lives on the tangent plane and we want to use the property (3.18) to collapse $\widetilde{K}_j \cap Q_{r(1-\sqrt{\varepsilon})}$ into $(\partial Q_{(1-\sqrt{\varepsilon})r}^d) \times \{0\}^{n-d}$. To this end let us define for every $j \in \mathbb{N}$, the following Lipschitz map:

$$\varphi_j(x) = \begin{cases} (x' + z'_{j,x}, x'') & \text{if } x \in R_{r(1-\sqrt{\varepsilon}), r} \\ x & \text{otherwise,} \end{cases}$$

with

$$z'_{j,x} := \min \left\{ 1, \frac{|x' - y'_j|}{\delta_j} \right\} \frac{(r - 4\|x''\|)_+}{r} \gamma_{j,x}(x' - y'_j),$$

where $\gamma_{j,x} > 0$ is such that $x' + \gamma_{j,x}(x' - y'_j) \in \partial Q_{(1-\sqrt{\varepsilon})r}^d \times \{0\}^{n-d}$ and $\|x''\|$ is defined in (3.13).

One can easily check that $\varphi_j \in D(0, r)$. Moreover, setting $\varphi_j(\widetilde{K}_j) =: K'_j$ we have that

$$K'_j \setminus Q_r = K_j \setminus Q_r$$

and

$$\mathcal{H}^d(K'_j \cap Q_{r(1-\sqrt{\varepsilon})}) = 0, \quad (3.19)$$

thanks to (3.15), since

$$\mathcal{H}^d\left(\partial Q_{(1-\sqrt{\varepsilon})r}^d \times \{0\}^{n-d}\right) = 0.$$

Since $\mathcal{P}(H)$ is a good class, by (1.1) there exists a sequence of competitors $(J_j)_{j \in \mathbb{N}} \subset \mathcal{P}(H)$ such that $J_j \setminus \overline{B}_{0,r} = K_j \setminus \overline{B}_{0,r}$ and $\mathcal{H}^d(J_j) = \mathcal{H}^d(K'_j) + o_j(1)$. Hence, thanks to (3.16) and (3.19) we get

$$\begin{aligned} \mathcal{H}^d(K_j) - \mathcal{H}^d(J_j) &\geq \mathcal{H}^d(K_j) - \mathcal{H}^d(K'_j) - o_j(1) = \mathcal{H}^d(K_j \cap Q_r) - \mathcal{H}^d(K'_j \cap Q_r) - o_j(1) \\ &\geq \mathcal{H}^d\left(K_j \cap Q_{r(1-\sqrt{\varepsilon})}\right) + \mathcal{H}^d\left(K_j \cap (R_{r,\varepsilon r} \setminus Q_{r(1-\sqrt{\varepsilon})})\right) + \\ &\quad - o_j(1) - (1 + C\sqrt{\varepsilon})\mathcal{H}^d\left(K_j \cap (R_{r,\varepsilon r} \setminus Q_{r(1-\sqrt{\varepsilon})})\right) \\ &\geq \mathcal{H}^d\left(K_j \cap Q_{r(1-\sqrt{\varepsilon})}\right) - C\sqrt{\varepsilon}\mathcal{H}^d\left(K_j \cap (R_{r,\varepsilon r} \setminus Q_{r(1-\sqrt{\varepsilon})})\right) - o_j(1). \end{aligned}$$

Passing to the limit as $j \rightarrow \infty$, and using (3.1), (3.2) and (3.10) we get that

$$\begin{aligned} \liminf_j \mathcal{H}^d(K_j) &\geq \liminf_j \mathcal{H}^d(J_j) + \mu(Q_{r(1-\sqrt{\varepsilon})}) - C\sqrt{\varepsilon}r^d \\ &\geq \liminf_j \mathcal{H}^d(J_j) + (\theta_0(1 - \sqrt{\varepsilon})^d - C\sqrt{\varepsilon})r^d. \end{aligned}$$

Since, for ε small, this is in contradiction with K_j be a minimizing sequence we finally conclude that $\theta(0) \geq 1$.

Step five: We prove that $\theta(x) \leq 1$ for every $x \in K$ such that the approximate tangent space to K exists. Arguing by contradiction, we assume that $\theta(x) = 1 + \sigma > 1$ for some x where K admits an approximate tangent plane T . As usually we assume that $x = 0$ and $T = \{y : y_{d+1}, \dots, y_n = 0\}$. By the monotonicity of the density established in Step 2, for every $\varepsilon > 0$ we can find $r_0 > 0$ such that

$$K \cap Q_r \subset R_{r,\varepsilon r}, \quad 1 + \sigma \leq \frac{\mu(Q_r)}{r^d} \leq 1 + \sigma + \varepsilon\sigma, \quad \forall r \leq r_0. \quad (3.20)$$

Let us also note that for every $r < r_0$ there exists j_0 such that

$$\mathcal{H}^d(K_j \cap Q_r) > \left(1 + \frac{\sigma}{2}\right)r^d, \quad \mathcal{H}^d((K_j \cap Q_r) \setminus R_{r,\varepsilon r}) < \frac{\sigma}{4}r^d, \quad \forall j \geq j_0, \quad (3.21)$$

Consider the map $P : \mathbb{R}^n \rightarrow \mathbb{R}^n \in \mathcal{D}(0, r)$ with $\text{Lip } P \leq 1 + C\sqrt{\varepsilon}$ defined in (3.14) which collapses $R_{(r(1-\sqrt{\varepsilon}), \varepsilon r)}$ onto the tangent plane. By exploiting the fact the $\mathcal{P}(H)$ is a good class we find that

$$\begin{aligned} \mathcal{H}^d(K_j \cap Q_r) - o_j(1) &\leq \underbrace{\mathcal{H}^d(P(K_j \cap R_{(1-\sqrt{\varepsilon})r, \varepsilon r}))}_{I_1} + \underbrace{\mathcal{H}^d(P(K_j \cap (R_{r,\varepsilon r} \setminus R_{(1-\sqrt{\varepsilon})r, \varepsilon r})))}_{I_2} \\ &\quad + \underbrace{\mathcal{H}^d(P(K_j \cap (Q_r \setminus R_{r,\varepsilon r})))}_{I_3}. \end{aligned}$$

By construction, $I_1 \leq r^d$, while, by (3.21), $\mathcal{H}^d(K_j \cap Q_r) > (1 + (\sigma/2))r^d$ and

$$I_3 \leq (\text{Lip } P)^d \mathcal{H}^d(K_j \cap (Q_r \setminus R_{r,\varepsilon r})) < (1 + C\sqrt{\varepsilon})^d \frac{\sigma}{4}r^d.$$

Hence, as $j \rightarrow \infty$,

$$\left(1 + \frac{\sigma}{2}\right)r^d \leq r^d + \liminf_{j \rightarrow \infty} I_2 + (1 + C\sqrt{\varepsilon})^d \frac{\sigma}{4}r^d,$$

that is,

$$\left(\frac{1}{2} - \frac{(1 + C\sqrt{\varepsilon})^d}{4}\right)\sigma \leq \liminf_{j \rightarrow \infty} \frac{I_2}{r^d}. \quad (3.22)$$

By (3.20), we finally estimate that

$$\begin{aligned} \limsup_{j \rightarrow \infty} I_2 &\leq (1 + C\sqrt{\varepsilon})^d \mu(Q_r \setminus Q_{(1-\sqrt{\varepsilon})r}) \\ &\leq (1 + C\sqrt{\varepsilon})^d \left((1 + \sigma + \varepsilon\sigma) - (1 + \sigma)(1 - \sqrt{\varepsilon})^d \right) r^d \end{aligned} \quad (3.23)$$

By choosing ε sufficiently small, (3.22) and (3.23) provide the desired contradiction. Hence $\theta \leq 1$ and, combining this with the previous step, $\mu = \mathcal{H}^d \llcorner K$.

Step six: We now show that the canonical density one rectifiable varifold associated K is stationary in $\mathbb{R}^n \setminus H$. In particular, applying Allard's regularity theorem, see [Sim83, Chapter 5], we will deduce that there exists an \mathcal{H}^d -negligible closed set $\Sigma \subset K$ such that $\Gamma = K \setminus \Sigma$ is a real analytic manifold. Since being a stationary varifold is a local property, to prove our claim it is enough to show that for every ball $B \subset\subset \mathbb{R}^n \setminus H$ we have

$$\mathcal{H}^d(K) \leq \mathcal{H}^d(\phi(K)) \quad (3.24)$$

whenever ϕ is a diffeomorphism such that $\text{spt}\{\phi - \text{Id}\} \subset B$. Indeed, by exploiting (3.24) with $\phi_t = \text{Id} + tX$, $X \in C_c^1(B)$ we deduce the desired stationarity property.

To prove (3.24) we argue as in [DGM14, Theorem 7]. Given $\varepsilon > 0$ we can find $\delta > 0$ and a compact set $\hat{K} \subset K \cap B$ with $\mathcal{H}^d(K \setminus \hat{K}) \cap B < \varepsilon$ such that K admits an approximate tangent plane $\pi(x)$ at every $x \in \hat{K}$,

$$\sup_{x \in \hat{K}} \sup_{y \in B_{x,\delta}} |\nabla \phi(x) - \nabla \phi(y)| \leq \varepsilon, \quad \sup_{x \in \hat{K}} \sup_{y \in \hat{K} \cap B_{x,\delta}} d(\pi(x), \pi(y)) < \varepsilon, \quad (3.25)$$

where d is a distance on $G(d)$, the d -dimensional Grassmanian. Moreover, denoting by $S_{x,r}$ the set of points in $B_{x,r}$ at distance at most εr from $x + \pi(x)$, then $K \cap B_{x,r} \subset S_{x,r}$ for every $r < \delta$ and $x \in \hat{K}$. By Besicovitch covering theorem we can find a finite disjoint family of closed balls $\{\bar{B}_i\}$ with $B_i = B_{x_i, r_i} \subset B \subset\subset \mathbb{R}^n \setminus H$, $x_i \in \hat{K}$, and $r_i < \delta$, such that $\mathcal{H}^d(\hat{K} \setminus \bigcup_i B_i) < \varepsilon$. By exploiting the construction of Step four, we can find $j(\varepsilon) \in \mathbb{N}$ and maps $P_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\text{Lip}(P_i) \leq 1 + C\sqrt{\varepsilon}$ and $P_i = \text{Id}$ on B_i^c , such that, for a certain $X_i \subset S_i = S_{x_i, \varepsilon r_i}$,

$$\begin{aligned} P_i(X_i) &\subset B_i \cap (x_i + \tau(x_i)), \\ \mathcal{H}^d\left(P_i((K_j \cap B_i) \setminus X_i)\right) &\leq C\sqrt{\varepsilon} \omega_d r_i^d, \quad \forall j \geq j(\varepsilon). \end{aligned} \quad (3.26)$$

Exploiting (3.25), (3.26), the area formula and that $\omega_d r_i^d \leq \mathcal{H}^d(K \cap B_i)$ (by the monotonicity formula), and setting $\alpha_i = \mathcal{H}^d((K \setminus \hat{K}) \cap B_i)$, we get (denoting with J_d^π the d -dimensional tangential jacobian with respect to the plane π and by J_d^K the one with respect to K)

$$\begin{aligned} \mathcal{H}^d(\phi(P_i(K_j \cap X_i))) &= \int_{P_i(K_j \cap X_i)} J_d^{\pi(x_i)} \phi(x) d\mathcal{H}^d(x) \leq (J^{\pi(x_i)} \phi(x_i) + \varepsilon) \omega_d r_i^d \\ &\leq (J_d^{\pi(x_i)} \phi(x_i) + \varepsilon) \mathcal{H}^d(K \cap B_i) \leq (J_d^{\pi(x_i)} \phi(x_i) + \varepsilon) (\mathcal{H}^d(\hat{K} \cap B_i) + \alpha_i) \\ &\leq \int_{\hat{K} \cap B_i} (J_d^K \phi(x) + 2\varepsilon) d\mathcal{H}^d(x) + ((\text{Lip } \phi)^d + \varepsilon) \alpha_i \\ &= \mathcal{H}^d(\phi(\hat{K} \cap B_i)) + 2\varepsilon \mathcal{H}^d(K \cap B_i) + ((\text{Lip } \phi)^d + \varepsilon) \alpha_i, \end{aligned} \quad (3.27)$$

where in the last identity we have used the area formula and the injectivity of ϕ . Since $P_i = \text{Id}$ on B_i^c , $\phi = \text{Id}$ on B^c , $B_i \subset B$ and the balls B_i are disjoint, the map $\tilde{\phi}$ which is equal to ϕ on $B \setminus \bigcup_i B_i$, equal to the identity on B^c and equal to $\phi \circ P_i$ on B_i is well defined. Moreover, by (3.27), we get

$$\mathcal{H}^d(\tilde{\phi}(K_j)) \leq \mathcal{H}^d(\phi(K)) + C\varepsilon$$

where C depends only on K . By exploiting the definition of good class, we get that

$$\mathcal{H}^d(K) \leq \mathcal{H}^d(\tilde{\phi}(K_j)) + \eta_j \leq \mathcal{H}^d(\phi(K)) + C\varepsilon + o_j(1).$$

Letting $j \rightarrow \infty$ and $\varepsilon \rightarrow 0$ we obtain (3.24).

Step seven: We finally address the dimension of the singular set. Recall that, by monotonicity, the density function

$$\Theta^d(K, x) = \lim_{r \rightarrow 0} \frac{\mathcal{H}^d(K \cap B_{x,r})}{\omega_d r^d}$$

is everywhere defined in $\mathbb{R}^n \setminus H$ and equals 1 \mathcal{H}^d -almost everywhere in K . Fixing $x \in K$ and a sequence $r_k \downarrow 0$, the monotonicity formula, the stationarity of $\mathcal{H}^d \llcorner K$ and the compactness theorem for integral varifolds [All72, Theorem 6.4] imply that (up to subsequences)

$$\mathcal{H}^d \llcorner \left(\frac{K-x}{r_k} \right) \rightarrow V \quad \text{locally in the sense of varifolds,} \quad (3.28)$$

with

- (a) V is a stationary integral varifold: in particular $\Theta^d(\|V\|, y) \geq 1$ for $y \in \text{spt}(V)$;
- (b) V is a cone, namely $(\delta_\lambda)_\# V = V$, where $\delta_\lambda(x) = \lambda x$, $\lambda > 0$;
- (c) $\Theta^d(\|V\|, 0) = \Theta^d(K, x) \geq \Theta^d(\|V\|, y)$ for every $y \in \mathbb{R}^n$.

Recall that the tangent varifold V depends (in principle) on the sequence (r_k) . We denote by $\text{TanVar}(K, x)$ the (nonempty) set of all possible limits V as in (3.28) varying among all sequences along which (3.28) holds. Given a cone W we set

$$\text{Spine}(W) := \{y \in \mathbb{R}^n : \Theta^d(\|W\|, y) = \Theta^d(\|W\|, 0)\} : \quad (3.29)$$

by [Alm00, 2.26] $\text{Spine}(W)$ is a vector subspace of \mathbb{R}^n , see also [Whi97]. We can stratify K in the following way: for every $k = 0, \dots, n$ we let

$$A_k := \{x \in K : \text{for all } V \in \text{TanVar}(K, x), \dim \text{Spine}(V) \leq k\}.$$

Clearly $A_0 \supset \dots \supset A_{d+1} = \dots = A_n = \emptyset$; moreover it holds: $\dim_{\mathcal{H}} A_k \leq k$, see [Alm00, 2.28]. In order to prove our claim, we need to show that $A_d \setminus A_{d-1} \subset K \setminus \Sigma$, where Σ as in Step six is the singular set of K , namely that every point in K having at least one tangent cone of maximal dimension d must be regular (note that, as the example of complex varieties shows, this is not true in general for a stationary varifold).

First, note that if $x \in A_d \setminus A_{d-1}$ and $V \in \text{TanVar}(K, x)$ satisfies $\dim \text{Spine}(V) = d$, then $V = \mathcal{H}^d \llcorner \text{Spine}(V)$: indeed up to a rotation $\text{spt}(V) = \text{Spine}(V) \times \Gamma$, where Γ is a cone in \mathbb{R}^{n-d} . If $\Gamma \neq \{0\}$ then $\Theta^d(\|V\|, 0) > \Theta^d(\|V\|, y)$ for any $y \in \text{Spine}(V) \setminus \{0\}$, which contradicts (3.29). Hence by (3.29) and (c), $\Theta^d(\|V\|, 0)$ is an integer and $V = \Theta^d(K, x) \mathcal{H}^d \llcorner \text{Spine}(V)$ (see also [Alm00, Theorem 2.26 (4)]). Second, the density lower bound $\Theta^d(K, \cdot) \geq 1$ and (3.28) imply that for every $\varepsilon > 0$

$$\frac{K-x}{r} \subset U_\varepsilon(\text{Spine}(V)) \quad \text{and} \quad \mathcal{H}^d \left(\frac{K-x}{r} \cap Q_{0,1} \right) \leq (1+\varepsilon) \Theta^d(K, x)$$

if r is sufficiently small. By arguing as in Step five (i.e. roughly speaking comparing K with $P(K)$ in $Q_{x,r}$, where P is the squeezing map (3.14) although one has to rigorously get through the minimizing sequence K_j) we obtain

$$\mathcal{H}^d \left(\frac{K-x}{r} \cap Q_{0,1} \right) \leq (1 + C\sqrt{\varepsilon}).$$

Letting $r \downarrow 0$ thanks to (3.28) we obtain $\|V\|(Q_{0,1}) \leq (1 + C\sqrt{\varepsilon})$, implying $\Theta^d(K, x) = 1$. We therefore fall into the hypotheses of Allard's regularity Theorem [All72, Regularity Theorem, Section 8], $K \cap Q_{x, \frac{r}{2}}$ is a real analytic submanifold. Equivalently $x \notin \Sigma$. \square

4. PROOF OF THEOREMS 1.5 AND 1.8

In this Section we prove Theorem 1.5 and 1.8. With Theorem 1.3 at hand, the proofs are quite similar to the corresponding ones in [DGM14] (see Theorems 4 and 7 there), hence we limit ourselves to provide a short sketch underlying only the main differences.

Proof of Theorem 1.5. We start by proving that $\mathcal{F}(H, \mathcal{C})$ is a good class in the sense of Definition 1.2: let $\tilde{K} \in \mathcal{F}(H, \mathcal{C})$, $x \in K$, $r \in (0, \text{dist}(x, H))$ and $\varphi \in \mathcal{D}(x, r)$. We show that $\varphi(\tilde{K}) \in \mathcal{F}(H, \mathcal{C})$ arguing by contradiction: assume that $\gamma(S^{n-d}) \cap \varphi(\tilde{K}) = \emptyset$ for some $\gamma \in \mathcal{C}$ and, without loss of generality, suppose also that $\gamma(S^{n-d}) \cap (\tilde{K} \setminus B_{x,r}) = \emptyset$. By Definition 1.1 there exists a sequence

$$(\varphi_j) \subset \mathcal{D}(x, r) \quad \text{such that} \quad \lim_j \|\varphi_j - \varphi\|_{C^0} = 0.$$

Since $\gamma(S^{n-d})$ is compact and $\varphi_j = \text{Id}$ outside $B_{x,r}$, for j sufficiently large $\gamma(S^{n-d}) \cap \varphi_j(\tilde{K}) = \emptyset$; moreover φ_j is invertible, hence $\varphi_j^{-1}(\gamma(S^{n-d})) \cap \tilde{K} = \emptyset$. But the property for φ_j of being isotopic to the identity implies $\varphi_j^{-1} \circ \gamma \in \mathcal{C}$, which contradicts $V \in \mathcal{F}(H, \mathcal{C})$. This proves (a).

Given a minimizing sequence $(K_j) \subset \mathcal{F}(H, \mathcal{C})$ which consists of rectifiable sets, we can therefore find a set K with the properties stated in Theorem 1.3. In order to conclude (b), namely that $K \in \mathcal{F}(H, \mathcal{C})$, we refer to [DGM14, Theorem 4(b)]: the proof is the same. \square

Proof of Theorem 1.8. As already observed in Remark 1.7, $\mathcal{A}(H, K_0)$ is a good class and we can therefore apply Theorem 1.3. We thus know that $\mathcal{H}^d \llcorner K_j \xrightarrow{*} \mu = \mathcal{H}^d \llcorner K$ and that K is a smooth set away from H and from a relatively closed set Σ of dimension less or equal than $(d-1)$. The conclusion of the proof can now be obtained by repeating verbatim Steps 4 and 6 in the proof of Theorem 7 in [DGM14]. \square

BIBLIOGRAPHY

- [All72] W. K. Allard. On the first variation of a varifold. *Ann. Math.*, 95:417–491, 1972.
- [All75] W. K. Allard. On the first variation of a varifold: boundary behaviour. *Ann. Math.*, 101:418–446, 1975.
- [Alm68] F. J. Jr Almgren. Existence and regularity almost everywhere of solutions to elliptic variational problems among surfaces of varying topological type and singularity structure. *Ann. Math.*, 87:321–391, 1968.
- [Alm76] F. J. Jr. Almgren. Existence and regularity almost everywhere of solutions to elliptic variational problems with constraints. *Mem. Amer. Math. Soc.*, 4(165):viii+199 pp, 1976.
- [Alm86] F. Almgren. Deformations and multiple-valued functions. Geometric measure theory and the calculus of variations. pages 29–130, 1986.
- [Alm00] F. J. Jr. Almgren. *Almgren's big regularity paper. Q-valued functions minimizing Dirichlet's integral and the regularity of area-minimizing rectifiable currents up to codimension 2*, volume 1 of *World Scientific Monograph Series in Mathematics*. World Scientific Publishing Co., Inc., River Edge, NJ, 2000. xvi+955 pp. pp. With a preface by Jean E. Taylor and Vladimir Scheffer.
- [Dav13] G. David. Regularity of minimal and almost minimal sets and cones: J. Taylor's theorem for beginners. In *Analysis and geometry of metric measure spaces*, volume 56 of *CRM Proc. Lecture Notes*, pages 67–117. Amer. Math. Soc., Providence, RI, 2013.
- [Dav14] G. David. Should we solve Plateau's problem again? In C. Fefferman; A. D. Ionescu; D.H. Phong; S. Wainger, editor, *Advances in Analysis: the legacy of Elias M. Stein*. Princeton University Press, 2014.

- [DS00] G. David and S. Semmes. Uniform rectifiability and quasiminimizing sets of arbitrary codimension. pages viii+132 pp., 2000.
- [De54] E. De Giorgi. Su una teoria generale della misura $(r - 1)$ -dimensionale in uno spazio ad r -dimensioni. *Ann. Mat. Pura Appl. (4)*, 36:191–213, 1954.
- [De55] E. De Giorgi. Nuovi teoremi relativi alle misure $(r - 1)$ -dimensionali in uno spazio ad r -dimensioni. *Ricerche Mat.*, 4:95–113, 1955.
- [DeL08] C. De Lellis. *Rectifiable sets, densities and tangent measures*. Zurich Lectures in Advanced Mathematics. European Mathematical Society, Zürich, 2008. vi+127 pp.
- [DGM14] C. De Lellis, F. Ghiraldin, and F. Maggi. A direct approach to Plateau’s problem. *JEMS*. To appear. URL [arXiv:1408.4047](https://arxiv.org/abs/1408.4047).
- [Fed69] H. Federer. *Geometric measure theory*, volume 153 of *Die Grundlehren der mathematischen Wissenschaften*. Springer-Verlag New York Inc., New York, 1969. xiv+676 pp pp.
- [FF60] H. Federer and W. H. Fleming. Normal and integral currents. *Ann. Math.*, 72:458–520, 1960.
- [Har14] J. Harrison. Soap film solutions to Plateau’s problem. *J. Geom. Anal.*, 24(1):271–297, 2014. ISSN 1050-6926.
- [Hir94] M. W. Hirsch. *Differential topology*, volume 33 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. ISBN 0-387-90148-5. x+222 pp. Corrected reprint of the 1976 original.
- [HP13] J. Harrison and H. Pugh. Existence and soap film regularity of solutions to Plateau’s problem. 2013. [arXiv:1310.0508](https://arxiv.org/abs/1310.0508).
- [Pre87] D. Preiss. Geometry of measures in \mathbf{R}^n : distribution, rectifiability, and densities. *Ann. of Math. (2)*, 125(3):537–643, 1987. ISSN 0003-486X.
- [Rei60] E. R. Reifenberg. Solution of the Plateau problem for m -dimensional surfaces of varying topological type. *Acta Math.*, 104:1–92, 1960.
- [Sim83] L. Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis*. Australian National University, Centre for Mathematical Analysis, Canberra, 1983. vii+272 pp.
- [Whi97] Brian White. Stratification of minimal surfaces, mean curvature flows, and harmonic maps. *J. Reine Angew. Math.*, 488:1–35, 1997.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH, SWITZERLAND
E-mail address: guido.dephilippis@math.uzh.ch

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH, SWITZERLAND
E-mail address: antonio.derosa@math.uzh.ch

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH, SWITZERLAND
E-mail address: francesco.ghiraldin@math.uzh.ch