The Logic of Counting Query Answers

Hubie Chen

Departamento LSI, Universidad del País Vasco and IKERBASQUE, Basque Foundation for Science E-20018 San Sebastián, Spain Stefan Mengel CNRS CRIL UMR 8188 France

Abstract—We consider the problem of counting the number of answers to a first-order formula on a finite structure. We present and study an extension of first-order logic in which algorithms for this counting problem can be naturally and conveniently expressed, in senses that are made precise and that are motivated by the wish to understand tractable cases of the counting problem.

I. INTRODUCTION

A. Overview

The computational problem of evaluating a logical formula on a finite relational structure is of central interest in database theory and logic. In the context of database theory, this problem is often referred to as *query evaluation*, as it models the posing of a query to a database, in a well-acknowledged way: the formula represents the query, and the structure represents the database. We refer to the results of such an evaluation as *answers*; logically, these are the satisfying assignments of the formula on the structure. The particular case of this problem where the formula is a sentence is known as *model checking*. In the sequel, we assume by default that (unless otherwise specified) all formulas and sentences under discussion are firstorder and relational.

This article concerns the problem of *counting query answers*: given a first-order formula and a finite structure, output the *number* of answers (for previous studies, see for example the works [1], [2], [3], [4]). This problem is a counting version of query evaluation, and generalizes model checking, which can be viewed as the particular case thereof where one is given a sentence and structure, and wants to decide if the number of answers is 1 or 0, corresponding to whether or not the empty assignment is satisfying. Motivation for studying this counting problem stems both from basic and fundamental interest, and from application scenarios: all practical query languages supported by database management systems have a counting operator, and it has indeed been argued [2] that database queries with counting are at the basis of decision support systems that handle large data volume.

With a first-order formula ϕ in hand, if one is interested in counting the *number* of answers to ϕ on given structures, it is natural to inquire if there is a language or logic in which one can directly express the mapping that provides, for each structure, the number of answers to ϕ . Such a logic could serve as a target language into which first-order formulas of interest (in the mentioned sense) could be compiled, and then optimized, rewritten, and evaluated. This article presents and studies such a logic, \sharp -logic, wherein the evaluation of a sentence on a structure yields an integer value. From the database-theoretic viewpoint, our presentation of \sharp -logic amounts to the introduction of a query language designed particularly for counting answers. We show that \sharp -logic enjoys and balances the following properties.

- Expressivity. In a sense made precise, #-logic allows for the expression of known efficient algorithms for tractable cases of the counting query answers problem. Moreover, this expression is (in our view) direct and clean, and illustrates that #-logic captures precisely the key computational primitives required by these algorithms; this capture, in turn, justifies the particular definition of #logic.
- Optimizability. Minimizing a crucial measure known as width can be performed computably in an expressive fragment of #-logic; this amounts to the fragment supporting an optimal form of query optimization, relative to this quantity.

Our hope is that this article will contribute to and invigorate a broader investigation of query languages for counting answers. Such an investigation could address issues such as the identification of desirable theoretical properties of such query languages, and techniques for performing query rewriting, optimization, and evaluation.

B. Background: complexity

As has been previously articulated in the literature, a typical situation in the database setting is the evaluation of a relatively short formula on a relatively large structure. Consequently, it has been argued that, in measuring the time complexity of query evaluation tasks, one could reasonably allow a slow (non-polynomial-time) preprocessing of the formula, so long as the desired evaluation can be performed in polynomial time following the preprocessing [5], [6]. Relaxing polynomial-time computation to allow arbitrary preprocessing of a parameter of a problem instance yields, in essence, the notion of *fixed*parameter tractability. This notion is at the core of and is the primary tractability notion in *parameterized complexity* theory, which provides a taxonomy for classifying problems where each instance has an associated parameter. Following this motivation, whenever the problem of counting query answers (or the model checking problem) is considered using parameterized complexity, in this article, the formula is taken to be the parameter.

The problem of counting query answers is well-known to be computationally intractable. It is possible to restrict this problem by considering restricted classes of queries, and then trying to understand which classes of queries are computationally well-behaved in that they give rise to a tractable case of the general problem. Precisely, for a class Φ of first-order formulas, define $count(\Phi)$ to be the problem where an instance is a formula $\phi \in \Phi$ paired with a finite structure **B**, and the output is the number of answers of ϕ on **B**. We hence have a family of problems, one problem $count(\Phi)$ for each such formula class Φ , and one can inquire which of these problems are tractable (and which are not). We will also have cause to consider the case of model checking, so, when Φ is a class of first-order sentences, define $MC(\Phi)$ to be the *model checking* problem where an instance is a sentence $\phi \in \Phi$ paired with a finite structure B, and the output is yes or no depending on whether or not **B** satisfies ϕ . The complexity of the problem family $MC(\Phi)$ has been considered in numerous papers, such as [7], [8], [9], [10], [11], [12], [13].

C. Background: width

Width is a syntactic measure of logical formulas. The width of a first-order formula ϕ is defined as the maximum number of free variables over all subformulas of ϕ . In studies of the problem family MC(Φ), width has emerged as a crucially relevant measure; we now explain how.

Say that a class Φ of first-order formulas has *bounded* width if there exists a constant $k \ge 1$ such that each $\phi \in \Phi$ has width at most k. It is now well-known that bounded width sentence classes are computationally desirable for model checking, made precise as follows.

Observation 1.1: [14], [15] Suppose that Φ is a class of sentences having bounded width. The computational problem $MC(\Phi)$ is polynomial-time decidable, via the algorithm that (given a sentence and a structure) simply computes the set of satisfying assignments for each subformula, inductively.

As suggested above, it is known that a problem is fixedparameter tractable if, after performing preprocessing on the parameter, an instance can be resolved in polynomial time [6, Theorem 1.37]. For a problem $MC(\Phi)$, one can readily observe that if each sentence $\phi \in \Phi$ can be algorithmically translated to a sentence lying in a class having bounded width, then $MC(\Phi)$ is fixed-parameter tractable as a consequence of Observation 1.1. This can be formalized as follows.

Observation 1.2: [11] Suppose that Φ is a sentence class. The following condition, which we will refer to as the *classical* condition, is sufficient for the problem $MC(\Phi)$ to be fixedparameter tractable: there exists a sentence class Φ' having bounded width and an algorithm f that computes, for each $\phi \in \Phi$, a logically equivalent sentence $f(\phi)$ that is in Φ' .

Research on the problem family $MC(\Phi)$ has succeeded in obtaining classifications on classes Φ of bounded arity where the quantifiers and connectives are restricted (see for examples [7], [10], [11]). An example relevant to the present article is a study [11] of existential positive logic (by which we mean the positive fragment of first-order logic consisting of formulas built from atoms, \land , \lor , and \exists); making crucial use of a hardness result by Grohe [7], this work observed that when Φ is a class of existential positive sentences having bounded arity, the problem MC(Φ) is fixed-parameter tractable if and only if the classical condition applies to Φ . (Here and elsewhere in our discussion, we assume the standard complexity-theoretic hypothesis that FPT \neq W[1].) That is, the sufficient condition for fixed-parameter tractability identified by Observation 1.2 is the *exclusive* explanation for fixed-parameter tractability, in the described setting of existential positive logic. Let us remark that the query preprocessing algorithm f here (in Observation 1.2) is related to and akin to the database notion of a query optimizer that computes a query execution plan.

Existential positive logic is a natural fragment of first-order logic [16] and is studied heavily in database theory. Existential positive formulas include and are semantically equivalent to so-called *unions of conjunctive queries*, also known as *selectproject-join-union queries*, which have been argued to be the most common database queries [17]. Recently, the present authors generalized the mentioned dichotomy on existential positive sentence classes, by presenting a classification theorem [18] describing the fixed-parameter tractable problems of the form count(Φ), where Φ is a bounded arity class of existential positive formulas. The classification theorem is in fact a trichotomy theorem, which demonstrates that the studied problems count(Φ) can exhibit three types of complexity behavior.

D. Contributions

As a means of introducing our contributions, we here wish to highlight a conceptual point: the applicability of the above *classical condition* indicates that for the model checking problem, *first-order logic itself can be used as a model of computation in which desirable, efficient algorithms can be expressed.* This condition posits the existence of an algorithm that translates first-order sentences to a polynomial-time evaluable format; and, the particular format used therein is that of a first-order sentence! Let us highlight that here, logic can be viewed as playing two complementary roles: on the one hand, the computational problems of interest are phrased directly in terms of logic; on the other hand, appropriate algorithmic solutions to this problem are themselves describable by logical sentences.

Inspired by this perspective of *logic as a useful model of computation*, the present work was motivated by the desire to develop a logic that could serve as a useful model of computation for the problem of counting query answers— analogously to how first-order logic itself serves as a useful model of computation for the model checking problem.

Let us point out some desiderata that such a logic ought to fulfill. First, recall that in the problem of counting query answers, an instance is a first-order formula paired with a structure, and the output is the *number* of answers. The hope, then, would be to be able to translate a first-order formula ϕ to a sentence ψ in the logic such that evaluating ψ on a structure returns the number of answers to ϕ on the structure; hence, in the logic, the evaluation of a sentence on a structure ought to return a *numerical quantity*, instead of a propositional value as in usual first-order logic. Second, the logic must accommodate the fact that, while the classical condition explains all tractable cases of model checking in existential positive logic (in the sense made precise), there are classes Φ of formulas for which the problem count(Φ) is fixed-parameter tractable, but on which the classical condition does not hold—in the sense that Φ does not have bounded width, even if each formula therein may be replaced with a logically equivalent one.¹

In this article, we introduce and study a logic, which we call \sharp -*logic* and which possesses the sought-after characteristics just described. In \sharp -logic, the evaluation of a \sharp -*sentence* (a type of formula in \sharp -logic) on a structure returns an integer value.

1) \sharp -logic, a preview: Syntactically, \sharp -logic consists of \sharp -formulas; each \sharp -formula ϕ has an associated set of free variables, denoted by free(ϕ), Let **B** be a structure, let ϕ be a \sharp -formula over the signature of **B**, and let h : free(ϕ) \rightarrow *B* be an assignment. Semantically, evaluating ϕ with respect to **B** and *h* returns an integer value, as opposed to a propositional value (as for a fo-formula). To present and discuss the semantics of \sharp -logic, we will notationally use $[\mathbf{B}, \phi]$ to denote the mapping that takes an assignment h : free(ϕ) \rightarrow *B* to the corresponding integer value (that is, the integer value provided to the triple consisting of ϕ , **B**, and h).

To offer the reader a feel for the syntax and semantics of \sharp -logic, we provide a discussion of some example formulas.

Example 1.3: Define $\phi(x, y, z)$ to be the formula $E(x, y) \land F(x, z)$. The first type of \sharp -formula is a *casting* of a foformula; define ψ to be $C(E(x, y), \{x, y\})$, which is a \sharp formula with free variables $\{x, y\}$. Let **B** be a structure, and let $h : \{x, y\} \to B$ be an assignment; $[\mathbf{B}, \psi](h)$ is equal to 1 or 0 depending on whether or not $\mathbf{B}, h \models E(x, y)$ (respectively). Once one has a \sharp -formula, it is possible to define a further \sharp -formula by *projecting* free variables. For example, $P\{y\}\psi$ is a \sharp -formula with free variables $\{x\}$. When $g : \{x\} \to B$ is an assignment, the value $[\mathbf{B}, P\{y\}\psi](g)$ is the number of extensions $h : \{x, y\} \to B$ of g such that $\mathbf{B}, h \models E(x, y)$. In an analogous fashion, one may define ψ' to be the \sharp -formula $C(F(x, z), \{x, z\})$; then, when $g : \{x\} \to B$ is an assignment, the value $[\mathbf{B}, P\{z\}\psi'](g)$ is the number of extensions $h' : \{x, z\} \to B$ of g such that $\mathbf{B}, h' \models F(x, z)$.

Now, let $g : \{x\} \to B$ be an assignment. Observe that the number of extensions $h^+ : \{x, y, z\} \to B$ of g such that $\mathbf{B}, h^+ \models E(x, y) \land F(x, z)$ is equal to the product of $[\mathbf{B}, P\{y\}\psi](g)$ and $[\mathbf{B}, P\{z\}\psi'](g)$. A product connective \times is provided by \sharp -logic, and said product is equal to $[\mathbf{B}, (P\{y\}\psi) \times (P\{z\}\psi')](g)$. Finally, $\theta = P\{x\}((P\{y\}\psi) \times (P\{z\}\psi'))$ is a \sharp -formula with no free variables; letting \emptyset denote the empty assignment, $[\mathbf{B}, \theta](\emptyset)$ will be equal to the sum, over all assignments $g : \{x\} \to B$, of $[\mathbf{B}, (P\{y\}\psi) \times (P\{z\}\psi')](g)$, which is equal to the number of assignments $f : \{x, y, z\} \to B$ such that $\mathbf{B}, f \models \phi$. Phrased in terminology that will be defined precisely, the \sharp -formula θ represents the fo-formula $\phi(x, y, z)$.

2) \sharp -logic, features: From the discussed viewpoint of bounded width as an explanation for the tractability of model checking, the relationship of the counting query answers problem to \sharp -logic is strongly analogous to the relationship of model checking to usual first-order logic. The following parallel of Observation 1.1 holds. (Note that the *width* of a formula in \sharp -logic will be defined in a natural way.)

Observation 1.4: Suppose that Ψ is a class of \sharp -sentences having bounded width. Then, the computational problem of evaluating a sentence $\psi \in \Psi$ on a finite structure is polynomial-time computable. (See Proposition 3.2 for a precise statement and further information.)

As expressed, one purpose of \sharp -logic is to allow for the translation of a first-order formula ϕ to a \sharp -sentence ψ such that ψ represents ϕ in that, for any structure **B**, the number of answers to ϕ on **B** is equal to the quantity that results from evaluating ψ on **B**. The following is an immediate consequence of the previous observation, and a parallel of Observation 1.2.

Observation 1.5: Suppose that Φ is a first-order formula class. The following condition, which we will refer to as the counting condition, is sufficient for the problem count(Φ) to be fixed-parameter tractable: there exists a \sharp -sentence class Ψ having bounded width and an algorithm f that computes, for each $\phi \in \Phi$, a \sharp -sentence $f(\phi)$ that is in Ψ and that represents ϕ .

Of course, Observation 1.5 is only of interest if the *counting condition* possesses *explanatory power*, that is, only if this condition allows one to explain the fixed-parameter tractability of problems $count(\Phi)$ having interest. We in fact show that, in the context of existential positive queries, the *counting condition* has maximal explanatory power:

Theorem 1.6: Let Φ be any class of existential positive queries having bounded arity. If $count(\Phi)$ is tractable, the counting condition applies to Φ . (See Theorem 4.6 for a precise statement.)

That is, the counting condition is the *exclusive* explanation for the tractability of count(Φ) in this existential positive setting, providing an analog to the result that the classical condition is the exclusive explanation for the tractability of $MC(\Phi)$ in the existential positive setting. On a conceptual level, we view this result as strong evidence that, for the problem of counting query answers, \sharp -logic is a useful, expressive model of computation in which relevant, efficient algorithms can be presented. This result is obtained as an immediate consequence of two theorems:

 We show that when such a problem count(Φ) is tractable, then there exists a bounded width class Ψ of #-sentences such that each φ ∈ Φ has a representation in Ψ (Theorem 4.4).

¹ An example of such a class is $\Theta = \{\theta_n \mid n \ge 1\}$ where $\theta_n(x_1, \ldots, x_n) = U_1(x_1) \land \cdots \land U_n(x_n)$. The number of answers for θ_n on any structure **B** is straightforwardly verified to be $\prod_{i=1}^n |U_i^{\mathbf{B}}|$. Given θ_n and **B**, this quantity can indeed be computed in polynomial time; however, it can be verified that the classical condition does not apply to Θ , in the mentioned sense.

• We prove that there is a *minimization algorithm* that, given an existential positive formula, computes a representation of minimum width (Theorem 4.5).

The latter theorem, which we view as a key contribution in and of itself, can be read as demonstrating that \sharp -logic is wellcharacterized and well-understood as a model of computation: conceiving of a \sharp -sentence representation of an existential positive formula as a computational procedure for counting query answers, this theorem provides a minimization algorithm that always outputs an *optimal* procedure for a given existential positive formula, where optimality here is measured using width.

In short, our presentation and study of \sharp -logic forwards the discussed use of logic as a means for expressing computationally desirable procedures; in particular, \sharp -logic allows for the direct expression of procedures for counting query answers.

3) Counting homomorphisms: The problem of counting the number of homomorphisms from a given *source* structure **A** to a given *target* structure **B** arises and has been studied in numerous contexts [19], [20]. This problem can be viewed as the special case of counting query answers where the formula is a quantifier-free conjunction of atoms; there is indeed a correspondence that allows one to pass from a source structure **A** to such a formula $\phi_{\mathbf{A}}$ which originates from the classical work [21] (and which is explained in Section II). This problem is now well-known to be polynomial-time tractable under a constant treewidth bound on the permitted source structures; indeed, the corresponding algorithm, which performs dynamic programming over a tree decomposition of **A**, has received a textbook treatment [22, Section 5.3].

We discuss how, from a tree decomposition for a structure **A**, one can compute, in polynomial time, a \sharp -sentence ψ that has width at most the width of the given decomposition (plus one), and that represents **A** in the sense that evaluating ψ on an arbitrary structure **B** always yields the number of homomorphisms from **A** to **B** (Example 3.4). Combining this result with Observation 1.4, we obtain that the algorithm of this well-known polynomial-time tractability result can be expressed in \sharp -logic (Proposition 3.5). Indeed, we believe that the resulting \sharp -sentences accurately, faithfully, and cleanly describe the execution of this algorithm.

4) A dual perspective on a classical theorem of Lovász: Fix a relational signature τ ; in the following discussion, all structures are finite and on τ . Let **B** be a structure, let $\operatorname{str}[\tau]$ denote the class of finite structures on τ , and let $L(\mathbf{B})$ be the vector from $\mathbb{Q}^{\operatorname{str}[\tau]}$ that maps a structure $\mathbf{A} \in \operatorname{str}[\tau]$ to the number of homomorphisms from \mathbf{A} to \mathbf{B} . A classical theorem of Lovász states that, for any two structures \mathbf{B} , \mathbf{B}' , it holds that $L(\mathbf{B}) = L(\mathbf{B}')$ iff \mathbf{B} and \mathbf{B}' are isomorphic [23].

When one is concerned with homomorphisms from one structure **A** to another structure **B**, sometimes, the structure **A** is referred to as the *left-hand* structure and the structure **B** is referred to as the *right-hand* structure. The vectors studied by Lovász indicate, for a structure **B**, the number of homomorphisms coming *from* each possible left-hand side structure. One can naturally formulate a dual vector, as follows. For any

structure **A**, define $R(\mathbf{A})$ to be the vector from $\mathbb{Q}^{\operatorname{str}[\tau]}$ that maps a structure $\mathbf{B} \in \operatorname{str}[\tau]$ to the number of homomorphisms from **A** to **B**. That is, the vector $R(\mathbf{A})$ indicates, for a structure **A**, the number of homomorphisms *to* each possible right-hand side structure.

Our previous work [18, Theorem 5.4] implied a dual of Lovász's theorem, namely, that for any two structures A, A', it holds that $R(\mathbf{A}) = R(\mathbf{A}')$ iff **A** and **A**' are isomorphic. In the present work, we prove and use a natural generalization of this fact, namely, that for any finite sequence A_1, \ldots, A_k of pairwise non-isomorphic structures, the vectors $R(\mathbf{A}_i)$ are linearly independent (Theorem 7.1). (We view these vectors as over the rational numbers \mathbb{Q} so as to be able to properly discuss linearly independence.) This linear independence theorem is used as a key tool to establish the correctness of our minimization algorithm. We believe that it should play an important role in future studies of counting query answers, and that the techniques and concepts that its proof requires may be of independent and future interest. Indeed, to prove this theorem, we extend notions and techniques from the work of Lovász [23]; for example, we introduce and crucially use a notion of multivariate polynomial associated to a primitive positive formula.

E. Discussion

Logics with counting mechanisms have been considered in finite model theory and descriptive complexity; one wellknown example is the counting logic studied by Immerman and Lander [24]. A typical motivation in these areas for studying such logics is the desire to extend first-order logic in order to capture properties not expressible in first-order logic. This motivation contrasts somewhat with ours here; our objective is to introduce logics that allow for the relatively direct expression of useful algorithms for the problem of counting query answers. We believe that it could be of interest to try to understand the relationship (if any) between existing counting logics and the logics studied in the present work.

We wish to emphasize that, as regards our present motivations, our logic² trades off expressivity and computability properties in an extremely desirable fashion. On the one hand, the algorithms for the tractable cases of $count(\Phi)$ (where Φ is existential positive) can be expressed in our logic, as described above; on the other hand, our *minimization algorithm* described above evidences that the measure of *width* can be computably minimized in our logic, and is thus in a certain sense well-characterized.

Previous work established that there is no algorithm for minimizing width in positive first-order logic [25, Section 5]. As a consequence, there is no algorithm for minimizing width in any logic that (1) includes positive first-order logic as a fragment and (2) where a width minimization algorithm would imply a width minimization algorithm for positive firstorder logic. Thus, such a logic would provably not exhibit the identified expressivity-computability tradeoff that our logic

²Precisely, the #EP-formulas, the fragment of our logic that we focus on.

enjoys. To the best of our knowledge, our width minimization algorithm is the first such algorithm for a logic with a form of counting mechanism; again, we view this as one contribution of this article.

We believe that our introduction of \sharp -logic may open up further research directions. One particular question for future research that we may pose is whether or not there are Ehrenfeucht-Fraïssé style games for proving inexpressibility in bounded width fragments of our logic.

II. PRELIMINARIES

A. Logic

We assume basic familiarity with the syntax and semantics of first-order logic. In this article, we focus on relational firstorder logic where equality is not built-in to the logic. Hence, each *vocabulary/signature* under discussion consists only of relation symbols. We assume structures under discussion to be *finite* (that is, have finite universe); nonetheless, we sometimes describe structures as *finite* for emphasis. We use the letters **A**, **B**, ... to denote structures, and the corresponding letters *A*, *B*, ... to denote their respective universes. When **A**, **B** are structures over the same signature τ , a *homomorphism* from **A** to **B** is a mapping $h : A \to B$ such that, for each $R \in \tau$ and each tuple $(a_1, \ldots, a_k) \in R^{\mathbf{A}}$, it holds that $(h(a_1), \ldots, h(a_k)) \in R^{\mathbf{B}}$.

We use the term *fo-formula* to refer to a first-order formula. An *ep-formula* (short for *existential positive formula*) is a foformula built from *atoms* (by which we refer to predicate applications of the form $R(v_1, \ldots, v_k)$, where R is a relation symbol and the v_i are variables), conjunction (\wedge), disjunction (\vee), and existential quantification (\exists). A *pp-formula* (short for *primitive positive formula*) is an ep-formula where disjunction does not occur. An fo-formula is *prenex* if it has the form $Q_1v_1 \ldots Q_nv_n\theta$ where θ is quantifier-free, that is, if all quantifiers occur in the front of the formula. The set of free variables of a formula ϕ is a *sentence* if free(ϕ) = \emptyset . We define an *ep-sentence* to be an ep-formula that is a sentence, and define *fo-sentence* and *pp-sentence* similarly.

We now present some definitions and conventions that are not totally standard. A primary concern in this article is in counting satisfying assignments of fo-formulas on a finite structure. The count is sensitive to the set of variables over which assignments are considered; and, we will sometimes want to count relative to a set of variables that is strictly larger than the set of free variables. Hence, we will often associate with each fo-formula ϕ a set V of variables called the *liberal variables*, denoted by $lib(\phi)$, for which it is required that $\operatorname{lib}(\phi) \supseteq \operatorname{free}(\phi)$. We generally assume that the variables in $lib(\phi) \setminus free(\phi)$ are not used in ϕ . To indicate that V is the set of liberal variables of ϕ , we often use the notation $\phi(V)$; we also use $\phi(v_1, \ldots, v_n)$, where the v_i are a listing of the liberal variables. Relative to a formula $\phi(V)$, when **B** is a structure, we will use $\phi(\mathbf{B})$ to denote the set of assignments $f: V \to B$ such that **B**, $f \models \phi$. We call an fo-formula ϕ free if free(ϕ) $\neq \emptyset$, and *liberal* if lib(ϕ) is defined and lib(ϕ) $\neq \emptyset$.

Example 2.1: Consider the formula

$$\phi(x, y, z) = E(x, y) \lor F(y, z).$$

Define $\psi(x, y, z) = E(x, y)$ and $\psi'(x, y, z) = F(y, z)$. The notation is intended to indicate that

$$\operatorname{lib}(\phi) = \operatorname{lib}(\psi) = \operatorname{lib}(\psi') = \{x, y, z\}.$$

In the context of studying ϕ , it is natural to define $\operatorname{lib}(\psi)$ and $\operatorname{lib}(\psi')$ to be $\{x, y, z\}$; under these definitions, it holds that $\phi(\mathbf{B}) = \psi(\mathbf{B}) \cup \psi'(\mathbf{B})$, but in general this would not hold in the case that (say) $\operatorname{lib}(\psi)$ was defined as $\{x, y\}$ (which set is equal to free (ψ)).

B. pp-formulas

It is well-known [21] that there is a correspondence between prenex pp-formulas and relational structures. In particular, each prenex pp-formula $\phi(S)$ (on signature τ) with $\operatorname{lib}(\phi) = S$ may be viewed as a pair (\mathbf{A}, S) consisting of a structure \mathbf{A} (on τ) and a set S; the universe A of A is the union of S with the variables appearing in ϕ , and the following condition defines the relations of A: for each $R \in \tau$, a tuple $(a_1, \ldots, a_k) \in A^k$ is in $R^{\mathbf{A}}$ if and only if $R(a_1, \ldots, a_k)$ appears in ϕ . In the other direction, such a pair (\mathbf{A}, S) can be viewed as a prenex pp-formula $\phi(S)$ where all variables in $A \setminus S$ are quantified and the atoms of ϕ are defined according to the above condition. A basic known fact [21] that we will use is that when $\phi(S)$ is a pp-formula corresponding to the pair (\mathbf{A}, S) , **B** is an arbitrary structure, and $f : S \rightarrow B$ is an arbitrary map, it holds that $\mathbf{B}, f \models \phi(S)$ if and only if there is an extension f' of f that is a homomorphism from A to B. We will freely interchange between the structure view and the usual notion of a prenex pp-formula. For a prenex pp-formula specified as a pair (\mathbf{A}, S) , we typically assume that $S \subseteq A$.

Example 2.2: Consider the pp-formula that is given as $\phi(u, v, w, x) = \exists y(E(u, v) \land F(w, y))$. To convert ϕ to a structure **A**, we take the universe A of **A** to be the union of lib(ϕ) with the variables appearing in ϕ , so $A = \{u, v, w, x, y\}$. The relations of **A** are as defined above, so $E^{\mathbf{A}} = \{(u, v)\}$ and $F^{\mathbf{A}} = \{(w, y)\}$. The resulting pair representation of ϕ is $(\mathbf{A}, \{u, v, w, x\})$.

C. Graphs

Throughout the paper, all graphs under discussion should be assumed to be undirected by default.

To every prenex pp-formula (\mathbf{A}, S) we assign a graph whose vertex set is $A \cup S$ and where two vertices are connected by an edge if they appear together in a tuple of a relation of \mathbf{A} . A prenex pp-formula (\mathbf{A}, S) is called *connected* if its graph is connected. A prenex pp-formula (\mathbf{A}', S') is a *component* of a prenex pp-formula (\mathbf{A}, S) over the same signature τ if there exists a set C that forms a connected component of the graph of (\mathbf{A}, S) , where $S' = S \cap C$, and for each relation $R \in \tau$, a tuple (a_1, \ldots, a_k) is in $R^{\mathbf{A}'}$ if and only if $(a_1, \ldots, a_k) \in$ $R^{\mathbf{A}} \cap C^k$.

Note that when this holds, the graph of (\mathbf{A}', S') is the connected component of the graph of (\mathbf{A}, S) on vertices C.

We will use the fact that, if $\phi(V)$ is a prenex pp-formula and $\phi_1(V_1), \ldots, \phi_k(V_k)$ is a list of its components, then for any finite structure **B**, it holds that $|\phi(\mathbf{B})| = \prod_{i=1}^{k} |\phi_i(\mathbf{B})|$.

Example 2.3: Consider the formula ϕ from Example 2.2. The connected components of the graph of ϕ can be readily verified to be $\{u, v\}$, $\{w, y\}$, and $\{x\}$. Hence, the pp-formula ϕ has 3 components, which can be readily verified to be $\phi_1(u, v) = E(u, v)$, $\phi_2(w) = \exists y F(w, y)$, and $\phi_3(x) = \top$; here, \top denotes the empty conjunction.

D. Treewidth

We give some basic facts about tree decompositions and treewidth; see for example [6] for more details.

A tree decomposition of a graph G = (V(G), E(G)) is a pair $(T, (B_t)_{t \in V(T)})$ where T is a tree and $(B_t)_{t \in V(T)}$ is a family of subsets of V(G) such that (1) for every $v \in V(G)$, the set $\{t \in V(T) \mid v \in B_t\}$ is non-empty and connected in T, and (2) for every edge $uv \in E(G)$, there is a $t \in V(T)$ such that $u, v \in B_t$. We also denote B_t using the notation B(t). The width of a tree decomposition $(T, (B_t)_{t \in V(T)})$ is defined as max $\{|B_t| : t \in V(T)\} - 1$. The treewidth tw(G) of G is the minimum width over all the tree decompositions of G. Computing tree decompositions of minimal width is fixedparameter tractable parameterized by the treewidth [26].

A tree decomposition is called *nice* if its tree T is rooted and every $t \in V(T)$ is of one of the following types:

- (leaf) t has no children and |B(t)| = 1.
- (introduce) t has one child t' and B(t) = B(t') ∪ {v} for a vertex v ∈ V\B(t').
- (forget) t has one child t' and $B(t) = B(t') \setminus \{v\}$ for a vertex $v \in B(t')$.
- (join) t has two children t_1, t_2 with $B(t) = B(t_1) = B(t_2)$.

It is well-known that a width k tree decomposition of G can be converted to a width k nice tree decomposition, in polynomial time.

III. *‡-LOGIC*

In this section, we present the syntax and semantics of \sharp -logic, as well as some associated terminology. Syntactically, \sharp -logic consists of \sharp -formulas; each \sharp -formula ϕ has an associated set of free variables, denoted by $\text{free}(\phi)$, as well as an associated set of *closed* variables, denoted by $\text{closed}(\phi)$. (At this point, the reader may wish to recall the preview of \sharp -logic presented in Section I-D1.)

A. Syntax

We define *#*-formulas inductively, as follows.

- C(φ, L) is a #-formula if φ is a fo-formula and L ⊇ free(φ).
 Define free(C(φ, L)) = L and closed(C(φ, L)) = Ø.
- PVφ is a ♯-formula if φ is a ♯-formula and V is a set of variables with V ∩ closed(φ) = Ø.
 Define free(PVφ) = free(φ)\V and closed(PVφ) = V ∪ closed(φ).

- EVφ is a ♯-formula if φ is a ♯-formula and V is a set of variables with V ∩ (free(φ) ∪ closed(φ)) = Ø.
 Define free(EVφ) = V ∪ free(φ) and closed(EVφ) = closed(φ).
- $\phi \times \phi'$ is a \sharp -formula if ϕ and ϕ' are \sharp -formulas with free(ϕ) = free(ϕ') and closed(ϕ) \cap closed(ϕ') = \emptyset . Define free($\phi \times \phi'$) = free(ϕ) and closed($\phi \times \phi'$) = closed(ϕ) \cup closed(ϕ').
- φ + φ' is a #-formula if φ and φ' are #-formulas with free(φ) = free(φ').
 Define free(φ + φ') = free(φ) and closed(φ + φ') = closed(φ) ∪ closed(φ').
- n is a ♯-formula if n ∈ Z.
 Define free(n) = Ø and closed(n) = Ø.

A formula $C(\phi, L)$ can be thought of as the casting of a fo-formula ϕ into a \sharp -formula; the P quantifier can be thought of as projecting or closing variables; and the E quantifier can be thought of as expanding the set of free variables. The connectives \times and + perform the usual arithmetic operations. We remark that, for each \sharp -formula ϕ , it holds that free (ϕ) and closed (ϕ) are disjoint; this is straightforwardly verified by induction.

Let ψ be a \sharp -formula. We say that a \sharp -formula θ is a \sharp subformula of ψ if θ is used in the inductive formation of ψ . We say that a fo-formula θ is a fo-subformula of ψ if ψ contains a subformula $C(\phi, L)$ where θ is a subformula of ϕ .

A subformula of ψ is a \sharp -subformula or fo-subformula of ψ . We define width(ψ) to be the maximum of $|\text{free}(\theta)|$ over all subformulas θ of ψ , and \sharp -width(ψ) to be the maximum of $|\text{free}(\theta)|$ over all \sharp -subformulas θ of ψ . We say that ψ is a \sharp -sentence if $\text{free}(\psi) = \emptyset$.

We define a $\sharp PP$ -formula to be a \sharp -formula where, in each \sharp -subformula of the form $C(\phi, L)$, ϕ is a pp-formula; the notion of $\sharp EP$ -formula is defined analogously. We define a $\sharp PP$ -sentence to be a $\sharp PP$ -formula that is a \sharp -sentence, and we define a $\sharp EP$ -sentence similarly.

B. Semantics

We define the semantics of our logic. For each structure **B**, each \sharp -formula ψ on the vocabulary of **B**, and each assignment $h : \text{free}(\psi) \rightarrow B$, we define $[\mathbf{B}, \psi](h)$ recursively, as follows.

- When C(φ, L) is a ♯-formula,
 [B, C(φ, L)](h) = 1 if B, h ⊨ φ;
 [B, C(φ, L)](h) = 0 otherwise.
- When $PV\phi$ is a \sharp -formula, $[\mathbf{B}, PV\phi](h) = \sum_{h'} [\mathbf{B}, \phi](h'),$ where the sum is over all extensions h': free $(\phi) \cup V \to B$ of h.
- When EVφ is a ♯-formula,
 [B, EVφ](h) = [B, φ](h ↾ free(φ)).
- When $\phi \times \phi'$ is a \sharp -formula, $[\mathbf{B}, \phi \times \phi'](h) = [\mathbf{B}, \phi](h) \cdot [\mathbf{B}, \phi'](h).$

- When $\phi + \phi'$ is a \sharp -formula, $[\mathbf{B}, \phi + \phi'](h) = [\mathbf{B}, \phi](h) + [\mathbf{B}, \phi'](h).$
- When n is a \sharp -formula, $[\mathbf{B}, n](h) = n$.

We consider two \sharp -formulas ϕ , ϕ' with free $(\phi) = \text{free}(\phi')$ to be *logically equivalent* if for each structure **B**, it holds that $[\mathbf{B}, \phi] = [\mathbf{B}, \phi']$. A \sharp -sentence ψ represents or is a representation of a fo-formula $\phi(V)$ if for each finite structure **B**, it holds that $|\phi(\mathbf{B})| = [\mathbf{B}, \psi](\emptyset)$, where \emptyset is the empty assignment. For simplicity, when ψ is a \sharp -sentence, we will typically write $[\mathbf{B}, \psi]$ in place of $[\mathbf{B}, \psi](\emptyset)$. We will use the term \sharp PP-representation to refer to a representation that is a \sharp -PP-formula, and define \sharp EP-representation similarly.

We make the basic observation that each fo-formula has a representation.

Proposition 3.1: For each fo-formula $\phi(V)$, the \sharp -sentence $PVC(\phi, V)$ is a representation of $\phi(V)$.

We also observe that, when a constant width bound is assumed, evaluation of \sharp -sentences can be performed in polynomial time.

Proposition 3.2: For each $k \ge 1$, there exists a polynomialtime algorithm that, given a finite structure **B** and a \sharp -sentence ψ having width $(\psi) \le k$, computes [**B**, ψ].

Proof. For each subformula θ of ψ and each mapping h: free $(\theta) \rightarrow B$, the algorithm computes $[\mathbf{B}, \theta](h)$ in the case that θ is a \sharp -subformula, and determines whether or not $\mathbf{B}, h \models \theta$ in the case that θ is a fo-subformula. This computation is performed inductively over the subformulas of ψ , and in the case that θ is a \sharp -subformula, the just-given semantics are used. \Box

C. Examples

Example 3.3: Let us define $\phi(x_0, x_1, x_2, y_0, y_1, y_2)$ as the formula $\phi_0 \land \phi_1 \land \phi_2$, where $\phi_i = \exists z_i T_i(x_i, x_{i+1}, y_i, z_i)$ for each $i \in \{0, 1, 2\}$; here, the quantity i + 1 appearing in x_{i+1} is computed modulo 3. These formulas are over the vocabulary $\{T_0, T_1, T_2\}$ having three relation symbols, each of arity 4.

Define $\psi_i = P\{y_i\}C(\phi_i, \{x_0, x_1, x_2, y_i\})$ for each $i \in \{0, 1, 2\}$. Observe that when **B** is a structure and $h : \{x_0, x_1, x_2\} \rightarrow B$ is a map, $[\mathbf{B}, \psi_i](h)$ gives the number of extensions $h' : \{x_0, x_1, x_2, y_i\} \rightarrow B$ of h satisfying ϕ_i on **B**. We have free $(\psi_0) =$ free $(\psi_1) =$ free $(\psi_2) = \{x_0, x_1, x_2\}$ and closed $(\psi_i) = \{y_i\}$. Now consider $\psi = (\psi_0 \times \psi_1) \times \psi_2$. It can be verified that, for a structure **B** and a map $h : \{x_0, x_1, x_2\} \rightarrow B$, $[\mathbf{B}, \psi](h)$ gives the number of extensions $h' : \{x_0, x_1, x_2, y_0, y_1, y_2\} \rightarrow B$ of h satisfying ϕ on **B**. It follows that the \sharp -sentence $\theta = P\{x_0, x_1, x_2\}\psi$ is a representation of ϕ . Since

$$3 = |\text{free}(\psi)| = |\text{free}(\psi_0)| = |\text{free}(\psi_1)| = |\text{free}(\psi_2)|,$$

we obtain that the representation θ has width equal to

$$\max(3, \mathsf{width}(\psi_0), \mathsf{width}(\psi_1), \mathsf{width}(\psi_2)) = 4.$$

As a further remark, consider, as an example, the subformula $C(\phi_0, \{x_0, x_1, x_2, y_0\})$ of ψ_0 . It holds that $x_2 \notin \text{free}(\phi_0)$,

and so $E\{x_2\}C(\phi_0, \{x_0, x_1, y_0\})$ is a \sharp -formula and is logically equivalent to $C(\phi_0, \{x_0, x_1, x_2, y_0\})$.

Example 3.4: Consider a prenex pp-formula ϕ without quantifiers whose structure view has the form (\mathbf{A}, A) . Suppose that (T, (B(t))) is a width k tree decomposition of the graph of (\mathbf{A}, A) ; without loss of generality, we may assume that this tree decomposition is nice, and that the root node r of T has $B(r) = \emptyset$. We explain how to give a representation of ϕ having width $\leq k + 1$.

Say that a node u of T is *below* a node t of T if t occurs on the unique simple path from u to the root of T (this is understood to hold in particular when u = t). When t is a node of T, define $B(\leq t)$ to be the union of B(u) over all nodes u that are below t.

We show that, for each node t of T, there exists a \sharp -formula ψ_t such that:

- free $(\psi_t) = B(t)$,
- $\operatorname{closed}(\psi_t) = B(\leqslant t) \setminus B(t)$, and
- for any structure D, the value [D, ψ_t](h) is equal to the number of extensions h': B(≤ t) → D of h such that (D, h') satisfies each atom R(a₁,...,a_k) of φ whose variables all fall into a bag B(u), with u below t.

We give a \sharp -formula depending on the type of the node t; we use the notation from the definition of nice tree decomposition.

- (introduce) ψ_t = (Evψ_{t'}) × C(α₁, B(t)) × ··· × C(α_m, B(t)) where the α_i are the atoms of φ whose variables fall into B(t). Note that the presence of the C(α_i, B(t)) ensures that [**D**, ψ_t](h) = 0 if (**D**, h) does not satisfy all of the α_i.
- (leaf) ψ_t is defined as in the previous case, except $(Ev\psi_{t'})$ is omitted from the product.
- (forget) $\psi_t = P\{v\}\psi_{t'}$.
- (join) $\psi_t = \psi_{t_1} \times \psi_{t_2}$. In this case, $\psi_{t_1} \times \psi_{t_2}$ is a \sharp -formula since free $(\psi_{t_1}) = B(t_1) = B(t_2) = \text{free}(\psi_{t_2})$, and, by the definition of tree decomposition, it holds that the sets $B(\leqslant t_1) \setminus B(t_1) = \text{closed}(\psi_{t_1})$ and $B(\leqslant t_2) \setminus B(t_2) = \text{closed}(\psi_{t_2})$ are disjoint.

The desired representation is ψ_r . The claim on the width of ψ_r holds, for we have the following: each \sharp -subformula ψ' of ψ_r has free $(\psi') = B(w)$ for a node w of T, and each fosubformula of ψ_r is either \top or an atom whose variables fall into a bag B(t).

From the discussion in Example 3.4, we obtain the following.

Proposition 3.5: Let $k \ge 1$. Consider the problem of computing, given a pair (\mathbf{A}, \mathbf{B}) of relational structures (over the same signature) where \mathbf{A} has treewidth $\le k$, the number of homomorphisms from \mathbf{A} to \mathbf{B} . This problem can be solved by the polynomial-time algorithm that computes a nice, width k tree decomposition (using a known polynomial-time algorithm for this task); computes, from \mathbf{A} , the representation ψ_r given

by Example 3.4; and, invokes the algorithm of Proposition 3.2 to compute $[\mathbf{B}, \psi_r]$.

IV. MAIN THEOREMS

A. Statements

The treewidth of a prenex pp-formula ϕ , denoted here by $\mathsf{tw}(\phi)$, is defined as the treewidth of the graph of ϕ . The following notions are adapted from [4], [18]. Let (\mathbf{A}, S) be a prenex pp-formula with graph G. An \exists -component of (\mathbf{A}, S) is a graph of the form G[W'] where there exists $W \subseteq A$ that is a connected component of $G[A \setminus S]$ and W' is the union of W with all vertices in S having an edge to W. Define contract (\mathbf{A}, S) to be the graph on vertices S obtained by starting from G[S] and adding an edge between any two vertices that appear together in an \exists -component of (\mathbf{A}, S) .³

Example 4.1: Let $\phi(u_1, u_2, u_3, u_4)$ be the pp-formula

$$\exists x_1 \exists x_2 \exists x_3 (R(u_1, x_1, x_2) \land S(u_2, x_2) \land T(u_2, x_3) \land U(u_3, x_3) \land P(u_3, u_4)).$$

The graph G of ϕ has vertex set

$$V = \{u_1, u_2, u_3, u_4, x_1, x_2, x_3\}$$

and edge set

$$E = \{\{u_1, x_1\}, \{u_1, x_2\}, \{x_1, x_2\}, \{u_2, x_2\}, \{u_2, x_3\}, \{u_3, x_3\}, \{u_3, u_4\}\}.$$

Note that, if we were to view ϕ as a pair (A, S), the structure A would have universe V and we would have $S = \operatorname{lib}(\phi) =$ $\{u_1, u_2, u_3, u_4\}$. There are two connected components of $G[A \setminus S]$, namely, $W_1 = \{x_1, x_2\}$ and $W_2 = \{x_3\}$. (Note that, in contrast, G itself is connected.) The \exists -components of ϕ are thus $W'_1 = W_1 \cup \{u_1, u_2\}$ and $W'_2 = W_2 \cup \{u_2, u_3\}$. The graph $contract(\phi)$ is the graph on vertices $S = \{u_1, u_2, u_3, u_4\}$ obtained by starting from G[S], which has the single edge $\{u_3, u_4\}$, and adding the edges $\{u_1, u_2\}$ and $\{u_2, u_3\}$. A core of (\mathbf{A}, S) is a prenex pp-formula $(c(\mathbf{A}), S)$ where c is an A-endomorphism fixing each $s \in S$ that has minimum image size. By $c(\mathbf{A})$, we mean the structure with universe c(A) and where $R^{c(\mathbf{A})} = \{c(t) \mid t \in R^{\mathbf{A}}\}$; here, c(t) denotes the tuple obtained by applying c to each entry of t. It is known that any core of a prenex pp-formula ϕ is logically equivalent to ϕ [21]. We speak of *the core* of a pp-formula, as it is unique up to isomorphism; this follows from the basic theory of cores [27].

A class Φ of prenex pp-formulas satisfies the *contraction condition* if the class containing each graph contract(ψ), where ψ is the core of a formula in Φ , has bounded treewidth; Φ satisfies the *tractability condition* if it satisfies the contraction condition and the cores of formulas in Φ have bounded treewidth. Previous work showed that, for bounded arity Φ , count(Φ) is fixed-parameter tractable if Φ satisfies the tractability condition; interreducible with the parameterized clique problem if Φ satisfies the contraction condition but not the tractability condition; and as hard as the parameterized counting clique problem otherwise (see [4] for a precise statement). We first study representations of pp-formulas, obtaining the following theorems.

Theorem 4.2: Let Φ be a class of prenex pp-formulas.

- The class Φ satisfies the tractability condition if and only if there exists k ≥ 1 such that each formula in Φ has a #PP-representation φ' such that width(φ') ≤ k.
- The class Φ satisfies the contraction condition if and only if there exists k ≥ 1 such that each formula in Φ has a #PP-representation φ' such that #-width(φ') ≤ k.

Theorem 4.3: There exists an algorithm that, given a prenex pp-formula ϕ , outputs a \sharp PP-representation ψ of ϕ of minimum width.

Building on this understanding of pp-formulas, we are then able to achieve general versions of these theorems for epformulas. Previous work showed that for any class Φ of epformulas, there exists a class Φ^+ of prenex pp-formulas such that the problems $count(\Phi)$ and $count(\Phi^+)$ are interreducible (see [18] for a precise statement), and hence (for example) whether or not the tractability condition holds on Φ^+ determines whether or not $count(\Phi)$ is fixed-parameter tractable. For the purposes of this extended abstract, it is sufficient to know that the class Φ^+ is essentially defined from Φ in the following way: for each $\phi \in \Phi$, it is shown that the function $|\phi(\cdot)|$, which maps a finite structure **B** to $|\phi(\mathbf{B})|$, can be written as a polynomial (over the integers) in unknowns $|\phi_1(\cdot)|, \ldots, |\phi_m(\cdot)|$ where the ϕ_i are pp-formulas; the class Φ^+ is defined to contain all such formulas ϕ_i arising in this way. (In general, the mentioned polynomial makes use of negative integers; this is a reason why our definition of #logic allows arbitrary integers, and not just natural numbers.) For more information, we refer the reader to [18].

Theorem 4.4: Let Φ be a class of ep-formulas.

- The class Φ⁺ satisfies the tractability condition if and only if there exists k ≥ 1 such that each formula in Φ has a #EP-representation φ' having width(φ') ≤ k.
- The class Φ⁺ satisfies the contraction condition if and only if there exists k ≥ 1 such that each formula in Φ has a #EP-representation φ' having #-width(φ') ≤ k.

Theorem 4.5: There exists an algorithm that, given an epformula ϕ , outputs a $\sharp \text{EP}$ -representation ψ of ϕ of minimum width.

The following is a consequence of the previous two theorems.

Theorem 4.6: Let Φ be a class of ep-formulas. If Φ^+ satisfies the tractability condition, then there exists $k \ge 1$ and an algorithm that, given a formula $\phi \in \Phi$, computes a $\sharp \text{EP-}$ representation ϕ' of ϕ having width $(\phi') \le k$.

Proof. The algorithm is that provided by Theorem 4.5. The claim on the width follows immediately from the first part of Theorem 4.4.

B. Overviews of the proofs

In order to prove the main theorems just presented, we will develop several tools spanning Sections V to VII before finally proving the main results in Section VIII. Here, we offer the

³ Note that in previous articles [4], [18] this graph was defined in terms of the core of the given formula.

reader guidance by introducing the pieces of the puzzle and explaining how they fit together.

As formulated above, the tractability condition is defined in part using the contraction condition, and thus it appears to consist of two independent parts. In a first step, in Section V, we simplify the situation by introducing a new notion that we call quantifier-aware width of a pp-formula. It is defined to be the minimal width of certain restricted tree decompositions of a pp-formula. We show that having bounded quantifier-aware width is equivalent to the tractability condition (Lemma 5.2); as a consequence of this lemma, after Section V, we will not have to deal with the tractability condition directly anymore and can work with the conceptually cleaner notion of quantifier-aware width. We then go on to show that quantifieraware width can be computed in a fixed-parameter fashion (Lemma 5.3) which is an important building block for our minimization algorithm. Finally, we prove that quantifieraware width of a pp-formula is essentially equal to the width of an optimal representation by a *PP*-formula of a particularly simple type, which we call basic (Lemma 5.4). Together with the rest of the results of Section V, this gives an important connection between pp-formulas and #PP-formulas and thus bridges the gap between ordinary first-order logic and #-logic.

Section VI gives insights into the structure of #EP-formulas by showing that we can always turn one into an equivalent weighted sum of basic #PP-formulas. This allows us to leverage most of the results of Section V to general #EP-formulas.

The perhaps most subtle but very important contribution to the proofs of the main theorem comes from the results of Section VII. In this section, we consider sums of the form $\sum_{i=1}^{m} c_i |\phi_i(V_i)|$ where each c_i is a non-zero rational constant and the ϕ_i are pp-formulas which are pairwise not counting equivalent. We call such terms *linear combinations*; each naturally maps a finite structure **B** to the value $\sum_{i=1}^{m} c_i |\phi_i(\mathbf{B})|$. The main result of Section VII is the *independence theorem* (Theorem 7.1) which states that for every linear combination there is a structure on which the linear combination evaluates to a non-zero value.

The independence theorem has concrete applications throughout the proofs of the main results in Section VIII. The reasoning is roughly as follows: we assign to a formula two different linear combinations having two different desirable properties (for example, small width and small number of summands) but computing the same value on every finite structure. Applying Theorem 7.1 on the difference of these linear combinations, we obtain that in fact both linear combinations are the same (up to *counting equivalence* of the ϕ_i , a notion of equivalence to be defined). Consequently, both linear combinations have the same properties and thus in particular have both of the two desirable properties. This then allows to reason about the properties of the original formula we started with.

The proofs of the main results in Section VIII use the above tools in a rather black-box fashion. Thus the reader is invited to first skip the proofs in Sections V to VII and see how everything fits together in Section VIII before reading the proofs of the individual pieces.

V. QUANTIFIER-AWARE WIDTH

In this section, we introduce a new width measure of ppformulas which we call *quantifier-aware width* and show that it is related to the width of $\sharp PP$ -formulas.

We here assume all tree decompositions of pp-formulas to be nice. So let $(T, (B(t))_{t\in T})$ be a nice tree decomposition of a pp-formula ϕ . For every variable x of ϕ let top(x) be the vertex t of T that is highest in T such that $x \in B(t)$. We call a tree decomposition of ϕ quantifier-aware if for every \exists -component C of ϕ and for all $x \in V(C) \setminus \text{free}(\phi)$ and all $y \in V(C) \cap \text{free}(\phi)$, we have that top(y) is on the path from top(x) to the root of T. We call the quantifier-aware width of a pp-formula ϕ , denoted by $qaw(\phi)$, the minimal treewidth of a quantifier-aware tree decomposition of ϕ plus 1.

Remark 5.1: The quantifier-aware width can be arbitrarily higher than their treewidth. To see this consider the formula $\phi = \exists z \bigwedge_{i \in [n]} E(x_i, z)$. The primal graph of ϕ is a star, so it has treewidth 1. We claim that the quantifier-aware width of ϕ is n + 1. To see this, observe first that the free variables x_1, \ldots, x_n must appear above top(z) in a bag of any quantifier-aware tree decomposition. But since $x_i z$ is an edge in the primal graph for every *i*, the variable x_i must also appear in a common bag with *z* and consequently also in top(z). Thus top(z) must contain n + 1 variables.

We now demonstrate properties of quantifier-aware width that will be used to establish our main theorems.

We first show that $qaw(\phi)$ is, in a sense, characterized by $tw(\phi)$ and $tw(contract(\phi))$, for every pp-formula ϕ . Consequently, quantifier-aware treewidth will allow us to characterize tractable classes of pp-formulas for counting.

Lemma 5.2: For every pp-formula ϕ we have

$$\max\{\mathsf{tw}(\phi),\mathsf{tw}(\mathsf{contract}(\phi))\} + 1$$
$$\leq \mathsf{qaw}(\phi) \leq \mathsf{tw}(\phi) + \mathsf{tw}(\mathsf{contract}(\phi)) + 1.$$

We now give a lemma that shows how to compute the quantifier-aware treewidth, which will allow us to compute prime PP-formulas of optimal width.

Lemma 5.3: The computation of quantifier-aware tree decompositions of pp-formulas having minimal width is fixedparameter tractable, when the parameter is taken as the quantifier-aware width.

We call a $\sharp PP$ -formula ϕ basic if it does not contain + nor subformulas of the form n, where $n \in \mathbb{Z}$. The following lemma demonstrates that basic $\sharp PP$ -formulas correspond very closely to pp-formulas.

Lemma 5.4:

- a) There exists an algorithm that, given a basic #PP-sentence φ', computes a pp-formula φ that φ' represents, such that width(φ') ≥ qaw(φ), and #-width(φ') ≥ tw(contract(φ)) + 1.
- b) There exists an algorithm that, given a pp-formula φ, computes a basic #PP-sentence φ' that represents φ, such that width(φ') ≤ qaw(φ), and #-width(φ') ≤ tw(contract(φ)) + 1.

We call a $\sharp PP$ -formula *constant* if it is only constructed from constants in \mathbb{Z} , \times , and P- and E-quantifiers. We call a $\sharp PP$ -formula *flat* if it is of the form $\sum_{i \in [\ell]} \psi_i \times \phi_i$ where the ψ_1, \ldots, ψ_n are constant and $\phi_1, \ldots, \phi_\ell$ are basic $\sharp PP$ formulas.

The main result of this section is the following normalization lemma.

Lemma 6.1: There exists an algorithm that computes, for a given #EP-formula ϕ , a logically equivalent flat #PP-formula ϕ' such that width $(\phi') \leq$ width (ϕ) .

We prove a sequence of lemmas to aid us.

Lemma 6.2: There exists an algorithm that computes, for a given #EP-formula ϕ of the form $C(\psi, L)$, a logically equivalent #PP-formula ϕ' such that width $(\phi') \leq$ width (ϕ) .

We call an \sharp -formula +-free if it does not contain +.

Lemma 6.3: There exists an algorithm that computes, for a given #EP-formula ϕ , a logically equivalent #PP-formula ϕ' of the form $\sum_{i=1}^{s} \phi_i$ where the ϕ_i are +-free such that width $(\phi') \leq$ width (ϕ) .

Proof. The proof is by straightforward induction on the structure of ϕ , pushing all occurences of + up in the formula; for instance, one proves that $PV(\psi_1 + \psi_2)$ is logically equivalent to $PV\psi_1 + PV\psi_2$. The base case $\phi = C(\psi, L)$ is Lemma 6.2.

The proof of the following lemma is by a straightforward induction.

Lemma 6.4: There exists an algorithm that computes, for a given constant $\sharp PP$ -formula ϕ , a logically equivalent $\sharp PP$ formula $\phi' = EV_1PV_2n$ with $n \in \mathbb{Z}$ such that width $(\phi') \leq$ width (ϕ) .

Lemma 6.5: There exists an algorithm that computes, for a given +-free \sharp PP-formula ϕ , a logically equivalent \sharp PPformula $\phi' = \psi_1 \times \psi_2$ where ψ_1 is constant and ψ_2 is basic such that width $(\phi') \leq$ width (ϕ) .

Proof. The proof is again straightforward induction in the style of Lemma 6.3. We consider only the the case of *P*-quantifiers which is the only case that is not completely clear from the definition.

So let $\phi = PV\phi'$ where $\phi' = \phi_1 \times \phi_2$ such that ϕ_1 is constant and ϕ_2 is basic. Note that by Lemma 6.4 we may assume that $\phi_1 = PV_1EV_2n$ for some $n \in \mathbb{Z}$. We claim that ϕ is logically equivalent to $\phi'_1 \times PV\phi_2$ where $\phi'_1 = PV_1E(V_2 \setminus V)$. To see this, consider a structure **B** and an assignment *h* to ϕ . Then $[\mathbf{B}, PV\phi'](h) = \sum_{h'} [\mathbf{B}, \phi_1 \times \phi_2](h') = \sum_{h'} ([\mathbf{B}, \phi_1](h') \cdot [\mathbf{B}, \phi_2](h'))$ where the *h'* are as in the definition. Now for an arbitrary assignment *h''* to ϕ' , $[\mathbf{B}, \phi_1](h'') = [\mathbf{B}, \phi_1](h') = [\mathbf{B}, \phi'_1](h)$ for all *h'*. Consequently, we have

$$[\mathbf{B}, PV\phi'](h) = [\mathbf{B}, \phi_1](h'') \cdot \sum_{h'} [\mathbf{B}, \phi_2](h')$$
$$= [\mathbf{B}, \phi_1'](h) \cdot \sum_{h'} [\mathbf{B}, \phi_2](h')$$
$$= [\mathbf{B}, \phi_1'](h) \cdot [\mathbf{B}, PV\phi_2](h)$$
$$= [\mathbf{B}, \phi_1' \times PV\phi_2](h).$$

Proof. (of Lemma 6.1) First use Lemma 6.3 to turn ϕ into a sum of +-free \sharp PP-formulas. After this, apply Lemma 6.5 to each of the summands.

VII. INDEPENDENCE THEOREM

In this section, we establish a key tool for reasoning about #EP-formulas.

Define two fo-formulas $\phi(V)$, $\phi'(V')$ to be *counting equivalent* if, for each structure **B**, it holds that $|\phi(\mathbf{B})| = |\phi'(\mathbf{B})|$; note that a decidable characterization of counting equivalence on pp-formulas is known ([18, Theorem 5.4]).

In the scope of this article, define a *linear combination* to be an expression of the form $\sum_{i=1}^{m} c_i |\phi_i(V_i)|$, where $m \ge 1$, each c_i is a non-zero rational number, and the $\phi_i(V_i)$ are pp-formulas that are pairwise not counting equivalent. Here, the notation $|\phi_i(V_i)|$ is intended to indicate the function that maps each finite structure **D** to the value $|\phi_i(\mathbf{D})|$. So, each linear combination ℓ naturally induces a mapping $\ell(\cdot)$ from finite structures to \mathbb{Q} , namely, the map given by $\ell(\mathbf{D}) =$ $\sum_{i=1}^{m} c_i |\phi_i(\mathbf{D})|$. The following theorem will be key for our understanding of equivalence of $\sharp \text{EP-formulas}$.

Theorem 7.1: (Independence theorem) For any linear combination ℓ , there exists a finite structure **D** such that $\ell(\mathbf{D}) \neq 0$.

We devote the rest of this section to proving this theorem. By multiplying all values c_i by a multiple of their denominators, we can and will assume that each value c_i is an integer.

In order to establish this theorem, we will make use of the following lemma.

Lemma 7.2: (Lovász [23]) For each univariate polynomial p with positive integer coefficients and each finite structure **B**, there exists a finite structure $p(\mathbf{B})$ such that, for each connected liberal pp-formula ϕ , it holds that $|\phi(p(\mathbf{B}))| = p(|\phi(\mathbf{B})|)$.

The following lemma shows that certain pp-formulas can be controlled independently of each other.

Lemma 7.3: Let $\phi_1(S_1), \ldots, \phi_n(S_n)$ be connected liberal pp-formulas (over signature τ) that are pairwise not counting equivalent. Then for every $m \ge 2$, there exist structures $(\mathbf{B}_{(a_1,\ldots,a_n)} | (a_1,\ldots,a_n) \in [m]^n)$ and injective functions $f_1,\ldots,f_n: [m] \to \mathbb{N}$ such that for each $(a_1,\ldots,a_n) \in [m]^n$ and each $i \in [n]$, it holds that $|\phi_i(\mathbf{B}_{(a_1,\ldots,a_n)})| = f_i(a_i)$.

Moreover, when **A** is any structure on which $|\phi_i(\mathbf{A})| > 0$ for each $i \in [n]$, each structure $\mathbf{B}_{(a_1,...,a_n)}$ can be chosen in the form $\mathbf{A} \times \mathbf{E}$, where **E** is a structure such that $|\phi(\mathbf{E})| > 0$ for all pp-formulas ϕ over τ . **Proof.** As each $\phi_i(S_i)$ is connected and liberal, the result [18, Theorem 5.14] ensures that there exists a structure C' such that the values $|\phi_i(\mathbf{C}')|$ are pairwise different, and where $|\phi(\mathbf{C}')| > 0$ for all pp-formulas ϕ . By taking a sufficiently large power P of C', we may obtain that for the structure $\mathbf{C} = \mathbf{C}'^P \times \mathbf{A}$, the values $c_i = |\phi_i(\mathbf{C})|$ are pairwise different. For each $(a_1, \ldots, a_n) \in [m]^n$, define $p_{(a_1, \ldots, a_n)}$ to be a univariate polynomial over the rationals that evaluates to 0 at 0, and to a_i at c_i (for each $i \in [n]$). Define D to be the absolute value of the product of all denominators of coefficients in the defined polynomial. Set $p'_{(a_1, \ldots, a_n)} = D \cdot p_{(a_1, \ldots, a_n)}$; each such polynomial has integer coefficients. Next, set $p_{(a_1, \ldots, a_n)}^-$ to be the restriction of $p'_{(a_1, \ldots, a_n)}$ to be $p'_{(a_1, \ldots, a_n)} + 2\sum_{(a_1, \ldots, a_n) \in [m]^n} (-p_{(a_1, \ldots, a_n)}^-)$. Now, for each $(a_1, \ldots, a_n) \in [m]^n$, define the structure $\mathbf{B}_{(a_1, \ldots, a_n)}$ as $p''_{(a_1, \ldots, a_n)}(\mathbf{C})$; for each $i \in [n]$, we have

$$|\phi_i(\mathbf{B}_{(a_1,\dots,a_n)})| = |\phi_i(p''_{(a_1,\dots,a_n)}(\mathbf{C}))| = p''_{(a_1,\dots,a_n)}(c_i);$$

the second equality here holds by the Lemma 7.2. From these equalities and the definitions of $p_{(a_1,...,a_n)}$ and $p''_{(a_1,...,a_n)}$, it is straightforward to verify that the defined structures have the desired property. Our claim concerning each $B_{(a_1,...,a_n)}$ having the form $\mathbf{A} \times \mathbf{E}$ holds, as $p''_{(a_1,...,a_n)}(0) = 0$ (for each $(a_1,...,a_n) \in [m]^n$), implying that the structures $\mathbf{B}_{(a_1,...,a_n)}$ provided can be obtained in the form $\mathbf{C} \times \cdot$, which has the form $\mathbf{A} \times \cdot$.

We now introduce a highly useful notion, that of *component* polynomial. Fix a set V of liberal variables. Denote by \mathcal{E} the set of counting equivalence classes of liberal connected pp-formulas (with liberal variables from V). A component polynomial q is a multivariate polynomial with integer coefficients over variables $\{X_e \mid e \in \mathcal{E}\}$. For any finite structure **B**, we define the value of q evaluated on **B**, denoted by $q[[\mathbf{B}]]$, as the integer value obtained by evaluating q when each X_e is given the value $|\phi_e(\mathbf{B})|$, for a formula $\phi_e \in e$. The following is our main theorem on component polynomials.

Theorem 7.4: When q is a non-zero component polynomial, there exists a finite structure **B** such that $q[[\mathbf{B}]] \neq 0$. Moreover, when $\phi_1(S_1), \ldots, \phi_n(S_n)$ are representatives of the equivalence classes $e_1, \ldots, e_n \in \mathcal{E}$ whose corresponding variables X_{e_i} appears in q, the structure **B** may be picked as a structure of the form provided by Lemma 7.3.

In order to establish this theorem, we will make use of the following known fact concerning multivariate polynomials; see for example [28, Lemma 2.1] for a proof.

Proposition 7.5: Let $p(x_1, \ldots, x_n)$ be a multivariate polynomial in n variables over a field F. For each $i \in [n]$, let d_i denote the degree of p as a polynomial in x_i , and suppose that $T_i \subseteq F$ is a set of size $d_i + 1$ or greater. Then, if p is not the zero polynomial, there exists a point $(t_1, \ldots, t_n) \in T_1 \times \cdots \times T_n$ such that $p(t_1, \ldots, t_n) \neq 0$.

Proof. (Theorem 7.4) Let $\phi_1(S_1), \ldots, \phi_n(S_n)$ be as described in the theorem statement. Let $m \ge 2$ be a value that exceeds the degree of each of the variables

 X_{e_1}, \ldots, X_{e_n} in q, and apply Lemma 7.3 to obtain structures $(\mathbf{B}_{(a_1,\ldots,a_n)} \mid (a_1,\ldots,a_n) \in [m]^n)$ and the corresponding functions $f_1, \ldots, f_n : [m] \to \mathbb{N}$. Evaluating q on these structures amounts to evaluating q when the variables (X_{e_1},\ldots,X_{e_n}) are given values in $f_1([m]) \times \cdots \times f_n([m])$. By Proposition 7.5, q must evaluate to a non-zero value on one of these structures.

Proof. (Theorem 7.1) Denote ℓ by $\sum_{i=1}^{m} c_i |\phi_i(V_i)|$ and let $(\mathbf{A}_1, V_1), \ldots, (\mathbf{A}_m, V_m)$ be the pairs corresponding to the formulas $\phi_1(V_1), \ldots, \phi_m(V_m)$. By rearranging the indices, we may assume for the sake of notation that $\mathbf{A}_1, \ldots, \mathbf{A}_k$ are homomorphically equivalent structures (where $k \in [m]$) and that for no i with $k < i \leq m$ does \mathbf{A}_i have a homomorphism to \mathbf{A}_1 .

For any structure B, it holds that one of the values $|\phi_1(\mathbf{B})|, \ldots, |\phi_k(\mathbf{B})|$ is non-zero if and only if all of them are. Now, for each i, define $\phi_i(V_i)$ from $\phi_i(V_i)$ by removing non-liberal components, that is, by removing each atom whose variables are all in a non-liberal component. Note that for every $i \in [k]$ and for every structure **B** such that $|\phi_i(\mathbf{B})| > 0$, we have that $|\hat{\phi}_i(\mathbf{B})| = |\phi_i(\mathbf{B})|$. Since the ϕ_i are pairwise not counting equivalent, it follows that the ϕ_i are pairwise not counting equivalent. For each formula $\phi_i(V_i)$, by considering its liberal connected components, we may define r_i to be a component polynomial which is a product of variables from $\{X_e \mid e \in \mathcal{E}\}$ such that $|\phi_i(\mathbf{B})| = r_i[\mathbf{B}]$ for all finite structures **B**. The products r_1, \ldots, r_k are pairwise distinct, so $r = c_1 r_1 + \cdots + c_k r_k$ is a non-zero component polynomial. By applying Lemma 7.3 with $A = A_1$ and then invoking Theorem 7.4, we obtain a finite structure D of the form $\mathbf{A} \times \mathbf{C}$ such that $r[[\mathbf{D}]] \neq 0$. Since there is by assumption a homomorphism from A_i to A and there is by Theorem 5.4 a homomorphism from \mathbf{A}_i to \mathbf{C} , we have $|\phi_i(\mathbf{D})| > 0$ for every $i \in [k]$. Consequently, $|\phi_i(\mathbf{D})| = |\phi_i(\mathbf{B})|$ by the observation from above. Since no structure A_i with $k < i \leq m$ maps homomorphically to A, we have $|\phi_{k+1}(\mathbf{D})| = \cdots =$ $|\phi_m(\mathbf{D})| = 0$ and hence $\ell(\mathbf{D}) = r[\mathbf{D}] \neq 0$.

VIII. PROOFS OF MAIN THEOREMS

The following lemma shows that if we are only interested in PP-representations of pp-formulas, we may restrict ourselves to basic PP-formulas.

Lemma 8.1: Let ϕ be a pp-formula and let ϕ' be a \sharp PP-representation of ϕ . Then there is a basic \sharp PP-sentence ϕ'' that is also a \sharp PP-representation of ϕ such that width $(\phi'') \leq$ width (ϕ') and \sharp -width $(\phi'') \leq \sharp$ -width (ϕ') .

Proof. (sketch) Using the normalization result given by Lemma 6.1 and the results of Section V, we can find numbers $c'_i \in \mathbb{Z}$ and pp-formulas ϕ''_i such that for all structures **B**

$$[\mathbf{B}, \phi'] = \sum_{i=1}^{\ell'} c'_i |\phi''_i(\mathbf{B})|, \qquad (1)$$

where the ϕ_i'' are pairwise not counting equivalent and have $qaw(\phi_i'') \leq width(\phi')$. Note that (1) is a linear combination.

Now note that $[\mathbf{B}, \phi'] = |\phi(\mathbf{B})|$, which gives another linear combination. If follows with Theorem 7.1 that (1) consists only of one summand with coefficient 1. Now applying the results of Section V yields the result.

Proof. (of Theorem 4.2 and Theorem 4.3; sketch) Observe that, by Lemma 8.1, we may restrict attention to basic \sharp PP-sentence. Then both statements can be shown with the results of Section V in a rather straightforward fashion.

Proof. (of Theorem 4.4; sketch) Let first Φ^+ satisfy the tractability condition. Then there is a constant k such that for all cores ψ of pp-formulas in Φ^+ we have $tw(\psi) \leq k$ and $tw(contract(\psi)) + 1 \leq k$. With the results of Section V it follows that $qaw(\psi) \leq 2k$ for such cores ψ . As discussed prior to the statement of Theorem 4.4, we can express, for each $\phi \in \Phi$, the function $|\phi(\cdot)|$ as a polynomial in unknowns $|\psi(\cdot)|$ with $\psi \in \Phi^+$ [18]. We can then substitute every term $|\psi'(\cdot)|$ by a \sharp PP-formula of width at most 2k with the results of Section V. This yields a \sharp PP-formula of width at most 2k.

For the other direction, assume the existence of a constant k such that each ep-formula in Φ has a $\sharp \text{EP-representation } \phi'$ with width $(\phi') \leq k$. Let ψ be a pp-formula from Φ^+ . We will show that ψ is equivalent to a pp-formula of quantifier-aware width at most k which completes the proof with Section V.

We first choose $\phi \in \Phi$ that witnesses $\psi \in \Phi^+$. Let ϕ' be the $\sharp \mathsf{EP}$ -representation of ϕ of width at most k. We first construct a linear combination $\ell_1(.)$ as in (1). Note that all summands of $\ell_1(.)$ have quantifier-aware width at most k as in the proof of Lemma 8.1.

We then construct a second representation of $|\phi(\mathbf{B})|$ as a linear combination $\ell_2(.)$. As in the first direction, for every structure we can express $|\phi(\mathbf{B})|$ as a polynomial in unknowns of the form $|\theta(\mathbf{B})|$. Arithmetic simplifications and elimination of counting equivalent terms gives the second linear combination $\ell_2(.)$ computing $|\phi(\mathbf{B})|$. We then argue that a term of the form $c \cdot |\psi(\mathbf{B})|$ must appear in $\ell_2(.)$.

We have that ℓ_1 and ℓ_2 compute $|\phi(\mathbf{B})|$ and are hence equal. With Theorem 7.1 it follows that ℓ_1 contains a term that is counting equivalent to ψ . Since all summands of $\ell_1(.)$ have quantifier-aware width at most k, the claim follows.

Proof. (of Theorem 4.5; sketch) Using the results of Section V and the normalization result of Lemma 6.1, ϕ can be turned into a linear combination as in (1). Moreover, starting with any such representation yields the same linear combination up to counting equivalence of the summands by Theorem 7.1. Now turning this linear combination into a \sharp PP-formula yields a \sharp PP-representation and minimizing the width of the summands with Theorem 4.3 gives a representation of optimal width.

REFERENCES

- R. Pichler and S. Skritek, "Tractable counting of the answers to conjunctive queries," in *Proceedings of the 5th Alberto Mendelzon International* Workshop on Foundations of Data Management, 2011.
- [2] G. Greco and F. Scarcello, "Counting solutions to conjunctive queries: structural and hybrid tractability," in *Proceedings of the 33rd ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems*, 2014, pp. 132–143.

- [3] A. Durand and S. Mengel, "Structural tractability of counting of solutions to conjunctive queries," in *Proceedings of the 16th International Conference on Database Theory (ICDT 2013)*, 2013.
- [4] H. Chen and S. Mengel, "A trichotomy in the complexity of counting answers to conjunctive queries," in 18th International Conference on Database Theory, ICDT 2015, March 23-27, 2015, Brussels, Belgium, 2015, pp. 110–126.
- [5] C. Papadimitriou and M. Yannakakis, "On the Complexity of Database Queries," *Journal of Computer and System Sciences*, vol. 58, no. 3, pp. 407–427, 1999.
- [6] J. Flum and M. Grohe, *Parameterized Complexity Theory*. Springer, 2006.
- [7] M. Grohe, "The complexity of homomorphism and constraint satisfaction problems seen from the other side," *Journal of the ACM*, vol. 54, no. 1, 2007.
- [8] I. Adler and M. Weyer, "Tree-width for first order formulae," in CSL 2009, 2009, pp. 71–85.
- [9] H. Chen and V. Dalmau, "Decomposing quantified conjunctive (or disjunctive) formulas," in *LICS*, 2012.
- [10] H. Chen and D. Marx, "Block-sorted quantified conjunctive queries," in *ICALP*, 2013.
- [11] H. Chen, "On the complexity of existential positive queries," ACM Trans. Comput. Log., vol. 15, no. 1, 2014.
- [12] H. Chen and M. Müller, "One hierarchy spawns another: graph deconstructions and the complexity classification of conjunctive queries," in Joint Meeting of the Twenty-Third EACSL Annual Conference on Computer Science Logic (CSL) and the Twenty-Ninth Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), CSL-LICS '14, Vienna, Austria, July 14 - 18, 2014, 2014, pp. 32:1–32:10.
- [13] H. Chen, "The tractability frontier of graph-like first-order query sets," in Joint Meeting of the Twenty-Third EACSL Annual Conference on Computer Science Logic (CSL) and the Twenty-Ninth Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), CSL-LICS '14, Vienna, Austria, July 14 - 18, 2014, 2014, p. 31.
- [14] N. Immerman, "Upper and lower bounds for first-order expressibility," J. Comput. System Sci., vol. 25, p. 7698, 1982.
- [15] M. Y. Vardi, "On the complexity of bounded-variable queries." in PODS'95, 1995, pp. 266–276.
- [16] B. Rossman, "Homomorphism preservation theorems," J. ACM, vol. 55, no. 3, 2008.
- [17] S. Abiteboul, R. Hull, and V. Vianu, Foundations of Databases. Addison-Wesley, 1995.
- [18] H. Chen and S. Mengel, "Counting answers to existential positive queries: A complexity classification," in *Proceedings of the 35th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems*, 2016, pp. 315–326.
- [19] M. Grohe and M. Thurley, "Counting homomorphisms and partition functions," *Contemporary Mathematics*, vol. 558, pp. 252–301.
- [20] A. A. Bulatov, "The complexity of the counting constraint satisfaction problem," J. ACM, vol. 60, no. 5, p. 34, 2013.
- [21] A. K. Chandra and P. M. Merlin, "Optimal implementation of conjunctive queries in relational data bases," in *Proceedings of STOC'77*, 1977, pp. 77–90.
- [22] F. V. Fomin and D. Kratsch, *Exact Exponential Algorithms*, 1st ed. New York, NY, USA: Springer-Verlag New York, Inc., 2010.
- [23] L. Lovsz, "Operations with structures," Acta Mathematica Academiae Scientiarum Hungarica, vol. 18, no. 3-4, pp. 321–328, 1967.
- [24] N. Immerman and E. Lander, "Describing graphs: A first-order approach to graph canonization," in *Complexity Theory Retrospective*, A. Selman, Ed. Springer New York, 1990, pp. 59–81.
- [25] S. Bova and H. Chen, "The complexity of width minimization for existential positive queries," in *ICDT*, 2014, pp. 235–244.
- [26] H. L. Bodlaender, "A linear-time algorithm for finding tree-decompositions of small treewidth," *SIAM J. Comput.*, vol. 25, no. 6, pp. 1305–1317, 1996. [Online]. Available: http://dx.doi.org/10.1137/S0097539793251219
- [27] P. Hell and J. Nesetril, "The core of a graph," *Discrete Math.*, vol. 109, pp. 117–126, 1992.
- [28] N. Alon, "Combinatorial nullstellensatz," Comb. Probab. Comput., vol. 8, no. 1-2, pp. 7–29, Jan. 1999.
- [29] V. Dalmau, P. G. Kolaitis, and M. Y. Vardi, "Constraint Satisfaction, Bounded Treewidth, and Finite-Variable Logics," in *Constraint Pro*gramming '02, ser. LNCS, 2002.

IX. MATERIAL

A. Proof of Lemma 5.2

Proof.[of Lemma 5.2] The main idea of the proof is that forcing the free variables of an \exists -component to appear above the quantified variables has a very similar effect as connecting them to a clique in the construction of the contract(ϕ). So from a quantifier-aware decomposition we get a decomposition of contract(ϕ) by restricting to the tree decomposition to the free variables. For the other direction, we can add the quantified variables to a tree decomposition of contract(ϕ) in a straightforward way. We now give the details.

For the first inequality, observe first that any quantifieraware tree decomposition of ϕ is a tree decomposition of ϕ , so tw $(\phi) + 1 \leq qaw(\phi)$ is obvious. To prove the inequality tw(contract (ϕ)) + 1 $\leq qaw(\phi)$, let $(T, (B(t))_{t\in T})$ be a quantifier-aware tree decomposition of ϕ . Introduce for every \exists -component of ϕ a new vertex v_C . Then substitute in every B(t) every non-free variable x of ϕ by v_C where C is the \exists -component that contains x. Call the result B'(t). We claim that $(T, (B'(t))_{t\in T})$ is a tree decomposition of contract (ϕ) . To see this, note that by the same argument as in Remark 5.1 we have for every \exists -component C a bag that contains $(V(C) \cap \text{free}(\phi)) \cup \{v_C\}$.

For the second inequality, first compute a tree decomposition $(T, (B(t))_{t \in T})$ of the contraction contract (ϕ) . Note that for every \exists -component C of ϕ there is a bag $B(t^*)$ that contains $V(C) \cap \text{free}(\phi)$, because this variable set forms a clique in contract(ϕ) and it is well known that for every clique in a graph every tree decomposition must contain a bag that contains this clique completely. Now compute a tree decomposition $(T', (B'(t))_{t \in T'})$ for $G[V(C) \setminus \text{free}(\phi)]$, where G is the primal graph of ϕ . Then construct for every $t \in T'$ a new bag $B''(t) := B'(t) \cup (V(C) \cap \text{free}(\phi))$. Finally, connect T to T' by connecting an arbitrary vertex of T'to t^* . Doing this for every \exists -component yields a quantifieraware tree decomposition of ϕ . Moreover, the width of the decomposition is at most $tw(contract(\phi)) + tw(\phi) + 1$ which completes the proof.

B. Proof of Lemma 5.3

Proof. The idea of the proof is to add edges to the graph of ϕ in such a way that the treewidth of the resulting graph is exactly $qaw(\phi)$; then, we apply standard algorithms for computing treewidth. To this end, let ϕ be a pp-formula with primal graph G and $S := \text{free}(\phi)$. For each \exists -component C of ϕ choose a vertex $x_C \in V(C) \setminus \text{free}(\phi)$ and connect it to all vertices $y \in V(C) \cap \text{free}(\phi)$; moreover, connect the vertices in $V(C) \cap \text{free}(\phi)$ by a clique. Call the resulting graph G'. We will show that the minimum of tw(G') + 1 over the choices of the x_C is $qaw(\phi)$.

We first show that for every choice of the x_C we have $\operatorname{tw}(G') + 1 \ge \operatorname{qaw}(\phi)$. To see this, fix a tree decomposition $(T, (B(t))_{t\in T})$ of G'. Since $V(C)\setminus\operatorname{free}(\phi)$ is connected in G', the bags containing $V(C)\setminus\operatorname{free}(\phi)$ are contained in a subtree T' of T. Moreover, because $\{x_c\} \cup (V(C) \cap \operatorname{free}(\phi))$ is a clique in G', we know that $\{x_c\} \cup (V(C) \cap \operatorname{free}(\phi)) \subseteq B(t^*)$ for some t^* in T'. Since none of the vertices in $V(C) \setminus \operatorname{free}(\phi)$ have any neighbors outside of V(C), we may assume that t^* is the root of T'. Then it is easy to see that $(T, (B_t)_{t \in T})$ can be turned into a quantifier-aware tree decomposition: We only have to potentially add a new bag $B(t^{**}) := V(C) \cap \operatorname{free}(\phi)$ and a vertex t^{**} in the decomposition. Then connect t^{**} to t^* and its parent and delete the edge between t^* and its parent.

For the other direction, let $(T, (B(t))_{t\in T})$ be a quantifieraware tree decomposition of ϕ . We will show that it is also a tree decomposition of G' for a choice of the x_C . First note that by the same argument as before, the vertices of $V(C) \cap \text{free}(\phi)$ are contained in a subtree T' of T. Let x'_C be the only variable of $V(C) \setminus \text{free}(\phi)$ that is contained in B(r) where B(r) is the root of T'. Note that by the same argument as in Remark 5.1, we know that B(r) contains $V(C) \cap \text{free}(\phi)$. Thus B(r) covers all edges introduced in the construction of G' when choosing $x_C = x'_C$. Thus $(T, (B(t))_{t\in T})$ is indeed a tree decomposition of G' for the right choice of the x_C .

Since computing tree decompositions is fixed parameter tractable when parameterized by the treewidth (see e.g. [6]), the only problem left to solve is the right choice of the x_C . But since the quantified variables of the different \exists -components are independent, we can do this choice independently for every \exists component C as follows: Construct G'' by choosing a vertex $x_C \in V(C) \cap \text{free}(\phi)$ and proceed as in the construction of G'. Now for all other \exists -components C' connect $V(C') \cap \text{free}(\phi)$ to a clique and delete all variables in $V(C') \setminus \text{free}(\phi)$. Clearly, trying all potential choices of x_C lets us optimize the choice for C. Doing this for all \exists -components gives the desired choice and thus the optimal quantifier-aware tree decomposition. \Box

C. Proof of Lemma 5.4

Proof. The proof relies on the observation that the condition on top(x) in quantifier-aware tree decompositions corresponds closely to the fact that free variables can only be closed by a *P*-quantifier in \sharp PP-formulas after the contained pp-formula has been casted by a *C*-quantifier. With this in mind, a \sharp PPformula is transformed into a pp-formula by making use of and inducting on the \sharp PP-formula's structure (viewed as a tree). The other direction is similar.

a) Let ϕ' be a basic \sharp PP-sentence. By potentially renaming quantified variables, make sure that every variable in ϕ' is either free or quantified exactly once. We construct ϕ by deleting all *C*-, *E*- and *P*-quantifiers and substituting all \times by \wedge . Obviously, the result is a pp-formula. Note that for every subformula ψ' of ϕ' , we have an associated subformula ψ of ϕ . For every subformula ψ of ϕ we define $lib(\psi)$ to be the variables of ψ' that are not quantified in ψ . We claim that for all \sharp -subformulas ψ' and every assignment *h* to free(ψ'),

$$[\mathbf{B}, \psi'](h) = |\{h' : \operatorname{lib}(\psi) \to B \mid h' \text{ extends } h, (\mathbf{B}, h') \models \psi\}|$$
(2)

We show (2) by induction on the structure of basic $\sharp PP$ formulas. If $\psi' = C\psi''$ for a pp-formula ψ'' , then we actually have $\psi = \psi''$. Moreover, h assigns to values to all liberal variables of ψ , so both sides of (2) are 1 if and only if h satisfies ψ . If $\psi' = EV\psi''$ or $\psi' = PV\psi''$, we get (2) directly from the semantics of \sharp -formulas and induction. Finally, if $\psi' = \psi'_1 \times \psi'_2$, we have that $\operatorname{lib}(\psi_1) \cap \operatorname{lib}(\psi_2) \subseteq \operatorname{lib}(\psi)$. Therefore, $\operatorname{lib}(\psi_1)$ and $\operatorname{lib}(\psi_2)$ overlap only at variables where a mapping $h' : \operatorname{lib}(\psi) \to B$ is defined, and (2) follows easily.

It remains to show the inequalities of the width measures. To this end, consider the syntax tree T of ϕ' . For every node t of T, define $B(t) := \operatorname{free}(\phi'_t)$ where ϕ'_t is the subformula of ϕ' that has t as its root. Note that $(T, (B(t))_{t \in T})$ satisfies the connectivity condition and is thus a tree decomposition of ϕ of width width $(\phi') - 1$. Also, $(T, (B(t))_{t \in T})$ is quantifier-aware because in ϕ' existential quantification is only allowed in the pp-part in which all free variables of ϕ are still free. This shows width $(\phi') \ge \operatorname{qaw}(\phi)$. Now observe that by deleting all bags that contain quantified variables we end up with a tree decomposition for $\operatorname{contract}(\phi)$. This shows \sharp -width $(\phi) \ge \operatorname{tw}(\operatorname{contract}(\phi)) + 1$.

b) Let now ϕ be a pp-formula and let $(T, (B(t))_{t \in T})$ be a nice quantifier-aware tree decomposition of ϕ of width k-1. For every \exists -component C of ϕ , the vertices $V(C) \setminus \text{free}(\phi)$ all lie in the bags of a subtree T_C of T. Moreover, we may w.l.o.g. assume that the bags in T_C do not contain any vertices not in V(C). Finally, we have that the bag $B(r_C)$ where r_c is the root of T_C contains $V(C) \cap \text{free}(\phi)$, because $(T, (B(t))_{t \in T})$ is quantifier-aware. The results of [29] assure that there is a pp-formula ϕ_C of width k that is logically equivalent to the pp-formula that we get by restricting ϕ to the atoms that have all of their variables in V(C).

We now construct for every $t \in T$ such that B(t) does not contain any quantified variables of ϕ a basic $\sharp PP$ -sentence ϕ'_t . So let t be a node of t with the desired properties. Let atom(t)be the atoms of ϕ containing only variables in B(t) and set $\bar{\phi}_t := \prod_{\psi \in atom(t)} C\psi$. If t has no children, set $\phi_t := \bar{\phi}_t$.

If t has a child t' such that $B(t')\setminus \text{free}(\phi) \neq \emptyset$, then t has only that one child because $(T, (B(t))_{t\in T})$ is nice. Let C be the unique \exists -component of the variable in $B(t')\setminus \text{free}(\phi)$. We set $\phi_t := \overline{\phi}_t \times C\phi_C$.

If t has a child t' such that $B(t')\setminus \text{free}(\phi) = \emptyset$ and a variable x is forgotten when going from t' to t, then set $\phi_t := Px\phi_{t'}$.

If t has a child t' and a variable x is introduced when going from t' to t, then set $\phi_t := \overline{\phi}_t \times Ex\phi_{t'}$.

If t has two children t_1 and t_2 , then note that

$$B(t_1) \setminus \operatorname{free}(\phi) = B(t_2) \setminus \operatorname{free}(\phi) = \emptyset$$

Moreover, free (ϕ_{t_1}) = free (ϕ_{t_2}) . We define $\phi_t := \phi_{t_1} \times \phi_{t_2}$. Set $\phi' := P$ free $(\phi_r)\phi_r$ where r is the root of ϕ .

An easy induction along the construction of ϕ similar to that in a) shows that ϕ' does indeed compute the correct value for every structure **B**. Moreover, the width of ϕ' is at most k which completes the proof.

If we do not have a bound on $qaw(\phi)$ but only on $tw(contract(\phi))$, the same construction as above yields the the bound $tw(contract(\phi))+1 \ge \sharp$ -width (ϕ') . The only difference

is that we do not have to bound the width of the pp-formulas with [29]. \Box

X. PROOF OF LEMMA 6.2

Proof. In a first step, we transform ψ into a logically equivalent ep-formula $\psi^d = \bigvee_{i=1}^s \psi_i$ where the ψ_i are pp-formulas; this can be done without increasing width [11]. Then we claim that ϕ is logically equivalent to

$$\phi' = \sum_{J \subseteq [s], J \neq \emptyset} (EL(-1)^{|J|+1}) \prod_{i \in J} C(\psi_i, L).$$

First note that this is a well-formed \sharp -formula, because for all additions and multiplications the free variables of all operands are *L*. It remains to show that ϕ' is logically equivalent to ϕ . So fix **B** and $h: L \to B$.

If h does not satisfy ψ , then $[\mathbf{B}, \phi](h) = 0$. Since h does not satisfy any ψ_i , it is easy to see that $[\mathbf{B}, \phi](h) = 0$ as well.

Assume h satisfies ψ ; say that $\psi_1, \ldots, \psi_\ell$ are the disjuncts that it satisfies. By definition $[\mathbf{B}, \phi](h) = 1$. Also, $[\mathbf{B}, \phi'](h)$ is equal to

$$\sum_{J\subseteq[s],J\neq\emptyset} (-1)^{|J|+1} \prod_{i\in J} [\mathbf{B}, C(\psi_i, L)](h)$$

which in turn is equal to $\sum_{J \subseteq [\ell], J \neq \emptyset} (-1)^{|J|+1}$; this latter quantity is equal to the sum

$$\ell - \binom{\ell}{2} + \binom{\ell}{3} - \ldots \pm \binom{\ell}{\ell} = (1-1)^{\ell} + 1 = 1.$$

It is readily seen that the width of ϕ' is not bigger than that of ϕ .

A. Proofs for Section VII

Proof. [Explanation for Lemma 7.2] Let p be a univariate polynomial with positive integer coefficients and variable X. Fix a representation of p as a term with 1 and X as the inputs and where addition and multiplication are the operations. For a vocabulary τ , let \mathbf{I}_{τ} denote the τ -structure with universe $\{a\}$ and where each relation symbol $R \in \tau$ has $R^{\mathbf{I}_{\tau}} = \{(a, \ldots, a)\}$. For each structure \mathbf{B} over vocabulary τ , we define $p(\mathbf{B})$ as the τ -structure obtained by evaluating the representation of pby interpreting 1 as \mathbf{I}_{τ} , X as \mathbf{B} , addition as the disjoint union \oplus of two structures, and multiplication as the product of two structures.

Here, by the disjoint union $\mathbf{A} \oplus \mathbf{B}$ of two structures \mathbf{A} , \mathbf{B} , we mean the structure obtained as follows: rename the elements of \mathbf{A} , \mathbf{B} so that their universes A, B are disjoint, and then take the structure with universe $A \cup B$ where $R^{\mathbf{A} \oplus \mathbf{B}} = R^{\mathbf{A}} \cup R^{\mathbf{B}}$ for each relation symbol R. By the product $\mathbf{A} \times \mathbf{B}$ of two structures \mathbf{A}, \mathbf{B} , we mean the structure with universe $A \times B$ and where $R^{\mathbf{A} \times \mathbf{B}} = \{((a_1, b_1), \dots, (a_k, b_k)) \mid (a_1, \dots, a_k) \in R^{\mathbf{A}}, (b_1, \dots, b_k) \in R^{\mathbf{B}}\}.$

Lemma 7.2 can be proved by a straightforward induction on the structure of p, using the observations that $|\phi(\mathbf{I}_{\tau})| = 1$, $|\phi(\mathbf{D} \oplus \mathbf{D}')| = |\phi(\mathbf{D})| + |\phi(\mathbf{D}')|$, and $|\phi(\mathbf{D} \times \mathbf{D}')| = |\phi(\mathbf{D})| \cdot |\phi(\mathbf{D}')|$.

B. Full proofs for Section VIII

Remember that we call a \sharp PP-formula ϕ basic if it does not contain + nor subformulas of the form n, where $n \in \mathbb{Z}$. The following lemma shows that if we are only interested in \sharp PP-representations of pp-formulas, we may restrict ourselves to basic \sharp PP-formulas.

Lemma 10.1: Let ϕ be a pp-formula and let ϕ' be a \sharp PP-representation of ϕ . Then there is a basic \sharp PP-sentence ϕ'' that is also a \sharp PP-representation of ϕ such that width $(\phi'') \leq$ width (ϕ') and \sharp -width $(\phi'') \leq \sharp$ -width (ϕ') .

Proof. With Lemma 6.1 we may assume that ϕ' is flat, so let $\phi' = \sum_{i=1}^{\ell} \psi'_i \times \phi'_i$ where each ψ'_i is constant and each ϕ'_i is basic. As the \sharp PP-formulas ϕ'_i are basic, by Lemma 5.4 (a) we obtain that there are pp-formulas ϕ_i representing them, where for each finite structure **B** we have $|\phi_i(\mathbf{B})| = [\mathbf{B}, \phi'_i]$. Set c_i to be the value that the constant formula ψ'_i evaluates to. We have that, for every structure **B**, $[\mathbf{B}, \phi'] = \sum_{i=1}^{\ell} c_i |\phi_i(\mathbf{B})|$, where $c_i \in \mathbb{Z}$ and ϕ_i is a pp-formula.

Now combine the summands of counting equivalent ppformulas to get a linear combination with

$$[\mathbf{B}, \phi'] = \sum_{i=1}^{\ell'} c'_i |\phi''_i(\mathbf{B})|, \qquad (3)$$

where the ϕ''_i are pairwise not counting equivalent pp-formulas and $c'_i \in \mathbb{Z} \setminus \{0\}$. Note that for all ϕ''_i we have by Lemma 5.4 that $qaw(\phi''_i) \leq width(\phi')$ and $tw(contract(\phi''_i)) + 1 \leq$ \sharp -width(ϕ').

Since ϕ' is a \sharp PP-representation of ϕ , we have $[\mathbf{B}, \phi'] = |\phi(\mathbf{B})|$ for all structures **B**. So, we have two linear combinations, $|\phi(\mathbf{B})|$ and that of (3) that evaluate to the same value. By taking the difference and invoking Theorem 7.1, we obtain that these two linear combinations are equal, up to counting equivalence of the pp-formulas. Since the linear combination $|\phi(\mathbf{B})|$ only has one summand with coefficient 1, it follows that the linear combination in (3) consists only of one summand with coefficient 1. Let $|\psi(\mathbf{B})|$ be that summand. We have that ψ and ϕ are counting equivalent. Now we apply Lemma 5.4 (b) on input ψ to construct a formula with the desired properties.

Another ingredient that we will use in the proofs of the main theorems is a syntactic characterization of counting equivalence. To this end, we say that two pp-formulas (\mathbf{A}, V) , (\mathbf{A}', V') over the same signature are *renaming equivalent* if there exist surjections $h_1 : V \to V'$ and $h_2 : V' \to V$ that can be extended to homomorphisms $\bar{h} : \mathbf{A} \to \mathbf{A}'$ and $\bar{h'} : \mathbf{A}' \to \mathbf{A}$, respectively.

A crucial result of [18] is that renaming equivalence is a syntactic characterization of counting equivalence.

Theorem 10.2: [18, Theorem 5.4] Any two pp-formulas $\phi_1(V), \phi_2(V')$ are counting equivalent if and only if they are renaming equivalent.

We will make use of the following easy corollary.

Corollary 10.3: Let $\phi(V)$ and $\phi'(V')$ be counting equivalent pp-formulas. Then there is a pp-formula ψ that is logically equivalent to ϕ such that $qaw(\psi) \leq qaw(\phi')$.

Proof. By Theorem 10.2, we have that ϕ and ϕ' are renaming equivalent. Construct $\psi(V)$ from ϕ' by renaming variables so that $\phi(V)$ and $\psi(V)$ are renaming equivalent via the identity mapping $i : V \to V$ on V. We claim that ψ and ϕ are logically equivalent, that is, for any structure **B** and any mapping $f : V \to B$, it holds that **B**, $f \models \psi$ iff **B**, $f \models \phi$.

We prove the forward direction; the backward direction is analogous. Let $[\mathbf{A}_{\psi}, V]$ and $[\mathbf{A}_{\phi}, V]$ be the structure views of the pp-formulas ψ and ϕ , respectively. Suppose that $\mathbf{B}, f \models \psi$. By the work of Chandra and Merlin [21] discussed in the preliminaries, we obtain that there is an extension $g : A_{\psi} \rightarrow B$ of f that is a homomorphism from \mathbf{A}_{ψ} to \mathbf{B} . Since ϕ and ψ are renaming equivalent via the identity mapping i on V, there exists a homomorphism h from \mathbf{A}_{ϕ} to \mathbf{A}_{ψ} that extends i. By composing h and g, we obtain an extension of f that is a homomorphism from \mathbf{A}_{ϕ} to \mathbf{B} . By the work of Chandra and Merlin [21] discussed in the preliminaries, we obtain that $\mathbf{B}, f \models \phi$.

Proof. [of Theorem 4.2] We start with the second statement. Let first Φ satisfy the contraction condition. Then there is a constant k such that for all cores ϕ of pp-formulas in Φ we have tw(contract(ϕ)) + 1 \leq k. But then Lemma 5.4 yields basic \sharp PP-representations ϕ' with \sharp -width(ϕ') \leq k.

Now assume there is a constant k such that every formula ϕ in Φ has a \sharp PP-representation ϕ' such that \sharp -width $(\phi') \leq k$. By Lemma 10.1 we may assume that ϕ' is basic. Then by Lemma 5.4 we obtain a pp-formula ϕ'' that is counting equivalent to ϕ and such that we have tw $(contract(\phi'')) + 1 \leq k$. As ϕ and ϕ'' are counting equivalent, by Theorem 10.2, they are renaming equivalent. Consequently, the treewidth of contract(\cdot) applied to the cores of ϕ and ϕ'' , is the same. Since passing to the core does not increase the treewidth of contract(\cdot), the treewidth of contract(\cdot) applied to the core of ϕ is $\leq k - 1$. By Corollary 10.3, we have that ϕ is logically equivalent to a formula ψ with tw $(contract(\psi)) + 1 \leq k$. Consequently, Φ satisfies the contraction condition.

For the first statement, let first Φ satisfy the tractability condition. Then there is a constant k such that for all cores ϕ of pp-formulas in Φ we have $\mathsf{tw}(\phi) \leq k$ and $\mathsf{tw}(\mathsf{contract}(\phi)) + 1 \leq k$. It follows that $\mathsf{qaw}(\phi) \leq 2k$ by Lemma 5.2. Then Lemma 5.4 yields a basic $\sharp\mathsf{PP}$ -representation ϕ' with $\mathsf{width}(\phi') \leq 2k$.

Now assume there is a constant k such that every formula ϕ in Φ has a \sharp PP-representation ϕ' such that width $(\phi') \leq k$. We may again assume that ϕ' is basic. Then Lemma 5.4 gives a pp-formula ϕ'' that is counting equivalent to ϕ such that $qaw(\phi) \leq k$. Using Corollary 10.3 it follows that ϕ is logically equivalent to a formula ψ with $qaw(\phi) \leq k$. Now applying Lemma 5.2 shows that Φ satisfies the tractability condition. \Box

Proof. [of Theorem 4.3] By Lemma 10.1 we may assume that the desired \sharp PP-representation is basic. It then suffices to minimize the width of the factor that is not constant. To do so, we translate to a pp-formula with Lemma 5.4 (a), then minimize the width with Lemma 5.3 and finally translate back to a basic \sharp PP-formula with Lemma 5.4 (b).

In what follows, we will use the following lemma that summarizes the main construction of the proof of Theorem 3.1 in [18].

Lemma 10.4: For every set Φ of ep-formulas there is a set Φ^+ of pp-formulas such that the following holds: For every $\phi \in \Phi$ there are pp-formulas $\phi_1, \ldots, \phi_\ell \in \Phi^+$, pp-sentences $\psi_1, \ldots, \psi_t \in \Phi^+$ and non-zero integers c_1, \ldots, c_l such that the following holds for every structure **B**:

- If **B** satisfies one of the sentences ψ_i , then $|\phi(\mathbf{B})| = |B|^{|\text{free}(\phi)|}$.
- Otherwise, we have $|\phi(\mathbf{B})| = \sum_{i=1}^{\ell} c_i |\phi_i(\mathbf{B})|$.

Moreover, the ϕ_i are pairwise not counting equivalent and there is no homomorphism from any sentence ψ_i to any ψ_j for $i \neq j$ and no homomorphism to any ϕ_j .

Finally, for every $\phi' \in \Phi^+$ we have that ϕ' appears as a ϕ_i or ψ_i as above for a $\phi \in \Phi$.

Proof. [of Theorem 4.4] First suppose that Φ^+ satisfies the tractability condition. Then there is a constant k such that for all cores ψ of pp-formulas in Φ^+ we have tw $(\psi) \leq k$ and tw(contract (ψ)) + 1 $\leq k$. With Lemma 5.2 it follows that qaw $(\psi) \leq 2k$ for the cores ψ of the pp-formulas in Φ . Then Lemma 10.4 provides pp-formulas $\phi_1, \ldots, \phi_\ell \in \Phi^+$, coefficients c_1, \ldots, c_ℓ , and pp-sentences $\psi_1, \ldots, \psi_t \in \Phi^+$ such that the following holds for every structure **B**: If **B** satisfies any sentence ψ_i , then $|\phi(\mathbf{B})| = |B|^{|\text{free}(\phi)|}$; otherwise, $|\phi(\mathbf{B})| = \sum_{i=1}^{\ell} c_i |\phi_i(\mathbf{B})|$. Let $|\text{free}(\phi)| = r$. Then $|\phi(\mathbf{B})|$ is equal to $(\prod_{i=1}^{t} (1 - |\psi_i(\mathbf{B})|))(\sum_{i=1}^{\ell} c_i |\phi_i(\mathbf{B})|) + |B|^r (1 - \prod_{i=1}^{t} (1 - |\psi_i(\mathbf{B})|))$. Now let $\phi'_1, \ldots, \phi'_\ell$ be the \sharp -formulas we get by applying Lemma 5.4 on $\phi_1, \ldots, \phi_\ell$ and then multiplying respectively by the \sharp -formulas

$$E$$
free $(\phi)c_1, \ldots, E$ free $(\phi)c_\ell$

that correspond to the coefficients c_i . By Lemma 5.4, we have that the width of each ψ'_i is bounded above by 2k.

Then set ψ to be $Plib(\phi)$ applied to the sum of

$$(E \operatorname{free}(\phi)(1 - C(\bigvee_{i=1}^{t} \psi_i, \emptyset))) \times (\sum_{i=1}^{\ell} \phi'_i)$$

and

$$E \operatorname{free}(\phi) C(\bigvee_{i=1}^{t} \psi_i, \emptyset).$$

Clearly, ψ is a $\sharp \text{EP}$ -representation of ϕ . Moreover, the width of ψ is the maximal width of the ψ_i and the ϕ'_i , which is bounded above by 2k.

The first direction of the second item can be proved completely analogously using tw(contract(.)) and #width of qaw and width.

For the other direction, assume that there is a constant k such that every ep-formula in Φ possesses a $\sharp \text{EP}$ -representation ϕ' with width $(\phi) \leq k$. Let ψ be a pp-formula from Φ^+ . We will show that ψ is equivalent to a pp-formula of quantifier-aware width at most k which completes the proof with Lemma 5.4.

We first choose $\phi \in \Phi$ such that ψ appears as a ϕ_i or ψ_i as in Lemma 10.4. Let ϕ' be the $\sharp \text{EP}$ -representation of ϕ of width at most k. With Lemma 6.1 we may assume that ϕ' is flat, i.e., it has the form $\sum_{i=1}^{\ell} \theta'_i \times \phi'_i$, where the θ'_i are constant and the ϕ'_i are basic. As in the proof of Lemma 10.1, this yields for every **B**

$$|\phi(\mathbf{B})| = \sum_{i=1}^{\ell} \bar{c}_i |\bar{\phi}_i(\mathbf{B})|, \qquad (4)$$

where the ϕ_i are pairwise not counting equivalent. Moreover, $qaw(\bar{\phi}_i) \leq k$.

We will now construct a second representation of $|\phi(\mathbf{B})|$ as a linear combination. To this end, let $|\text{lib}(\phi)| = r$ and let $\phi_1, \ldots, \phi_\ell, \psi_1, \ldots, \psi_t$ and c_1, \ldots, c_ℓ be as in Lemma 10.4. Then as before, for every structure $|\phi(\mathbf{B})| = \prod_{i=1}^t (1 - |\psi_i(\mathbf{B})|)(\sum_{i=1}^\ell c_i |\phi_i(\mathbf{B})|) + |B|^r (1 - \prod_{i=1}^t (1 - |\psi_i(\mathbf{B})|)).$ Now multiplying the righthand side out, we get that $|\phi(\mathbf{B})|$ can be expressed as a weighted sum of terms of the form $|\phi_i(\mathbf{B})| \cdot \prod_{j \in J} |\psi_j(\mathbf{B})|$ and of the form $|B|^r \prod_{j \in J} |\psi_j(\mathbf{B})|$. These terms are equivalent to

$$|(\phi_i \wedge \bigwedge_{j \in J} \psi_j)(\mathbf{B})|$$

and

$$|(\psi^r \wedge \bigwedge_{j \in J} \psi_j)(\mathbf{B})|,$$

since the ψ_j are sentences; here, ψ^r is the true formula with liberal variables $\operatorname{lib}(\phi)$.

Now combine counting equivalent summands to get a linear combination

$$\phi(\mathbf{B})| = \sum_{i=1}^{\ell} c'_i |\phi'_i(\mathbf{B})|.$$
(5)

We claim that the linear combination in (5) contains c. $|\psi(\mathbf{B})|$ or $c \cdot |\psi^r \wedge \psi(\mathbf{B})|$ for some $c \neq 0$ as a summand. To see this, consider first the case that ψ is not a sentence. Note that in this case $c \cdot \psi(\mathbf{B})$ appears in the weighted sum from Lemma 10.4 and thus (for $J = \emptyset$) also in the sum we get before combining counting equivalent summands. Moreover, ψ is not counting equivalent to any other summand ψ' in this sum. To see this, note first that by Lemma 10.4 there is no homomorphism from any of the ψ_i to ψ . Moreover, the ϕ_i are pairwise not counting equivalent. Thus $c|\psi|$ appears in (5) in this case. If ψ is one of the sentences ψ_i , then $\psi^r \wedge \psi$ is contained in the sum before combining counting equivalent summands (for $J = \{i\}$). Obviously, $\psi^r \wedge \psi$ is not counting equivalent to any conjunction that contains a non-sentences ψ_i . Moreover, since there are no homomorphism between $\psi = \psi_i$ and ψ_i by Lemma 10.4, $\psi^r \wedge \psi$ is not counting equivalent to any other summand, so it must be contained in (5).

It follows that (4) and (5) give two linear combinations that are equal for every structure **B**. Using Theorem 7.1 it follows that (4) contains a summand $c \cdot |\bar{\psi}|$ that is counting equivalent to ψ . Moreover, $qaw(\bar{\psi}) \leq k$. Invoking Corollary 10.3, it follows that ψ is logically equivalent to a formula with quantifieraware width at most k.

Again, the second direction of the second item follows completely analogously. $\hfill \Box$

Proof. [of Theorem 4.5] The algorithm first comptues a #EPrepresentation ψ of the input $\phi \in \Phi$, such as that given by Proposition 3.1. By Lemma 6.1 we may assume that ψ is flat, i.e., it has the form $\psi = \sum_{i=1}^{\ell} \psi_i \times \phi_i$ where the ψ_i are constant and the ϕ_i are basic. Note that width $(\psi) = \max_{i \in \ell} (\text{width}(\phi_i))$.

Note that every flat #EP-representation of ϕ can be turned into a linear combination as in (3). Moreover, starting with any such representation yields the same linear combination up to counting equivalence of the summands by Theorem 7.1. Now turning this linear combination into a flat #PP-formula, one obtains a #PP-representation; minimizing the width of the summands with Theorem 4.3 gives a representation of optimal width.