# Ground states for a coupled nonlinear Schrödinger system

Filipe Oliveira\*

Department of Mathematics, FCT-UNL-CMA Universidade Nova de Lisboa Caparica Campus, 2829-516, Portugal

#### Abstract

We study the existence of ground states for the coupled Schrödinger system

$$\begin{cases} -\Delta u + u &= |u|^{2q-2}u + b|v|^{q}|u|^{q-2}u \\ -\Delta v + \omega^{2}v &= |v|^{2q-2}v + b|u|^{q}|v|^{q-2}v \end{cases}$$
(1)

in  $\mathbb{R}^n$ , for  $\omega \ge 1$ , b > 0 (the so-called "attractive case") and q > 1  $(q < \frac{n}{n-2} \text{ if } n \ge 3)$ . We improve for several ranges of  $(q, n, \omega)$  the known results concerning the existence of positive ground state solutions to (1) with non-trivial components. In particular, we prove that for 1 < q < 2 such ground states exist in all dimensions and for all values of  $\omega$ , which constitutes a drastic change of behaviour with respect to the case  $q \ge 2$ . Furthermore, for q > 2 and in the one-dimensional case n = 1, we improve the results in [14].

**Keywords:** Non-trivial ground states; Coupled nonlinear Schrödinger Systems; Nehari Manifold.

AMS Subject Classification: 35J20, 35J50, 35J60

#### 1 Introduction

In this paper we consider the system

$$\begin{cases} -\Delta u + \lambda_1 u = |u|^{2q-2}u + b|v|^q |u|^{q-2}u \\ -\Delta v + \lambda_2 v = |v|^{2q-2}v + b|u|^q |v|^{q-2}v, \end{cases}$$
(2)

<sup>\*</sup>e-mail address: fso@fct.unl.pt

with  $u, v : \mathbb{R}^n \to \mathbb{R}$   $(n \ge 1), q > 1, b > 0$  and  $\lambda_1, \lambda_2 > 0$ , which appears in several physical contexts, namely in nonlinear optics (see [1] and the references therein).

By rescaling the x variable and/or inverting the roles of u and v, it is easy to see that (2) can be reduced, without loss of generality, to the system

$$\begin{cases} -\Delta u + u &= |u|^{2q-2}u + b|v|^{q}|u|^{q-2}u \\ -\Delta v + \omega^{2}v &= |v|^{2q-2}v + b|u|^{q}|v|^{q-2}v, \qquad \omega \ge 1. \end{cases}$$
(3)

In the last years, this system has been extensively studied by many authors (see for instance [2], [11], [12], [17]). In particular, in [4] and [5] the authors studied the case q = 2 and n = 2, 3, proving the existence of a constant  $\Lambda > 0$ depending on  $\omega$  such that for  $b < \Lambda$  the system (3) admits a non-trivial radial solution  $(u, v) \neq (0, 0)$  (with u, v > 0 if b > 0). The authors also showed the existence of another constant  $\Lambda' \ge \Lambda$  such that for  $b > \Lambda'$  the system possesses a radial ground state solution  $W_* = (u_*, v_*)$   $(u_*, v_* > 0)$ , in the sense that  $W_*$  minimizes the energy functional associated to (3) among all solutions in  $(u, v) \in H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n) \setminus \{(0, 0)\}$ . In [9] Ikoma and Tamaka showed that for  $0 < b < \min\{\Lambda, 1\}$ , the solutions found in [4],[5] are in fact also least energy solutions.

In [14], following some of the ideas presented in [15], the authors proved the existence of a radial non-trivial ground state solution  $(u^*, v^*)$   $(u^*, v^* \ge 0)$ for every b > 0 and for (q, n) satisfying

$$1 < q < \begin{cases} +\infty & if \quad n = 1, 2\\ \frac{n}{n-2} & if \quad n \ge 3. \end{cases}$$

$$\tag{4}$$

Furthermore, it is shown that for

$$b \ge \mathcal{C}_{\omega,n,q} := \frac{1}{2} \left[ 1 + \frac{n}{2} \left( 1 - \frac{1}{q} \right) + \frac{1}{w^2} \left( 1 - \frac{n}{2} \left( 1 - \frac{1}{q} \right) \right) \right]^q \omega^{2q - n(q-1)} - 1 \quad (5)$$

there exists a ground state  $(u^*, v^*)$  with  $u^*, v^* > 0$ .

In the present paper we will prove the existence of a positive radial decreasing ground state solution to (3) for all (q, n) satisfying the condition (4). Exploring this radial decay, we improve the constant  $C_{\omega,n,q}$  derived in [14] for all q > 1 and large  $\omega$  in the case n = 1 and for all 1 < q < 2 in any dimension, in fact replacing it by 0 in the latter case. When dealing with the system (3) it is often necessary to treat the case n = 1 separately due to the lack of compactness of the injection  $H^1_d(\mathbb{R}) \hookrightarrow L^q(\mathbb{R}), q > 2$ , where  $H^1_d(\mathbb{R})$  denotes the space of the radially symmetric functions of  $H^1(\mathbb{R})$ . This lack of compactness is, in a sense, a consequence of the inequality

$$|u(x)| \le C|x|^{\frac{1-n}{2}} ||u||_{H^1(\mathbb{R}^n)}$$
(6)

for  $u \in H^1_d(\mathbb{R})$ . Indeed, (6) gives no decay in the case n = 1. However, if u is also radially decreasing, it is easy to establish that

$$|u(x)| \le C|x|^{-\frac{n}{2}} ||u||_{L^2(\mathbb{R}^n)},$$

which provides decay in all space dimensions, hence compacity by applying the classical Strauss' compactness lemma ([16]). Hence, putting

$$H^1_{rd}(\mathbb{R}^n) = \{ u \in H^1_d(\mathbb{R}^n) : u \text{ is radially decreasing} \},\$$

we get the compactness of the injection  $H^1_{rd}(\mathbb{R}^n) \hookrightarrow L^q(\mathbb{R}^n)$  for all  $n \geq 1$ (see the Appendix of [3] for more details). We will use this fact to present a unified approach for the problem of the energy minimization of (3), valid in all space dimensions.

Before stating our results more precisely, and following the functional settings in [4], [5] and [14], let us introduce a few notations: we denote by  $\|\cdot\|_q$  the standard  $L^q(\mathbb{R}^n)$  norm and, for  $(u, v) \in E := H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n)$ , we put

$$||(u,v)||_{\dot{\omega}}^2 := ||u||^2 + ||v||_{\dot{\omega}}^2 := ||u||_2^2 + ||\nabla u||_2^2 + \omega^2 ||v||_2^2 + ||\nabla v||_2^2.$$

We introduce the energy functional associated to (3),

$$I(u,v) := \frac{1}{2} \|(u,v)\|_{\dot{\omega}}^2 - \frac{1}{2q} \Big( \|u\|_{2q}^{2q} + \|v\|_{2q}^{2q} + 2b\|uv\|_q^q \Big),$$

noticing that (u, v) is a solution of (3) if and only if  $\nabla I(u, v) = 0$ . We will study the minimization problem

$$\inf\{I(u,v): (u,v) \in \mathcal{N}\},\tag{7}$$

where the so-called Nehari manifold  $\mathcal{N}$  is defined by

$$\mathcal{N} := \{ (u, v) \in H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n) : (u, v) \neq (0, 0), \nabla I(u, v) \perp (u, v) \},\$$

that is,  $(u, v) \in \mathcal{N}$  if and only if  $(u, v) \neq (0, 0)$  and

$$\tau(u,v) := \langle \nabla I(u,v), (u,v) \rangle_{L^2} = \|(u,v)\|_{\dot{\omega}}^2 - \left( \|u\|_{2q}^{2q} + \|v\|_{2q}^{2q} + 2b\|uv\|_q^q \right) = 0.$$

As pointed out in [4] for the case q = 2, we notice that

$$\langle \nabla \tau(u,v), (u,v) \rangle_{L^2} = 2 \| (u,v) \|_{\dot{\omega}}^2 - 2q \Big( \|u\|_{2q}^{2q} + \|v\|_{2q}^{2q} + 2b \|uv\|_q^q \Big),$$

and, if  $(u, v) \in \mathcal{N}$ ,

$$\langle \nabla \tau(u,v), (u,v) \rangle_{L^2} = 2(1-q) ||(u,v)||^2_{\dot{\omega}} < 0$$
 (8)

which shows that  $\mathcal{N}$  is locally smooth.

Furthermore, it is easy to check that  $[h_1, h_2]$ Hess  $\tau_{(0,0)} {}^t[h_1, h_2] > 0$  for all  $(h_1, h_2) \neq (0, 0)$ : (0, 0) is a strict minimizer of  $\tau$ , hence an isolated point of the set  $\{\tau(u, v) = 0\}$ , implying that  $\mathcal{N}$  is a complete manifold. Finally, any critical point of I constrained to  $\mathcal{N}$  is a critical point of I. Indeed, let us consider  $(u, v) \in \mathcal{N}$  a critical point of I constrained to  $\mathcal{N}$ . There exists a Lagrange multiplier  $\lambda$  such that  $\nabla I(u, v) = \lambda \nabla \tau(u, v)$ . By taking the  $L^2$  scalar product with (u, v),

$$\langle \nabla I(u,v), (u,v) \rangle_{L^2} = \lambda \langle \nabla \tau(u,v), (u,v) \rangle_{L^2},$$

that is, in view of (8),  $0 = \lambda(2-2q) ||(u,v)||^2_{\dot{\omega}}$ , hence  $\lambda = 0$  and  $\nabla I(u,v) = 0$ .

Putting  $E_{rd} = H_{rd}^1 \times H_{rd}^1$  the cone of symmetric radially decreasing nonnegative functions of E, we will prove the following result:

**Theorem 1.1** Let  $n \geq 1$  and q > 1, with  $q \leq \frac{n}{n-2}$  if n > 3. There exists a minimizing sequence  $(u_n, v_n) \in E_{rd}$  for the minimization problem (7). Furthermore,  $(u_n, v_n) \rightarrow (u_*, v_*) \in E_{rd}$  strongly in  $H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n)$ . In particular

$$I(u_*, v_*) = \min_{\mathcal{N}} I(u, v) = \min_{\mathcal{N} \cap E_{rd}} I(u, v)$$
  
= min{ $I(u, v) : (u, v) \neq (0, 0) \text{ and } \nabla I(u, v) = 0$ }. (9)

Concerning the existence of ground states with non-trivial components, we will show:

**Theorem 1.2** Let  $n \ge 1$  and 1 < q < 2, with  $q < \frac{n}{n-2}$  if  $n \ge 3$ . Then for all b > 0 there exists a ground state solution  $(u, v) \in E_{rd}$  to (3) with u > 0 and v > 0.

**Theorem 1.3** Let n = 1 and  $q \ge 2$ . If

$$b \ge \mathcal{D}_{\omega,q} = \frac{2^q - 1}{2} \omega^{1 + \frac{q}{2}} - \frac{1}{2} \omega^{-\frac{q}{2}}$$
(10)

there exists a ground state solution  $(u, v) \in E_{rd}$  to (3) with u > 0 and v > 0.

Notice that

$$\mathcal{D}_{\omega,q} < \mathcal{C}_{\omega,1,q} = \frac{1}{2} \left( \frac{3}{2} - \frac{1}{2q} + \frac{1}{\omega^2} \left( \frac{1}{2} + \frac{1}{2q} \right) \right)^q \omega^{1+q} - 1$$

at least for large values of  $\omega$ .

#### 2 Proof of Teorem 1.1

We begin by observing that for  $(f,g) \in E$ ,  $(f,g) \neq (0,0)$ , with  $\tau(f,g) \leq 0$ , there exists  $t \in ]0,1]$  such that  $(tf,tg) \in \mathcal{N}$ . Indeed, if  $\tau(f,g) = 0$ , we choose t = 1. If  $\tau(f,g) < 0$  we simply notice that

$$\tau(tf, tg) = t^2 \Big( \|(f, g)\|_{\dot{\omega}}^2 - t^{2q-2} (\|f\|_{2q}^{2q} + \|g\|_{2q}^{2q} + 2b\|fg\|_q^q) \Big) := t^2 T_{f,g}(t),$$

with  $T_{f,g}(0) > 0$  and  $T_{f,g}(1) < 0$ . Also, we notice that if  $(f,g) \in \mathcal{N}$ ,

$$I(f,g) = \left(\frac{1}{2} - \frac{1}{2q}\right) \|(f,g)\|_{\dot{\omega}} = \left(\frac{1}{2} - \frac{1}{2q}\right) \left(\|f\|_{2q}^{2q} + \|g\|_{2q}^{2q} + 2b\|fg\|_{q}^{q}\right).$$
(11)

We now take a minimizing sequence  $(u_n, v_n) \in \mathcal{N}$  for the problem

$$m = \inf\{I(u, v) : (u, v) \in \mathcal{N}\}.$$

From (11), it is clear that  $m \ge 0$  and that  $(u_n, v_n)$  is bounded in E.

We put  $u_n^*$  and  $v_n^*$  the decreasing radial rearrangements of  $|u_n|$  and  $|v_n|$  respectively. It is well-known that this rearrangement preserves the  $L^p$  norm  $(1 \le p \le +\infty)$ . Furthermore, the Pólya-Szegö inequality

$$\|\nabla f^*\|_2 \le \|\nabla |f|\|_2$$

in addition with the inequality  $\|\nabla f\|_2 \leq \|\nabla f\|_2$  (see [13]) shows that

$$\|(u_n^*, v_n^*)\|_{\dot{\omega}}^2 \le \|(u_n, v_n)\|_{\dot{\omega}}^2.$$

On the other hand, the Hardy-Littlewood inequality

$$\int |fg| \le \int f^*g^*$$

combined with the monotonicity of the map  $\lambda \to \lambda^q$  (see for instance [8] for details) yields  $||fg||_q \leq ||f^*g^*||_q$  and, finally,

$$\tau(u_n^*, v_n^*) \le \tau(u_n, v_n) = 0.$$

Next, let  $t_n \in [0, 1]$  such that  $(t_n u_n^*, t_n v_n^*) \in \mathcal{N}$ . We obtain

$$I(t_n u_n^*, t_n v_n^*) = t_n^2 \left(\frac{1}{2} - \frac{1}{2q}\right) \|(u_n^*, v_n^*)\|_{\dot{\omega}}^2 \le \left(\frac{1}{2} - \frac{1}{2q}\right) \|(u_n, v_n)\|_{\dot{\omega}}^2 = I(u_n, v_n)$$

and we obtained a minimizing sequence  $(t_n u_n^*, t_n v_n^*)$  in  $E_{rd}$ , denoted again, in what follows, by  $(u_n, v_n)$ . Since this sequence is bounded in  $H^1(\mathbb{R}^n)$ , up to a subsequence,  $(u_n, v_n) \rightarrow (u_*, v_*)$  in  $H^1(\mathbb{R}^n)$  weak. Also, since the injection  $E_{rd} \rightarrow L^{2q}(\mathbb{R}^n)$  is compact, up to a subsequence,  $(u_n, v_n) \rightarrow (u_*, v_*)$ in  $L^{2q}(\mathbb{R}^n)$  strong.

in  $L^{2q}(\mathbb{R}^n)$  strong. Hence, since  $||u_n||_{2q}^{2q} + ||v_n||_{2q}^{2q} + 2b||u_nv_n||_q^q \to ||u_*||_{2q}^{2q} + ||v_*||_{2q}^{2q} + 2b||u_*v_*||_q^q$ , we deduce that

 $\tau(u_*, v_*) \le \liminf \tau(u_n, v_n) = 0.$ 

Once again, let  $t \in [0, 1]$  such that  $(tu_*, tv_*) \in \mathcal{N}$ .

$$m \leq I(tu_*, tv_*) = t^2 \left(\frac{1}{2} - \frac{1}{2q}\right) ||(u_*, v_*)||_{\dot{\omega}}^2$$
$$\leq \left(\frac{1}{2} - \frac{1}{2q}\right) \liminf ||(u_n, v_n)||_{\dot{\omega}}^2 \leq \liminf I(u_n, v_n) = m.$$

This implies that  $(tu_*, tv_*)$  is a minimizer. In particular, all inequalities above are in fact equalities: t = 1,  $(u_*, v_*) \in \mathcal{N}$ ,  $||(u_*, v_*)||_{\dot{\omega}} = \lim ||(u_n, v_n)||_{\dot{\omega}}$ ,  $||u_n||_{H^1} \to ||u_*||_{H^1}$ ,  $||v_n||_{H^1} \to ||v_*||_{H^1}$  and  $(u_n, v_n) \to (u_*, v_*)$  in  $H^1(\mathbb{R}^n)$ strong.

Finally, it is clear that  $(u_*, v_*)$  is a ground state: if  $(w_1, w_2) \neq (0, 0)$  is a critical point of I such that  $I(w_1, w_2) < I(u_*, v_*)$ , taking once again  $w_1^*$ and  $w_2^*$  the decreasing radial rearrangements of  $|w_1|$  and  $|w_2|$ , there exists  $t \in ]0, 1]$  such that  $(tw_1^*, tw_2^*) \in \mathcal{N}$  and  $I(tw_1^*, tw_2^*) \leq I(w_1, w_2)$ , which leads to a contradiction. This completes the proof of Theorem 1.1.

## 3 Ground states with non-trivial components

Let  $(u_*, v_*) \in E_{rd}$  the ground state mentionned in Theorem 1.1. If  $v_* = 0$ ,  $u_* = u_0$  is the unique positive radially symmetric solution of the elliptic equation  $-\Delta u + u = u^{2q-1}$  (see [10]).

Also, if  $u_* = 0$ ,  $v_* = v_0$  is the unique positive radially symmetric solution of  $-\Delta v + \omega^2 v = v^{2q-1}$ , which relates to  $u_0$  by the relation  $v_0(x) = \omega^{\frac{1}{q-1}} u_0(\omega x)$ . Hence, to show the existence of a ground state with nontrivial components, we only have to exhibit an element  $(f,g) \in \mathcal{N} \cap E_{rd}$ ,  $f \neq 0$ ,  $g \neq 0$ , such that

$$I(f,g) \le \min\{I(u_0,0), I(0,v_0)\}.$$
(12)

Since  $I(u_0, 0) = \left(\frac{1}{2} - \frac{1}{2q}\right) \|u_0\|_{2q}^{2q}$ ,  $I(0, v_0) = \omega^{\frac{2q}{q-1}-n} \left(\frac{1}{2} - \frac{1}{2q}\right) \|u_0\|_{2q}^{2q}$  and  $\frac{2q}{q-1} - n > 0$ , for  $\omega \ge 1$  the inequality (12) reduces to

$$I(f,g) \le I(u_0,0).$$
 (13)

We first compute x > 0 such that  $(f, g) := (xu_0, x\theta v_0) \in \mathcal{N}$ , where  $\theta > 0$  will be chosen later (see [6] and [7] for a recent application of a related technique to the Schrödinger-KdV system):

$$\tau(f,g) = x^2 \|(u_0,\theta v_0)\|_{\dot{\omega}}^2 - x^{2q} \Big( \|u_0\|_{2q}^{2q} + \theta^{2q} \|v_0\|_{2q}^{2q} + 2b\theta^q \|u_0 v_0\|_q^q \Big) = 0.$$

Since

$$\|\theta v_0\|_{\dot{\omega}}^2 = \omega^{2+\frac{2}{q-1}-n} \|\theta u_0\|_2^2 + \omega^{2+\frac{2}{q-1}-n} \|\theta \nabla u_0\|_2^2 = \omega^{\frac{2q}{q-1}-n} \theta^2 \|u_0\|^2$$

and

$$\|v_0\|_{2q}^{2q} = \omega^{\frac{2q}{q-1}-n} \|u_0\|_{2q}^{2q},$$

we obtain

$$x^{2q-2} = \frac{(1+\theta^2\omega^{\frac{2q}{q-1}-n})\|u_0\|^2}{(1+\theta^{2q}\omega^{\frac{2q}{q-1}-n})\|u_0\|^{2q}_{2q} + 2b\theta^q\|u_0v_0\|^q_q} = \frac{1+\theta^2\omega^{\frac{2q}{q-1}-n}}{1+\theta^{2q}\omega^{\frac{2q}{q-1}-n} + 2b\theta^q\frac{\|u_0v_0\|^q_q}{\|u_0\|^{2q}_{2q}}}$$

Since  $u_0$  is radial and nonincreasing and  $\omega \geq 1$ ,

$$\|u_0 v_0\|_q^q = \omega^{\frac{q}{q-1}} \int u_0^q(x) u_0^q(\omega x) dx \le \omega^{\frac{q}{q-1}} \int u_0^q(x) u_0^q(x) dx = \omega^{\frac{q}{q-1}} \|u_0\|_{2q}^{2q}.$$

Also,

$$\|u_0 v_0\|_q^q \ge \omega^{\frac{q}{q-1}} \int u_0^{2q}(\omega x) dx = \omega^{\frac{q}{q-1}-n} \|u_0\|_{2q}^{2q}$$

Hence, we obtain

$$\frac{1+\theta^2\omega^{\frac{2q}{q-1}-n}}{1+\theta^{2q}\omega^{\frac{2q}{q-1}-n}+2b\theta^q\omega^{\frac{q}{q-1}}} \le x^{2q-2} \le \frac{1+\theta^2\omega^{\frac{2q}{q-1}-n}}{1+\theta^{2q}\omega^{\frac{2q}{q-1}-n}+2b\theta^q\omega^{\frac{q}{q-1}-n}}$$
(14)

and

$$I(f,g) = x^2 \left(\frac{1}{2} - \frac{1}{2q}\right) \|(u_0, \theta v_0)\|_{\dot{\omega}}^2 = x^2 \left(\frac{1}{2} - \frac{1}{2q}\right) (1 + \theta^2 \omega^{\frac{2q}{q-1} - n}) \|u_0\|^2.$$

The condition (13) then becomes  $x^2(1 + \theta^2 \omega^{\frac{2q}{q-1}-n}) \leq 1$ . In view of (14), a sufficient condition is

$$\frac{(1+\theta^2\omega^{\frac{2q}{q-1}-n})^q}{1+\theta^{2q}\omega^{\frac{2q}{q-1}-n}+2b\theta^q\omega^{\frac{q}{q-1}-n}} \le 1,$$

that is,

$$b \ge \frac{(1+\theta^2 \omega^{\frac{2q}{q-1}-n})^q - 1 - \theta^{2q} \omega^{\frac{2q}{q-1}-n}}{2\theta^q \omega^{\frac{q}{q-1}-n}}.$$

We now put  $\theta^2 = \epsilon^2 \omega^{n - \frac{2q}{q-1}}$ , for  $\epsilon > 0$ , obtaining the condition

$$b \ge \frac{(1+\epsilon^2)^q - 1}{2\epsilon^q} \omega^{q - \frac{n}{2}(q-2)} - \frac{1}{2} \epsilon^q \omega^{(\frac{n}{2}-1)q}.$$

For 1 < q < 2,  $\lim_{\epsilon \to 0} \frac{(1 + \epsilon^2)^q - 1}{2\epsilon^q} = 0$ .

Hence, the arbitrary value of  $\epsilon$  establishes the sufficient condition b > 0.

For n = 1, putting  $\epsilon = 1$ , we obtain the bound

$$b \ge \frac{2^q - 1}{2} \omega^{1 + \frac{q}{2}} - \frac{1}{2} \omega^{-\frac{q}{2}},\tag{15}$$

as stated in Theorem 1.3. We finish by making a few remarks:

**Remark 3.1** For  $\omega = 1$  and  $\theta = 1$ , we obtain, for all  $n \ge 1$ , the bound  $2^{q-1} - 1$  which is known to be optimal for  $q \ge 2$ , in the sense that for  $b < 2^{q-1} - 1$  all ground states of (3) have one null component (see [14], Theorem 2.5).

**Remark 3.2** The bound in (15) can be slightly improved for large values of  $\omega$  by replacing the quantity  $\frac{2^q - 1}{2}$  by the minimum of  $\frac{(1 + \epsilon^2)^q - 1}{2\epsilon^q}$  for  $\epsilon > 0$ .

**Remark 3.3** For  $n \ge 4$  we have 1 < q < 2, hence the problem of the existence of ground states with non-trivial components is completely solved for these spatial dimensions.

Acknowledgment: This article was partially supported by Fundação para a Ciência e Tecnologia, through contract PEst-OE/MAT/UI0297/2015.

### References

- N. Akhmediev and A. Ankiewicz, Partially coherent solitons on a finite background, Phys. Rev. Lett. 82, 26-61, 1999.
- [2] T. Bartsch and Z.Q. Wang, Note on ground states of nonlinear Schrödinger systems, J. Partial Diff. Eq. 19, 200-207, 2006.
- [3] H. Berestycki and P. L. Lions, Nonlinear Scalar Field Equations, I: Existence of a Ground State, A.R.M.A. 82, 313-345, 1983.
- [4] A. Ambrosetti and E. Colorado, Bound and ground states of coupled nonlinear Schrödinger equations, C.R. Acad. Sci. Paris, Ser. 1 342, 453-458, 2006.
- [5] A. Ambrosetti and E. Colorado, Standing waves of some coupled nonlinear Schrödinger equations, J. London Math. Soc. 75 (1), 67-82, 2007.
- [6] E. Colorado, On the existence of bound and ground states for some coupled nonlinear Schrödinger–Kortewegde Vries equations, arXiv:1411.7283, 2014.
- [7] E. Colorado, Existence of Bound and Ground States for a System of Coupled Nonlinear Schrödinger-KdV Equations, arXiv:1410.7638, 2014.
- [8] H. Hardy, J. E. Littlewood and G. Polya, Inequalities, Cambridge University Press, 1952.
- [9] N.Ikoma and N.Tanaka, A local mountain pass type result for a system of nonlinear Schrödinger equations, Calc.Var. Partial Diff. Equ. 40, 449-480, 2011.
- [10] M.K. Kwong, Uniqueness of positive solutions of  $-\Delta u + u = u^p$  in  $\mathbb{R}^N$ , Arch. Rat. Mech. Anal. 105, 243-266, 1989.
- [11] T.C. Lin and J. Wei, Ground state of N coupled nonlinear Schrödinger equations in  $\mathbb{R}^n$ ,  $n \leq 3$ , Comm. Math. Phys. 255, 629-653, 2005.

- [12] T.-C. Lin and J. Wei, Erratum: Ground state of N coupled nonlinear Schrödinger equations in  $\mathbb{R}^n$ ,  $n \leq 3$ , Comm. Math. Phys. 277, 573-576, 2008.
- [13] P. L. Lions, The concentration-compactness principle in the calculus of variations, Part 1, Ann. Inst. H. Poincaré 1, 109-145, 1984.
- [14] L.A. Maia, E.Montefusco and B. Pellacci, Positive solutions for a weakly coupled nonlinear Schrödinger System, J. Diff. Eq. 229, 743-767, 2006.
- [15] P.H. Rabinowitz, On a class of nonlinear Schrödinger equations, Z. Angew. Math. Phys. 43, 270-291, 1992.
- [16] W. A. Strauss, Existence of solitary waves in higher dimensions, Comm. Math. Phys. 55, 149-162, 1977.
- [17] B. Sirakov, Least energy solitary waves for a system of nonlinear Schrödinger equations in  $\mathbb{R}^n$ , Comm. Math. Phys. 271, 199-221, 2007.