

Fixed-parameter Tractable Distances to Sparse Graph Classes

Jannis Bulian and Anuj Dawar

University of Cambridge Computer Laboratory

April 1, 2024

We show that for various classes $\mathcal{C}$ of sparse graphs, and several measures of distance to such classes (such as edit distance and elimination distance), the problem of determining the distance of a given graph G to $\mathcal{C}$ is fixed-parameter tractable. The results are based on two general techniques. The first of these, building on recent work of Grohe et al. establishes that any class of graphs that is slicewise nowhere dense and slicewise first-order definable is FPT. The second shows that determining the elimination distance of a graph G to a minor-closed class $\mathcal{C}$ is FPT.

1 Introduction

The study of parameterized algorithmics for graph problems has thrown up a large variety of structural parameters of graphs. Among these are parameters that measure the *distance* of a graph G to a class $\mathcal{C}$ in some way. The simplest such measures are those that count the number of vertices or edges that one must delete (or add) to G to obtain a graph in $\mathcal{C}$. A common motivation for studying such parameters is that if a problem one wishes to solve is tractable on the class $\mathcal{C}$, then the distance to $\mathcal{C}$ provides an interesting parameterization of that problem (called *distance to triviality* by Guo et al. [14]). Other examples of this include the study of modulators to graphs of bounded tree-width in the context of kernelization (see [9, 11]) or the parameterizations of colouring problems (see [17]). On the other hand, determining the distance of an input graph G to a class $\mathcal{C}$ is, in general, a computationally challenging problem in its own right. Such problems have also been extensively studied with a view to establishing their complexity when parameterized by the distance. A canonical example is the problem of determining the size of a minimum vertex cover in a graph G , which is just the vertex-deletion distance of G to the class of edge-less graphs. More generally, Cai [3] studies the parameterized

complexity of distance measures defined in terms of addition and deletion of vertices and edges to hereditary classes $\mathcal{C}$. Counting deletions of vertices and edges gives a rather simple notion of distance, and many more involved notions have also been studied. Classic examples include the crossing number of a graph which provides one notion of distance to the class of planar graphs or the treewidth of a graph which can be seen as a measure of distance to the class of trees. Another recently introduced measure is *elimination distance*, defined in [2] where it was shown that graph isomorphism is FPT when parameterized by elimination distance to a class of graphs of bounded degree.

In this paper we consider the fixed-parameter tractability of a variety of different notions of distance to various different classes $\mathcal{C}$ of sparse graphs. We establish two quite general techniques for establishing that such a distance measure is FPT. The first builds on the recent result of Grohe et al. [13] which shows that the problem of evaluating first-order formulas on any *nowhere dense* class of graphs is FPT with the formula as parameter. We extract from their proof of this result a general statement about the fixed-parameter tractability of definable sparse classes. To be precise, we show that parameterized problems that are both *slicewise nowhere dense* and *slicewise first-order definable* (these terms are defined precisely below) are FPT. As an application of this, it follows that if $\mathcal{C}$ is a nowhere dense class of graphs that is definable by a first-order formula, then the parameterized problem of determining the distance of a graph G to $\mathcal{C}$ is FPT, for various notions of distance that can be themselves so defined. In particular, we get that various forms of edit distance to classes of bounded-degree graphs are FPT (a result established by Golovach [12] by more direct methods). Another interesting application is obtained by considering elimination distance of a graph G to the class $\mathcal{C}$ of empty graphs. This is nothing other than the *tree-depth* of G . While elimination distance to a class $\mathcal{C}$ is in general not first-order definable, it is in the particular case where $\mathcal{C}$ is the class of empty graphs. Thus, we obtain as an application of our method the result that tree-depth is FPT, a result previously known from other algorithmic meta theorems (see [18, Theorem 17.2]). The method of establishing that a parameterized problem is FPT by establishing that it is slicewise nowhere dense and slicewise first-order definable appears to be a powerful method of some generality which will find application beyond these examples.

Our second general method specifically concerns elimination distance to a minor-closed class $\mathcal{C}$. We show that this measure is fixed-parameter tractable for any such $\mathcal{C}$, answering an open question posed in [2]. Note that while a proper minor-closed class is always nowhere dense, it is not generally first-order definable (for instance, neither the class of acyclic graphs nor the class of planar graphs is), and elimination distance to such a class is also not known to be first-order definable. Thus, our results on the tractability of slicewise first-order definable classes do not apply here. Instead, we build on work of Adler et al. [1] to show that from a finite list of the forbidden minors characterising $\mathcal{C}$, we can compute the set of forbidden minors characterising the graphs at elimination distance k to $\mathcal{C}$. Adler et al. show how to do this for apex graphs, from which one immediately obtains the result for graphs that are k deletions away from $\mathcal{C}$. To extend this to elimination distance k , we show how we can construct the forbidden minors for the closure of a minor closed class under disjoint unions.

In Section 2 we present the definitions necessary for the rest of the paper. Section 3 establishes our result for slicewise first-order definable and slicewise nowhere dense problems and gives some applications of the general method. Section 4 establishes that the problem of determining elimination distance to any minor-closed class is FPT. Some open questions are discussed in Section 5.

2 Preliminaries

First-order logic. We assume some familiarity with first-order logic for Section 3. A *(relational) signature* σ is a finite set of relation symbols, each with an associated arity. A σ -*structure* A consists of a set $V(A)$ and for each k -ary relation symbol $R \in \sigma$ a relation $R(A) \subseteq V(A)^k$. Our structures will mostly be (coloured) graphs, so $\sigma = \{E\}$ or $\sigma = \{E, C_1, C_2, \dots, C_r\}$ where E is binary and the C_i are unary relation symbols. A graph G is then a σ -structure with vertex set $V(G)$, edge relation $E(G)$, and colours $C_i(G)$.

A first-order formula φ is recursively defined by the following rules:

$$\varphi := R(x_1, \dots, x_r) \mid x = y \mid \neg\varphi \mid \varphi \vee \psi \mid \exists x.\varphi.$$

We also use the following abbreviations:

$$\varphi \wedge \psi := \neg(\neg\varphi \vee \neg\psi), \quad \forall x.\varphi := \neg\exists x.\neg\varphi.$$

The *quantifier rank* of a formula φ is the nesting depth of quantifiers in φ . For a more detailed presentation we refer to Hodges [15].

Parameterized Complexity. Parameterized complexity theory is a two-dimensional approach to the study of the complexity of computational problems. We find it convenient to define problems as classes of structures rather than strings. A *problem* $Q \subseteq \text{str}(\sigma)$ is an (isomorphism-closed) class of σ -structures given some signature σ . A *parameterization* is a function $\kappa : \text{str}(\sigma) \rightarrow \mathbb{N}$. We say that Q is *fixed-parameter tractable* with respect to κ if we can decide whether an input $A \in \text{str}(\sigma)$ is in Q in time $O(f(\kappa(A)) \cdot |x|^c)$, where c is a constant and f is some computable function. For a thorough discussion of the subject we refer to the books by Downey and Fellows [6], Flum and Grohe [8] and Niedermeier [20].

A parameterized problem (Q, κ) is *slicewise first-order definable* if there is a computable function f such that:

- for all $i \in \mathbb{N}$, we have that $f(i) = \varphi_i \in \text{FO}[\sigma]$;
- a σ -structure A with $\kappa(A) \leq i$ is in Q if and only if $A \models \varphi_i$.

Slicewise definability of problems in a logic was introduced by Flum and Grohe [7].

Graph theory. A *graph* G is a set of vertices $V(G)$ and a set of edges $E(G) \subseteq V(G) \times V(G)$. We assume that graphs are loop-free and undirected, i.e. that E is irreflexive and symmetric. We mostly follow the notation in Diestel [5]. For a set $S \subseteq V(G)$ of vertices, we write $G \setminus S$ to denote the subgraph of G induced by $V(G) \setminus S$.

Let $r \in \mathbb{N}$. An r -independent set in a graph G is a set of vertices of G such that their pairwise distance is at least r .

A graph H is a *minor* of a graph G , written $H \preceq G$, if there is a map, called the *minor map*, that takes each vertex $v \in V(H)$ to a tree T_v that is a subgraph of G such that for any $u \neq v$ the trees are disjoint, i.e. $T_u \cap T_v = \emptyset$, and such that for every edge $uv \in E(H)$ there are vertices $u' \in T_u, v' \in T_v$ such that $u'v' \in E(G)$. A class of graphs $\mathcal{C}$ is *minor-closed* if $H \preceq G \in \mathcal{C}$ implies $H \in \mathcal{C}$.

The *set of minimal excluded minors* $M(\mathcal{C})$ is the set of graphs in the complement of $\mathcal{C}$ such that for each $G \in M(\mathcal{C})$ all proper minors of G are in $\mathcal{C}$. By the Robertson-Seymour Theorem [21] the set $M(\mathcal{C})$ is finite for every minor-closed class $\mathcal{C}$. It is a consequence of this theorem that membership in a minor-closed class can be tested in $O(n^3)$ time.

Let $r \in \mathbb{N}$. A minor H of G is a *depth- r minor* of G if there is a minor map that takes vertices in H to trees that have radius at most r . A class of graphs $\mathcal{C}$ is *nowhere dense* if for every $r \in \mathbb{N}$ there is a graph H_r such that for no $G \in \mathcal{C}$ we have $H_r \preceq_r G$. A nowhere dense class of graphs $\mathcal{C}$ is called *effectively nowhere dense* if there is a computable function f from integers to graphs such that for no $G \in \mathcal{C}$ and no r we have $f(r) \preceq_r G$. We are only interested in effectively nowhere dense classes so we simply use the term *nowhere dense* to mean effectively nowhere dense.

We say that a parameterized graph problem (Q, κ) is *slicewise nowhere dense* if there is a computable function h from pairs of integers to graphs such that for all $i \in \mathbb{N}$, we have for no $G \in \{H \in Q \mid \kappa(H) \leq i\}$ and r that $h(i, r) \preceq_r G$. We will call h the *parameter function* of Q .

For a class of graphs $\mathcal{C}$ we denote the closure of $\mathcal{C}$ under taking disjoint unions by $\overline{\mathcal{C}}$. We say that a graph G is an *apex graph* over a class $\mathcal{C}$ of graphs if there is a vertex $v \in V(G)$ such that the graph $G \setminus \{v\} \in \mathcal{C}$. The class of all apex graphs over $\mathcal{C}$ is denoted $\mathcal{C}^{\text{apex}}$.

A graph G has *deletion distance* k to a class $\mathcal{C}$ if there are k vertices $v_1, \dots, v_k \in V(G)$ such that $G \setminus \{v_1, \dots, v_k\} \in \mathcal{C}$.

The *elimination distance* of a graph G to a class $\mathcal{C}$ defined as follows:

$$ed_{\mathcal{C}}(G) := \begin{cases} 0, & \text{if } G \in \mathcal{C}; \\ 1 + \min\{ed_{\mathcal{C}}(G \setminus v) \mid v \in V(G)\}, & \text{if } G \notin \mathcal{C} \text{ and } G \text{ is connected}; \\ \max\{ed_{\mathcal{C}}(H) \mid H \text{ a connected component of } G\}, & \text{otherwise.} \end{cases}$$

3 A general method for editing distances

In this section we establish a general technique for showing that certain definable parameterized problems on graphs are FPT. As an application, we show that certain natural

distance measures to sparse graph classes are **FPT**. To be precise, we show that if a parameterized problem is both slicewise first-order definable and slicewise nowhere dense, then it is **FPT**. In particular, this implies that if we have a class $\mathcal{C}$ that is first-order definable and nowhere dense and the distance measure we are interested in is also first-order definable (that is to say, for each k there is a formula that defines the graphs at distance k from $\mathcal{C}$), then the problem of determining the distance is **FPT**. More generally, if we have a parameterized problem (Q, κ) that is slicewise nowhere dense and slicewise first-order definable, and a measure of distance to it is definable in the sense that for any values of k and d , there is a first-order formula defining the graphs at distance d to the class $\{G \mid G \in Q \text{ and } \kappa(G) \leq k\}$, then the problem of deciding whether a graph has distance at most d to this class is **FPT** parameterized by $d + k$. In particular, this yields the result of Golovach [12] as a consequence.

The method is an adaption of the main algorithm in Grohe *et al.* [13]. Since the proof is essentially a modification of their central construction, rather than give a full account, we state the main results they prove and explain briefly how the proofs can be adapted for our purposes. For a full proof, this section is best read in conjunction with the paper [13]. Section 3.1 gives an overview of the key elements of the construction from [13] and the elements from it which we need to extract for our result. Section 3.2 then gives our main result and Section 3.3 derives some consequences for distance measures.

3.1 Evaluating Formulas on Nowhere Dense Classes

The key result of [13] is:

Theorem 3.1 (Grohe *et al.* [13], Theorem 1.1). *For every nowhere dense class $\mathcal{C}$ and every $\epsilon > 0$, every property of graphs definable in first-order logic can be decided in time $O(n^{1+\epsilon})$ on $\mathcal{C}$.*

We first give a sketch of the algorithm from Theorem 3.1 with an emphasis on the changes needed for our purposes. We refer to [13] for several definitions and results.

The algorithm developed in the proof of Theorem 3.1 uses a locality-based approach, similar to that used by Frick and Grohe [10] to show that first-order evaluation is **FPT** on graphs of local bounded treewidth and developed in [4] for application to graph classes with locally excluded minors. The idea is that any first-order formula φ is, by Gaifman's theorem, equivalent to a Boolean combination of *local* formulae, that is formulae that assert the existence of neighbourhoods satisfying certain conditions. In classes of sparse graphs where the size (or other parameter) of neighbourhoods of a given radius can be bounded, this yields an efficient evaluation algorithm.

In nowhere dense classes of graphs, we cannot in general bound the size of neighbourhoods. For example, the class of apex graphs is nowhere dense, but a graph may contain a vertex whose neighbourhood is the whole graph. However, nowhere dense classes are *quasi-wide* [18], which means that we can remove a small (i.e. parameter-dependent) set of vertices (the bottleneck) to ensure that there are many vertices that are far away from each other. Grohe *et al.* [13] use this approach to iteratively transform the input graph into a coloured graph where key bottleneck vertices are removed and vertices are

coloured to keep relevant information. At the same time the formula φ to be evaluated is also transformed so that it can be evaluated on the modified graph. This procedure terminates on nowhere dense classes of graphs within a constant number of steps.

A key data structure used in the algorithm is a neighbourhood cover, i.e. a collection of connected subgraphs, called clusters, so that each neighbourhood of a vertex is contained in one of the clusters. The radius of a cover is the maximum radius of any of its clusters. The degree of a vertex in a cover is the number of clusters the vertex is contained in. An important result from [13] is that graphs from a nowhere dense class allow for small covers and that such a cover can be efficiently computed.

Theorem 3.2 (Grohe *et al.* [13], Theorem 6.2). *Let $\mathcal{C}$ be a nowhere dense class of graphs. There is a function f such that for all $r \in \mathbb{N}$ and $\epsilon > 0$ and all graphs $G \in \mathcal{C}$ with $n \geq f(r, \epsilon)$ vertices, there exists an r -neighbourhood cover of radius at most $2r$ and maximum degree at most n^ϵ and this cover can be computed in time $f(r, \epsilon) \cdot n^{1+\epsilon}$. Furthermore, if $\mathcal{C}$ is effectively nowhere dense, then f is computable.*

In this theorem, f is a function of r and ϵ and depends on the class $\mathcal{C}$ in the sense that it is determined, for an effectively nowhere dense $\mathcal{C}$ by its parameter function. To be precise, the algorithm needs to order the vertices of G to witness a weak colouring number of less than n^ϵ . The weak colouring number is an invariant of the graph that is guaranteed to be low for graphs from a nowhere dense class. The time bound $f(r, \epsilon) \cdot n^{1+\epsilon}$ is obtained using an algorithm for this from Nešetřil and Ossona de Mendez [19].

While the algorithm of [13] assumes that the input graph G comes from the class $\mathcal{C}$, we can say something more. For a fixed nowhere dense class $\mathcal{C}$, where we know the parameter function h , we can, given G , r and ϵ , compute a bound on the running time of the algorithm from Theorem 3.2. By running the algorithm to this bound, we have the following as a direct consequence of the proof of Theorem 3.2.

Lemma 3.3. *There is a computable function f and an algorithm A which given any graph G with n vertices and for any $r \in \mathbb{N}$ and $\epsilon > 0$ either computes an r -neighbourhood cover of radius at most $2r$ and maximum degree at most n^ϵ or determines that $G \notin \mathcal{C}$.*

At the core of the proof of Theorem 3.1 is the Rank-Preserving Locality Theorem. Given a neighbourhood cover $\mathcal{X}$ in a graph G , the algorithm iteratively removes bottleneck vertices and adds colours to the neighbourhoods of removed vertices to obtain a coloured graph denoted $G \star_{\mathcal{X}}^{q+1} q$. Here q is an integer parameter obtained from the first-order formula φ that we wish to evaluate in G . At the same time, φ is transformed into a formula $\hat{\varphi}$ that is (a) in the expanded signature of the $G \star_{\mathcal{X}}^{q+1} q$; and (b) in a logic FO^+ which enriches FO by allowing us to assert distances between vertices without the need for quantifiers. This ensures that the *local* sentence $\hat{\varphi}$ has the same quantifier rank as φ , giving us the Rank-Preserving Locality Theorem below. In the following statement, a $(q+1, r)$ -independence sentence is a formula asserting the existence of a distance- r -independent set of size $q+1$ of a particular colour.

Theorem 3.4 (Rank-Preserving Locality Theorem, Grohe *et al.* [13], Theorem 7.5). *For every $q \in \mathbb{N}$ there is an r such that for every FO -formula $\varphi(x)$ of quantifier rank q*

there is an FO^+ -formula $\hat{\varphi}(x)$, which is a Boolean combination of $(q+1, r)$ -independence sentences and atomic formulas, such that for any graph G every r -neighbourhood cover $\mathcal{X}$ of G , and every $v \in V(G)$,

$$G \models \varphi(v) \iff G \star_{\mathcal{X}}^{q+1} q \models \hat{\varphi}(v).$$

Furthermore, $\hat{\varphi}$ is computable from φ , and r is computable from q .

An important tool for constructing $G \star_{\mathcal{X}}^{q+1} q$ is a game characterisation of nowhere dense classes. The game has three parameters: ℓ, m, r . In the (ℓ, m, r) Splitter game two players Connector and Splitter play against each other. In each round Connector chooses a vertex u , and Splitter has to respond with a set A of vertices of size at most m in the r -neighbourhood of u . In the next round the graph is the neighbourhood of u with the vertices from A removed. If the graph is empty, Splitter wins. If Connector survives for more than ℓ rounds, she wins. Grohe *et al.* [13] prove [13, Theorem 4.2] that if $\mathcal{C}$ is a nowhere dense class, then there are ℓ, m such that Splitter has a winning strategy on the $(\ell, m, 2r)$ Splitter game on every graph in $\mathcal{C}$.

The Splitter's strategy on a graph G (which can be efficiently computed) is the essential tool in the construction of $G \star_{\mathcal{X}}^{q+1} q$. The inductive procedure used to compute $G \star_{\mathcal{X}}^{q+1} q$ from G is outlined in [13, Proof of Theorem 8.1]. We note that the termination of the algorithm depends on the length of the game – which is bounded by a constant since $\mathcal{C}$ is nowhere dense. The strategy to compute Splitter's moves is described in [13, Remark 4.3]. Since the run time of the algorithm to compute $G \star_{\mathcal{X}}^{q+1} q$ only depends on q and the length of the Splitter game and we can compute this in advance, we can once again extract the fact that if we start with an arbitrary graph G , we can efficiently *either* transform it into $G \star_{\mathcal{X}}^{q+1} q$ *or* determine that it is not in the class $\mathcal{C}$. This is summed up in the following lemma.

Lemma 3.5. *Let $\mathcal{C}$ be a nowhere dense class of graphs. There is an algorithm that runs in time $O(q)$ which given a graph G returns $G \star_{\mathcal{X}}^{q+1} q$ or determines that $G \notin \mathcal{C}$.*

Theorem 3.4 reduces the problem of evaluating a formula of first-order logic to deciding a series of distance- r -independent set problems. So, the final ingredient is to show that this is tractable. Formally, the problem is defined as follows:

DISTANCE INDEPENDENT SET

Input: A graph G and $k, r \in \mathbb{N}$.

Parameter: $k + r$

Problem: Does G contain an r -independent set of size k ?

The problem is shown to be **FPT** on nowhere dense classes of graphs [13, Theorem 5.1]. Since the runtime of the algorithm depends on the length of the Splitter game and Splitter's strategy, and this can be bounded in advance, [13, Theorem 5.1] can be restated as follows:

Lemma 3.6. *Let $\mathcal{C}$ be a nowhere dense class of graphs. Then there is an algorithm A and a computable function f such that for every $\epsilon > 0$ A runs in time $f(\epsilon, r, k)$ and either solves the DISTANCE INDEPENDENT SET problem or determines that $G \notin \mathcal{C}$.*

This is all we need to evaluate $\hat{\varphi}$ on $G \star_{\mathcal{X}}^{q+1} q$, which is equivalent to evaluating φ on G by Theorem 3.4.

3.2 Deciding Definable nowhere dense Problems

The main result of [13] establishes that checking whether $G \models \varphi$ is FPT when parameterized by φ provided that G comes from a known nowhere dense class $\mathcal{C}$. Thus, the formula is arbitrary, but the graphs come from a restricted class. In Section 3.1 above we give an account of this proof from which we can extract the observation that the algorithm can be modified to work for an arbitrary input graph G with the requirement that the algorithm *may* simply reject the input if G is not in $\mathcal{C}$. This suggests a tractable way of deciding $G \models \varphi$ provided that φ defines a nowhere dense class. Now the graphs is arbitrary, but the formula comes from a restricted class. We formalise the result in the following theorem:

Theorem 3.7. *Let (Q, κ) be a problem that is slice-wise first-order definable and slice-wise nowhere dense. Then (Q, κ) is fixed-parameter tractable.*

Proof. In the following, for ease of exposition, we assume that an instance of the problem consists of a graph G and $\kappa(G) = i$ for some positive integer i .

Step 1: Compute φ and the parameters function. Since (Q, κ) is slice-wise first-order definable, we can compute from i a first-order formula φ which defines the class of graphs $C_i = \{H \mid H \in Q \text{ and } \kappa(H) \leq i\}$. Moreover, since (Q, κ) is slice-wise nowhere dense, we can compute from i an algorithm that computes the parameter function h for C_i .

Step 2: Obtain $\hat{\varphi}$ from φ . By the Rank-Preserving Locality Theorem (Theorem 3.4), we can compute from φ the formula $\hat{\varphi}$ and a radius r .

Step 3: Find a small cover $\mathcal{X}$ for G . By Lemma 3.3, we can either find a cover $\mathcal{X}$ for G , or reject if the algorithm determine that $G \notin C_i$.

Step 4: Simulate Splitter game to compute $G \star_{\mathcal{X}}^{q+1} q$. By Lemma 3.5 we obtain $G \star_{\mathcal{X}}^{q+1} q$ or reject if the algorithm determines that $G \notin C_i$.

Step 5: Evaluate $\hat{\varphi}$ on $G \star_{\mathcal{X}}^{q+1} q$. Finally we evaluate $\hat{\varphi}$ on $G \star_{\mathcal{X}}^{q+1} q$. To do this, we need to solve the distance independent set problem. We can do this by Lemma 3.6.

Since evaluating $\hat{\varphi}$ on $G \star_{\mathcal{X}}^{q+1} q$ is equivalent to evaluating φ on G this allows us to decide whether $G \in Q$.

□

3.3 Applications

In this Section we discuss some applications of Theorem 3.7 that demonstrate its power. We begin by considering simple edit distances.

Edit Distances A graph G has *deletion distance* k to a class $\mathcal{C}$ if there exists a set S of k vertices in G so that $G \setminus S \in \mathcal{C}$. Suppose (Q, κ) is a parameterized graph problem. We define the problem of deletion distance to Q as follows:

DELETION DISTANCE TO Q

Input: A graph G and $k, d \in \mathbb{N}$.

Parameter: $k + d$

Problem: Does G contain a set S of k vertices so that $\kappa(G \setminus S) \leq d$ and $G \setminus S \in Q$?

In many of the examples below, we define formulas of first-order logic by *relativisation*. For convenience, we define the notion here.

Definition 3.8. Let φ and $\psi(x)$ be first-order formulas, where ψ has a distinguished free variable x . The relativisation of φ by $\psi(x)$, denoted $\varphi^{[x.\psi]}$ is the formula obtained from φ by replacing all subformulas of the form $\exists v \varphi'$ in φ by $\exists v(\psi[v/x] \wedge \varphi')$, and all subformulas of the form $\forall v \varphi'$ in φ by $\forall v(\psi[v/x] \rightarrow \varphi')$. Here $\psi[v/x]$ denotes the result of replacing the free occurrences of x in ψ with v in a suitable way avoiding capture.

The key idea here is that if $\varphi^{[x.\psi]}$ is true in a graph G if φ is true in the subgraph of G induced by the vertices that satisfy $\psi(x)$. Note that the variable x that is free in ψ is bound in $\varphi^{[x.\psi]}$. Other variables that appear free in ψ remain free in $\varphi^{[x.\psi]}$. We stress this as it is needed in Proposition 3.11 where nested relativisations are used.

Proposition 3.9. *If (Q, κ) is slice-wise nowhere dense and slice-wise first-order definable then DELETION DISTANCE TO Q is FPT.*

Proof. It suffices to show that DELETION DISTANCE TO Q is also slice-wise nowhere dense and slice-wise first-order definable. For the latter, note that if φ_i is the first-order formula that defines the class of graphs $\mathcal{C}_i = \{G \mid \kappa(G) \leq i \text{ and } G \in Q\}$, then the class of graphs at deletion distance k to $\mathcal{C}_i$ is given by:

$$\exists w_1, \dots, w_k \varphi_i^{[x.\theta_k]}$$

where $\theta_k(x)$ is the formula $\bigwedge_{1 \leq i \leq k} x \neq w_i$.

To see that DELETION DISTANCE TO Q is also slice-wise nowhere dense, let h be the parameter function for Q . If the graph $h(i, r)$ has m vertices, then K_m is not a depth- r -minor of any graph in $\mathcal{C}_i$. Then a graph which has deletion distance k to $\mathcal{C}_i$ cannot have K_{m+k} as a depth- r -minor. Indeed, suppose $K_{m+k} \preceq_r G$ and $G \setminus S \in \mathcal{C}_i$ where S is a set of k vertices. Vertices from S can appear in the images of at most k vertices from K_{m+k} under the minor map. Thus, this minor map also witnesses that $K_m \preceq_r G \setminus S$, a contradiction. $\square$

Instead of deleting vertices, we can also consider editing the graph by adding or deleting edges. It is easily seen that we can modify a first-order formula φ to define the class of graphs G that can be made to satisfy φ by k edge additions or deletions. An added or deleted edge is identified by a pair of vertices $uv \in E(G)$. If we want to add the edge uv we can replace all occurrences of $E(w_1, w_2)$ in φ by:

$$(w_1 = u \wedge w_2 = v) \vee (w_1 = v \wedge w_2 = u) \vee E(w_1, w_2).$$

Similarly we delete an edge uv by replacing all occurrences of $E(w_1, w_2)$ in φ by:

$$(w_1 \neq u \vee w_2 \neq v) \wedge (w_1 \neq v \vee w_2 \neq u) \wedge E(w_1, w_2).$$

Thus, an analogue of Proposition 3.9 is obtained for any combination of vertex and edge deletions and additions. Golovach [12] proved that editing a graph to degree d using at most k edge additions/deletions is FPT parameterized by $k + d$. Since the class of graphs of degree d is first-order definable and nowhere dense for any d , the result also follows directly from Theorem 3.7.

Tree-depth. Tree-depth is a graph parameter that lies between the widely studied parameters vertex cover number and tree width. It has interesting connections to nowhere dense graph classes. It is usually defined as follows:

Definition 3.10. The *tree-depth* of a graph G , written $td(G)$, is defined as follows:

$$td(G) := \begin{cases} 0, & \text{if } V(G) = \emptyset; \\ 1 + \min\{td(G \setminus v) \mid v \in V(G)\}, & \text{if } G \text{ is connected;} \\ \max\{td(H) \mid H \text{ a component of } G\}, & \text{otherwise.} \end{cases}$$

Note that a graph has tree-depth k if and only if it has elimination distance k to the class of empty graphs. So one can think of elimination distance as a natural generalisation of tree-depth.

It is known that the problem of determining the tree-depth of graph is FPT, with tree-depth as the parameter (see [18, Theorem 7.2]). We now give an alternative proof of this, using Theorem 3.7. It is clear that for any k , the class of graphs of tree-depth at most k is nowhere dense. We show below that it is also first-order definable.

Proposition 3.11. *For each $k \in \mathbb{N}$ there is a first-order formula φ_k such that a graph G has tree-depth k if and only if $G \models \varphi_k$.*

Proof. We use the fact that in a graph of tree-depth less than k , there are no paths of length greater than 2^k . This allows us, in the inductive definition of tree-depth above, to replace the condition of connectedness (which is not first-order definable) with a first-order definable condition on vertices at distance at most 2^k .

Let $\text{dist}_d(u, v)$ denote the first-order formula with free variables u and v that is satisfied by a pair of vertices in a graph G if, and only if, they have distance at most d in G . Note that the formula $\text{dist}_d^{[x.x \neq w]}(u, v)$ is then a formula with three free variables u, v, w which defines those u, v which have a path of length d in the graph obtained by deleting the vertex w .

We can now define the formula φ_k by induction. Only the empty graph has tree-depth 0, so $\varphi_0 := \neg \exists v(v = v)$.

Suppose that φ_k defines the graphs of tree-depth at most k , let

$$\theta_k := (\forall u, v \text{ dist}_{2^{k+1}}(u, v)) \wedge \exists w(\varphi_k^{[x.x \neq w]}).$$

The formula θ_k defines the connected graphs of tree depth at most $k+1$. Indeed, the first conjunct ensures that the graph is connected as no pair of vertices has distance greater than 2^{k+1} and the second conjunct ensures we can find a vertex w whose removal yields a graph of tree-depth at most k .

We can now define the formula φ_{k+1} as follows.

$$\varphi_{k+1} := (\forall u, v \text{ dist}_{2^{k+1}+1}(u, v) \rightarrow \text{dist}_{2^{k+1}}(u, v)) \wedge \forall w \theta_k^{[x.\text{dist}_{2^{k+1}}(w, x)]}.$$

The formula asserts that there are no pairs of vertices whose distance is strictly greater than 2^{k+1} and that for every vertex w , the formula θ_k holds in its connected component, namely those vertices which are at distance at most 2^{k+1} from w . $\square$

While the proof of Proposition 3.9 shows that deletion distance to any slice-wise first-order definable class is also slice-wise first-order definable, Proposition 3.11 shows that elimination distance to the particular class of empty graphs is slice-wise first-order definable. It does not establish this more generally for elimination distance to any slice-wise nowhere dense class.

4 Elimination distance to classes characterised by excluded minors

In this section we show that determining the elimination distance of a graph to a minor-closed class $\mathcal{C}$ is FPT when parameterized by the elimination distance. More generally, we formulate the following parameterized problem where the forbidden minors of $\mathcal{C}$ are also part of the parameter.

ELIMINATION DISTANCE TO EXCLUDED MINORS

Input: A graph G , a natural number $k \in \mathbb{N}$ and a set of graphs M

Parameter: $k + \sum_{G \in M} |G|$

Problem: Does G have elimination distance k to the class $\mathcal{C}$ characterised by $M(\mathcal{C}) = M$?

It is not difficult to show that the class of graphs which have elimination distance k to a minor-closed class $\mathcal{C}$ is also a minor-closed class. Indeed, this can be seen directly from an alternative characterisation of elimination distance that we establish below. The characterisation is in terms of the iterated closure of $\mathcal{C}$ under the operation of disjoint unions and taking the class of apex graphs. We introduce a piece of notation for this in the next definition. Recall that we write $\mathcal{C}^{\text{apex}}$ for the class of all the apex graphs over $\mathcal{C}$, and that we write $\overline{\mathcal{C}}$ for the closure of $\mathcal{C}$ under disjoint unions.

Definition 4.1. For a class of graphs $\mathcal{C}$, let $\mathcal{C}_0 := \mathcal{C}$, and $\mathcal{C}_{i+1} := \overline{\mathcal{C}_i^{\text{apex}}}$.

We show next that the class $\mathcal{C}_k$ is exactly the class of graphs at elimination distance k from $\mathcal{C}$.

Proposition 4.2. *Let $\mathcal{C}$ be a class of graphs and $k \geq 0$. Then $\mathcal{C}_k$ is the class of all graphs with elimination distance at most k to $\mathcal{C}$.*

Proof. We prove this by induction. Only the graphs in $\mathcal{C}$ have elimination distance 0 to $\mathcal{C}$, so the statement holds for $k = 0$.

Suppose the statement holds for k . If $G \in \mathcal{C}_{k+1}$, then G is a disjoint union of graphs $G_1, \dots, G_s$ from $\mathcal{C}_k^{\text{apex}}$, so we can remove at most one vertex from each of the G_i and obtain a graph in $\mathcal{C}_k$. Thus the elimination distance of G to $\mathcal{C}_k$ is 1, and by induction the elimination distance to $\mathcal{C}$ is $k + 1$. Conversely, if G has elimination distance $k + 1$ to $\mathcal{C}$, then we can remove a vertex from each component of G to obtain a graph G' with elimination distance k to $\mathcal{C}$. Using the induction hypothesis each component of G' is in $\mathcal{C}_k$, and thus $G \in \mathcal{C}_{k+1}$. $\square$

It is easy to see that if $\mathcal{C}$ is a minor-closed class of graphs then so is $\mathcal{C}_k$ for any k . Indeed, it is well-known that $\mathcal{C}^{\text{apex}}$ is minor-closed for any minor-closed $\mathcal{C}$, so we just need to note that $\overline{\mathcal{C}}$ is also minor-closed. But it is clear that if H is a minor of a graph G that is the disjoint union of graphs $G_1, \dots, G_s$, then H itself is the disjoint union of (possibly empty) minors of $G_1, \dots, G_s$. Thus, the class of graphs of elimination distance at most k to a minor-closed class $\mathcal{C}$ is itself minor-closed. We next show that we can construct the set of its minimal excluded minors from the corresponding set for $\mathcal{C}$.

To obtain $M(\mathcal{C}_k)$, we need to iteratively compute $M(\mathcal{C}^{\text{apex}})$ and $M(\overline{\mathcal{C}})$ from $M(\mathcal{C})$. Adler et al. [1] show that from the set of minimal excluded minors $M(\mathcal{C})$ of a class $\mathcal{C}$, we can compute $M(\mathcal{C}^{\text{apex}})$:

Theorem 4.3 ([1], Theorem 5.1). *There is a computable function that takes the set of graphs $M(\mathcal{C})$ characterising a minor-closed class $\mathcal{C}$ to the set $M(\mathcal{C}^{\text{apex}})$.*

We next aim to show that from $M(\mathcal{C})$ we can also compute $M(\overline{\mathcal{C}})$. Together with Theorem 4.3 this implies that from $M(\mathcal{C})$ we can compute $M(\mathcal{C}_k)$, the set of minimal excluded minors for the class of graphs with elimination distance k to $\mathcal{C}$.

We begin by characterising minor-closed classes that are closed under disjoint unions in terms of the connectedness of their excluded minors.

Lemma 4.4. *Let $\mathcal{C}$ be a class of graphs closed under taking minors. Then $\mathcal{C}$ is closed under taking disjoint unions iff each graph in $M(\mathcal{C})$ is connected.*

Proof. Let $\mathcal{C}$ be a minor-closed class of graphs, and let $M(\mathcal{C}) = \{H_1, \dots, H_s\}$ be its set of minimal excluded minors.

Suppose each of the graphs in $M(\mathcal{C})$ is connected. Let $H \in M(\mathcal{C})$ and let $G = G_1 \oplus \dots \oplus G_r$ be the disjoint union of graphs $G_1, \dots, G_r \in \mathcal{C}$. Because H is connected, we have that $H \preceq G$ if and only if $H \preceq G_i$ for one $1 \leq i \leq r$. So, since all the $G_i \in \mathcal{C}$, we have $H \not\preceq G$ and thus $G \in \mathcal{C}$. This shows that $\mathcal{C}$ is closed under taking disjoint unions.

Conversely assume one of the graphs $H \in M(\mathcal{C})$ is not connected and let $A_1, \dots, A_t$ be its components. Then $A_1, \dots, A_t \in \mathcal{C}$, since each A_i is a proper minor of H , and H is minor-minimal in the complement of $\mathcal{C}$. However, $A_1 \oplus \dots \oplus A_t = H \notin \mathcal{C}$. $\square$

Definition 4.5. For a graph G with connected components $G_1, \dots, G_r$, let $\mathcal{H}$ denote the set of *connected* graphs H with $V(H) = V(G)$ and such that the subgraph of H induced by $V(G_i)$ is exactly G_i . We define the *connection closure* of G to be the set of

all minimal (under the subgraph relation) graphs in $\mathcal{H}$. The connection closure of a set of graphs is the union of the connection closures of the graphs in the set.

Note that if G has e edges and m components, then any graph in the connection closure of G has exactly $e + m - 1$ edges. This is because it has G as a subgraph and in addition $m - 1$ edges corresponding to a tree on m vertices connecting the m components.

Lemma 4.6. *Let $\mathcal{C}$ be a minor-closed class of graphs. Then $M(\overline{\mathcal{C}})$ is the set of minor-minimal graphs in the connection closure of $M(\mathcal{C})$.*

Proof. Let $\mathcal{C}$ be a minor-closed class of graphs, with $M(\mathcal{C})$ its set of minimal excluded minors, and let $\hat{M}$ be the connection closure of $M(\mathcal{C})$.

Let G be a graph such that $\hat{H} \not\preceq G$ for all $\hat{H} \in \hat{M}$. Suppose for a contradiction that G is not a disjoint union of graphs from $\mathcal{C}$. Then there is a component G' of G that is not in $\mathcal{C}$ and therefore there is a graph $H \in M(\mathcal{C})$ such that $H \preceq G'$. We show that one of the graphs in the connection closure of H is a minor of G' .

Let $\{w_1, \dots, w_s\}$ be the vertex set of H and consider the image $T_1, \dots, T_s$ of the minor map from H to G' . Let T be a minimal subtree of G' that contains all of the T_i . Such a tree must exist since G' is connected. Let $\hat{H}$ be the graph with the same vertex set as H , and an edge between two vertices w_i, w_j whenever either $w_i w_j \in E(H)$ or when there is a path between T_{w_i} and T_{w_j} in T that is disjoint from any T_{w_k} with $w_i \neq w_k \neq w_j$. We claim that $\hat{H}$ is in the connection closure of H . By construction, $\hat{H}$ is connected and contains all components of H as disjoint subgraphs, so we only need to argue minimality. $\hat{H}$ has no vertices besides those in H so no graph obtained by deleting a vertex would contain all components of H as subgraphs. To see that no edge of $\hat{H}$ is superfluous, we note it has exactly $e + m - 1$ edges and thus no proper subgraph could be connected and have all components of H as disjoint subgraphs. By the construction $\hat{H} \preceq G' \preceq G$, so by the transitivity of the minor relation we have that $\hat{H} \preceq G$.

Conversely let G be an arbitrary graph and assume that $\hat{H} \in \hat{M}$ and $\hat{H} \preceq G$. Because $\hat{H}$ is connected, there is a connected component G' of G such that $\hat{H} \preceq G'$. Now there must be a graph $H \in M(\mathcal{C})$ such that $\hat{H}$ is in the connection closure of H , and since H is a subgraph of $\hat{H}$, $H \preceq \hat{H}$. Then, by the transitivity of the minor relation, $H \preceq G'$ and thus $G' \notin \mathcal{C}$. Therefore G is not a disjoint union of graphs from $\mathcal{C}$. $\square$

Now we can put everything together and prove our main theorem:

Theorem 4.7. *There is a computable function which takes a set M of excluded minors characterising a minor-closed class $\mathcal{C}$ and $k \geq 0$ to the set $M(\mathcal{C}_k)$.*

Proof. The proof is by induction. For $k = 0$, the set of minimal excluded minors of $\mathcal{C}_0$ is $M(\mathcal{C}_0) = M(\mathcal{C})$, which is given. For $k > 0$, we have that $\mathcal{C}_k = \overline{\mathcal{C}_{k-1}}^{\text{apex}}$. By the induction hypothesis we can compute $M(\mathcal{C}_{k-1})$, by Theorem 4.3 we can compute $M(\mathcal{C}_{k-1}^{\text{apex}})$ and using Lemma 4.6 we can compute the connection closure of $M(\mathcal{C}_{k-1}^{\text{apex}})$ to obtain $M(\overline{\mathcal{C}_{k-1}^{\text{apex}}}) = M(\mathcal{C}_k)$. $\square$

So by the Robertson-Seymour Theorem we have the following:

Corollary 4.8. *Let $\mathcal{C}$ be a minor-closed graph class. Then the problem ELIMINATION DISTANCE TO EXCLUDED MINORS is FPT.*

5 Conclusion

We are motivated by the study of the fixed-parameter tractability of edit distances in graphs. Specifically, we are interested in edit distances such as the number of vertex or edge deletions, as well as more involved measures like elimination distance. Aiming at studying general techniques for establishing tractability, we establish an algorithmic meta-theorem showing that any slicewise first-order definable and slicewise nowhere dense problem is **FPT**. This yields, for instance, the tractability of counting the number of vertex and edge deletions to a class of bounded degree. As a second result, we establish that determining elimination distance to any minor-closed class is **FPT**, answering an open question of [2].

A natural open question raised by these two results is whether elimination distance to the class of graphs of degree d is **FPT**. When d is 0, this is just the tree-depth of a graph, and this case is covered by our first result. For positive values of d , it is not clear whether elimination distance is first-order definable. Indeed, a more general version of the question is whether for any nowhere dense and first-order definable $\mathcal{C}$, elimination distance to $\mathcal{C}$ is **FPT**.

Another interesting case that seems closely related to our methods, but is not an immediate consequence is that of classes that are given by first-order interpretations from nowhere dense classes of graphs. For instance, consider the problem of determining the deletion distance of a graph to a disjoint union of complete graphs. This problem, known as the cluster vertex deletion problem is known to be **FPT** (see [16]). The class of graphs that are disjoint unions of cliques is first-order definable but certainly not nowhere dense and so the method of Section 3 does not directly apply. However, this class is easily shown to be interpretable in the nowhere dense class of forests of height 1. Can this fact be used to adapt the methods of Section 3 to this class?

References

- [1] I. Adler, M. Grohe, and S. Kreutzer, *Computing excluded minors*, SODA '08: Proceedings of the nineteenth annual ACM-SIAM symposium on Discrete algorithms, SIAM, January 2008.
- [2] J. Bulian and A. Dawar, *Graph isomorphism parameterized by elimination distance to bounded degree*, Parameterized and Exact Computation - 9th International Symposium, IPEC 2014, Wroclaw, Poland, September 10-12, 2014. Revised Selected Papers, 2014, pp. 135–146.
- [3] L. Cai, *Fixed-parameter tractability of graph modification problems for hereditary properties*, Inf. Process. Lett. **58** (1996), 171–176.
- [4] A. Dawar, M. Grohe, and S. Kreutzer, *Locally excluding a minor*, Proc. 22nd IEEE Symp. on Logic in Computer Science, 2007, pp. 270–279.
- [5] R. Diestel, *Graph Theory*, Springer, January 2000.
- [6] R. G. Downey and M. R. Fellows, *Parameterized Complexity*, Springer, October 2012.
- [7] J. Flum and M. Grohe, *Fixed-Parameter Tractability, Definability, and Model-Checking.*, SIAM J. Comput. **31** (2001), no. 1, 113–145.
- [8] ———, *Parameterized Complexity Theory*, Springer, May 2006.
- [9] F. V. Fomin, D. Lokshtanov, N. Misra, and S. Saurabh, *Planar F-deletion: Approximation, kernelization and optimal FPT algorithms*, 53rd Annual IEEE Symposium on Foundations of Computer Science, FOCS, 2012, pp. 470–479.
- [10] M. Frick and M. Grohe, *Deciding First-Order Properties of Locally Tree-Decomposable Graphs*, Automata, Languages and Programming, Springer, January 1999, pp. 331–340.
- [11] J. Gajarský, P. Hliněný, J. Obdržálek, S. Ordyniak, F. Reidl, P. Rossmanith, F. Sanchez Villaamil, and S. Sikdar, *Kernelization using structural parameters on sparse graph classes*, Algorithms - ESA 2013 - 21st Annual European Symposium, 2013, pp. 529–540.
- [12] P. A. Golovach, *Editing to a Graph of Given Degrees*, Parameterized and Exact Computation, Springer, September 2014, pp. 196–207.
- [13] M. Grohe, S. Kreutzer, and S. Siebertz, *Deciding first-order properties of nowhere dense graphs*, STOC '14: Proceedings of the 46th Annual ACM Symposium on Theory of Computing, ACM, May 2014.

- [14] J. Guo, F. Hüffner, and R. Niedermeier, *A Structural View on Parameterizing Problems: Distance from Triviality*, Parameterized and Exact Computation, Springer, 2004, pp. 162–173.
- [15] W. Hodges, *A Shorter Model Theory*, Cambridge University Press, 1997.
- [16] F. Hüffner, C. Komusiewicz, H. Moser, and R. Niedermeier, *Fixed-parameter algorithms for cluster vertex deletion*, Theory Comput. Syst. **47** (2010), 196–217.
- [17] D. Marx, *Parameterized coloring problems on chordal graphs*, Theor. Comput. Sci. **351** (2006), 407–424.
- [18] J. Nešetřil and P. Ossona de Mendez, *Sparsity - graphs, structures, and algorithms*, Springer, 2012.
- [19] J. Nešetřil and P. Ossona de Mendez, *Grad and classes with bounded expansion II. Algorithmic aspects*, European Journal of Combinatorics **29** (2008), no. 3, 777–791.
- [20] R. Niedermeier, *Invitation to Fixed-Parameter Algorithms*, Oxford University Press, February 2006.
- [21] N. Robertson and P. D. Seymour, *Graph minors. XX. wagner’s conjecture*, J. Comb. Theory, Ser. B **92** (2004), 325–357.