

# Scalar Glueball in a Top-Down Holographic Approach to QCD\*

DENIS PARGANLIJA

Institut für Theoretische Physik, Technische Universität Wien, Wiedner  
Hauptstr. 8-10, 1040 Vienna, Austria

Identification of glueballs – bound states of gauge bosons in Quantum Chromodynamics (QCD) – is a very important open question in dynamics of the strong interaction. The search for the glueball ground state, carrying scalar quantum numbers, poses a particular challenge due to the existence of *(i)* several candidates for its realisation in the physical spectrum and *(ii)* inevitable mixing of the pure glueball state with those comprised of quarks. In this article, I discuss implications of an approach in holographic QCD where, among others, the mass and the two-pion decay of the pure scalar glueball can be studied.

PACS numbers: 11.25.Tq, 12.38.Lg, 12.39.Mk, 13.25.Jx, 14.40.Be

## 1. Introduction

The non-Abelian nature of Quantum Chromodynamics (QCD) – the theory of strong interaction – gives rise to the expectation that its gauge bosons, the gluons, form composite objects denoted as glueballs [1]. These states would have access to various quantum numbers  $J^{PC}$ , where  $J$  denotes the total spin,  $P$  the parity and  $C$  the charge conjugation; the corresponding spectrum in the QCD Yang-Mills sector has been determined in numerical simulations [2, 3] but the identification of glueballs in experimental data has proven to be a challenge, particularly in the  $J^{PC} = 0^{++}$  (scalar) channel.

There are at least two reasons to study glueball states. Firstly, their masses are generated solely by the strong interaction; no influence of the Higgs mechanism – providing, e.g., quarks with a current mass – is present in the

---

\* Presented at EEF70 – Workshop on Unquenched Hadron Spectroscopy: Non-Perturbative Models and Methods of QCD vs. Experiment at the occasion of Eef van Beveren’s 70th birthday, 1-5 September 2014, University of Coimbra, Portugal

glueball mass generation. Secondly, due to their structure glueballs must possess integer spin assigning them to mesons; our understanding of the physical meson spectrum would not be complete without glueballs.

Glueball ground state is scalar [4]; listings of the Particle Data Group (PDG) [5] contain five isospin-zero scalar states in the energy region below 1 GeV:  $f_0(500)$  or  $\sigma$ ,  $f_0(980)$ ,  $f_0(1370)$ ,  $f_0(1500)$  and  $f_0(1710)$ . While all of them possess the quantum numbers of the ground state, there are strong reasons to focus on resonances above 1 GeV as candidates for the scalar glueball: (i) simulations in lattice QCD determine the ground-state mass at between approximately 1.65 GeV and 1.8 GeV [2, 3]; (ii) various effective approaches to low-energy QCD arrive at an analogous result for the mass while describing the overall dynamics correctly [6]. Complication is that  $f_0$  resonances will most certainly have contributions not only from the (pure) glueball states but also from those containing quarks ( $\bar{q}q$  [7],  $\bar{q}\bar{q}qq$  [8] and others). This leads to various issues in both theory and experiment [9, 10] and represents the main reason why the scalar glueball has still not been clearly identified.

In this article, the question of the scalar glueball is tackled by a holographic approach to non-perturbative QCD. Such approaches are based on the idea of a duality between strongly coupled quantum gauge field theories and weakly coupled supergravity/superstring theories in one dimension higher, pursuing Maldacena's groundbreaking conjecture of a complete equivalence between the supergravity limit of type-IIB string theory on an  $AdS_5 \times S^5$  space and the large- $N$  limit of an  $\mathcal{N} = 4$  supersymmetric and conformal  $U(N)$  gauge theory on its boundary (AdS/CFT correspondence [11]). In Ref. [12], Witten proposed an analogous duality in type-IIA string theory, with supersymmetry and conformality not present in line with their absence in QCD. The supersymmetry is broken by compactification on a circle ( $S^1$ ); for a vanishing circle radius, the 5-dimensional pure Yang-Mills theory is reduced to a 4-dimensional one. However, the supergravity approximation requires finite circle radius (whose inverse is defined as the Kaluza-Klein mass  $M_{KK}$ ), and also a large coupling. Thus constructed holographic approaches are referred to as *top-down* models [13, 14]; there are also more phenomenological *bottom-up* constructions – see Ref. [15] and references therein. This article describes the implications of the top-down Witten-Sakai-Sugimoto model [14] for the glueball spectroscopy; more details can be found in Ref. [16].

## 2. The Model and Its Implications

While Witten's model contained only gauge fields, the novel feature of Witten-Sakai-Sugimoto model is the inclusion of chiral quarks introduced by  $N_f$  (number of flavours) probe D8 and anti-D8 branes [inducing  $U(N_f) \times U(N_f)$  chiral symmetry] that extend along all dimensions of the 10-dimensional space with the exception of a (Kaluza-Klein) circle. The branes are usually antipodally separated with regard to this  $S^1$ . The space geometry, however, is such that the branes and antibranes merge at a certain point – interpreted as realisation of the chiral-symmetry breaking.

Up to a Chern-Simons term, the corresponding action for D8-branes reads

$$S_{\text{D8}} = -T_{\text{D8}} \text{Tr} \int d^9x e^{-\Phi} \sqrt{-\det(\tilde{g}_{MN} + 2\pi\alpha' F_{MN})} \quad (1)$$

where  $T_{\text{D8}} = (2\pi)^{-8} l_s^{-9}$  (and  $l_s^2 = \alpha'$ , with  $\alpha'$  the string coupling),  $g_{MN}$  is the metric of the D-brane world volume,  $\Phi$  is the dilaton field and  $F_{MN}$  a field strength tensor whose components are, upon dimensional reduction, identified as meson fields of interest. No backreaction of the Witten-model background to D8-branes is considered; consequently  $N_f$  is fixed and, as can be argued [14], significantly smaller than the number of colours (large- $N_c$  limit).

The above action is expanded up to the second order in fields as

$$S_{\text{D8}}^{(2)} = -\kappa \text{Tr} \int d^4x \int_{-\infty}^{\infty} dZ \left[ \frac{1}{2} K^{-\frac{1}{3}} \eta^{\mu\rho} \eta^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} + M_{\text{KK}}^2 \eta^{\mu\nu} F_{\mu Z} F_{\nu Z} \right] \quad (2)$$

where  $\kappa = \lambda N_c / (216\pi^3)$  [16],  $\lambda = g_{\text{YM}}^2 N_c$  is the 't Hooft coupling (and  $g_{\text{YM}}$  the 4-dimensional coupling),  $Z$  is the holographic radial coordinate (and  $K = 1 + Z^2$ ) and  $\eta^{\mu\nu}$  is the flat metric  $\text{diag}(-, +, +, +)$ . The Kaluza-Klein mass  $M_{\text{KK}}$  sets the model scale; beside the scale, the model contains only one unknown quantity: the coupling  $\lambda$ . They are usually determined such that the mass of the rho meson and the pion decay constant correspond to their physical values. This yields  $M_{\text{KK}} = 949$  MeV and  $\lambda = 16.63$ ; alternative methods for their determination do not alter model conclusions [16].

Once  $M_{\text{KK}}$  and  $\lambda$  are known, the model describes various experimental quantities surprisingly well [14]. Glueballs are of interest in this article; building on the work of Ref. [17], the following masses in the scalar channel are obtained:

$$\begin{aligned} M_{G_E} &= 855 \text{ MeV}; \quad M_{G_E^*} = 2168 \text{ MeV} \\ M_{D_E} &= 1487 \text{ MeV}; \quad M_{D_E^*} = 2358 \text{ MeV}. \end{aligned} \quad (3)$$

Two types of glueball states are present: one, denoted with  $G_D$ , is predominantly dilaton while the other, denoted with  $G_E$ , involves a graviton polarisation in a fourth spatial dimension, unlike the dilaton, and has therefore been termed *exotic* [18]. Excited states are denoted with an asterisk. We observe that the exotic ground state is approximately 50% lighter than the expectation from lattice QCD; the dilaton mode mass is only approximately 15% smaller than the lattice result. Numerical simulations also indicate the mass of the first excited scalar state to be  $\simeq 2600$  MeV with errors amounting to  $\simeq 300$  MeV [2] (although mass corrections in the unquenched case may be substantial [3]); the excited dilaton state is within errors consistent with the lattice result while the excited exotic mode is approximately 15% too light. Hence already from the mass results the indication is that the exotic mode could be discarded while the dilaton mode appears compatible with simulations of the Yang-Mills sector of QCD.

This is corroborated by the decay ratios of the modes. The corresponding Lagrangians are obtained by inserting 10-dimensional metric fluctuations (whose explicit forms together with further details are presented in Ref. [16]) into the action for D8 branes and integrating over the bulk coordinates (see also Ref. [19]); ratios of decay widths  $\Gamma$  and the respective masses  $M$  read

$$\begin{aligned}\Gamma_{G_E \rightarrow \pi\pi}/M_{G_E} &= 0.092; \quad \Gamma_{G_E^* \rightarrow \pi\pi}/M_{G_E^*} = 0.149 \\ \Gamma_{G_D \rightarrow \pi\pi}/M_{G_D} &= 0.009; \quad \Gamma_{G_D^* \rightarrow \pi\pi}/M_{G_D^*} = 0.011.\end{aligned}\tag{4}$$

The two main candidates for the scalar glueball are the  $f_0(1500)$  and  $f_0(1710)$  resonances (see Ref. [10] and references therein). Experimental data imply  $\Gamma_{f_0(1500) \rightarrow \pi\pi}/M_{f_0(1500)} = 0.025 \pm 0.003$  [5] and  $\Gamma_{f_0(1710) \rightarrow \pi\pi}/M_{f_0(1710)} \simeq 0.009 - 0.017$  [20]. The exotic mode is thus too broad, again indicating that its interpretation as a physical state is uncertain; contrarily, the ratio  $\Gamma/M$  for the dilaton mode is within the experimental interval for  $f_0(1710)$ .

Similar is true for the excited states:  $\Gamma_{G_E^* \rightarrow \pi\pi}$  is larger than  $\Gamma_{G_D^* \rightarrow \pi\pi}$  and nicely comparable to widths of  $f_0$  states near and above 2 GeV; however, the full decay width of  $G_E^*$  having contributions from  $2K$ ,  $2\eta$ ,  $4\pi$  and other channels is unphysically large ( $\sim 1$  GeV, see Ref. [16]). Contrarily,  $\Gamma_{G_D^* \rightarrow \pi\pi}$  is small but the full decay width of  $G_D^*$  is of the order of 460 MeV [16] and thus significantly closer to the data [5] whose current uncertainties unfortunately do not allow for a clear identification of an excited scalar glueball.

### 3. Summary and Outlook

In this article, a top-down holographic approach to low-energy QCD – Witten-Sakai-Sugimoto model – has been presented and its implications for

phenomenology of scalar glueballs have been discussed. The model offers two sets of glueball states, a dilaton and an *exotic* mode that, unlike the dilaton, involves a graviton polarisation in a fourth spatial dimension. The exotic ground state has a mass approximately 50% smaller than the value expected in lattice QCD; its  $2\pi$  decay width is substantially larger than that of the two prime candidates for the scalar glueball, the resonances  $f_0(1500)$  and  $f_0(1710)$ . Contrarily, the mass of the dilaton mode ( $= 1487$  MeV) is quite close to masses of both mentioned resonances; its  $2\pi$  decay width is within the experimental range for  $f_0(1710)$ , which therefore appears to be the preferred candidate for the glueball ground state. In the excited channel, the exotic state is unphysically broad while the dilaton width  $\sim 460$  MeV is close to the (still ambiguous) data on  $f_0$  states near/above 2 GeV. Nonetheless, further pursuit of glueball dynamics in holography is called for, particularly in light of expectations from the planned PANDA experiments at FAIR [21].

#### 4. Acknowledgments

I am grateful to F. Br  nner and A. Rebhan for collaboration and to D. Bugg and S. Janowski for extensive discussions. I am also grateful to the Workshop Organisers for their generous support of my participation. This work is supported by the Austrian Science Fund FWF, project no. P26366.

#### REFERENCES

- [1] A. Salomone, J. Schechter and T. Tudron, Phys. Rev. D **23**, 1143 (1981); C. Rosenzweig, A. Salomone and J. Schechter, Nucl. Phys. B **206**, 12 (1982) [Erratum-ibid. B **207**, 546 (1982)]; A. A. Migdal and M. A. Shifman, Phys. Lett. B **114**, 445 (1982); H. Gomm and J. Schechter, Phys. Lett. B **158**, 449 (1985); R. Gomm, P. Jain, R. Johnson and J. Schechter, Phys. Rev. D **33**, 801 (1986).
- [2] C. J. Morningstar and M. J. Peardon, Phys. Rev. D **60**, 034509 (1999) [hep-lat/9901004]; Y. Chen *et al.*, Phys. Rev. D **73**, 014516 (2006) [hep-lat/0510074]; M. Loan, X. Q. Luo and Z. H. Luo, Int. J. Mod. Phys. A **21**, 2905 (2006) [hep-lat/0503038].
- [3] E. Gregory, A. Irving, B. Lucini, C. McNeile, A. Rago, C. Richards and E. Rinaldi, JHEP **1210**, 170 (2012) [arXiv:1208.1858 [hep-lat]].
- [4] G. B. West, Phys. Rev. Lett. **77**, 2622 (1996) [hep-ph/9603316].
- [5] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. C, **38**, 090001 (2014).
- [6] F. E. Close and A. Kirk, Eur. Phys. J. C **21**, 531 (2001) [hep-ph/0103173]; F. Giacosa, T. Gutsche and A. Faessler, Phys. Rev. C **71**, 025202 (2005) [hep-ph/0408085]; F. Giacosa, T. Gutsche, V. E. Lyubovitskij and A. Faessler,

- Phys. Lett. B **622**, 277 (2005) [hep-ph/0504033]; F. Giacosa, T. Gutsche, V. E. Lyubovitskij and A. Faessler, Phys. Rev. D **72**, 094006 (2005) [hep-ph/0509247]; H. Y. Cheng, C. K. Chua and K. F. Liu, Phys. Rev. D **74**, 094005 (2006) [hep-ph/0607206]; V. Mathieu, N. Kochelev and V. Vento, Int. J. Mod. Phys. E **18**, 1 (2009) [arXiv:0810.4453 [hep-ph]]; S. Janowski, D. Parganlija, F. Giacosa and D. H. Rischke, Phys. Rev. D **84**, 054007 (2011) [arXiv:1103.3238 [hep-ph]]; S. Janowski, F. Giacosa and D. H. Rischke, Phys. Rev. D **90**, no. 11, 114005 (2014) [arXiv:1408.4921 [hep-ph]].
- [7] D. Parganlija, F. Giacosa and D. H. Rischke, Phys. Rev. D **82**, 054024 (2010) [arXiv:1003.4934 [hep-ph]]; D. Parganlija, P. Kovacs, G. Wolf, F. Giacosa and D. H. Rischke, Phys. Rev. D **87**, 014011 (2013) [arXiv:1208.0585 [hep-ph]].
- [8] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977); A. H. Fariborz, R. Jora and J. Schechter, Phys. Rev. D **72**, 034001 (2005) [arXiv:hep-ph/0506170]; F. Giacosa, Phys. Rev. D **75**, 054007 (2007) [hep-ph/0611388]; A. Heinz, S. Struber, F. Giacosa and D. H. Rischke, Phys. Rev. D **79**, 037502 (2009) [arXiv:0805.1134 [hep-ph]]; A. Heinz, S. Struber, F. Giacosa and D. H. Rischke, Acta Phys. Polon. Supp. **3**, 925 (2010) [arXiv:1006.5393 [hep-ph]].
- [9] D. V. Bugg, Phys. Rept. **397**, 257 (2004) [hep-ex/0412045]; E. Klempt and A. Zaitsev, Phys. Rept. **454**, 1 (2007) [arXiv:0708.4016 [hep-ph]].
- [10] D. Parganlija, J. Phys. Conf. Ser. **426**, 012019 (2013) [arXiv:1211.4804 [hep-ph]]; D. Parganlija, J. Phys. Conf. Ser. **503**, 012010 (2014) [arXiv:1312.2830 [hep-ph]].
- [11] J. M. Maldacena, Int. J. Theor. Phys. **38**, 1113 (1999) [Adv. Theor. Math. Phys. **2**, 231 (1998)] [hep-th/9711200].
- [12] E. Witten, Adv. Theor. Math. Phys. **2**, 505 (1998) [hep-th/9803131].
- [13] I. R. Klebanov and M. J. Strassler, JHEP **0008**, 052 (2000) [hep-th/0007191]; J. Babington, J. Erdmenger, N. J. Evans, Z. Guralnik and I. Kirsch, Phys. Rev. D **69**, 066007 (2004) [hep-th/0306018]; M. Kruczenski, D. Mateos, R. C. Myers and D. J. Winters, JHEP **0405**, 041 (2004) [hep-th/0311270].
- [14] T. Sakai and S. Sugimoto, Prog. Theor. Phys. **113**, 843 (2005) [hep-th/0412141]; T. Sakai and S. Sugimoto, Prog. Theor. Phys. **114**, 1083 (2005) [hep-th/0507073]; T. Imoto, T. Sakai and S. Sugimoto, Prog. Theor. Phys. **124**, 263 (2010) [arXiv:1005.0655 [hep-th]]; F. Brünner, D. Parganlija and A. Rebhan, Acta Phys. Polon. Supp. **7**, no. 3, 533 (2014) [arXiv:1407.6914 [hep-ph]]; A. Rebhan, arXiv:1410.8858 [hep-th].
- [15] S. S. Afonin, Int. J. Mod. Phys. A **25**, 5683 (2010) [arXiv:1001.3105 [hep-ph]].
- [16] F. Brünner, D. Parganlija and A. Rebhan, arXiv:1501.07906 [hep-ph].
- [17] R. C. Brower, S. D. Mathur and C. I. Tan, Nucl. Phys. B **587**, 249 (2000) [hep-th/0003115].
- [18] N. R. Constable and R. C. Myers, JHEP **9910**, 037 (1999) [hep-th/9908175].
- [19] K. Hashimoto, C. -ITan and S. Terashima, Phys. Rev. D **77**, 086001 (2008) [arXiv:0709.2208 [hep-th]].
- [20] D. Parganlija, arXiv:1208.0204 [hep-ph].
- [21] M. F. M. Lutz *et al.* [PANDA Collaboration], arXiv:0903.3905 [hep-ex].