Special Relativity in a Discrete Quantum Universe

Alessandro Bisio,^{1, 2, *} Giacomo Mauro D'Ariano,^{1, 2, †} and Paolo Perinotti^{1, 2, ‡}

¹Universita degli Studi di Pavia, Dipartimento di Fisica, QUIT Group ²Istituto Nazionale di Fisica Teorica e Nucleare, Sezione di Pavia

The hypothesis of a discrete fabric of the universe—the "Planck scale"—is always on stage, since it solves mathematical and conceptual problems in the infinitely small. However, it clashes with special relativity, which is designed for the continuum. Here we show how the clash can be overcome within a discrete quantum theory where the evolution of fields is described by a quantum cellular automaton. The reconciliation is achieved by defining the change of observer as a change of representation of the dynamics, without any reference to space-time. We use the relativity principle, i.e. the invariance of dynamics under change of inertial observer, to identify a change of inertial frame with a symmetry of the dynamics. We consider the full group of such symmetries, and recover the usual Lorentz group in the relativistic regime of low energies, while at the Planck scale the covariance is nonlinearly distorted.

I. INTRODUCTION

Is the world continuous or discrete? Richard Feynman [1, 2] motivated a discrete universe as the only way it can be simulated by its own constituents, which means by a quantum computer. Einstein himself considered a discrete space-time as a possibility, however, he complained about the lack of an appropriate mathematical framework [3]. Usually we dismiss discreteness on the basis of a mathematical convenience of continuos theories. But the continuum leads to still unsolved mathematical problems in the infinitely small, problems that do not arise in the discrete. The discrete, on the other hand, seems to raise a problem: the disagreement with Einstein's special relativity.

The debate about the clash between a discrete space and Lorentz symmetry has been recently renewed because in some approaches to quantum gravity (such as Regge Calculus [4], spin-foam [5], causal sets, [6]) the fundamental description of space-time is a discrete structure, to which the continuum is only an approximation. The scale of this discreteness is the Planck length, which is amazingly small-the Planck length compared to a meter is like the electron radius compared to the size of our galaxy.

Why the Lorentz transformations would not work with a discrete space-time? The objection is that a discrete space-time would not be invariant under the Lorentz group, even if we take it as discrete. Such a point, raised more than sixty years ago, was disproved for d = 3 space dimensions in Ref. [7]. However, it was shown that the minimum admissible boost would be huge: 0.866 times the speed of light! It seems therefore that there is no way for the reconciliation of Einstein's special relativity with a discrete fabric of space-time. The issue is, however, a false problem, originating from the unnecessary requirement of enforcing a covariance designed for the continuum. The right point of view is to take the Lorentz covariance only as an approximate symmetry, and recovering it in the regime where the discreteness looks continuous. This is similar to what happens for a crystalline medium, which looks isotropic at large scales, whereas instead it is highly anisotropic at microscopic scales. The smaller is the crystal structure, the more accurate is the continuum symmetry: think that the Planck length is 10^{-25} Å!

The right point of view is thus to consider the continuum as an approximation of the discrete when observed at very large scales. It is thus conceptually legitimate that the Lorentz transformations are actually distorted at the tiny Planck scale. An example of such a distorted Lorentz symmetry is that of *doubly special relativity* [8– 10], where the distorted Lorentz transformations, in addition to the speed of light, preserve also an energy scale.

Here we show that if we take only the very essence of the relativity principle—the invariance of the physical law under change of *inertial representation*—we get nonlinear Lorentz transformations, which happen to be of the same kind as those of doubly special relativity. In the continuum description the reference frame is a Cartesian coordinate system—what is called "position representation" in quantum theory. Other representations of the dynamics are given in terms of constants of motion—such as the momentum or energy representations—and these provide a viable notion of inertial frame in a world made of countably many quantum systems.

II. RESULTS

We take the pragmatic point of view that quantum theory is more efficient a description of our world than classical theory, pretty much like Kepler laws for the planet orbits are more efficient than Ptolemaic epicycles. We therefore consider the most general quantum discrete theory, which is a *quantum cellular automaton* [1, 11, 12]. This consists of infinitely many quantum systems (qubits, Fermionic or Bosonic fields), whose evolution occurs in

^{*}alessandro.bisio@unipv.it

[†]dariano@unipv.it

[‡]paolo.perinotti@unipv.it




FIG. 1: The green surface represents the dispersion relation in Eq. (8) where we fixed $k_z = 0$ ($\omega_{\mathbf{k}} = \arccos \lambda^{\pm}(\mathbf{k})$ have the same plot). The red surface is the usual relativistic dispersion relation $\omega_{\mathbf{k}}^2 = k_x^2 + k_y^2$. Notice that the two surfaces get closer approaching the origin for $\mathbf{k} \to 0$.

discrete steps, and which interact *locally*, namely every system interacts with a bounded number of other systems. A consequence of locality is that signals propagate at finite speed over the interacting network. For our purposes it is sufficient to consider a single particle, and technically this simplifies the automaton to be linear in the quantum field. Moreover we require that the dynamics is reversible, hence it is described by a unitary matrix: this is what is called *Quantum Walk* [13, 14].

In the discrete context universality of the physical law corresponds to the homogeneity of the evolution. This implies that the network of interactions represents a group G of "translations", moving from a system to another interacting with it: this is a so-called *Cayley graph* of G (see Methods). Moreover, if we want to reproduce the physical situation of flat space where parallel transports along two different paths end up in the same result, then the "translation" group must be Abelian. This allows us to make use the Fourier transform. Upon denoting by $|\mathbf{k}\rangle$ the eigenvector of the translations corresponding to wave-vector \mathbf{k} , we write the quantum walk unitary operator A as follows

$$A = \int_{\mathsf{B}} \mathrm{d}\,\mathbf{k}\,A_{\mathbf{k}} \otimes |\mathbf{k}\rangle \langle \mathbf{k}|,\tag{1}$$

where B denotes the Brillouin zone of the interaction lattice, and A_k is a unitary *s* dimensional matrix, with *s* finite. As proved in Ref. [15], the easiest nontrivial quantum walk has s = 2. The eigenvalues of *A* are obtained by solving the eigenvalue equation

$$A_{\mathbf{k}}\psi(\omega, \mathbf{k}) = e^{i\omega}\psi(\omega, \mathbf{k}) \tag{2}$$

which can be rewritten in relativistic notation as follows

$$n_{\mu}(k)\sigma^{\mu}\psi(k) = 0, \qquad (3)$$

where we introduced the four-vectors $k = (\omega, \mathbf{k}), n(k) = (\sin \omega, \mathbf{n}(\mathbf{k})), \text{ and } \sigma = (I, \sigma), \text{ with } \sigma = (\sigma_x, \sigma_y, \sigma_z) \text{ denoting the Pauli matrices, and the vector } \mathbf{n}(\mathbf{k}) \text{ are defined}$

by

$$\mathbf{n}(\mathbf{k}) \cdot \boldsymbol{\sigma} := \frac{i}{2} (A_{\mathbf{k}} - A_{\mathbf{k}}^{\dagger}). \tag{4}$$

The eigenvalues can be collected into two functions $\omega^{\pm}(\mathbf{k})$ called *dispersion relations*. In this scenario the constants of motions are \mathbf{k} and ω^{\pm} , hence a change of representation corresponds to a map $k \mapsto k'(k)$. Now the principle of relativity corresponds to the requirement that the eigenvalue equation (3) is preserved under a change of representation as follows

$$n_{\mu}(k)\sigma^{\mu} = \tilde{\Gamma}_{k}^{-1} n_{\mu}(k')\sigma^{\mu} \Gamma_{k}, \qquad (5)$$

where Γ_k , Γ_k are invertible matrices.

Eq. (5) translates the relativity principle for the QW evolution: the dynamics is left invariant by a change of observer.

The simplest example of change of observer is the one given by the trivial relabeling k' = k and by the matrices $\Gamma_k = \tilde{\Gamma}_k = e^{i\lambda(\mathbf{k})}$, where $\lambda(\mathbf{k})$ is an arbitrary real function of \mathbf{k} . When $\lambda(\mathbf{k})$ is a linear function we recover the usual group of translations. The set of changes of representation $k \mapsto k'(k)$ for which Eq. (5) holds are a group, which is the largest group of symmetries of the dynamics.

If to the general assumptions defining the quantum walk we just add that of isotropy, it turns out that there are only two admissible quantum walks [15], which in the small wave-vector regime give exactly the two Weyl equations for the left and right massless Fermion. Indeed, with the above assumptions the only possible lattice is the body centered cubic one, and modulo local unitary equivalence the two admissible quantum walks are

$$A_{\mathbf{k}}^{\pm} := \lambda^{\pm}(\mathbf{k})I - i\mathbf{n}^{\pm}(\mathbf{k}) \cdot \boldsymbol{\sigma}^{\pm}, \qquad (6)$$

where

$$\mathbf{n}^{\pm}(\mathbf{k}) := \begin{pmatrix} s_x c_y c_z \pm c_x s_y s_z \\ c_x s_y c_z \mp s_x c_y s_z \\ c_x c_y s_z \pm s_x s_y c_z \end{pmatrix},$$

$$\lambda^{\pm}(\mathbf{k}) := (c_x c_y c_z \mp s_x s_y s_z),$$

$$c_\alpha := \cos(k_\alpha/\sqrt{3}), \ s_\alpha := \sin(k_\alpha/\sqrt{3}), \ \alpha = x, y, z,$$

(7)

where $\sigma^+ = \sigma$ and $\sigma^- = \sigma^T$, with T denoting the transposed matrix. The dispersion relations are given by

$$n_{\mu}^{\pm}(k)n^{\mu\pm}(k) = 0, \tag{8}$$

and are plotted in Fig. 1.

In the small wave-vector regime $\mathbf{k} \sim \mathbf{k}_0 = (0, 0, 0)$ one has $n(k) \sim k$, recovering the usual relativistic dispersion relation. The Weyl equations can be also recovered in the neighborhood of the wavevectors $\mathbf{k}_1 = \frac{\pi}{2}(1, 1, 1)$, $\mathbf{k}_2 = -\frac{\pi}{2}(1, 1, 1)$, $\mathbf{k}_3 = -\frac{\pi}{2}(1, 0, 0)$. The mapping between the vectors \mathbf{k}_i exchange chirality of the particle




FIG. 2: The distortion effects of the Lorentz group for the discrete Planck-scale theory represented by the quantum walk in Eq. (6). Left figure: the orbit of the wavevectors $\mathbf{k} = (k_x, 0, 0)$, with $k_x \in \{.05, .2, .5, 1, 1.7\}$ under the rotation around the z axis. Right figure: the orbit of wavevectors with $|\mathbf{k}| = 0.01$ for various directions in the (k_x, k_y) plane under the boosts with $\boldsymbol{\beta}$ parallel to \mathbf{k} and $|\boldsymbol{\beta}| \in [0, \tanh 4]$.

and double the particles to four species in total. Therefore we have four different particles–two left-handed and two right-handed–namely the discreteness also doubles the particles, which is the well known phenomenon of Fermion doubling [16]. In the following the term "small wavevector" will denote the neighborhoods of the vectors $\mathbf{k}_i \ i = 0, \dots 3$.

We now show that the group of symmetries of the dynamics of the quantum walks in Eq. (6) contains a nonlinear representation of the Poincaré group, which exactly recovers the usual linear one in the small wavevector regime. For any arbitrary non vanishing function f(k) we can introduce the four-vector

$$p^{(f)} = \mathcal{D}^{(f)}(k) := f(k)n(k)$$
 (9)

and rewrite the eigenvalue equation (3) as follows

$$p_{\mu}^{(f)}\sigma^{\mu}\psi(k) = 0.$$
 (10)

Upon denoting the usual Lorentz transformation by L_{β} for a suitable f (an example is provided in the supplemental material) the Brillouin zone splits into four regions B_i $i = 1, \ldots, 4$ centered around \mathbf{k}_i $i = 0, \ldots, 3$, such that the composition

$$\mathcal{L}_{\boldsymbol{\beta}}^{(f)} := \mathcal{D}^{(f)-1} L_{\boldsymbol{\beta}} \mathcal{D}^{(f)} \tag{11}$$

is well defined on each region separately (see Methods). The four invariant regions corresponding to the four different massless Fermionic particles show that the Wigner notion of "particle" as invariant of the Poincaré group survives in a discrete world, consistent with a physical interpretation of the Fermion-doubled particles. For fixed function f the maps $\mathcal{L}^{(f)}_{\beta}$ provide a non-linear representation of the Lorentz group [9, 10, 17]. In Figs. 2 and 3 we show the numerical evaluation of some wavevector




FIG. 3: The green surface represents the orbit of the wavevector $\mathbf{k} = (0.3, 0, 0)$ under the full rotation group SO(3).

orbits under subgroups of the nonlinear Lorentz. The distortion effects due to underlying discreteness are evident at large wavevectors and boosts. The relabeling $k \to k'(k) = \mathcal{L}_{\beta}^{(f)}(k)$ satisfies (5) with $\Gamma_k = \Lambda_{\beta}$ and $\tilde{\Gamma}_k = \tilde{\Lambda}_{\beta}$ for the right-handed particles, and $\Gamma_k = \tilde{\Lambda}_{\beta}$ and $\tilde{\Gamma}_k = \Lambda_{\beta}$ for the left-handed particles, with Λ_{β} and $\tilde{\Lambda}_{\beta}$ being the $(0, \frac{1}{2})$ and $(\frac{1}{2}, 0)$ representation of the Lorentz group, independently on k in each pertaining region.

For varying f, we obtain a much larger group, including infinitely many copies of the nonlinear Lorentz one. In the small wave-vector regime the whole group collapses to the usual linear Lorentz group for each particle.

Up to now we have analyzed what happens with massless particles. A simple way to obtain the Dirac equation is to pair an automaton in Eq. (6) with its adjoint into a direct sum, as in Ref. [15], thus leading to a new automaton giving the Dirac equation in the small wave-vector regime. A relevant feature of the discreteness is that because of unitarity the mass parameter is upper bounded [18]. Now if one derives the full symmetry group of the dynamics as we have done for the two automata in Eq. (6) one discovers that the group is a nonlinear representation of the de Sitter group SO(1,4) with infinite cosmological constant, with the rest mass of the particle playing the role of the additional coordinate (see Methods). It is noticeable that even for pure boosts the rest-mass is involved in the transformation. For rest-mass much smaller than the upper bound and for pure boosts one recovers the previous nonlinear Lorentz group for zero-mass.

III. CONCLUSION

We have seen what happens of the Lorentz group in a quantum world that is discrete. The main point is to abandon the idea of enforcing the exact Lorentz symmetry on the discrete, but instead to consider the symmetry as an approximate one that holds only in the small wavevector and small mass regime. But the natural question is now: how small? According to the common opinion the scale of discreteness a is identified with the Planck scale. In terms of the maximum wavevector k_M in the Brillouin zone, one has $k_M = \frac{\sqrt{3}\pi}{a}$. In the small wavevector regime we recover the simple relations [18] $c = \frac{a}{\sqrt{3}\pi}$ and $\hbar = \mu ac$, with c, \hbar , and τ denoting the speed of light, the Planck constant, and the time-step, respectively. Then the maximum mass μ of the quantum walk is the Planckmass. A way of deriving μ and a heuristically is to keep literally the argument of taking the mass of the particle bounded in order to keep the Compton wave-length λ_C larger than the Schwartschild radius. Noticeably for $m = \mu$ the dispersion relation is constant, namely with no propagation of information, a situation reminiscent of a micro black hole [19]. Remarkably general relativity enters the present quantum digital framework also through the unforeseeable appearance of the De Sitter symmetry group, which connects different Dirac particle mass values. Are these only coincidences? The dream is that it is a new route to quantum gravity.

The crucial question is now what can be actually seen experimentally. Recently, experimental tests of Planckscale phenomenology have been proposed [20–22]. In particular, the modification to the usual dispersion relations can in principle be detected in observation of gamma-ray bursts from deep-space events, where billions of light-year of distance can sufficiently amplify the weak vacuum dispersive behavior due to discreteness [23]. In our context this can be proved with the free electromagnetic field derived as the two-particle sector of the quantum walk in Eq. (6) [24]. This possibility reconnects with the recent analysis of data [25] from Fermi-LAT concluding that the observations set an upper-bound to the scale of discreteness which is smaller than the Planck scale a by a factor 2.8. The analysis of Ref. [25] can be refined with a complete theoretical derivation based on Ref. [24] and on the results presented in this letter. This would also take into account the possibility of a compensating effect due to the phenomenon of relative locality [26]. In short relative locality is the phenomenon due to the nonlinearity of the Lorentz transformations, which generalizes the relativity of simultaneity to relativity of the full space-time coincidence of events. The separation of events under boost is amplified by the difference of their frequency domain. Indeed the Fermi-LAT observation is based on a predicted time-delay between two events with a huge difference in frequency, which could then be compensated by the relative-locality effect. The fully fledged discrete theory given here, derived from very general principles, allows for a thorough quantitative evaluation that takes into account both the dispersive vacuum and nonlinear Lorentz transformations.

Acknowledgments

This work has been supported by the Templeton Foundation under the project ID# 43796 A Quantum-Digital Universe.

- R. Feynman, International journal of theoretical physics 21, 467 (1982).
- [2] A. J. Hey, ed., Feynman and Computation (Perseus Books, Reading, 1998).
- [3] J. Stachel, in From Quarks to Quasars: Philosophical Problems of Modern Physics, edited by R. G. Colodny and A. Coffa (University of Pittsburgh Press, Pittsburgh, 1986) pp. 349–81.
- [4] G. Immirzi, Nuclear Physics B-Proceedings Supplements 57, 65 (1997).
- [5] A. Perez, Classical and Quantum Gravity **20**, R43 (2003).

- [6] L. Bombelli, J. Lee, D. Meyer, and R. Sorkin, Physical review letters 59, 521 (1987).
- [7] A. Schild, Phys. Rev. 74 (1947).
- [8] G. Amelino-Camelia, Physics Letters B **510**, 255 (2001).
- [9] G. Amelino-Camelia, International Journal of Modern Physics D 11, 35 (2002).
- [10] J. Magueijo and L. Smolin, Physical Review Letters 88, 190403 (2002).
- [11] B. Schumacher and R. Werner, Arxiv preprint quantph/0405174 (2004).
- [12] P. Arrighi, V. Nesme, and R. Werner, Journal of Com-

puter and System Sciences 77, 372 (2011).

- [13] D. Meyer, Journal of Statistical Physics 85, 551 (1996).
- [14] A. Ambainis, E. Bach, A. Nayak, A. Vishwanath, and J. Watrous, in *Proceedings of the thirty-third annual ACM* symposium on Theory of computing (ACM, 2001) pp. 37– 49.
- [15] G. M. D'Ariano and P. Perinotti, Phys. Rev. A 90, 062106 (2014).
- [16] L. Susskind, Phys. Rev. D 16, 3031 (1977).
- [17] G. Amelino-Camelia and T. Piran, Physical Review D 64, 036005 (2001).
- [18] G. M. D'Ariano, Physics Letters. Section A: General, Atomic and Solid State Physics 376, 697 (2012).
- [19] S. W. Hawking, Phys. Rev. D 53, 3099 (1996).
- [20] M. Moyer, Scientific American (2012).
- [21] C. Hogan, Physical Review D 85 (2012).
- [22] I. Pikovski, M. Vanner, M. Aspelmeyer, M. Kim, and C. Brukner, Nature Physics **331**, 393 (2012).
- [23] G. Amelino-Camelia, J. Ellis, N. E. Mavromatos, D. V. Nanopoulos, and S. Sarkar, Nature **393**, 763 (1998).
- [24] A. Bisio, G. M. D'Ariano, and P. Perinotti, Annals of Physics , http://dx.doi.org/10.1016/j.aop.2016.02.009 (2016).
- [25] V. Vasileiou, J. Granot, T. Piran, and G. Amelino-Camelia, Nat Phys 11, 344 (2015).
- [26] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman, and L. Smolin, Phys. Rev. D 84, 084010 (2011).
- [27] G. M. D'Ariano, M. Erba, P. Perinotti, and A. Tosini, arXiv preprint arXiv:1511.03992 (2015).

SUPPLEMENTAL INFORMATION

IV. QUANTUM WALK FROM PRINCIPLES

Quantum Walks (QWs) describe the evolution of a quantum particle over a lattice. The dynamics is assumed to be *reversible*, hence the QW will be represented by a unitary transformation. By denoting with G the set of the lattice's points we can conveniently introduce the Hilbert space $\ell^2(G)$ and the orthonormal basis $|\mathbf{x}\rangle$ which corresponds to the position of the particle. If we associate the Hilbert space \mathbb{C}^s to the internal degrees of freedom of the particle, the QW is then a unitary operator on $\ell^2(G) \otimes \mathbb{C}^s$.

a. Locality The QW evolution is assumed to be local i.e. information propagates through the lattice at a bounded speed. Given a lattice, let $N_{\mathbf{x}}$ be the set of nearest neighbors of the site \mathbf{x} . If the particle is localized at site \mathbf{x} , then, after one step of the QW, it must be localized within a finite set $N_{\mathbf{x}}$. Such a locality condition introduces a notion of causal cone in the lattice.

b. Homogeneity The evolution is assumed to be homogeneous, i.e. requirement that all the sites are equivalent. The QW evolution should not not allow one to discriminate between two points \mathbf{x} and \mathbf{x}' . This requirement (see Ref. [1] for a full derivation and Appendix A for a shorter review) implies that the set og points G is a group and the lattice is a possible Cayley Graph of this group.

c. Isotropy The assumption of isotropy translate the requirement that there is no priviledged direction on the lattice. The mathematical translation of this requirement requires the existence of a group of permutation that act on the generators of the group G that can be faithfully represented on the internal degrees of freedom (see Ref. [1] for a full derivation and the Appendix A for a shorter review)

d. Flat and curved space The above sketched framework encompass a broad variety of dynamics. In particular, depending on the properties of the group G, we can have a quantum dynamics on a generally curved space. If we are interested to make contact with special relativity, it is natural to restrict the scenario to QWs corresponding to dynamics on the flat Minkovski spacetime. This requirement correspond to assume the group G to be virtually abelian, i.e. G has an abelian subgroup of finite order. We can further restrict ourselves to the case in which G is abelian without any loss of generality. The price to pay for this restriction is to add additional internal degrees of freedom (see Ref. [2] and Appendix A for a more complete discussion).

e. Fourier analysis If the group G is abelian it is convenient to study the dynamics in the Fourier transform basis $|\mathbf{k}\rangle := (2\pi)^{\frac{3}{2}} \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot\mathbf{x}} |\mathbf{x}\rangle$. Since homogeneity condition implies that the QW commutes with the translations on the lattice, in the Fourier basis the QW operator can be written as

$$A = \int_{B} \mathrm{d}^{3}\mathbf{k} \, |\mathbf{k}\rangle \langle \mathbf{k}| \otimes A_{\mathbf{k}} \tag{12}$$

where *B* denotes the first Brillouin zone of the underlying lattice. The unitary constraint implies that $A_{\mathbf{k}}$ is unitary for every $\mathbf{k} \in B$ and the locality assumption implies that $A_{\mathbf{k}}$ is a matrix polynomial in $e^{i\mathbf{h}\cdot\mathbf{k}}$. Notice that due to the discreteness of the lattice the QW is band-limited in \mathbf{k} . The spectrum $\{e^{i\omega_{\mathbf{k}}^{(i)}}\}$ of the operator $A_{\mathbf{k}}$, and expecially its *dispersion relation* that is the expression of the phases $\omega_{\mathbf{k}}^{(i)}$ as functions of \mathbf{k} , plays a crucial role in the analysis of the QW dynamics.

f. Weyl Quantum Walk If the dimension of the Hilbert space of the internal degrees of freedom is s = 2 and the group G is \mathbb{Z}^3 the requirements of *locality*, homogeneity and isotropy then [1] the QWs can only be defined over the body-centered-cubic lattice and they are equivalent (up to a local change of basis) to the following two QWs:

$$A_{\mathbf{k}}^{\pm} := \lambda^{\pm}(\mathbf{k})I - i\mathbf{n}^{\pm}(\mathbf{k}) \cdot \boldsymbol{\sigma}^{\pm}, \qquad (13)$$
$$\mathbf{n}^{\pm}(\mathbf{k}) := \begin{pmatrix} s_{x}c_{y}c_{z} \pm c_{x}s_{y}s_{z} \\ c_{x}s_{y}c_{z} \mp s_{x}c_{y}s_{z} \\ c_{x}c_{y}s_{z} \pm s_{x}s_{y}c_{z} \end{pmatrix},$$

$$\lambda^{\pm}(\mathbf{k}) := (c_x c_y c_z \mp s_x s_y s_z), \qquad (14)$$
$$c_{\alpha} := \cos(k_{\alpha}/\sqrt{3}), \ s_{\alpha} := \sin(k_{\alpha}/\sqrt{3}), \ \alpha = x, y, z.$$

The Pauli matrices $\sigma^+ = \sigma$ are the usual ones, while the $\sigma^- = \sigma^T$ are their transposed ones, and $\mathbf{k} \in \mathsf{B}$ where B denotes the Brillouin zone of the BCC lattice.

V. NONLINEAR LORENTZ TRANSFORMATIONS FOR THE WEYL QUANTUM WALK

In this section we give the explicit construction of the non-linear representation of the Lorentz group on the set of the solutions of the Weyl QW dynamics. The set of solution will be split into four regions, each one of them carrying a non-linear deformation of the Lorentz group.

Let us consider the splitting of the Brillouin zone ${\sf B}$

$$B'_{0} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}/\sqrt{3}) > 0 \}, B'_{1} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}/\sqrt{3}) > 0 \}, B'_{2} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}/\sqrt{3}) < 0 \}, B'_{3} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}/\sqrt{3}) < 0 \},$$
(15)

with $\mathsf{B} = \bigcup_{i=0}^{3} \mathsf{B}'_{i}$ up to a nullmeasure set, and let us denote with $\mathbf{n}^{(i)}(\mathbf{k})$ the restriction of $\mathbf{n}(\mathbf{k})$ to B'_{i} . Notice

that we dropped the label \pm denoting the two different Weyl walk, since the results holds the same in the two cases. We now denote with U the unit open ball in \mathbb{R}^3 (\overline{U} denotes its closure) and with H the subset of U defined as follows

$$\mathsf{H} := \{ \mathbf{m} \in \mathsf{U}s.t.m_x = \pm m_z, 2m_x^2 + 2m_y^2 \ge 1 \}.$$
 (16)

We then consider the regions

$$\mathsf{B}_i := \mathbf{n}^{(i)-1}(\mathsf{U} \setminus \mathsf{H}) \tag{17}$$

and the function $f(\omega, \mathbf{k})$ defined as follows:

$$f(\omega, \mathbf{k}) = g(\mathbf{n}(\mathbf{k})) =$$

$$= \tilde{g}(r, \theta, \phi) := 1 + r \int_{0}^{r} ds \left(\frac{1}{a(s)} + \frac{1}{b(s, \theta, \phi)} \right)$$

$$a(r) := 1 - r^{2},$$

$$b(r, \theta, \phi) := (\cos^{2}(\phi) - \sin^{2}(\phi))^{2} +$$

$$+ (\frac{1}{2} - r^{2}(1 - \cos^{2}(\theta)\sin^{2}(\phi)))^{2}$$
(18)

where we used the spherical coordinates $n_x(\mathbf{k}) = r \cos \theta \cos \phi$, $n_y(\mathbf{k}) = r \sin \theta$, $n_z(\mathbf{k}) = r \cos \theta \sin \phi$. Finally, we define the maps $\mathcal{D}^{(i)}$ as

$$\mathcal{D}^{(i)}: \Sigma_{i} \to \mathsf{\Gamma}_{0}, \quad \mathcal{D}^{(i)}: \begin{pmatrix} \omega \\ \mathbf{k} \end{pmatrix} \mapsto f(\omega, \mathbf{k}) \begin{pmatrix} \sin \omega \\ \mathbf{n}^{(i)}(\mathbf{k}) \end{pmatrix},$$

$$\Sigma_{i}:= \{(\omega, \mathbf{k}) \text{ s.t. } \mathbf{k} \in \mathsf{B}_{i}, \sin^{2} \omega - |\mathbf{k}|^{2} = 0\},$$

$$\mathsf{\Gamma}_{0}:= \{p \in \mathbb{R}^{4} \text{ s.t. } p_{\mu}p^{\mu} = 0\}.$$
(19)

One can prove (see Appendix) that the maps $\mathcal{D}^{(i)}$ define an analytic diffeomorphism between Σ_i and Γ_0 . As a consequence, the composition

$$\mathcal{L}_{\beta}^{(i)}: \Sigma_i \to \Sigma_i \qquad \mathcal{L}_{\beta}^{(i)}:= \mathcal{D}^{-1} \circ L_{\beta} \circ \mathcal{D}$$
 (20)

is a well defined nonlinear representation of the Lorentz group on the set Σ_i . Since the union of the B_i sets coincides with the whole (up to a null measure set) Brillouin zone, we have that the collection of the maps $\mathcal{L}_{\beta}^{(i)}$ provide a notion of Lorentz transformation for any (up to a null measure set) solution of the Weyl QW dynamics.

VI. DYNAMICAL SYMMETRIES OF DIRAC QUANTUM WALK

The Dirac quantum walk is obtained by performing the direct-sum of two Weyl walks and introducing offdiagonal blocks in such a way that the obtained matrix is a well defined quantum walk. As proved in Ref. [1], there are only two admissible walk

$$D^{\pm} = \begin{pmatrix} nA^{\pm} & imI\\ imI & nA^{\pm} \end{pmatrix}$$

$$0 \le n, m \le 1 \quad n^2 + m^2 = 1$$
(21)

The eigenvalue equation of the Dirac QW can be written as

$$[p_{\mu}^{(f)}(\omega, \mathbf{k}, m)\gamma^{\mu} - mI]\psi(\omega, \mathbf{k}, m) = 0, \qquad (22)$$

where γ^{μ} are the Dirac γ matrices in the Weyl representation, and m is then interpreted as the particle mass. Due to the explicit dependence of $p_{\mu}^{(f)}$ from m the covariance under change of reference cannot leave the value of minvariant. In such case the dispersion relation resorts to the conservation of the de Sitter norm

$$\sin^2 \omega - (1 - m^2) |\mathbf{n}(\mathbf{k})|^2 - m^2 = 0.$$
 (23)

The group leaving Eq. (23) invariant is the De Sitter group SO(1, 4). In the limit of $m \ll 1$, the usual Lorentz symmetry is recovered. The analysis of De Sitter covariance of Eq. (22) will be given in a forthcoming publication.

Appendix A: 3-dimensional Quantum Walks and the Weyl Quantum Walk

We consider Quantum Walks (QWs) that describe the evolution of a quantum particle over a lattice of \mathbb{R}^3 . The dynamics is assumed to be reversible, hence the QW will be represented by a unitary transformation. We consider *Bravais lattices*, i.e. lattices that are generated by linear independent discrete translations. By denoting with G the set of the lattice's points we can conveniently introduce the Hilbert space $\ell^2(G)$ and the orthonormal basis $|\mathbf{x}\rangle$ which corresponds to the position of the particle. The internal degrees of freedom of the particle are described by the Hilbert space \mathbb{C}^s .

The QW evolution is assumed to be *local* i.e. information propagates through the lattice at a bounded speed. Given a Bravais lattice, let $N_{\mathbf{x}}$ be the set of nearest neighbors of the site \mathbf{x} . If the particle is localized at site \mathbf{x} , then, after one step of the QW, it must be localized within the set $N_{\mathbf{x}}$. The set $N_{\mathbf{x}}$ corresponds to set of vectors $S := {\mathbf{h}}$ such that $N_{\mathbf{x}} = {\mathbf{y} \text{ s.t. } \mathbf{y} = \mathbf{x} + \mathbf{h}, \mathbf{h} \in S}$ Such a locality condition introduces a notion of causal cone in the lattice.

The QW evolution is also assumed to be *homogeneous*, i.e. it must commute with the discrete traslations. Let $T_{\mathbf{h}}$ be the translation operators on $\ell^2(G)$, such that i.e. $T_{\mathbf{h}}|\mathbf{x}\rangle = |\mathbf{x} + \mathbf{h}\rangle$ These assumption implies that the QW evolution can be represented by a unitary operator on $\ell^2(G) \otimes \mathbb{C}^s$ as follows:

$$A := \sum_{\mathbf{h} \in S} T_{\mathbf{h}} \otimes A_{\mathbf{h}} \tag{A1}$$

where $A_{\mathbf{h}}$ are $s \times s$ matrices and $T_{\mathbf{h}}$ is the translations that connect the site \mathbf{x} with its nearest neighbor $\mathbf{x} + \mathbf{h}$.

The QW dynamics is conveniently study in the Fourier transform basis $|\mathbf{k}\rangle := (2\pi)^{\frac{3}{2}} \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot\mathbf{x}} |\mathbf{x}\rangle$, in which the

QW operator A becomes

$$A = \int_{B} \mathrm{d}^{3}\mathbf{k} \, |\mathbf{k}\rangle \langle \mathbf{k}| \otimes A_{\mathbf{k}}, \qquad A_{\mathbf{k}} := \sum_{\mathbf{h} \in S} e^{-i\mathbf{k} \cdot \mathbf{h}} A_{\mathbf{h}}, \tag{A2}$$

where B denotes the first Brillouin zone of the underlying lattice and it is defined by the following linear constraints:

$$B := \bigcap_{i} \{ \mathbf{k} \in \mathbb{R}^{3} | -\pi | \tilde{\mathbf{h}}_{i} |^{2} \le | \mathbf{k} \cdot \tilde{\mathbf{h}}_{i} | \le \pi | \tilde{\mathbf{y}}_{i} |^{2} \}$$

$$\tilde{\mathbf{h}}_{i} \cdot \mathbf{h}_{j} = \delta_{ij}, \quad \mathbf{h}_{j} \in S.$$
 (A3)

The unitary constraint implies that $A_{\mathbf{k}}$ is unitary for every $\mathbf{k} \in B$ and the locality assumption implies that $A_{\mathbf{k}}$ is a matrix polynomial in $e^{i\mathbf{h}\cdot\mathbf{k}}$. Notice that due to the discreteness of the lattice the QW is band-limited in \mathbf{k} . The spectrum $\{e^{i\omega_{\mathbf{k}}^{(i)}}\}$ of the operator $A_{\mathbf{k}}$, and expecially its dispersion relation that is the expression of the phases $\omega_{\mathbf{k}}^{(i)}$ as functions of \mathbf{k} , plays a crucial role in the analysis of the QW dynamics.

It was shown in Ref.[1] that, when the dimension of the Hilbert space of the internal degrees of freedom is s = 2, the additional assumption of *isotropy*[3] of the evolution implies that the QWs can only be defined over the body-centered-cubic lattice and they are equivalent (up to a local change of basis) to the following two QWs:

$$A_{\mathbf{k}}^{\pm} := \lambda^{\pm}(\mathbf{k})I - i\mathbf{n}^{\pm}(\mathbf{k}) \cdot \boldsymbol{\sigma}^{\pm}, \qquad (A4)$$

where we define

$$\mathbf{n}^{\pm}(\mathbf{k}) := \begin{pmatrix} s_x c_y c_z \pm c_x s_y s_z \\ c_x s_y c_z \mp s_x c_y s_z \\ c_x c_y s_z \pm s_x s_y c_z \end{pmatrix},$$

$$\lambda^{\pm}(\mathbf{k}) := (c_x c_y c_z \mp s_x s_y s_z),$$

$$c_{\alpha} := \cos(k_{\alpha}/\sqrt{3}), \ s_{\alpha} := \sin(k_{\alpha}/\sqrt{3}), \ \alpha = x, y, z.$$
(A5)

The Pauli matrices $\sigma^+ = \sigma$ are the usual ones, while the $\sigma^- = \sigma^T$ are their transposed ones, and $\mathbf{k} \in \mathsf{B}$ where B denotes the Brillouin zone of the BCC lattice (see Fig. 4). In the limit $|\mathbf{k}| \ll 1$ we have

$$\mathbf{n}^{\pm}(\mathbf{k}) \sim \frac{\mathbf{k}}{\sqrt{3}}, \quad A^{\pm}_{\mathbf{k}} \sim \exp[-i\frac{\mathbf{k}}{\sqrt{3}} \cdot \boldsymbol{\sigma}^{\pm}],$$

corresponding to the evolution of the Weyl's equation
with the rescaling
$$\frac{\mathbf{k}}{\sqrt{3}} \rightarrow \mathbf{k}$$
. Since the A^+ and A^- re-
produce the dynamics of the Weyl equation in the limit
 $|\mathbf{k}| \ll 1$, we refer to them as Weyl QW.

For the sake of simplicity, in the following we will consider only one Weyl QW, i.e. we define $A_{\mathbf{k}} := A_{\mathbf{k}}^+$, the choice of Pauli matrices is the usual one $\boldsymbol{\sigma} := \boldsymbol{\sigma}^+$, and we similarly drop all the \pm superscripts. This choice is completely painless since all the results of this paper can be easily adapted to the choice $A_{\mathbf{k}} = A_{\mathbf{k}}^-$. In order to simplify the notation we also adopt the convention

$$k_i \to \frac{k_i}{\sqrt{3}}$$
 (A7)

(A6)




FIG. 4: *Top*: Body centered cubic lattice. Each pair of nearest neighbor is connected by a blue link. Each point of the lattice has 8 nearest neighbors, e.g. the red points are the nearest neighbors of the yellow one. *Bottom*: Brillouin zone B of the BCC lattice. The zone is a rhombic dodecahedron in which the opposite faces are identified.

in order to get rid of the annoying $\frac{1}{\sqrt{3}}$ factor. With this choice we have that in the $|\mathbf{k}| \ll 1$ limit, $\mathbf{n}(\mathbf{k}) \simeq \mathbf{k}$.

Appendix B: Deformed relativity and Weyl quantum walk

In this section we prove that the construction in Eq. is a well defined deformed Lorentz symmetry for each set Σ_i Let us define

$$\mathcal{D} = \mathcal{N} \circ \mathcal{P}, \qquad \mathcal{P} : (\omega, \mathbf{k}) \mapsto (\omega, \mathbf{n}(\mathbf{k}))$$
(B1)
$$\mathcal{N} : \begin{pmatrix} \omega \\ \mathbf{m} \end{pmatrix} \mapsto g(\omega, \mathbf{m}) \begin{pmatrix} \sin \omega \\ \mathbf{m} \end{pmatrix}$$

where we also assumed $f(\omega, \mathbf{k}) = g(\omega, \mathbf{n}(\mathbf{k}))$.

We now need to study separately the properties of the two maps \mathcal{P} and \mathcal{N} .

1. Study of the map n(k)

In this subsection we study the analytical properties of the map \mathcal{P} , which, according to Eq. (B1) resorts to the map **n**. The analysis will proceed through the determination of the largest subdomains $\{\mathbf{B}'_i\}$ of invertibility of the map **n**. We first prove that on the closure $\overline{\mathbf{B}}_i$ of each domain the map is surjective on the closed unit sphere $\overline{\mathbf{U}}$. Then, we determine the geometry of the ranges $\mathbf{n}(\mathbf{B}'_i)$, showing that they are homotopic to a solid genus-two torus.

Let us denote by B the Brillouin zone of the center cubic lattice. B, upon a proper identification of its boundary points (see Fig. 4) is a compact 3-dimensional manifold. The Jacobian $J_{\mathbf{n}}(\mathbf{k})$ of the map $\mathbf{n}(\mathbf{k})$ is given by

$$J_{\mathbf{n}}(\mathbf{k}) := \det[\partial_i n_j(\mathbf{k})] = \cos(2k_y)\lambda(\mathbf{k}), \qquad (B2)$$

and it vanishes on the set

$$F = G \cup X,$$

$$X := \{ \mathbf{k} \in \mathsf{B} | \cos(2k_y) = 0 \},$$

$$G := \{ \mathbf{k} \in \mathsf{B} | \lambda(\mathbf{k}) = 0 \}.$$
(B3)

Since $\nabla \lambda(\mathbf{k}) \neq 0$ for all \mathbf{k} such that $\lambda(\mathbf{k}) = 0$, the implicit function theorem guarantees that G is a well defined 2dimensional surface. In the following we will denote by $\{\mathsf{B}'_i\}$ (*i* ranging in some set) the disjoint connected subsets of $\mathsf{B} \setminus \mathsf{F}$, thus

$$\mathsf{B} \setminus \mathsf{F} = \bigcup_{i} \mathsf{B}'_{i}, \quad \mathsf{B}'_{i} \cap \mathsf{B}'_{j} = \emptyset \text{ for } i \neq j.$$
(B4)

For each *i* the set B'_i is open and we denote as $\overline{B'_i}$ its closure and as $\partial B'_i$ its boundary.

Now let us denote with $\overline{U} \subset \mathbb{R}^3$ the closed unit-radius ball, and with S^2 the sphere of radius 1 in \mathbb{R}^3 . Moreover, let us define the parametric curves

$$\mathbf{e}_{\pm}(t) := \frac{1}{\sqrt{2}} (\sin(t), \cos(t), \pm \sin(t))^T \qquad (B5)$$

and the sets

$$\begin{aligned} \mathsf{Q}_a &:= \mathsf{U} \setminus (\mathbf{e}_+(\mathsf{T}_1) \cup \mathbf{e}_-(\mathsf{T}_2)) \\ \mathsf{Q}_b &:= \mathsf{U} \setminus (\mathbf{e}_+(\mathsf{T}_2) \cup \mathbf{e}_-(\mathsf{T}_1)) \\ \mathsf{T}_1 &:= (-\frac{\pi}{2}, \frac{\pi}{2}) \\ \mathsf{T}_2 &:= (-\pi, -\frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]. \end{aligned} \tag{B6}$$

Given all the definition introduced in this section, we have then the following result:

Lemma 1 There are four different B'_i regions, determined by the following conditions

$$B'_{0} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}) > 0 \}, B'_{1} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}) > 0 \}, B'_{2} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}) < 0 \}, B'_{3} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}) < 0 \}.$$
(B7)

For each *i*, let $\mathbf{n}^{(i)}(\mathbf{k})$ denote the restriction of the map $\mathbf{n}(\mathbf{k})$ to the set B'_i . Then $\mathbf{n}^{(i)}(\mathbf{k})$ defines a diffeomorphism between B'_i and its image $\mathbf{n}^{(i)}(\mathsf{B}'_i)$ and we have

$$\mathbf{n}^{(0)}(\mathsf{B}'_0) = \mathbf{n}^{(2)}(\mathsf{B}'_2) = \mathsf{Q}_a$$

$$\mathbf{n}^{(1)}(\mathsf{B}'_1) = \mathbf{n}^{(3)}(\mathsf{B}'_3) = \mathsf{Q}_b.$$
 (B8)

The proof of this result is rather involved and can be found in Appendix C. The B'_i regions are plotted in Fig. 5 The most important consequence of this result is that, for each *i*, the set $\mathbf{n}(B'_i)$ (see Fig. 6) i) coincides with U except a null measure set and ii) it is homeomorphic to a genus two torus.

2. Study of the map \mathcal{N}

Since for all *i* the region $\mathbf{n}(\mathsf{B}'_i)$ has a nontrivial topology, the set $\{(\omega, \mathbf{m}) \text{ s.t. } |\omega| \leq \frac{\pi}{2}, \mathbf{m} \in \mathbf{n}(\mathsf{B}'_i), \sin^2 \omega - |\mathbf{m}|^2 = 0\}$ cannot be diffeomorphic to any Lorentzinvariant region of \mathbb{M}^4 . A possible way to change the topology of $\mathbf{n}(\mathsf{B}'_i)$ is to exclude the set $\mathsf{H} \subseteq \overline{\mathsf{U}}$ (as it is shown in Fig. 6) of vectors \mathbf{m} satisfying the following inequalities

$$\begin{cases} m_x = \pm m_z, \\ 2m_x^2 + m_y^2 \le 1, \\ 2m_x^2 + 2m_y^2 \ge 1. \end{cases}$$
(B9)

Then, the set $U \setminus H$ is topologically trivial and we have $U \setminus H \subset \mathbf{n}(\mathsf{B}'_i)$ for all *i*. Let us now consider the function $\mathcal{N} : (\omega, \mathbf{m}) \mapsto (p_0, \mathbf{p}) = g(\omega, \mathbf{m})(\sin \omega, \mathbf{m})$ restricted to the set

$$\mathsf{N} := \{(\omega, \mathbf{m}) \text{ s.t } \mathbf{m} \in \mathsf{U} \setminus \mathsf{H}, |\omega| \le \frac{\pi}{2}, \sin^2 \omega - |\mathbf{m}|^2 = 0\}.$$
(B10)

As shown in Appendix D, it is possible to define the function $g(\omega, \mathbf{m})$ is such that \mathcal{N} defines a diffeomorphism between N and the null mass-shell

$$\Gamma_0 := \{ p \in \mathbb{M}^4, \text{ s.t. } p^\mu p_\mu = 0 \}.$$
 (B11)

and that its Jacobian matrix at the origin is 0, i.e

$$J_{\mathcal{N}}(\mathbf{0}) = I. \tag{B12}$$

Finally, for each *i* we denote by B_i the counter-image of the set $\mathsf{U} \setminus \mathsf{H}$ under the map $\mathbf{n}^{(i)}$ and by $\mathcal{D}^{(i)}$ the composition

$$\mathcal{D}^{(i)} : \mathbf{\Sigma}_{i} \to \mathbf{\Gamma}_{0} \qquad \mathcal{D}^{(i)} := \mathcal{N} \circ \mathcal{P}^{(i)} \qquad (B13)$$
$$\mathcal{P}^{(i)} : \mathbf{\Sigma}_{i} \to \mathbf{N} \qquad \mathcal{P}^{(i)} : \begin{pmatrix} \omega \\ \mathbf{k} \end{pmatrix} \mapsto \begin{pmatrix} \omega \\ \mathbf{n}^{(i)}(\mathbf{k}) \end{pmatrix}$$
$$\mathcal{N} : \mathbf{N} \to \mathbf{\Gamma}_{0} \qquad \mathcal{N} : \begin{pmatrix} \omega \\ \mathbf{m} \end{pmatrix} \mapsto g(\omega, \mathbf{m}) \begin{pmatrix} \sin \omega \\ \mathbf{m} \end{pmatrix}$$
$$\mathbf{\Sigma}_{i} := \{(\omega, \mathbf{k}) \text{ s.t. } \mathbf{k} \in \mathbf{B}_{i}, \sin^{2} \omega - |\mathbf{k}|^{2} = 0\}.$$




FIG. 5: (Colors online). Top left figure: surfaces $\lambda(\mathbf{k}) = 0$ in Eq. (A5) (yellow) and $\cos(2k_y) = 0$ (red planes) inside the Brillouin zone (transparent). Top middle figure: B_0 zone (red X-shaped). Top right figure: B_0 (red) and B_1 (blue). Bottom left to right: B_1 , B_2 , B_3 . Bottom right: region B_1 represented in a properly translated Brillouin zone. In this paper the Lorentz transformations are those that leave the dispersion relations of the Weyl QW invariant, and act on the Weyl spinor independently of the wave-vector. In such way they are nonlinear in (ω, \mathbf{k}) and linear over the Weyl spinor. Therefore the Lorentz group acts as a group of diffeomorphisms over the Brillouin zone B. The four domains $B_i \subset B$ are Lorentz invariant (up to a null-measure set, see Fig. 6). More precisely a point (ω, \mathbf{k}) with $\mathbf{k} \in B_i$ and $\sin^2 \omega - |\mathbf{n}(\mathbf{k}')|^2 = 0$ is mapped to a point (ω', \mathbf{k}') with $\sin^2 \omega' - |\mathbf{n}(\mathbf{k}')|^2 = 0$ and $\mathbf{k}' \in B_i$. Moreover, the map **n** maps each B_i into the same set (up to null

to a point (ω, \mathbf{k}) with $\sin^{-}\omega - |\mathbf{n}(\mathbf{k})|^{-} = 0$ and $\mathbf{k} \in B_i$. Moreover, the map \mathbf{n} maps each B_i into the same set (up to null measure set: see Fig. 6). Since the kinematics of a wavevector \mathbf{k} depends only on the vector $\mathbf{n}(\mathbf{k})$, we can conclude that the B_i regions are kinematically equivalent and they can be interpreted as four different massless Weyl Fermions. Because of the identification of the boundary points in the Brillouin zone, all the B_i regions have the same X-shape as B_0 . This is evident in the bottom right figure, in which we see that the region B_1 (in red), when represented in a properly translated Brillouin zone (in blue), has the same X-shape as the region B_0 . Considering the identification of the boundary points of the Brillouin zone in Fig. 4, one realizes that the opposite arms of the X are glued together, resulting in a solid double-torus (genus-two). This result is rigorously proved in the text where we show that the B_i regions are diffeomorphic to a solid ball pierced by two arches of ellipses (Fig. 6).

For each *i*, the map $\mathcal{D}^{(i)}$ is an analitic diffeomorphism between the region Σ_i and the Lorentz invariant set Γ_0 which satisfies the condition $J_{\mathcal{D}_i}(\mathbf{0}) = I$. Then the composition

$$\mathcal{L}_{\beta}^{(i)}: \Sigma_i \to \Sigma_i \qquad \mathcal{L}_{\beta}^{(i)}:= \mathcal{D}^{-1} \circ L_{\beta} \circ \mathcal{D} \qquad (B14)$$

is a well defined nonlinear representation of the Lorentz group on the set Σ_i . Since the union of the B_i sets coincides with the whole (up to a null measure set) Brillouin zone, we have that the collection of the maps $\mathcal{L}_{\beta}^{(i)}$ provide a notion of Lorentz transformation for any (up to a null measure set) solution of the Weyl QW dynamics.

Appendix C: Proof of Lemma 1

In this section will give the proofs of the results contained in Lemma 1. Since the proof is quite involved, we split it into several pieces. Let us begin by defining the sets

$$\mathsf{Q}' := \overline{\mathsf{U}} \setminus \mathsf{R} \qquad \mathsf{R} := \mathsf{S}^2 \cup \mathsf{E}_+ \cup \mathsf{E}_- \,. \tag{C1}$$

Obviously Q' is open and connected, with $\overline{Q'} = \overline{U}$ and $\partial Q' = R$. We now prove some useful properties of the map **n**.




FIG. 6: (Colors online) Left figure: region Q_a . Right figure: H zone in red inside the unit ball. In the left figure, the tubes around the arches $\mathbf{e}_+(\mathsf{T})_1$ and $\mathbf{e}_-(\mathsf{T})_2$ emphasize the piercing of the ball by the one-dimensional holes along the elliptic arches $\mathbf{e}_+(\mathsf{T})_1$ and $\mathbf{e}_-(\mathsf{T})_2$. The region Q_a is clearly homeomorphic to a solid torus of genus two. Because of this non-trivial topological feature the set $\{(\omega, \mathbf{m}) \text{ s.t. } |\omega| \leq \frac{\pi}{2}, \mathbf{m} \in \mathbf{n}(\mathsf{B}_i), \sin^2 \omega - |\mathbf{m}|^2 = 0\}$ cannot be diffeomorphic to any Lorentz-invariant region of \mathbb{M}^4 . However it is possible to remove from the region Q_a a null-measure set such that the resulting topology is trivial. This can be done by removing the set H (red zones in the right figure), resulting in a star-shaped open set in \mathbb{R}^3 .

Sublemma 1 Let \mathbf{n}_i denote the restriction of the map \mathbf{n} to \mathbf{B}'_i . Then for each i we have that \mathbf{n}_i is a diffeomorphism between \mathbf{B}'_i and $\mathbf{n}(\mathbf{B}'_i)$

Proof. Since by definition $\mathbf{k} \in \mathsf{B}'_i \Rightarrow \mathbf{k} \notin \mathsf{F}$ we have $J_{\mathbf{n}}(\mathbf{k}) \neq 0$ for all $\mathbf{k} \in \mathsf{B}'_i$. Since B'_i is connected and \mathbf{n} is analytical, we have the thesis.

Sublemma 2 We have the following inclusions:

1. $\mathbf{n}(\overline{\mathsf{B}'_i}) \subseteq \overline{\mathsf{U}},$ 2. $\partial \mathbf{n}(\mathsf{B}'_i) \subseteq \mathsf{R}.$

Proof. Let us start with the proof of item 1. By explicit computation we have $|\mathbf{n}(\mathbf{k})|^2 = 1 - \lambda^2(\mathbf{k}) \leq 1$ which implies that the image of \mathbf{n} is contained in \overline{U} .

We now prove item 2. Thanks to Lemma 1 we have that $\mathbf{n}(\mathsf{B}'_i)$ is open. On the other hand, since \mathbf{n} is continuous and $\overline{\mathsf{B}'_i}$ is compact, we have that $\mathbf{n}(\overline{\mathsf{B}'_i})$ is compact and then it is closed. Then the trivial inclusion $\mathbf{n}(\mathsf{B}'_i) \subseteq \mathbf{n}(\overline{\mathsf{B}'_i})$ implies $\mathbf{n}(\mathsf{B}'_i) \subseteq \mathbf{n}(\overline{\mathsf{B}'_i})$. By definition we have $\mathbf{n}(\mathsf{B}'_i) = \mathbf{n}(\mathsf{B}'_i) \cup \partial \mathbf{n}(\mathsf{B}'_i) \subseteq \mathbf{n}(\mathsf{B}'_i)$. By definition we have $\mathbf{n}(\mathsf{B}'_i) = \mathbf{n}(\mathsf{B}'_i) \cup \partial \mathbf{n}(\mathsf{B}'_i)$ with $\mathbf{n}(\mathsf{B}'_i) \cap \partial \mathbf{n}(\mathsf{B}'_i) = \emptyset$ and $\mathbf{n}(\overline{\mathsf{B}'_i}) = \mathbf{n}(\mathsf{B}'_i) \cup \mathbf{n}(\partial\mathsf{B}'_i)$. Then the inclusion $\mathbf{n}(\mathsf{B}'_i) \subseteq$ $\mathbf{n}(\overline{\mathsf{B}'_i})$ implies $\partial \mathbf{n}(\mathsf{B}'_i) \subseteq \mathbf{n}(\partial\mathsf{B}'_i)$. Since $\partial\mathsf{B}'_i \subseteq \mathsf{F}$ we have $\mathbf{n}(\partial\mathsf{B}'_i) \subseteq \mathbf{n}(\mathsf{F})$. One can then verify by direct computation that $\mathbf{n}(\mathsf{F}) \subseteq \mathsf{R}$ thus proving the thesis.

We now recall a result of basic topology which will be useful in the following.

Sublemma 3 Let A and B be open sets such that $\overline{A} \subset \overline{B}$. Then there exists a point p such that $p \in \operatorname{int}\overline{B}$ and $p \notin \overline{A}$.

Proof. Let us suppose that $B \subseteq \overline{A}$. Since B is open and \overline{A} is closed, we have $\overline{B} \subseteq \overline{A}$ which contradicts the hypothesis.

The following result will be of crucial importance.

Sublemma 4 The following identity holds:

$$\overline{\mathbf{n}}(\mathsf{B}'_i) = \overline{\mathsf{U}}.\tag{C2}$$

Proof. First we prove the easiest inclusion $\overline{\mathbf{n}}(\mathsf{B}'_i) \subseteq \overline{\mathsf{U}}$. From item 1 of Sublemma 2 we have that $\mathbf{n}(\mathsf{B}'_i) \subseteq \mathbf{n}(\overline{\mathsf{B}'_i}) \subseteq \overline{\mathsf{U}}$ (the first inclusion is trivial). Reminding that $\mathbf{n}(\mathsf{B}'_i)$ is open we have $\overline{\mathbf{n}}(\mathsf{B}'_i) \subseteq \overline{\mathsf{U}}$.

We now prove that $\overline{U} \subseteq \overline{\mathbf{n}(\mathsf{B}'_i)}$. By contradiction, let us suppose that the strict inclusion $\overline{\mathbf{n}(\mathsf{B}'_i)} \subset \overline{\mathsf{U}}$ holds. Then, thanks to Sublemma 3, we find $p \in \mathsf{U}$ such that $p \notin \overline{\mathbf{n}(\mathsf{B}'_i)}$. Moreover we can find an open neighborhood N of p such that $\mathsf{N} \cap \overline{\mathbf{n}(\mathsf{B}'_i)} = \emptyset$ and then without loss of generality we can suppose that $p \in \mathsf{Q}'$. Since R has no interior points, $\mathbf{n}(\mathsf{B}'_i)$ cannot be included in R, whence $\mathbf{n}(\mathsf{B}'_i) \cap \mathsf{Q}'$ is not empty. Let us now fix a point $q \in \mathbf{n}(\mathsf{B}'_i) \cap \mathsf{Q}'$. Then, for any continuous path γ connecting p and q there exist t'such that $\gamma(t') \in \partial \mathbf{n}(\mathsf{B}'_i)$. From item 2 of Sublemma 2 we have $\gamma(t') \in \mathsf{R}$. Since this conclusion contradicts the fact that Q' is connected, we have proved the thesis.

As a consequence we have

Corollary 1 The following inclusion holds $Q' \subseteq n(B'_i)$.

Proof. From Lemma 4 we have $Q' \cup R = \mathbf{n}(B'_i) \cup \partial \mathbf{n}(B'_i)$. Reminding that $Q' \cap R = \emptyset = \mathbf{n}(B'_i) \cap \partial \mathbf{n}(B'_i)$ and the inclusion $\partial \mathbf{n}(B'_i) \subseteq R$, proved in Sublemma 2, we have the thesis. We now turn our attention to the regions B'_i . Our first objective is to determine how many different B'_i regions are. The answer is provided by the following result.

Sublemma 5 The regions B'_i are in one-to-one correspondence with the solution of the equation $|\lambda(\mathbf{k})|^2 = 1$.

Proof. We proved that the map \mathbf{n}_i defines a diffeomorphism between B'_i and the set $\mathbf{n}(\mathsf{B}'_i) \subseteq \overline{P}$ which includes the origin. Then, for each B'_i there exist a point \mathbf{k} such that $\mathbf{n}(\mathbf{k}) = 0$ and it is unique. Since $\mathbf{n}(\mathbf{k}) = 0$ if and only if $|\mathbf{n}(\mathbf{k})|^2 = |\lambda(\mathbf{k})|^2 - 1 = 0$ we have the thesis.

Thanks to this result it is sufficient to find the solutions of $|\lambda(\mathbf{k})|^2 = 1$ in the Brillouin zone. One can easily check that there are only 4 solutions and then 4 different regions B'_0, \ldots, B'_3 .

We can now prove Eq. (B7) of Lemma 1.

Sublemma 6 The region B'_i are given by

$$B'_{0} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}) > 0 \}, B'_{1} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}) > 0 \}, B'_{2} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) > 0, \cos(2k_{y}) < 0 \}, B'_{3} := \{ \mathbf{k} \in B | \lambda(\mathbf{k}) < 0, \cos(2k_{y}) < 0 \}.$$
(C3)

Proof. Let us denote with \tilde{B}_i the regions defined by the right hand sides of Eq. C3. One can immediately see that: i) the \tilde{B}_i are open sets, ii) the \tilde{B}_i are mutually disjoint and iii) the union of the \tilde{B}_i is the union of the B'_i . We now prove that for all \tilde{B}_i there exist a unique B'_j such that $\tilde{B}_i \subseteq B'_j$. This fact, together with the previous properties of the \tilde{B}_i gives $\tilde{B}_i = B'_i$

Clearly for all \tilde{B}_i we must have $\tilde{B}_i \subseteq B'_{j_1} \cup \cdots \cup B'_{j_k}$ for some $k \ge 1$ Let us suppose then that there exist \tilde{B}_i such that $\tilde{B}_i \subseteq B'_{j_1} \cup \cdots \cup B'_{j_k}$ with k strictly greater than 1. Since we have as many \tilde{B}_i as B'_i , there must exist a B'_i and two points $\mathbf{k}_a \in \tilde{B}_a$ and $\mathbf{k}_b \in \tilde{B}_a$ such that $\mathbf{k}_a, \mathbf{k}_b \in B'_i$. Since B'_i is connected there must exist a path connecting \mathbf{k}_a and \mathbf{k}_b that entirely lies within B'_i . On the other hand, since the \tilde{B}_i are disjoint, this path would cross the border of the \tilde{B}_a but this contradicts the fact that the border of the \tilde{B}_a are not included in B'_i .

Finally, we can give the complete characterization of the sets $\mathbf{n}(\mathsf{B}'_i)$. From Corollary 1 we have the inclusion $\mathsf{Q}' \subseteq \mathbf{n}(\mathsf{B}'_i)$. Since $|\mathbf{n}(\mathbf{k})| = 1 \iff \lambda(\mathbf{k}) = 0$, we know that the $S^2 \not\subset \mathbf{n}(\mathsf{B}'_i)$. It is easy to check that also the points $p_{\pm} := (0, \pm \frac{\sqrt{2}}{2}, 0)$ are not included in the set $\mathbf{n}(\mathsf{B}'_i)$. For any region $\mathbf{n}(\mathsf{B}'_i)$, we will determine which ones of the 8 open arches defined as

are included in $\mathbf{n}(\mathsf{B}'_i)$.

Let us consider the sets E_{j}^{+} . If we for some t we have $\mathbf{n}(\mathbf{k}) = \mathbf{e}_{+}(t)$ and $\cos(2k_{y}) \neq 0$ then it must be

$$k_x = \frac{\pi}{4} + n\frac{\pi}{2}$$
 $k_z = k_x + m\pi$ (C5)

for n and m integers. Eq. (C5) then implies

$$\lambda(\mathbf{k}) = (-1)^m \frac{1}{2} (\cos(k_y) - \sin(k_y)).$$
(C6)

From Eq. (C6) we have

$$\lambda(\mathbf{k}) > 0 \Rightarrow \begin{cases} m \text{ even } \wedge -\frac{3}{4}\pi < k_y < \frac{1}{4}\pi \\ m \text{ odd } \wedge \frac{1}{4}\pi < k_y < \frac{5}{4}\pi \end{cases}$$
(C7)

Then, if we assume $\mathbf{k} \in \mathsf{B}_0'$ we must have

$$m \text{ even }, -\frac{1}{4}\pi < k_y < \frac{1}{4}\pi$$
$$m \text{ odd }, \frac{3}{4}\pi < k_y < \frac{5}{4}\pi$$
(C8)

However, since the two sets of **k** are related by a translation of $(0, l\pi, l\pi), l \in \mathbb{Z}$ they actually represent the same set in the Brillouin zone. So it suffices to consider just the first set in Eq. (C8), that can be written as

$$k_x = \frac{1}{4}\pi + n\frac{1}{2}\pi \qquad \frac{1}{4}\pi < k_y < \frac{1}{4}\pi k_z = \frac{1}{4}\pi + n\frac{1}{2}\pi + m\pi = \frac{1}{4}\pi + n\frac{1}{2}\pi$$
(C9)

where we used Eq. (C5) and in the second equality in the second line of Eq. (C9) we used the hypothesis that m is even. Using again the fact that we identify points related by a translation of $(l\pi, 0, l\pi), l \in \mathbb{Z}$ we find just two inequivalent sets

$$\mathsf{Z}_1 := \begin{cases} k_x = \frac{1}{4}\pi \\ \frac{1}{4}\pi < k_y < \frac{1}{4}\pi \\ k_z = \frac{1}{4}\pi \end{cases} \quad \mathsf{Z}_2 := \begin{cases} k_x = -\frac{1}{4}\pi \\ \frac{1}{4}\pi < k_y < \frac{1}{4}\pi \\ k_z = -\frac{1}{4}\pi \end{cases}$$

It is now easy to show that the images of this two sets under the map \mathbf{n} are $\mathbf{n}(\mathsf{Z}_1) = \mathsf{E}_2^+$ and $\mathbf{n}(\mathsf{Z}_2) = \mathsf{E}_4^+$. By applying an analogous line of reasoning one can prove all the following inclusions

$$(E_{2}^{+} \cup E_{4}^{+} \cup E_{1}^{-} \cup E_{3}^{-}) \subset \mathbf{n}(B_{0}^{\prime})
(E_{1}^{+} \cup E_{3}^{+} \cup E_{2}^{-} \cup E_{4}^{-}) \not\subset \mathbf{n}(B_{0}^{\prime})
(E_{2}^{+} \cup E_{4}^{+} \cup E_{1}^{-} \cup E_{3}^{-}) \subset \mathbf{n}(B_{2}^{\prime})
(E_{1}^{+} \cup E_{3}^{+} \cup E_{2}^{-} \cup E_{4}^{-}) \not\subset \mathbf{n}(B_{1}^{\prime})
(E_{1}^{+} \cup E_{3}^{+} \cup E_{2}^{-} \cup E_{4}^{-}) \subset \mathbf{n}(B_{1}^{\prime})
(E_{2}^{+} \cup E_{4}^{+} \cup E_{1}^{-} \cup E_{3}^{-}) \not\subset \mathbf{n}(B_{1}^{\prime})
(E_{1}^{+} \cup E_{3}^{+} \cup E_{2}^{-} \cup E_{4}^{-}) \subset \mathbf{n}(B_{3}^{\prime})
(E_{2}^{+} \cup E_{4}^{+} \cup E_{1}^{-} \cup E_{3}^{-}) \not\subset \mathbf{n}(B_{3}^{\prime}).$$
(C10)

This result completes the proof of Eq. (B8) of Lemma 1.

Appendix D: The function $g(\omega, \mathbf{m})$

In this section we now show how it is possible to define a function $g(\omega, \mathbf{m})$ such that the map \mathcal{N} defines a diffeomorphism between N and the null mass-shell Γ_0 . The purpose of the following construction is to reduce the region N to a star-shaped region \tilde{N} by removing a null measure region, and to define the function $g(\omega, \mathbf{m})$ in such a way that the map \mathcal{N} is invertible on \tilde{N} . Since multiplication by $g(\omega, \mathbf{m})$ rescales the four vector $(\sin \omega, \mathbf{m})$ without affecting its direction, in order to have an invertible map \mathcal{N} it is sufficient to ensure that $g(\omega, \mathbf{m})$ is radially monotonic versus \mathbf{m} , namely $g(\omega, r\mathbf{m}_0)$ must be monotonic versus r.

Let us denote E_{\pm} the ellipses defined by the parametric equations (B5). We define the polynomials

$$h_{\mathsf{U}}(r,\theta,\phi) := 1 - r^{2},$$

$$h_{\mathsf{E}}(r,\theta,\phi) := (\cos^{2}(\phi) - \sin^{2}(\phi))^{2} + (\frac{1}{2} - r^{2}(1 - \cos^{2}(\theta)\sin^{2}(\phi)))^{2},$$
(D1)

where we used the spherical coordinates $m_x = r \cos \theta \cos \phi$, $m_y = r \sin \theta$, $m_z = r \cos \theta \sin \phi$. Clearly we have $h_{\mathsf{U}}(\mathbf{m}), h_{\mathsf{E}}(\mathbf{m}) > 0 \ \forall \mathbf{n} \in, h_{\mathsf{U}}(\mathbf{m}) = 0 \Leftrightarrow \mathbf{m} \in \mathsf{S}^2$, $h_{\mathsf{E}}(\mathbf{m}) = 0 \Leftrightarrow \mathbf{m} \in \mathsf{E}_+ \cup \mathsf{E}_-$ and $h_{\mathsf{U}}, h_{\mathsf{E}}$ are analytic on $\mathsf{U} \setminus \mathsf{H}$. Since $\mathsf{U} \setminus \mathsf{H}$ is star-shaped we can define

$$\tilde{g}(r,\theta,\phi) := r \int_0^r ds \left(\frac{1}{h_{\mathsf{U}}(s,\theta,\phi)} + \frac{1}{h_{\mathsf{E}}(s,\theta,\phi)} \right). \quad (\mathrm{D2})$$

The condition $h_{\mathsf{U}}(\mathbf{m}), h_{\mathsf{E}}(\mathbf{m}) > 0 \ \forall \mathbf{n} \in \mathsf{U} \setminus \mathsf{H}$ implies that the function $\tilde{g}(r, \theta, \phi)$ is radially monotonic on $\mathsf{U} \setminus \mathsf{H}$.

- G. M. D'Ariano, P. Perinotti, Phys. Rev. A 90, 062106 (2014).
- [2] G. M. D'Ariano, M. Erba, P. Perinotti, A. Tosini, arXiv preprint arXiv:1511.03992.
- [3] Within this context, the isotropy condition, meaning that all the direction on the lattice are equivalent, is stated as follows. First we require that the set S is the disjoint union of two subsets $S_+ \cup S_- = S$ such that if $\mathbf{h} \in S_+$

Since $h_{\mathsf{U}}(\mathbf{m})$ are $h_{\mathsf{E}}(\mathbf{m})$ are analytic on $\mathsf{U} \setminus \mathsf{H}$ we have that $\tilde{g}(r, \theta, \phi)$ is analytic on $(\mathsf{U} \setminus \mathsf{H}) \setminus \mathbf{0}$. Moreover, since $\tilde{g}(r, \theta, \phi)$ is even in r we have that $\tilde{g}(\mathbf{m})$ is analytic on the whole domain $\mathsf{U} \setminus \mathsf{H}$. Finally it is easy to check that $g(\mathbf{m})$ diverges to $+\infty$ as \mathbf{m} approaches the boundary of $\mathsf{U} \setminus \mathsf{H}$, and that $(\nabla g)(\mathbf{0}) = \mathbf{0}$. Let us define

$$g(\mathbf{m}) := \tilde{g}(\mathbf{m}) + 1. \tag{D3}$$

We now check that, with this definition of the map $g(\mathbf{m})$, the map \mathcal{N} defines an analytic diffeomorphism between $U \setminus H$ and Γ_0 with the property $J_{\mathcal{N}}(\mathbf{0}) = I$. Clearly \mathcal{N} is analytic in N so we just need to prove that it gives a bijection between N and Γ_0 . Let us fix a versor **j** in \mathbb{R}^3 . Then, in the **j** direction we have

$$\mathcal{N}(\omega, r\mathbf{j}) = g(r, \theta_{\mathbf{j}}, \phi_{\mathbf{j}}) \begin{pmatrix} \sin(\omega) \\ r\mathbf{j} \end{pmatrix}.$$
 (D4)

Since $g(r, \theta_{\mathbf{j}}, \phi_{\mathbf{j}})$ is monotone we clearly have that \mathcal{N} is injective. We now prove the surjectivity of $\mathcal{N}'(\omega, r) :=$ $\mathcal{N}(\omega, r\mathbf{j})$ on the set $\mathsf{K} := \{(p_0, p_1) \in \mathbb{R}^2 \text{ s.t. } p_0^2 - p_1^2 = 0\}$. Let us fix a point $p = (p_0, p_1) \in \mathsf{K}$. Since $g(r, \theta_{\mathbf{j}}, \phi_{\mathbf{j}})$ is monotone and surjective on $[1, +\infty)$ we can find a value r_p such that $r_p g(r_p, \theta_{\mathbf{j}}, \phi_{\mathbf{j}}) = p_1$. Clearly, since $|r_p| < 1$, also the equation $\sin^2(\omega_p) = |r_p|^2$ can be solved and then $\mathcal{N}'(\omega_p, r_p) = (p_0, p_1)$. Since the surjectivity of \mathcal{N} holds for any direction \mathbf{j} , we have that \mathcal{N} is a diffeomorphism between N and Γ_0 . Finally, since $g(\mathbf{0}) = 1$ and $\nabla g(\mathbf{0}) =$ $\mathbf{0}$, we have that the Jacobian of the map \mathcal{N} is the identity, which proves Eq. (B12). Obviously, our choice of the map $g(\mathbf{m})$ is not unique.

then $\mathbf{h} \in S_{-}$. Then, let L be a group automorphisms of the Bravais lattice which is transitive over the translations in S_{+} . We then say that A is isotropic if there exists a faithful representation U of L over \mathbb{C}^{s} such that one has

$$A = \sum_{\mathbf{h} \in S} T_{\mathbf{h}} \otimes A_{\mathbf{h}} = \sum_{\mathbf{h} \in S} T_{l(\mathbf{h})} \otimes U_{l} A_{\mathbf{h}} U_{l}^{\dagger}, \quad \forall l \in L.$$