

Topology of the cone of positive maps on qubit systems

Marek Miller^{*1} and Robert Olkiewicz^{†1}

¹Instytut Fizyki Teoretycznej, Uniwersytet Wrocławski, Poland

Abstract

An alternative, geometrical proof of a known theorem concerning the decomposition of positive maps of the matrix algebra $M_2(\mathbb{C})$ has been presented. The premise of the proof is the identification of positive maps with operators preserving the Lorentz cone in four dimensions, and it allows to decompose the positive maps with respect to those preserving the boundary of the cone. In addition, useful conditions implying complete positivity of a map of $M_2(\mathbb{C})$ have been given, together with a sufficient condition for complete positivity of an extremal Schwarz map of $M_n(\mathbb{C})$. Lastly, following the same geometrical approach, a description in topological terms of maps that are simultaneously completely positive and completely copositive has been presented.

Introduction

The theory of positive maps constitutes an increasingly popular subject of research, and its mathematical appeal matches only its ability to present constructive applications to the theory of quantum information and quantum entanglement [1, 2]. The classification of positive maps on $M_2(\mathbb{C})$, the algebra of square complex matrices of size 2, in physical terms representing a qubit, was a first turning point of the theory. The theorem proven by E. Størmer [3] and S. L. Woronowicz [4] says that every positive map $S : M_2(\mathbb{C}) \rightarrow M_n(\mathbb{C})$, where $n \leq 3$, can be decomposed into the sum: $S = \Lambda_1 + \Lambda_2 \circ t$, where both maps $\Lambda_1, \Lambda_2 : M_2(\mathbb{C}) \rightarrow M_n(\mathbb{C})$ are completely positive, possibly zero, and t stands for the transposition map of $M_2(\mathbb{C})$. As the structure of completely positive maps had been entirely understood thanks to the work of W. F. Stinespring [5] and M.-D. Choi [6], the theorem offered the full description of positive maps in this low-dimensional case. Hence, e.g. the structure of entanglement witnesses of two-qubit systems can be expressed concisely with the celebrated PPT criterion by A. Peres [7] and R., P., M. Horodecki [8].

In this paper, we present an alternative proof of the cited theorem. The premise of the proof is the identification of positive maps of $M_2(\mathbb{C})$ with operators preserving the four-dimensional Lorentz cone. We show that the decomposition of positive maps is rooted in the fact that every such operator is a convex combination of those additionally preserving the cone's topological boundary [9]. Thus, the proof of the theorem, contrary to the previous attempts based mainly on involved algebraic manipulation of matrices, provides a geometrical insight into the nature of the decomposition discussed above. It must be mentioned that a similar method of looking at positive maps and their application to the theory of quantum composite systems has been employed before (see e.g. [10]). In addition, we approach the conjecture stated by Robertson [11], who asks whether a positive extremal map that fulfils the Kadison-Schwarz inequality (1.1) is completely positive. With an additional assumption, we give the affirmative answer to this question for positive maps on matrix

^{*}marek.miller@ift.uni.wroc.pl

[†]robert.olkiewicz@ift.uni.wroc.pl

algebras. Last but not least, using that geometrical approach, we have been able to provide useful conditions for a positive map to be completely positive, as well as to describe the set of bistochastic maps that are simultaneously completely positive and completely copositive in terms of a norm-induced topology on the unit ball of the identified set of operators.

1 Preliminaries

Let $M_n = M_n(\mathbb{C})$ be the algebra of complex matrices of size n ; the algebra of real matrices will always be denoted explicitly by $M_n(\mathbb{R})$. We will use letters A, B, X , etc. to specify a matrix of M_n . The symbol $\mathbf{1}_n$, or simply $\mathbf{1}$, means the identity matrix of M_n . The norm of A , denoted by $\|A\|$, is understood as the operator norm of A as a linear map acting on \mathbb{C}^n . For $A \in M_n$, we denote its trace by $\text{Tr } A$; and by A^t and $A^* = \overline{A}^t$ its transpose and conjugate transpose, respectively. The commutator of two matrices A, B is defined as $[A, B] = AB - BA$. We say that a matrix A is positive-semidefinite, or simply positive, if $A = A^*$ and A has a non-negative spectrum.

A linear map $S: M_n \rightarrow M_n$ is said to be positive, indicated as $S \geq 0$, if for any $A \in M_n$ such that $A \geq 0$, we have $S(A) \geq 0$. The identity map of M_n is labelled I_n , or simply I . For a positive map S , its operator norm is given by $\|S\| = S(\mathbf{1})$. Any positive map is Hermitian, i.e. $S(X^*) = S(X)^*$, for all $X \in M_n$. We say that a positive map fulfils the Kadison-Schwarz inequality, if

$$S(X^*)S(X) \leq S(X^*X), \quad (1.1)$$

for any matrix $X \in M_n$. A positive map S that preserves the identity, i.e. $S(\mathbf{1}) = \mathbf{1}$, fulfils the Kadison-Schwarz inequality, but only for normal matrices X , that is the ones such that $XX^* = X^*X$ [12]. Following [11], we call a positive map that preserves the identity and fulfils the Kadison-Schwarz inequality – a Schwarz map. A map S is called bistochastic, if it is positive and preserves both the identity and trace, i.e. $S(\mathbf{1}) = \mathbf{1}$ and $\text{Tr } S(X) = \text{Tr } X$ for any $X \in M_n$. For $k \in \mathbb{N}$, a positive map S , such that the map $I_k \otimes S: M_k \otimes M_n \rightarrow M_k \otimes M_n$ is positive, is called k -positive. If a map is a k -positive map for every k , it is called completely positive. Similarly, a map is k -copositive, or completely copositive, if $I_k \otimes (S \circ t)$ is positive for some k , or for every k , respectively, where $t: A \mapsto A^t$, $A \in M_n$, is the transposition map. A positive map S is called decomposable, if it can be written in the form $S(A) = \Lambda_1(A) + \Lambda_2(A^t)$, where both maps Λ_1, Λ_2 are completely positive, possibly zero (see [6] for more details on completely positive maps). By an extremal positive map on M_n , we exclusively mean an extremal element of the cone of all positive maps on M_n , as defined below.

From now on, we focus on the algebra M_2 . We shall use the Greek indices μ, ν to denote the numbers $\mu, \nu = 0, 1, 2, 3$; whereas the Latin indices stand for $i, j = 1, 2, 3$. Let (σ_μ) , $\mu = 0, 1, 2, 3$; be a basis of M_2 such that $\sigma_0 = \frac{1}{\sqrt{2}}\mathbf{1}$ and σ_i , $i = 1, 2, 3$, are normalised Pauli matrices

$$\sigma_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.2)$$

A Hermitian element $\sigma(x)$ of M_2 , $\sigma(x) = \sum_{\mu=0}^3 x_\mu \sigma_\mu$, for $x = (x_\mu) = (x_0, \vec{x}) \in \mathbb{R}^4$, is positive if and only if $x \in L_4$, where $L_4 = \left\{ x \in \mathbb{R}^4 : x_0 \geq \sqrt{(x_1)^2 + (x_2)^2 + (x_3)^2} \right\}$. By $\vec{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$, we denote the 'spatial' part of the vector x , and by $\|\vec{x}\|$ – its Euclidean norm.

Let $S: M_2 \rightarrow M_2$ be a linear map. If S is Hermitian, then we can define a real matrix $\pi(S) \in M_4(\mathbb{R})$ by

$$\sigma(\pi(S)x) = S\sigma(x), \quad x \in \mathbb{R}^4. \quad (1.3)$$

Clearly, for two Hermitian maps S_1, S_2 , we have $\pi(S_1 S_2) = \pi(S_1) \pi(S_2)$ and $\pi(S^{-1}) = \pi(S)^{-1}$, whenever the map S^{-1} exists. The map S is positive, if and only if $\pi(S)$ maps L_4 into itself.

A set $\mathcal{P} \subset \mathbb{R}^n$ is a (convex) cone, if $\alpha x + \beta y \in \mathcal{P}$, whenever $x, y \in \mathcal{P}$, and $\alpha, \beta \geq 0$. An element $x \in \mathcal{P}$ is called extremal, if from the fact that $x - y \in \mathcal{P}$, for some vector $y \in \mathcal{P}$, follows that $y = \alpha x$,

$\alpha \geq 0$. We will denote by $\text{Ext } \mathcal{P}$ the set of extremal elements of the cone \mathcal{P} . The set of operators of $M_n(\mathbb{R})$ that map \mathcal{P} into itself will be denoted by $\Gamma(\mathcal{P})$. For any subset U of a linear space, by the convex hull of U we mean the set $\text{conv} U = \{tx + (1-t)y; x, y \in U, 0 \leq t \leq 1\}$ containing all convex combinations of the points in U . All the subsets of the set of positive maps specified above, i.e. the set of bistochastic, k -positive, k -copositive, completely positive, completely copositive and decomposable maps, are convex cones themselves.

We call the cone L_4 the Lorentz cone in \mathbb{R}^4 . The topological boundary ∂L_4 of L_4 is the set $\partial L_4 = \{x \in L_4 : x_0 = \|\vec{x}\|\}$. It is true that in this case $\partial L_4 = \text{Ext } L_4$. The symbol $\Theta(L_4) \subset \Gamma(L_4)$ will stand for the group of operators of $M_4(\mathbb{R})$ that map ∂L_4 onto itself. By [9, Theorem 2.4], for every $\tilde{A} \in \Theta(L_4)$, $\tilde{A} = r\tilde{O}$, where $r > 0$ and \tilde{O} belongs to the orthochronous Lorentz group $O^+(1, 3)$. Let $\rho : SL_2(\mathbb{C}) \rightarrow SO^+(1, 3)$ be the standard homomorphism (the spinor map) between the group of invertible complex matrices of M_2 with determinant equal to one and the *proper* orthochronous Lorentz group $SO^+(1, 3) \subset O^+(1, 3)$ (see e.g. [13] for a definition of the Lorentz group). By definition,

$$\sigma(\rho(V)x) = V^* \sigma(x) V, \quad (1.4)$$

for any $V \in SL_2(\mathbb{C})$ and $x \in \mathbb{R}^4$. Hence, for a map $S_V : M_2 \rightarrow M_2$, such that $S_V A = V^* A V$, $V \in SL_2(\mathbb{C})$, we have $\rho(V) = \pi(S_V)$. It is evident that every element of $O^+(1, 3)$, which does not belong to $SO^+(1, 3)$, can be written as a multiple ΛJ , where $\Lambda \in SO^+(1, 3)$ and J is a diagonal matrix, $J = \text{diag}(1, 1, -1, 1)$. (The particular form of the matrix J has been chosen only for the sake of convenience; the essential fact is that $J \in O^+(1, 3)$ and $\det J = -1$.)

2 Positive maps

We present the main result of the paper, which is a geometrical proof of the decomposition theorem for positive maps of M_2 .

Theorem 1. *Let $S : M_2 \rightarrow M_2$ be a positive map. Then $S = \Lambda_1 + \Lambda_2 \circ t$, where the maps $\Lambda_1, \Lambda_2 : M_2 \rightarrow M_2$ are completely positive and t stands for the transposition map of M_2 .*

Proof. The map S acts on the basis vectors σ_μ by $S\sigma_\mu = \sum_{\nu=0}^3 \pi(S)_{\mu\nu} \sigma_\nu$. Suppose at first that $\pi(S) \in \Theta(L_4)$. If $\pi(S) = \rho(V) \in SO^+(1, 3)$, $V \in SL_2(\mathbb{C})$, then $S\sigma(x) = \sum_{\mu=0}^3 x_\mu V^* \sigma_\mu V = V^* \sigma(x) V$, $x \in \mathbb{R}^4$, and the map S is completely positive, since every matrix $A \in M_2$ is a linear combination of Hermitian matrices. On the other hand, if $\pi(S) = \rho(V)J$, then

$$\begin{aligned} S\sigma(x) &= x_0 V^* V + x_1 V^* \sigma_1 V - x_2 V^* \sigma_2 V + x_3 V^* \sigma_3 V = \\ &= x_0 V^* V + x_1 V^* \sigma_1^t V + x_2 V^* \sigma_2^t V + x_3 V^* \sigma_3^t V = V^* \sigma(x)^t V, \end{aligned} \quad (2.1)$$

because $\sigma_2^t = -\sigma_2$. Hence, S is a composition of the transposition map and the one that is completely positive (i.e. S is completely copositive). Next, suppose that $\pi(S) \in \delta(L_4)$, where $\delta(L_4) = \{uw^t : u, w \in \partial L_4\}$, the set of rank-one operators that preserve ∂L_4 . It follows from [9], Lemma 3.2, that S is an extremal positive map such that $\text{rank} S = 1$, which implies that S must be completely positive.

If $\pi(S)$ is any matrix that preserves the Lorentz cone, we infer from [9], Theorem 4.5, that $\pi(S) \in \text{conv}(\Theta(L_4) \cup \delta(L_4))$. This deep geometrical result can be described by saying that every operator that preserves the Lorentz cone L_4 is a convex combinations of those operators that additionally preserve the boundary of the cone. By what has been said above, we are allowed to write

$$S = \Lambda_1 + \Lambda_2 \circ t, \quad (2.2)$$

where each Λ_1, Λ_2 is a completely positive map. \square

Remark 1. Theorem 1 provides an alternative proof of a known result [3, 4]. But whereas methods employed in those papers are based mainly on algebraic manipulations of matrices, the introduction of the map π has allowed us to consider geometrical properties of the relevant cone as a subset of \mathbb{R}^4 , in order to obtain the full characterisation of positive maps of M_2 . This characterisation could therefore be understood as having its source in the particular symmetry of the cone L_4 representing positive matrices of M_2 . In this simplest case, we were able to apply the deep result from [9], concerning the description of maps that preserve the cone L_4 , in hopes of transforming it back to the realm of linear maps of complex matrices. If we switch to the case of positive maps of M_3 , the cone of those vectors $x \in \mathbb{R}^9$ such that $\lambda(x) = \sum_{\mu=0}^8 x_\mu \lambda_\mu$ is a positive matrix of M_3 , where $(\lambda_\mu)_{\mu=0}^8$ is any linear space basis of M_3 , is no longer equal to the Lorentz cone K_9 . In fact, the geometry of the cone becomes highly non-trivial, rendering the analysis of operators that preserve it much more involved [14]. Moreover, since that cone is only a proper subset of K_9 , the Theorem 4.5 of [9] can no longer be applied.

A natural question that arises from the Theorem 1 is when a positive map of M_2 is completely positive. Checking that by means of the definition of complete positivity might be a tedious task, hence we propose the following result.

Proposition 1. *Let $S : M_2 \rightarrow M_2$ be a positive map preserving the identity. We consider the following conditions:*

1. *S is a Schwarz map and there exists a rank-one orthogonal projection P , for which $S(P)^2 = S(P)$.*
2. *There is a rank-one orthogonal projection P such that for every $X \in M_2$, $S([P, X]) = [S(P), S(X)]$.*
3. *S is completely positive.*

Then the following relation holds between the above conditions: $1 \Rightarrow 2 \Rightarrow 3$.

Proof. 1) \Rightarrow 2). Let $A_k = kP + iX$, for some $X = X^* \in M_2$ and $k \in \mathbb{N}$. Putting A_k into the Kadison-Schwarz inequality (1.1), we obtain

$$i[S(P), S(X)] - iS([P, X]) \leq \frac{1}{k} (S(X^2) - S(X)^2), \quad (2.3)$$

for every $k \in \mathbb{N}$. Hence

$$i[S(P), S(X)] - iS([P, X]) \leq 0. \quad (2.4)$$

By taking $A'_k = kP - iX$, and repeating essentially the same calculation, we get the reverse of (2.4), and thus

$$S([P, X]) = [S(P), S(X)], \quad (2.5)$$

for any Hermitian matrix X . Because (2.5) is linear in X , the equation holds for every matrix $X \in M_2$.

2) \Rightarrow 3). Let P be a rank-one orthogonal projection such that

$$S([P, X]) = [S(P), S(X)], \quad (2.6)$$

for every $X \in M_2$.

If S is a completely positive map, then for every unitary matrices $U, V \in M_2$ the map $\tilde{S} : A \mapsto U^* S(V^* A V) U$ is also completely positive. Therefore, we can assume without loss of generality that $P = P_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, and $S(P)$ is diagonal. Let $P_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$, $E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $E_{21} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Suppose that $S(P_1) = \lambda_1 P_1 + \lambda_2 P_2$, $0 \leq \lambda_2 \leq \lambda_1 \leq 1$. If $S(E_{12}) = 0$, then of course $S(E_{21}) = 0$ as well, and as a result, S maps M_2 into the Abelian algebra of diagonal matrices. Hence, S is completely positive. On the other hand, if $S(E_{12}) = \begin{pmatrix} w & z \\ x & y \end{pmatrix} \neq 0$, $w, z, x, y \in \mathbb{C}$, when we plug $X = E_{12}$ into (2.6), remembering that $\lambda_1 \geq \lambda_2$, we obtain that $x = y = w = 0$ and $\lambda_1 - \lambda_2 = 1$. It means that

$\lambda_1 = 1$ and $\lambda_2 = 0$, and $S(E_{12}) = zE_{12}$. Altogether, S acts on a matrix $X \in M_2$, $X = (x_{ij})_{i,j=1,2}$, by

$$S(X) = \begin{pmatrix} x_{11} & z x_{12} \\ \bar{z} x_{21} & x_{22} \end{pmatrix}. \quad (2.7)$$

Now, it is easy to see that S is completely positive. Indeed, it comes from positivity of S that $|z| \leq 1$. We write

$$S(X) = \begin{pmatrix} 1 & z \\ \bar{z} & 1 \end{pmatrix} \circ X, \quad (2.8)$$

where the symbol \circ in this case denotes the Hadamard product, i.e. the element-wise product of two matrices of the same dimension. Because $|z| \leq 1$, the matrix $\begin{pmatrix} 1 & z \\ \bar{z} & 1 \end{pmatrix}$ is positive. It is a known fact that a map $X \mapsto A \circ X$ is completely positive, if and only if A is a positive matrix (see e.g. Lemma 1 of [15] for a simple proof of this fact). \square

Example 1. One can easily show that the condition 1 of Proposition 1 is strictly stronger than the condition 2. Let S be a bistochastic map of M_2 such that

$$\pi(S) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad 0 < b < 1, \quad (2.9)$$

i.e. $S(\sigma_1) = b\sigma_1$ and $S(\sigma_2) = S(\sigma_3) = 0$. Then $S([P_0, X]) = [S(P_0), S(X)]$, where $P_0 = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$, and $X \in M_2$. However, there is no rank-one orthogonal projection P such that $S(P)^2 = S(P)$. Indeed, since $P_0 = \frac{1}{\sqrt{2}}(\sigma_0 + \sigma_1)$, $S(P_0) = \frac{1}{\sqrt{2}}(\sigma_0 + b\sigma_1)$. Let X be a Hermitian matrix, $X = \sigma(x)$, $x \in \mathbb{R}^4$, $S(X) = x_0\sigma_0 + bx_1\sigma_1$. Then clearly $[S(P_0), S(X)] = 0$. Moreover, $[P_0, \sigma(x)] = i(x_2\sigma_3 - x_3\sigma_2)$, hence $S([P_0, \sigma(x)]) = 0$. The condition 2 of Prop. 1 holds for any Hermitian X , therefore it holds also for any matrix $X \in M_2$. On the other hand, $P = \sigma(a)$, $a \in \mathbb{R}^4$, is a rank-one orthogonal projection, if and only if $a_0 = \|\vec{a}\| = \frac{1}{\sqrt{2}}$. Then $S(P) \neq 1$ and since $b < 1$, $S(P)$ cannot be a rank-one orthogonal projection, i.e. $S(P)^2 \neq S(P)$.

For extremal positive maps, the condition 1 of Proposition 1 could be proven sufficient for complete positivity in any dimension. At the same time, a more general problem posed by Robertson [11], whether there is a Schwarz map that is extremal as a positive map, but not 2-positive, remains open. It should be noted that from [16, Theorem 3.3] we know that every extremal 2-positive map is completely positive.

Theorem 2. *Let $S : M_n \rightarrow M_n$ be a Schwarz map that is extremal in the set of all positive maps. Suppose additionally, that there exist rank-one orthogonal projections $P, Q \in M_n$, for which $S(P) = Q$. Then S is completely positive.*

Proof. Let $(e_i)_{i=1}^n$ be the standard orthonormal basis in \mathbb{C}^n . Because we want to prove that S is completely positive, we can safely assume that $P = Q = P_n$, where $P_n = e_n e_n^*$ is the orthogonal projection on the one-dimensional space spanned by the vector $e_n = (0, 0, \dots, 0, 1) \in \mathbb{C}^n$. From the fact that S preserves the identity follows that $S(P_n^\perp) = P_n^\perp$, where $P_n^\perp = \mathbf{1}_n - P_n$. For a matrix $X \in M_n$, written in the block form $X = \begin{pmatrix} A & u \\ w^t & z \end{pmatrix}$, where $A \in M_{n-1}$, $u, w \in \mathbb{C}^{n-1}$ are column vectors, and $z \in \mathbb{C}$, we have

$$S(X) = \begin{pmatrix} \hat{S}_0(A) & S_1 u \\ (\bar{S}_1 w)^t & z \end{pmatrix}, \quad (2.10)$$

where $\hat{S}_0 : M_{n-1} \rightarrow M_{n-1}$ is necessarily a Schwarz map of M_{n-1} , and $S_1 \in M_{n-1}$. Indeed, since S is a Schwarz map, by [1, Proposition 2.1.5], $S(P_n X) = P_n S(X)$. Now, it follows that $S(P_n X P_n) = P_n S(X) P_n$, and similarly $S(P_n^\perp X P_n^\perp) = P_n^\perp S(X) P_n^\perp$, $S(P_n^\perp X P_n) = P_n^\perp S(X) P_n$, $S(P_n X P_n^\perp) = P_n S(X) P_n^\perp$, which shows that S must have the form (2.10).

If we take a one-dimensional orthogonal projection $P_u = uu^*$, where $u = (u_1, u_2, \dots, u_n) = (\vec{u}, u_n)$, $\vec{u} = (u_1, u_2, \dots, u_{n-1}) \in \mathbb{C}^{n-1}$, $u_n \in \mathbb{C}$, we have then

$$SP_u = S \begin{pmatrix} \vec{u}\vec{u}^* & \overline{u_n}\vec{u} \\ u_n\vec{u}^* & |u_n|^2 \end{pmatrix} = \begin{pmatrix} \hat{S}_0(\vec{u}\vec{u}^*) & \overline{u_n}S_1\vec{u} \\ u_n(S_1\vec{u})^* & |u_n|^2 \end{pmatrix}. \quad (2.11)$$

In the case when $u_n \neq 0$, taking the Schur complement (see [17, Theorem 1.12, p.34]), we have that $SP_u \geq 0$, if and only if

$$S_1\vec{u}\vec{u}^*S_1^* \leq \hat{S}_0(\vec{u}\vec{u}^*). \quad (2.12)$$

But that means that S is a sum of two positive maps that act on a one-dimensional orthogonal projection by:

$$SP_u = \begin{pmatrix} S_1\vec{u}\vec{u}^*S_1^* & \overline{u_n}S_1\vec{u} \\ u_n(S_1\vec{u})^* & |u_n|^2 \end{pmatrix} + \begin{pmatrix} \hat{S}_0(\vec{u}\vec{u}^*) - S_1\vec{u}\vec{u}^*S_1^* & 0 \\ 0 & 0 \end{pmatrix}. \quad (2.13)$$

Since S is extremal and $S(P_n) = P_n$,

$$SP_u = \begin{pmatrix} S_1\vec{u}\vec{u}^*S_1^* & \overline{u_n}S_1\vec{u} \\ u_n(S_1\vec{u})^* & |u_n|^2 \end{pmatrix} = \begin{pmatrix} S_1 & 0 \\ 0 & 1 \end{pmatrix} P_u \begin{pmatrix} S_1 & 0 \\ 0 & 1 \end{pmatrix}^*. \quad (2.14)$$

If we put $U = \begin{pmatrix} S_1 & 0 \\ 0 & 1 \end{pmatrix} \in M_n$, then by the fact that $S(P_n^\perp) = P_n^\perp$, we have that $UU^* = \mathbf{1}_n$, i.e. U is unitary, and $S(X) = UXU^*$, $X \in M_n$, which completes the proof. \square

Having presented geometrical arguments for the decomposition theorem of positive maps of M_2 in Theorem 1, we pass now to a proposition concerning bistochastic maps. We choose a tentative symbol Δ_∞ for the set of all bistochastic map of M_2 . Let also $\Delta_1 \subset \Delta_\infty$ denote the set of those bistochastic maps that are both completely positive and completely copositive at the same time. Using the same geometric picture of positive maps as operators preserving the Lorentz cone L_4 , we can show that the set Δ_1 is surprisingly large. It is worth mentioning that some related results could be found in [18].

Any matrix $Y \in M_n(\mathbb{R})$ admits the singular value decomposition: $Y = O_1 D O_2$, where O_1, O_2 are orthogonal matrices and $D = \text{diag}(t_1, t_2, \dots, t_n)$ is a diagonal matrix such that the *singular values* of Y are ordered: $t_1 \geq t_2 \geq \dots \geq t_n \geq 0$. The singular values of Y are the eigenvalues of the matrix $|Y| = (Y^t Y)^{1/2}$. In the space of operators $M_n(\mathbb{R})$, we distinguish two norms. The first one is the usual operator norm: $\|\cdot\|$; the second one is the trace norm given by $\|Y\|_1 = \sqrt{\text{Tr } Y^t Y}$. We denote the unit ball of $M_3(\mathbb{R})$ in the operator norm by: $K_\infty = \{Y \in M_3(\mathbb{R}) : \|Y\| \leq 1\}$; and the ball in the trace norm by: $K_1 = \{Y \in M_3(\mathbb{R}) : \|Y\|_1 \leq 1\}$.

Theorem 3. *There is a convex isomorphism F between the set of bistochastic maps of M_2 and the set K_∞ , $F(\Delta_\infty) = K_\infty$. Moreover, $F(\Delta_1) = K_1$.*

Proof. Let S be a bistochastic map. It is true that in that case

$$\pi(S) = \begin{pmatrix} 1 & \vec{0}^t \\ \vec{0} & Y \end{pmatrix}, \quad (2.15)$$

where $Y \in M_3(\mathbb{R})$ and $\vec{0} \in \mathbb{R}^3$ is a null column vector. We define $F(S) = Y$. From the definition (1.3) of the map π , it is evident that for $S_1, S_2 \in \Delta_\infty$, we have $F(\lambda S_1 + (1 - \lambda)S_2) = \lambda F(S_1) + (1 - \lambda)F(S_2)$, $0 \leq \lambda \leq 1$, and $F(S_1 S_2) = F(S_1)F(S_2)$. The map S is positive, if and only if $\pi(S)$ preserves the cone L_4 , which in this case is equivalent to the fact that $F(S) = Y \in K_\infty$. Moreover, it is easy to see that both F and F^{-1} are continuous mapping with respect to the topologies induced by the natural norms on Δ_∞ and K_∞ . Hence, $F(\Delta_\infty) = K_\infty$ and in addition, F is a homeomorphism.

Suppose now that $S \in \Delta_1$. Since S is completely positive, from [19, Theorem 1], we know that S is a convex combination of unitary maps: $X \mapsto U^* X U$, $U \in U(2)$. From the proof of Theorem 1,

for the particular form of $\pi(S)$ as in (2.15), it means that Y belongs to the convex hull of the group $\text{SO}(3)$. From [20], Corollary on p. 139, we infer that if $t_1 \geq t_2 \geq t_3 \geq 0$ are singular values of the matrix Y , then $Y \in \text{conv SO}(3)$, if and only if

$$t_1 \leq 1 \quad \text{and} \quad t_1 + t_2 - \text{sig}(\det Y) t_3 \leq 1. \quad (2.16)$$

S is also completely copositive, i.e. $S' = S \circ t$ is a completely positive bistochastic map. Let $F(S') = Y'$. Then $Y' = Y \text{diag}(1, -1, 1)$. Both matrices, Y and Y' , share the same singular values and $\det Y' = -\det Y$. Because $Y' \in \text{conv SO}(3)$, we have that $t_1 + t_2 + \text{sig}(\det Y) t_3 \leq 1$. Combining this with (2.16), since $\det Y = 0$, if and only if $t_3 = 0$, we obtain that the map S is both completely positive and completely copositive, if and only if $\|Y\|_1 = t_1 + t_2 + t_3 \leq 1$, which altogether means that $F(\Delta_1) = K_1$. \square

To summarise, we have presented a geometrical approach that proves itself useful in the investigation of properties of positive maps on M_2 . In particular, we have been able to show that the decomposition theorem for positive maps on M_2 is a direct consequence of the fact that every operator preserving the Lorentz cone is a convex combination of the ones that in addition preserve the boundary of the cone. These operators translate as completely positive and completely copositive maps, and therefore one is able to obtain an elegant and simple proof of the decomposition theorem that leads e.g. to the celebrated PPT criterion for separable states of two-qubit systems [7, 8]. The proof of Proposition 1, although it does not make use of this geometrical approach explicitly, is backed by the insight into the structure of the operators that preserve the Lorentz cone and have norm equal to one. Hence the assumption 1 in Proposition 1. Finally, Theorem 3 shows that the set of bistochastic maps that are completely positive and completely copositive is unexpectedly large and topologically equivalent to the unit ball of $M_3(\mathbb{R})$ with respect to the trace norm.

References

- [1] E. Størmer. *Positive Linear Maps of Operator Algebras*. Springer monographs in mathematics. Springer, 2012.
- [2] D. Chruściński and G. Sarbicki. Entanglement witnesses: construction, analysis and classification. *Journal of Physics A: Mathematical and Theoretical*, 47(48):483001, 2014.
- [3] E. Størmer. Positive linear maps of operator algebras. *Acta Mathematica*, 110(1):233–278, 1963.
- [4] S.L. Woronowicz. Positive maps of low dimensional matrix algebras. *Reports on Mathematical Physics*, 10(2):165–183, 1976.
- [5] W. F. Stinespring. Positive functions on C^* -algebras. *Proceedings of the American Mathematical Society*, 6(2):211–216, 1955.
- [6] M.-D. Choi. Completely positive linear maps on complex matrices. *Linear algebra and its applications*, 10(3):285–290, 1975.
- [7] A. Peres. Separability criterion for density matrices. *Physical Review Letters*, 77(8):1413–1415, 1996.
- [8] M. Horodecki, P. Horodecki, and R. Horodecki. Separability of mixed states: necessary and sufficient conditions. *Physics Letters A*, 223(1-2):1–8, 1996.
- [9] R. Loewy and H. Schneider. Positive operators on the n -dimensional ice cream cone. *Journal of Mathematical Analysis and Applications*, 49(2):375–392, 1975.
- [10] Jon Magne Leinaas, Jan Myrheim, and Eirik Ovrup. Geometrical aspects of entanglement. *Physical Review A*, 74(1):012313, 2006.

- [11] A. G. Robertson. Schwarz inequalities and the decomposition of positive maps on C^* -algebras. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 94, pages 291–296. Cambridge Univ Press, 1983.
- [12] M.-D. Choi. A Schwarz inequality for positive linear maps on C^* -algebras. *Illinois Journal of Mathematics*, 18(4):565–574, 1974.
- [13] G.L. Naber. *The Geometry of Minkowski Spacetime: An Introduction to the Mathematics of the Special Theory of Relativity*. Applied mathematics sciences. Dover, 1992.
- [14] K. Goyal, S. N. Simon, B. R. Singh, and S. Simon. Geometry of the generalized Bloch sphere for qutrit. *arXiv preprint arXiv:1111.4427*, 2011.
- [15] Adám Besenyei and Dénes Petz. Completely positive mappings and mean matrices. *Linear Algebra and its Applications*, 435(5):984–997, 2011.
- [16] M. Marciniak. On extremal positive maps acting between type I factors. *arXiv preprint arXiv:0812.2311*, 2008.
- [17] F. Zhang. *The Schur complement and its applications*. Springer, 2006.
- [18] Erling Størmer. A decomposition theorem for positive maps, and the projection onto a spin factor. *arXiv preprint arXiv:1308.3332*, 2013.
- [19] L. J. Landau and R. F. Streater. On Birkhoff’s theorem for doubly stochastic completely positive maps of matrix algebras. *Linear algebra and its applications*, 193:107–127, 1993.
- [20] F. Miranda, H. and R. C. Thompson. Group majorization, the convex hulls of sets of matrices, and the diagonal element-singular value inequalities. *Linear algebra and its applications*, 199:131–141, 1994.