

Quantum phase transition in the \mathbb{Z}_3 Kitaev-Potts model

Razieh Mohseninia,^{1,*} Saeed S. Jahromi,^{2,†} Laleh Memarzadeh,^{1,‡} and Vahid Karimipour^{1,§}

¹*Department of Physics, Sharif University of Technology, P.O. Box 11155-9161, Tehran, Iran.*

²*Department of Physics, K.N. Toosi University of Technology, P.O. Box 15875-4416, Tehran, Iran*

The stability of the topological order phase induced by the \mathbb{Z}_3 Kitaev model, which is a candidate for fault-tolerant quantum computation, against the local order phase induced by the 3-State Potts model is studied. We show that the low energy sector of the Kitaev-Potts model is mapped to the Potts model in the presence of transverse magnetic field. Our study relies on two high-order series expansion based on continuous unitary transformations in the limits of small- and large-Potts couplings as well as mean-field approximation. Our analysis reveals that the topological phase of the \mathbb{Z}_3 Kitaev model breaks down to the Potts model through a first order phase transition. We capture the phase transition by analysis of the ground state energy, one-quasiparticle gap and geometric measure of entanglement.

PACS numbers: 03.67.-a, 03.65.Vf, 05.30.Pr, 05.50.+q, 64.70.Tg.

I. INTRODUCTION

Quantum computers are much more powerful than the classical ones^{1,2}. However, a practical realization of such machines is still a big challenge ahead due to the fragility of the qubits and their decoherence arising from inevitable interactions with the environment. Over the years, several error correcting schemes have been developed³⁻⁵ to make the proper working of a quantum computer possible, despite such interactions. Unfortunately, error correcting methods are themselves error prone due to the imperfections of the very gate which are responsible for detecting and correcting errors. Even this can be taken into account by the clever techniques of fault-tolerant quantum computation⁶. Unfortunately the error threshold below which fault tolerant quantum computation is possible, is very low⁷. In order to overcome such problems, one can combine the main quantum feature of the quantum world, namely superposition of states, with the robustness of classical bits which is the result of a macroscopic number of very small entities, comprising each bit. This has led to the idea of topological quantum computation^{8,9}, where qubits are formed from degenerate ground states of a topologically ordered many body system. The important property that these degenerate ground states cannot be distinguished by local measurements, protects them against local errors arising from coupling with the environment, at least as long as temperature is sufficiently low (see below).

The prototype of such topological qubits is the toric code first introduced by Kitaev⁸. Since then other models have also been developed, notably the color code of Bombin and Martin-Delgado¹⁰. In these models, information is stored in the topologically degenerate ground states of the system and the computation is performed by braiding the quasiparticles (QP) of the model. Here one can use the gapped ground state and the robust nature of the topological phase of the model

to protect information against local errors. The only perturbations that cause logical error are those with length equal to the system size. It should be noted that this applies only to zero temperatures, since it has been shown in¹¹ and elaborated in¹²⁻¹⁴ that the proliferation of topological defects is capable of destroying topological order and hence quantum information encoded in such order at any finite temperature. Of course by tuning the parameters of such models¹² it is possible to increase the time scale for keeping the quantum information if the temperature is kept sufficiently low.

In his seminal paper, Kitaev showed that universal quantum computation is not possible with an Abelian group and in order to perform all universal gates of quantum computation, one needs to either resort to the non-Abelian case or use other procedures like magic state distillation^{15,16} or add some other non-topological resources like measurements to Abelian models as in¹⁷⁻²⁰. Aside from the ability of Abelian models to perform universal quantum computation in the above sense, their robustness and stability against external perturbations is still a crucial question which has to be investigated. Plenty of recent studies have been devoted for investigating such questions²¹⁻³¹. The motivation for such studies are not entirely based on quantum computation. In fact since topological order adds an entirely new paradigm for studying phase transition in condensed matter compared with the traditional one of symmetry breaking, it is very instructive to study various facets of this new phenomena. For example how topological order is destroyed under thermal fluctuation¹¹⁻¹³, how it gives way to local order in the presence of an external field like magnetic field²⁷ are just two such broad questions. Of particular interest to us is the question of how topological order gives way to local order in the absence of external field when a local ordering interaction is added. This question was first investigated in our previous work²¹, where we studied the competition of topological order and ferromagnetic order in a model which we called Kitaev-Ising model.

There it was shown that there is a critical coupling of Ising where topological order is replaced with ferromagnetic order. In that model we used a mean-field analysis and a simple calculation of Wilson-Loop operators to find the transition point.

In the context of condensed matter physics, it is important to understand the role of local degrees of freedom and their symmetry in topological order and its properties. It is mainly for such reasons, aside from the appeal of d -level states or qudits for quantum computation, that investigation of the above questions for general topological models for d -level states is important. For example it is very desirable to understand the competition of Z_d topological order and Z_d local ferromagnetic order. The latter is a phase possessed by the d -level Potts model where the Z_d symmetry is broken. The Potts model is a direct generalization of Ising model and has been extensively studied since its inception³⁷. While it is very desirable to perform such a study for general values of d , the analysis turns out to be very difficult. The 3-State Potts model is an exception in the sense that the Hamiltonian is much simpler, because it has an equidistant spectrum which brings much simplification in our analysis. It is therefore natural that Z_3 Potts model has been studied more intensively in the series of d -level Potts model, both in the classical context and in the new context in relation to topological order.

It has been shown that the Z_3 Kitaev model is more robust against temperature than the Z_2 one³⁰. The robustness of the model has also been studied in transverse magnetic field³¹ and it has been shown that the topological order transforms to the magnetic order via a first order phase transition. Here we want to extend these studies by studying the competition of Z_3 topological order induced by the Z_3 Kitaev model and the local order induced by the 3-State Potts model. We will see that the equidistant spectrum of the model leads to much technical simplification in the method we use, namely Perturbative Continuous Unitary Transformations (PCUT)^{32–35}. We will show that like the Kitaev-Ising model²¹, the Kitaev-Potts model also shows a first-order phase transition and the topological order break downs at a critical coupling of Potts. In contrast to the Kitaev-Ising model, here our study is more comprehensive in that we use three different methods for locating the transition point, namely we capture the phase transition by analysis of the ground state energy, one-quasiparticle gap and geometric measure of entanglement. The location of the transition turns out to be slightly different by these methods, and we provide arguments to show that with better approximations of these methods, this slight change will be removed.

The outline of the paper is as follows: In Sec. II and III, we briefly review the Z_d Kitaev and Potts models and

their essential features needed for our study. In Sec. IV, we show that the low energy sector of the Z_3 Kitaev-Potts model can be mapped to the 3-State Potts model in transverse magnetic field. We start our analysis of the mapped model in Sec. V by using mean field approximation and Perturbative Continuous Unitary Transformations method. By applying the PCUT method to the small Potts coupling limits, we calculate the one-quasiparticle (1-QP) gap and also the Geometric Measure of Entanglement. We further compute the ground-state energy in both small and large Potts couplings and capture the phase transition and the breakdown of the topological phase of the Kitaev model by analysis of the ground state energy and its derivatives, as well as the Geometric Measure of Entanglement and energy gap. Finally, Sec. VI is devoted to the conclusion.

II. Z_d KITAEV MODEL

Z_d Kitaev model is the generalization of the Kitaev model from Z_2 to Z_d group³⁶. The model can be defined on any two-dimensional lattice. In the present work, we consider the model on a square lattice on which the periodic boundary conditions are imposed on both sides, where the lattice becomes a torus. The lattice has N plaquettes, N vertices and $2N$ edges. The qudits live on the edges of the lattice. The Hamiltonian of the model consists of two kinds of operators, i.e. the plaquette and vertex operators. The operators are defined based on the generalized Pauli operators (acting on a qudit) as: $\sigma_x|j\rangle = |j+1 \bmod d\rangle$ and $\sigma_z|j\rangle = \omega^j|j\rangle$, $\omega = e^{2\pi i/d}$. These operators are not Hermitian and they don't square to I except for $d=2$. They further obey the following commutation relation: $\sigma_z\sigma_x = \omega\sigma_x\sigma_z$. The Hamiltonian of the model is given by the sum of the plaquette and vertex operators as:

$$H_{\text{Kitaev}} := -J \sum_s (A_s + A_s^\dagger) - K \sum_p (B_p + B_p^\dagger), \quad (1)$$

where s and p denote the stars (vertices) and plaquettes respectively. In order for the model to be exactly solvable, the A_s 's and B_p 's are defined such that they commute with each other. To this end an arbitrary direction is assigned to each edge and the A_s and B_p operators are defined as follows:

- $A_s := \prod_{i \in s} \sigma_{x,i}^{\pm 1}$, if the link's direction is inward, σ_x is applied, otherwise σ_x^{-1} is,
- $B_p := \prod_{i \in p} \sigma_{z,i}^{\pm 1}$, by starting from a link and moving counterclockwise, if each link's direction is the same as moving's direction σ_z is applied, otherwise σ_z^{-1} is. (Fig. 1)

These conventions lead to the commutativity of all A_s 's and B_p 's. Let us note that these arbitrary orientations do not have a physical significance, i.e. they lead to unitary




FIG. 1. (Color Online) The conventions for defining vertex and plaquette operators. $A_s := \prod_{i \in s} \sigma_{x,i}^{\pm 1}$, where the $+$ ($-$) corresponds to links with inward (outward) direction and $B_p := \prod_{i \in p} \sigma_{z,i}^{\pm 1}$, where the $+$ ($-$) corresponds to counter-clockwise (clockwise) direction of the links around plaquettes.

equivalent models. That is if we change the orientations on some links (L) in an arbitrary manner, we will end up with a model (\tilde{H}) which is iso-spectral with the original one (H). This unitary operator should transform σ_x to σ_x^\dagger and σ_z to σ_z^\dagger in the Hamiltonian:

$$\tilde{H} = U H U^\dagger, \quad U = \bigotimes_{i \in L} u_i, \quad (2)$$

where

$$u_i \sigma_{x,i} u_i^\dagger = \sigma_{x,i}^\dagger, \quad u_i \sigma_{z,i} u_i^\dagger = \sigma_{z,i}^\dagger, \quad (3)$$

It is straightforward to show that u is given by:

$$u = \sum_k |d-k\rangle \langle k|. \quad (4)$$

Note that there are $2N$ stabilizers in the Hamiltonian, but only $2N - 2$ of them are independent, because of the following two constraints on the torus:

$$\prod_s A_s = \prod_p B_p = I. \quad (5)$$

So there are d^2 degenerate ground states. The ground state is the state that is stabilized by all of the star and plaquette operators simultaneously and is equal to:

$$|\tilde{0}\tilde{0}\rangle := \prod_s (1 + A_s + A_s^2 + A_s^3 + \dots + A_s^{d-1}) |0\rangle^{\otimes 2N}. \quad (6)$$

In order to construct the other $d^2 - 1$ degenerate ground states, we define the following four string operators:

- $T_{z,1} = \prod_{i \in C_1} \sigma_{z,i}^{\pm 1}$, by starting from a link on the loop C_1 and moving on it, if each link's direction is the same as moving's direction σ_z is applied, otherwise σ_z^{-1} is,
- $T_{z,2} = \prod_{i \in C_2} \sigma_{z,i}^{\pm 1}$, by starting from a link on the loop C_2 and moving on it, if each link's direction is the same as moving's direction σ_z is applied, otherwise σ_z^{-1} is,




FIG. 2. (Color Online) Four nontrivial loops in a torus. C_1 and C_2 are in the real lattice, while \tilde{C}_1 and \tilde{C}_2 are in the dual lattice.

- $T_{x,1} = \prod_{i \in \tilde{C}_1} \sigma_{x,i}^{\pm 1}$, if the links's direction is the same as moving's direction on C_1 , σ_x is applied, otherwise σ_x^{-1} is,
- $T_{x,2} = \prod_{i \in \tilde{C}_2} \sigma_{x,i}^{\pm 1}$, if the links's direction is the same as moving's direction on C_2 , σ_x is applied, otherwise σ_x^{-1} is,

where C_1 , C_2 , \tilde{C}_1 and \tilde{C}_2 are shown in Fig. 2. All the degenerate ground states are given as:

$$|\tilde{i}\tilde{j}\rangle = T_{x,2}^j T_{x,1}^i |\tilde{0}\tilde{0}\rangle, \quad i, j = 0, 1, 2, \dots, d-1. \quad (7)$$

One can further check that:

$$T_{z,1} |\tilde{i}\tilde{j}\rangle = \omega^i |\tilde{i}\tilde{j}\rangle, \quad T_{z,2} |\tilde{i}\tilde{j}\rangle = \omega^j |\tilde{i}\tilde{j}\rangle. \quad (8)$$

III. POTTS MODEL

Considering any lattice of interest, the classical Potts model is defined by the following Hamiltonian³⁷:

$$H_{Potts} = - \sum_{i,j} \delta_{s_i, s_j}, \quad (9)$$

where the sum runs over the nearest neighbor sites of the lattice (i, j) and s_i takes d different values. One can take these values to be d different roots of unity. For $d=2$, $s_i = \{1, -1\}$, which reduces the Hamiltonian (9) to the renowned Ising model. This simplification is a consequence of the following definition for the delta function: $\delta_{s_i, s_j} = \frac{1}{2}(1 + s_i s_j)$. Such a definition can be extended to d -level and the analogue formula for the Potts model reads:

$$\delta_{s_i, s_j} = \frac{1}{2d} \sum_{r=0}^{d-1} ((s_i s_j^*)^r + (s_i^* s_j)^r), \quad (10)$$

where s_j^* is the complex conjugate of s_j . For general value of d , the model possesses Z_d symmetry $s_j \rightarrow \omega s_j$.

The quantum Potts model consists of d -level spins (qudits). The nearest neighbor spins interact with each other by the following Hamiltonian:

$$H_{Potts} = -\frac{1}{2d} \sum_{\langle i,j \rangle} \sum_{r=0}^{d-1} ((\sigma_{z,i} \sigma_{z,j}^\dagger)^r + (\sigma_{z,i}^\dagger \sigma_{z,j})^r). \quad (11)$$

The ground state of the system is the state where all spins are polarized in the same direction. So there are d degenerate ground states as follows:

$$|\tilde{i}\rangle := |i\rangle^{\otimes L}, \quad i = 0, 1, \dots, d-1, \quad (12)$$

where L is the number of spins.

IV. THE 3-STATE KITAEV-POTTS

Our aim is to study the phase transition of the Kitaev model in presence of the Potts interaction. As our series expansion technique, i.e. the PCUT method is only applicable to those models with equidistant spectrum, we restrict our study to the $d = 3$ or qutrits and show that Z_3 Kitaev and 3-State Potts models have equidistant spectrum. The full Hamiltonian of the \mathbb{Z}_3 Kitaev model perturbed by the Potts interaction is given by:

$$\begin{aligned} H &= H_{Kitaev} + \lambda H_{Potts} \\ &= -J \sum_s (A_s + A_s^\dagger) - K \sum_p (B_p + B_p^\dagger) \\ &\quad - \frac{\lambda}{6} \sum_{\langle i,j \rangle} \sum_{r=0}^2 ((\sigma_{z,i} \sigma_{z,j}^\dagger)^r + (\sigma_{z,i}^\dagger \sigma_{z,j})^r). \end{aligned} \quad (13)$$

where λ is the perturbation parameter and is a measure of the strength of the Potts interaction. The perturbed Kitaev Hamiltonian is no longer exactly solvable. This is due to the fact that the σ_z operators in the Potts model do not commute with vertex operators of the Kitaev model. However, the plaquette operators still commute with the full Hamiltonian (13). The ground state of the Hamiltonian (13) is therefore in the sector in which $B_p = 1$ for all plaquette operators. In this sector the Kitaev-Potts's Hamiltonian reduces to the following form:

$$\begin{aligned} H &= -J \sum_s (A_s + A_s^\dagger) \\ &\quad - \frac{\lambda}{6} \sum_{\langle i,j \rangle} \sum_{r=0}^2 ((\sigma_{z,i} \sigma_{z,j}^\dagger)^r + (\sigma_{z,i}^\dagger \sigma_{z,j})^r) - 2KN. \end{aligned} \quad (14)$$

In order to tackle the Hamiltonian (14), we first define the following new basis and rewrite the full Hamiltonian in this new basis:

$$|\mathbf{r}\rangle = |r_1, r_2, \dots, r_N\rangle := \prod_i A_i^{r_i} |0\rangle^{\otimes 2N}, \quad r_i = 0, 1, 2, \quad (15)$$




FIG. 3. (Color Online) In the mapped Kitaev-Potts Hamiltonian, the original lattice (the lattices with $\sqrt{N} * \sqrt{N}$ Plaquettes, wherein \sqrt{N} is even) is de-coupled into two sublattices shown with blue dots and red dashes.

where i runs over all the vertices of the lattice. This transformation was first defined in²¹ for the the Kitaev-Ising model. One can readily check that $B_p|\mathbf{r}\rangle = |\mathbf{r}\rangle$. We can therefore interpret Eq. (15) as the basis for the sector in which $B_p = 1$ for all p . Next, we determine the action of the different terms in the Hamiltonian (14) on this new basis:

$$A_s|\mathbf{r}\rangle = A_s \prod_i A_i^{r_i} |0\rangle^{\otimes 2N} = |r_1, r_2, \dots, r_s + 1, \dots, r_N\rangle, \quad (16)$$

so A_s in this basis acts like generalized Pauli operator, σ_x , which we denote by \hat{X} . Clearly, the action of A_s^\dagger can be regarded as \hat{X}^\dagger . In order to recast the Potts interaction in the new basis, we attach an arrow to each link. The direction of the arrows are illustrated in Fig. 2. The action of a Potts interaction term like $\sigma_{z,a} \sigma_{z,b}^\dagger$ in the new basis, is denoted by:

$$\begin{aligned} \sigma_{z,a} \sigma_{z,b}^\dagger |\mathbf{r}\rangle &= \sigma_{z,a} \sigma_{z,b}^\dagger A_1^{r_1} A_2^{r_2} A_3^{r_3} \prod_{i \neq 1,2,3} A_i^{r_i} |0\rangle^{\otimes 2N} \\ &= \omega^{r_1} \omega^{-r_3} |\mathbf{r}\rangle, \end{aligned} \quad (17)$$

which means that the Potts interaction term, $\sigma_{z,a} \sigma_{z,b}^\dagger$, commutes with all of the vertex operators except A_1 and A_3 , actually it acts like $\hat{Z}_1 \hat{Z}_3^\dagger$ in this new basis:

$$\sigma_{z,a} \sigma_{z,b}^\dagger \equiv \hat{Z}_1 \hat{Z}_3^\dagger. \quad (18)$$

It is straightforward to check that the following relations further hold in the new basis:

$$\begin{aligned} \sigma_{z,a}^\dagger \sigma_{z,b} &\equiv \hat{Z}_1^\dagger \hat{Z}_3, \\ \sigma_{z,c}^\dagger \sigma_{z,d} &\equiv \hat{Z}_1 \hat{Z}_3^\dagger, \\ \sigma_{z,c} \sigma_{z,d}^\dagger &\equiv \hat{Z}_1^\dagger \hat{Z}_3. \end{aligned} \quad (19)$$

Therefore the \mathbb{Z}_3 Kitaev-Potts model in the new basis is given by:

$$\tilde{H} = \tilde{H}_A + \tilde{H}_B - 2KN, \quad (20)$$

where

$$\tilde{H}_A = -J \sum_{i \in A} (\hat{X}_i + \hat{X}_i^\dagger) - \frac{\lambda}{3} \sum_{\langle i, j \rangle \in A} \sum_{r=0}^2 (\hat{Z}_i \hat{Z}_j^\dagger)^r + (\hat{Z}_j \hat{Z}_i^\dagger)^r, \quad (21)$$

and

$$\tilde{H}_B = -J \sum_{i \in B} (\hat{X}_i + \hat{X}_i^\dagger) - \frac{\lambda}{3} \sum_{\langle i, j \rangle \in B} \sum_{r=0}^2 (\hat{Z}_i \hat{Z}_j^\dagger)^r + (\hat{Z}_j \hat{Z}_i^\dagger)^r. \quad (22)$$

The Hamiltonian (20) is nothing but the sum of two Potts models in a transverse magnetic field, wherein the Potts interactions act on nearest neighbor vertices in the two de-coupled sublattices shown in Fig. 3.

The two de-coupled Hamiltonians are exactly the same. In the following sections we present our results for \tilde{H}_A . The results can be extended to the full Hamiltonian (20), without loss of generality. From now on to the end of the paper by H , we mean \tilde{H}_A , unless stated otherwise.

V. METHODS

In this section, we present the solution to the mapped model by applying the Mean field approximation and Perturbative Continuous Unitary Transformations method to the Hamiltonian (21).

A. Mean Field approximation

Suppose we are interested in finding the ground state energy of a given Hamiltonian. We therefore need to minimize $\langle \Psi | H | \Psi \rangle$ over all $|\Psi\rangle$'s in the Hilbert space. One approximation (Mean field approximation) would be that we do the minimization only over product states. But the Hamiltonian has translational symmetry so we can search for the minimum energy only over states of the form $|\Psi\rangle = |\Phi\rangle^{\otimes n}$, which have translational symmetry. For the mapped Hamiltonian n is the number of vertices in each sublattice, $n = \frac{N}{2}$.

$$E = \min_{|\Phi\rangle}^{\otimes n} \langle \Phi | H | \Phi \rangle^{\otimes n}, \quad (23)$$

where $|\Phi\rangle$ is a general one-qutrit state. For the mapped Hamiltonian,

$$H = -J \sum_i (\hat{X}_i + \hat{X}_i^\dagger) - \frac{\lambda}{3} \sum_{\langle i, j \rangle} \sum_{r=0}^2 (\hat{Z}_i \hat{Z}_j^\dagger)^r + (\hat{Z}_j \hat{Z}_i^\dagger)^r,$$

we have:

$$E_0 = -Jn(\langle \hat{X} \rangle + \langle \hat{X}^\dagger \rangle) - 4n\frac{\lambda}{3} \sum_{r=0}^2 (\langle \hat{Z}^r \rangle \langle \hat{Z}^{-r} \rangle). \quad (24)$$




FIG. 4. (Color Online) Ground state energy per site, ε_0 , as a function of x . The inset represents its first derivative. The sharp jump in the derivation of energy at $x_c \approx 0.115$ signals the first order quantum phase transition.

Taking $|\Phi\rangle = \sum_{i=0}^2 a_i |i\rangle$, we should minimize:

$$E_0 = -Jn \left(\sum_i (a_i a_{i+1}^* + a_i^* a_{i+1}) \right) - 4n\lambda \left(\sum_i |a_i|^4 \right). \quad (25)$$

It's easy to check that the above equation recasts into:

$$E_0 = -Jn(|a_0 + a_1 + a_2|^2 - 1) - 4n\lambda(|a_0|^4 + |a_1|^4 + |a_2|^4), \quad (26)$$

that has a permutation symmetry by changing the a_i 's indices from $(0, 1, 2)$ to $(\sigma(1), \sigma(2), \sigma(3))$. If the state that minimize this expression respects the full symmetry, it should be equal to $|\Phi\rangle = \frac{1}{\sqrt{3}}(|0\rangle + |1\rangle + |2\rangle)$ and the ground state energy per site is equal to:

$$\varepsilon_0 = -2J - \frac{4}{3}\lambda. \quad (27)$$

Setting $J = 1$, we see that Eq. (27) is analytic for any value of λ and one does not observe any phase transition. However, if some of the symmetries are broken and the ground state remains invariant under only one permutation, it will be in the form of $|\Phi\rangle = a_0|0\rangle + a_1|1\rangle + a_1|2\rangle$. By setting $a_0 = \sin \theta$, $a_1 = \frac{1}{\sqrt{2}} e^{i\alpha} \cos \theta$, the ground state energy per site reduces to:

$$\varepsilon_0 = -J(\cos^2 \theta + \sqrt{2} \sin 2\theta \cos \alpha) - 4\lambda(\sin^4 \theta + \frac{1}{2} \cos^4 \theta). \quad (28)$$

By minimizing ε_0 , the ground state as a function of perturbation parameter $x = \frac{2\lambda}{9J}$ (this explicit form of x is chosen because it makes the comparison between mean-field and PCUT results easier) is given as follows:

$$|\Phi\rangle = \begin{cases} \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) & \text{for } x < 0.115 \\ \sin \theta_x |0\rangle + \frac{\cos \theta_x}{\sqrt{2}} |1\rangle + \frac{\cos \theta_x}{\sqrt{2}} |2\rangle & \text{for } x > 0.115 \end{cases} \quad (29)$$

where,

$$x = \frac{\sin 2\theta_x - 2\sqrt{2} \cos 2\theta_x}{36 \sin 2\theta_x (\sin^2 \theta_x - \frac{1}{2} \cos^2 \theta_x)}. \quad (30)$$

The above relation shows that the nature of the ground state changes at a critical point $x_c = 0.115$, which can be a signal of the phase transition. Finally we analyze the case where the symmetry is fully broken and the ground state has the general form as $|\Phi\rangle = \sum_{i=0}^2 a_i |i\rangle$. In this case, we minimize the energy numerically. Figure 4 demonstrates the mean-field ground state energy per site as a function of $x = \frac{2\lambda}{9J}$. As we can see, there is a small kink in the ground state energy curve (a sharp jump in its derivative) at $x_c \approx 0.115$. So the results show that the ground state actually respects some permutation symmetries. This jump signals a first order quantum phase transition. In order to investigate the critical point more accurately, we resort to a more accurate approximation technique, i.e. the Perturbative Continuous Unitary Transformation method³²⁻³⁵ which is the subject of the next subsection.

B. Perturbative Continuous Unitary Transformations

In this section, we briefly review the Perturbative Continuous Unitary Transformation (PCUT) method and apply it to the small- and large-coupling limits of the problem.

The Continuous Unitary Transformation (CUT) method which was first introduced by Wegner³² in the framework of condensed matter theory, is basically used to diagonalize or block-diagonalize a given Hamiltonian by applying an infinite number of unitary operators to the initial Hamiltonian in a continuous fashion as:

$$H(\ell) = U^\dagger(\ell) H U(\ell), \quad (31)$$

where ℓ is the continuous flow parameter such that $H = H(\ell = 0)$ and $H_{\text{eff}} = H(\ell = \infty)$ is the (block-) diagonal Hamiltonian. The Hamiltonian is transformed by a unitary operator which its evolution is governed by:

$$\partial_\ell U(\ell) = -U(\ell) \eta(\ell). \quad (32)$$

In which $\eta(\ell)$ is the anti-Hermitian generator of the unitary transformation $U(\ell)$. Combining Eq. (31) and (32) together one can show that the initial Hamiltonian flows in the form of a differential commutator equation :

$$\partial_\ell H(\ell) = [\eta(\ell), H(\ell)]. \quad (33)$$

The method therefore requires the choice of a suitable generator for the unitary operators to obtain the desirable form of H_{eff} . Uhrig and Knetter introduced the quasiparticle (QP) conserving generator which is very

well suited for our purpose. We refer the interested reader to Refs^{33,34} for detailed discussions on QP conserving generator.

The perturbative version of the CUT method (PCUT), can be applied to the Hamiltonians of the form $H = Q + \lambda V$ where the first part of the Hamiltonian, Q , is diagonal with an equidistant spectrum bounded from below and the second part can be treated as a perturbation (λ is the expansion parameter). The method further requires that the perturbing part can be written in the form $V = \sum_{n=-N}^N T_n$, where T_n increments (decrements, if $n < 0$) the number of excitations (quasiparticles) by n such that $[Q, T_n] = n T_n$ ³³. Transforming the initial problem by using the QP conserving generator, the effective Hamiltonian is brought to the form that conserves the number of quasiparticles, $[H_{\text{eff}}, Q] = 0$. The energy spectrum of the system can therefore be extracted perturbatively by acting the H_{eff} on the ground state and the multi-particle sectors of the Hilbert space.

1. Small-coupling limit ($\lambda \ll J$)

In the following we discuss the procedure of applying the PCUT method to the small-coupling limit ($\lambda \ll J$) of the mapped Kitaev-Potts model, i.e. the Potts model in transverse magnetic field on the A or B sublattices of Fig. 3:

$$H = -J \sum_{i \in A} (\hat{X}_i + \hat{X}_i^\dagger) - \frac{\lambda}{3} \sum_{\langle i, j \rangle \in A} \sum_{r=0}^2 \left((\hat{Z}_i \hat{Z}_j^\dagger)^r + (\hat{Z}_j \hat{Z}_i^\dagger)^r \right). \quad (34)$$

The first term in Eq. (34) is an effective field term which is diagonal. Denoting the local vacuum of each site by $|0\rangle$, the elementary excitations of the model for $d = 3$ are two separate spin flips labeled by $|1\rangle$ and $|2\rangle$ which correspond to the eigenstates of X operator with ω and ω^{-1} eigenvalues, respectively. Let us note that either of the excitations cost an energy of $3J$. The elementary excitations are energetically indistinguishable. So the first term for $d = 3$ has an equidistant spectrum and can be regarded as Q for implementation of the PCUT, and the PCUT results obtained in 1-QP sector of the Hilbert space such as 1-QP gap are degenerate for both of the excitations.

The Potts interaction at the right side of Eq. (34) can be treated as a perturbation V which for $d = 3$ is denoted by:

$$V = -\frac{2\lambda}{3} \sum_{\langle i, j \rangle} \left((\hat{Z}_i \hat{Z}_j^\dagger) + (\hat{Z}_j \hat{Z}_i^\dagger) \right) + C, \quad (35)$$

where C is a constant. The perturbing part consists of two-body interactions and can change the number of excitations over the ground state of the effective field term

Q by $n = \{0, \pm 1, \pm 2\}$ when it acts on the bonds of the square lattice. Therefore the Hamiltonian (34) can be written as:

$$H = Q - x(T_2 + T_1 + T_0 + T_{-1} + T_{-2}), \quad (36)$$

where

$$Q = \sum_i \frac{-(\hat{X}_i + \hat{X}_i^\dagger) + 2I}{3}, \quad (37)$$

is the quasiparticle counting operator (I is the identity operator), $x = \frac{2\lambda}{9J}$ is the expansion parameter and T_n operators are given by:

$$\begin{aligned} T_{+2} &= \sum_{\langle i,j \rangle} |12\rangle_{i,j} \langle 00| + |21\rangle_{i,j} \langle 00|, \\ T_{+1} &= \sum_{\langle i,j \rangle} |22\rangle_{i,j} \langle 01| + |22\rangle_{i,j} \langle 10| \\ &\quad + |11\rangle_{i,j} \langle 02| + |11\rangle_{i,j} \langle 20|, \\ T_0 &= \sum_{\langle i,j \rangle} |12\rangle_{i,j} \langle 21| + |01\rangle_{i,j} \langle 10| + |02\rangle_{i,j} \langle 20| \\ &\quad + |21\rangle_{i,j} \langle 12| + |10\rangle_{i,j} \langle 01| + |20\rangle_{i,j} \langle 02|. \end{aligned} \quad (38)$$

From the hermiticity condition $T_n^\dagger = T_{-n}$, and absorbing the expansion parameter $-x$ in the definition of T_n operators, Eq. (36) is recast into:

$$H = Q + \sum_{n=-2}^{n=2} T_n. \quad (39)$$

Under the CUTs, the above Hamiltonian is continuously transformed with the flow parameter ℓ as:

$$H(\ell) = Q + \sum_{n=-2}^{n=2} T_n(\ell), \quad (40)$$

where we wish to reach a situation where $T_n(\ell = \infty) = 0$ for all $n \neq 0$. In order to fulfil this demand, we choose the quasiparticle conserving generator in the following form^{33,34}:

$$\eta(\ell) = T_{+2}(\ell) + T_{+1}(\ell) - T_{-1}(\ell) - T_{-2}(\ell). \quad (41)$$

With this choice of generator, the flow Eq. (33) can be written as:

$$\begin{aligned} \partial_\ell T_0(\ell) &= 2[T_{+2}(\ell), T_{-2}(\ell)] + 2[T_{+1}(\ell), T_{-1}(\ell)], \\ \partial_\ell T_{+2}(\ell) &= -2T_{+2}(\ell) + [T_{+2}(\ell), T_0(\ell)], \\ \partial_\ell T_{+1}(\ell) &= -T_{+1}(\ell) + 2[T_{+2}(\ell), T_{-1}(\ell)] + [T_{+1}(\ell), T_0(\ell)]. \end{aligned} \quad (42)$$

Let us stress that the Hamiltonian remains hermitian under the unitary transformation. We can therefore calculate $T_{-2}(\ell)$ and $T_{-1}(\ell)$ from the hermiticity condition. Solving the flow equation is still a very cumbersome task because there are an infinite number of terms in $T_n(\ell)$.




FIG. 5. (Color Online) 1-QP dispersion of the Kitaev-Potts model in the small coupling regime. The minimum of the dispersion is located at $K=(0,0)$.

However, we can tackle the problem by performing a perturbative expansion of the flow equation. We can therefore introduce the expansion of the T_n operators as³⁵:

$$T_n(\ell) = \sum_{i=1}^{\infty} T_n^{(i)}(\ell), \quad (43)$$

where i is the order of perturbation. Using this relation, the perturbative expansion of the flow equation is written as:

$$\begin{aligned} \partial_\ell T_0^{(k)}(\ell) &= 2 \sum_{j=1}^{k-1} [T_{+2}^{(j)}(\ell), T_{-2}^{(k-j)}(\ell)] + 2 \sum_{j=1}^{k-1} [T_{+1}^{(j)}(\ell), T_{-1}^{(k-j)}(\ell)], \\ \partial_\ell T_{+2}^{(k)}(\ell) &= -2T_{+2}^{(k)}(\ell) + \sum_{j=1}^{k-1} [T_{+2}^{(j)}(\ell), T_0^{(k-j)}(\ell)], \\ \partial_\ell T_{+1}^{(k)}(\ell) &= -T_{+1}^{(k)}(\ell) + 2 \sum_{j=1}^{k-1} [T_{+2}^{(j)}(\ell), T_{-1}^{(k-j)}(\ell)] \\ &\quad + \sum_{j=1}^{k-1} [T_{+1}^{(j)}(\ell), T_0^{(k-j)}(\ell)]. \end{aligned} \quad (44)$$

Solving this set of equations for the initial condition $T_n^{(i)}(\ell = 0) = \delta_{1,i} T_n$ and then taking the limit of $(\ell \rightarrow \infty)$, we can obtain H_{eff} . The effective QP conserving Hamiltonian for the small-coupling limit up to order 3 in perturbation parameter is obtained as:

$$\begin{aligned} H_{\text{eff}}^{(3)} &= Q - xT_0 + x^2[T_1, T_{-1}] + \frac{x^2}{2}[T_2, T_{-2}] \\ &\quad - \frac{x^3}{8} \left([T_2, [T_0, T_{-2}]] + [[T_2, T_0], T_{-2}] \right) \\ &\quad - \frac{x^3}{2} \left([T_1, [T_1, T_{-2}]] + [[T_2, T_{-1}], T_{-1}] \right) \\ &\quad - \frac{x^3}{2} \left([T_1, [T_0, T_{-1}]] + [[T_1, T_0], T_{-1}] \right). \end{aligned} \quad (45)$$




FIG. 6. (Color Online) The 1-QP gap of the Kitaev-Potts model for different DlogPadé approximants. The critical point highlighted by gray dashed line is consistent with the closure of the gap at $x_c \approx 0.129$. The inset demonstrates the scaling on the closure of the bare series of the gap for different orders of perturbation.

The ground state energy and 1-QP gap can be obtained by acting the effective hamiltonian on 0P and 1P sector of the Q. We have calculated the ground state energy and 1-QP energy gap of the system in the small-coupling limit up to order 8 in perturbation theory:

$$\epsilon_0^{\text{sc}} = -\frac{2}{3} - 2x^2 - x^3 - \frac{17x^4}{2} - \frac{847x^5}{36} - \frac{18407x^6}{144} - \frac{15290\lambda^7}{27} - \frac{995278817\lambda^8}{311040}, \quad (46)$$

$$\Delta^{\text{sc}} = 1 - 4x - 10x^2 - 5x^3 - \frac{1895x^4}{6} + \frac{14107x^5}{18} - \frac{3572759x^6}{216} + \frac{26566267x^7}{324} - \frac{85919559673x^8}{77760}. \quad (47)$$

Fig. 6 illustrates the 1-QP gap of the system in the small-coupling limit as a function of $x = \frac{2\lambda}{J}$. The bare and extrapolated series are well converged. Closure of the 1-QP gap occurs at $x_c \approx 0.129$. The inset of Fig. 6 further demonstrates scaling of the closure of the bare series of the gap for different orders of perturbation. As one can see, by increasing the order of perturbation the transformation point starts to become smaller until it converges to $x_c \approx 0.12$ which is fully consistent with the closure of DlogPadé extrapolants.

Existence of anyonic excitations in the system is one of the signatures of the topological order. As one can see from Fig. 6, the anyonic gap is open until in the vicinity of the transition point. It is therefore reasonable to point out that the topological order survives during the tuning of the perturbation until the critical point at which the anyons condenses due to the strength of the Potts interaction and the topological order breaks down^{38,39}.

Let us further note that the knowledge of the gap is not solely sufficient to determine the first- or second-order nature of the phase transition^{38,39} and one has to




FIG. 7. (Color Online) The ground state energy per site ϵ_0 as a function of θ . Solid line corresponds to the bare series of order 8. Symbols denote different Padé approximants.

analyze the ground state energy of the system and its derivatives⁴⁰ to capture the phase transition correctly. We therefore postpone further discussion on the phase transition to Sec. VB3 after we calculate the ground state energy series in large-coupling limits.

2. Large-coupling limit ($\lambda \gg J$)

Following our discussion, we now apply the PCUT method to the large-coupling limit of the problem ($\lambda \gg J$). For $J = 0$, the Hamiltonian is 3-State Potts model which is ferromagnetically ordered and has an equidistant spectrum thus satisfying the first condition for the PCUT method. The excitations of the model further correspond to the anti-ferromagnetic bonds of the square lattice. When $J \neq 0$, the effective field term in Eq. (34) can be considered as a perturbation which changes the number of excitations by $n = \{0, \pm 1, \pm 2, \pm 3, \pm 4\}$. We can therefore write the Hamiltonian of the model in the large-coupling limit in terms of T_n operators as:

$$H = Q - h \sum_{n=-4}^4 T_n, \quad (48)$$

where Q is the quasiparticle counting operator defined as:

$$Q = \frac{2I - \sum_{\langle i,j \rangle} (\hat{Z}_i \hat{Z}_j^\dagger) + (\hat{Z}_j \hat{Z}_i^\dagger)}{3}, \quad (49)$$

and $h = J/2\lambda$ is the expansion parameter. Using the PCUT method, we have calculated the ground state energy per site, ϵ_0 , in the large-coupling limit up to order 8 in perturbation parameter by acting the H_{eff} on the




FIG. 8. (Color Online) First derivative of the ground state energy per site ($\partial_\lambda \varepsilon_0$) as a function of θ . Solid line corresponds to the bare series of order 8. Symbols denote different Padé approximants.

0-QP sector of the Hilbert space:

$$\epsilon_0^{\text{lc}} = -\frac{8}{3} - \frac{h^2}{2} - \frac{h^3}{8} - \frac{19h^4}{672} - \frac{3h^5}{128} - \frac{12779h^6}{846720} - \frac{1052987h^7}{121927680} - \frac{458808396457h^8}{62768369664000}. \quad (50)$$

As we have already mentioned in the previous section, one can obtain a better understanding about the nature of the phase transition by analysis of the ground state energy results. We will therefore provide strong evidences for the first-order phase transition in the \mathbb{Z}_3 Kitaev-Potts model in the next section.

3. Analyzing series expansion results

To investigate the nature of the phase transition, we study the ground state energy per site of the system in the whole range of the expansion parameter by merging the small- and large-coupling results. Setting $\lambda = \sin \theta$ and $J = \cos \theta$, we can join the small- and large-coupling results to obtain a complete picture. Fig. 7 demonstrates the ground state energy per site ε_0 as a function of θ . The small- and large-coupling series cross each other at $\theta_c \approx 0.52$, giving rise to a kink in the ε_0 curve which is fully consistent with the first-order phase transition. The location of the kink is essentially the same for different Padé extrapolants and has strong agreement with the closure of the 1-QP gap ($x_c = \frac{2}{9} \tan \theta_c \approx 0.129$).

Using the Feynman-Hellman theorem, we have calculated the first derivative of ε_0 which is equivalent to the magnetization in statistical mechanics. Figs. 8 and 9 depict first and second derivatives of ε_0 for different θ values. The sharp jumps in the first derivative is a clear signature of the first-order phase transition⁴⁰. Let us further note that the first order nature of the




FIG. 9. (Color Online) Second derivative of the ground state energy per site ($\partial_\lambda^2 \varepsilon_0$) as a function of θ . Solid line corresponds to the bare series of order 8. Symbols denote different Padé approximants.

phase transition in the \mathbb{Z}_3 Potts model in transverse magnetic field has already been confirmed using series expansion combined with infinite projected entangled-pair state (IPEPS) method in Ref.³¹. The jump in the second derivative further confirms the location of the transition point which is exactly the same as that of the ground-state energy for the bare series and different Padé extrapolants.

C. Geometric Measure of Entanglement

In this section we calculate a measure of multipartite entanglement, Geometric Measure of Entanglement (GME)⁴¹, as another tool for capturing the phase transition. GME is a measure of multipartite entanglement in quantum many body systems which measures the distance, Hilbert Schmidt distance, between a given state, $|\Psi(x)\rangle$, and the closest product state, $|P\rangle$, as follows:

$$\text{GME} = -\log_2 \left(\max_{|P\rangle} |\langle P | \Psi(x) \rangle|^2 \right), \quad (51)$$

where the maximization is over all product states. This implies that the more entangled the states are, more distance from the set of product states, which is a convex set, they will have. In order to calculate the GME, we have to calculate the ground state of the mapped Hamiltonian, $|\Psi(x)\rangle$, by PCUT procedure up to a specific order of perturbation. The PCUT method transforms the initial Hamiltonian by a unitary transformation ($U(\infty)$) to a block-diagonal form (H_{eff}), the basis of the Hilbert space is rotated, such that H_{eff} commutes with Q . So they have the same eigenstates. It's also known that H_{eff} 's ground state is the vacuum state of Q with no excitations ($|\bar{0}\rangle$)⁴². The H_{eff} and the initial Hamiltonian, H , are unitary equivalent and their ground states are

related to each other as follows:

$$|\Psi(x)\rangle = U(\infty)|\tilde{0}\rangle. \quad (52)$$

In order to determine $U(\ell)$, we expand Eq. (32) as a function of the perturbation parameter x :

$$\partial_\ell U^{(k)}(\ell) = - \sum_{j=0}^{k-1} U^{(j)}(\ell) \eta^{(k-j)}(\ell), \quad (53)$$

where

$$\eta^{(k-j)}(\ell) = T_{+2}^{(k-j)}(\ell) + T_{+1}^{(k-j)}(\ell) - T_{-1}^{(k-j)}(\ell) - T_{-2}^{(k-j)}(\ell), \quad (54)$$

and k is the order of perturbation. By using Eqs. (44), and the initial condition $U^k(0) = 1\delta_{k,0}$, one can solve the above equation perturbatively up to second order in x . Taking the limit $\ell \rightarrow \infty$, the $U(\infty)$ is found to be:

$$\begin{aligned} U = & 1 + \frac{x}{2}(T_{+2} - T_{-2}) + x(T_{+1} - T_{-1}) \\ & + \frac{x^2}{2}(T_{+1} - T_{-1})^2 + \frac{x^2}{8}(T_{+2} - T_{-2})^2 \\ & + \frac{x^2}{4}[T_0, (T_{+2} + T_{-2})] + x^2[T_0, (T_{+1} + T_{-1})] \\ & + \frac{x^2}{6}(T_{+1}T_{+2} + T_{-1}T_{-2} + 2T_{+2}T_{+1} + 2T_{-2}T_{-1}) \\ & + \frac{x^2}{2}([T_1, T_{-2}] + [T_{-1}, T_2] - T_{-2}T_1 - T_2T_{-1}). \end{aligned} \quad (55)$$

The ground state of the initial Hamiltonian H is therefore given by:

$$\begin{aligned} |\Psi(x)\rangle = U|\tilde{0}\rangle = & (1 - \frac{4nx^2}{8})|\tilde{0}\rangle + \frac{x}{2} \sum_{4n} |1 \ 2\rangle \\ & + \frac{x^2}{4} \left(\sum_{4n} |1 \ 2\rangle + 2 \sum_{8n} \left| \begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix} \right\rangle + 2 \sum_{4n} |1 \ 0 \ 2\rangle \right) \\ & + \frac{x^2}{6} \left(2 \sum_{4n} \left| \begin{smallmatrix} 2 & 2 \\ 2 & 0 \end{smallmatrix} \right\rangle + 2 \sum_{4n} \left| \begin{smallmatrix} 1 & 1 \\ 1 & 0 \end{smallmatrix} \right\rangle \right. \\ & \left. + 2 \sum_{2n} |2 \ 2 \ 2\rangle + 2 \sum_{2n} |1 \ 1 \ 1\rangle \right) \\ & + \frac{x^2}{8} \left(\sum_{8n(2n-7)} \left| \begin{smallmatrix} 1 & 2 \\ 1 & 2 \end{smallmatrix} \right\rangle \right). \end{aligned} \quad (56)$$

This state is a superposition of some product states, these product states are eigenstates of Q . The first term, $|\tilde{0}\rangle$ refers to the ground state of Q with no excitations, which is the state of all spins in the \hat{X} eigenstate with $+1$ eigenvalue, $|0\rangle$. $|1 \ 2\rangle$ is the state of all the spins in $|0\rangle$ state, except for the two of them, which are nearest neighbour, and have the states of $|1\rangle$ and $|2\rangle$. The summation must be done over all the states of this kind, and one should note that in a lattice with n vertices the number of this kind of states are 2 times the number of bonds,




FIG. 10. (Color Online) GME as a function of x for $n=25$. Convexity of GME changes close to the $x_c \approx 0.16$. The inset further demonstrates the derivative of GME as a function of x . The sharp jump in the figure is a clear signature of phase transition.

i.e. $2 \times 2n = 4n$. Furthermore, $\left| \begin{smallmatrix} 1 & 0 \\ 0 & 2 \end{smallmatrix} \right\rangle$ and $|1 \ 0 \ 2\rangle$ denote the state with all spins in the $|0\rangle$ state, except for the two of them, which are next nearest neighbours and have the states of $|1\rangle$ and $|2\rangle$. The last term, $\left| \begin{smallmatrix} 1 & 2 \\ 1 & 2 \end{smallmatrix} \right\rangle$, also refers to the state with all spins in $|0\rangle$ state except for the four of them. The state of these four spins consist of two clusters of $|1 \ 2\rangle$, where these two clusters can be separate or not.

$|\Psi(x)\rangle$ is exact up to the second order in perturbation parameter x and is normalized. We have also calculated the ground state of the initial Hamiltonian up to the fourth order, numerically and calculated the GME. Since the ground state has translational symmetry, the closest product state also preserves this symmetry⁴³. We can therefore perform the maximization only over the states in the form $|P\rangle = |\phi\rangle^{\otimes n}$, where $|\phi\rangle = \cos(\theta)|0\rangle + e^{-i\alpha} \sin(\theta) \sin(\varphi)|1\rangle + e^{-i\beta} \sin(\theta) \cos(\varphi)|2\rangle$. In other word, the maximization is only over 4 parameters which makes the numerical calculation of the GME possible. Fig. 10 illustrates the geometric measure of entanglement as a function of x . As we can see, the convexity of the GME changes sign from positive to negative close to a critical point, $x_c \approx 0.16$. The sharp jump in the derivative of GME captures the phase transition more clearly, since it's known that for two-dimensional systems discontinuity in the derivative of multipartite entanglement leads to a quantum phase transition⁴⁴. The critical point is very close to the one obtained from analysis of the ground state energy and the gap i.e. $x_c \approx 0.129$. The slight difference is likely the consequence of the difference in orders of perturbation. This can be best inferred by reminding the scaling behavior of the closure of 1-QP gap (inset of Fig. 6) which predicts that increasing the order

of perturbation makes the result to become smaller and converge to ≈ 0.12 . The same fact should also holds here for the location of sharp jump in the derivative of the GME.

VI. CONCLUSION

The \mathbb{Z}_d Kitaev model is a system with a topologically ordered ground state which, when complemented with some other resources like magic state distillation^{15,16} or measurements²⁰, is well suited for the purpose of universal quantum computation without resorting to non-Abelian groups. It is therefore of great interest to study the stability and robustness of the topological phase of the model in the presence of external perturbations. In this paper, we studied the competition between the topological order induced by the \mathbb{Z}_3 Kitaev model and the local order induced by the 3-State Potts model on the square lattice. We showed that the Kitaev model in the presence of the Potts interaction

is mapped to Potts model in a transverse magnetic field. Using the high-order series expansion based on the continuous unitary transformations in the small- and large- Potts couplings, we showed that the topological phase breaks down to a non-topological phase with local order parameter through a first-order quantum phase transition at $x_c = \frac{2\lambda_c}{9J_c} \approx 0.129$. Our results were further in good agreement with the mean-field approximation results at $x_c \approx 0.115$. Computing the Geometric Measure of Entanglement also shows that the derivative of GME has a sharp jump very close to the critical point and also, the convexity of GME changes sign from positive to negative in this point.

VII. ACKNOWLEDGEMENTS

R.M. acknowledges R. Haghshenas for fruitful discussions.

* mohseninia@physics.sharif.ir

† s.jahromi@dena.kntu.ac.ir

‡ memarzadeh@sharif.edu

§ vahid@sharif.edu

- ¹ Shor, P. W. "Polynomial-Time Algorithms for Prime Factorization and Discrete Logarithms on a Quantum Computer". SIAM J.Sci.Statist.Comput., 26, 1997. [1](#)
- ² Nielsen, M. A. and Chuang, I. L. "Quantum Computation and Quantum Information". 2000. [1](#)
- ³ Shor, P. W. "Scheme for reducing decoherence in quantum computer memory". Physical review A, 52(4), R2493, (1995). [1](#)
- ⁴ Steane, A. M. "Error correcting codes in quantum theory". Physical Review Letters, 77(5), 793, (1996).
- ⁵ Gottesman, D. "A Class of Quantum Error-Correcting Codes Saturating the Quantum Hamming Bound". Phys.Rev. A54 (1996) 1862. [1](#)
- ⁶ Gottesman, D. "A Theory of Fault-Tolerant Quantum Computation". Phys. Rev. A57, 127 (1998). [1](#)
- ⁷ Preskill, J. lecture notes in Quantum Computation, <http://www.theory.caltech.edu/people/preskill/ph229/notes/chap7.pdf>. [1](#)
- ⁸ Kitaev, A. Y. "Fault-tolerant quantum computation by anyons". Annals of Physics, 303(1), 2-30, (2003). [1](#)
- ⁹ Dennis, E., Kitaev, A., Landahl, A., Preskill, J. "Topological quantum memory". Journal of Mathematical Physics, 43(9), 4452-4505, (2002). [1](#)
- ¹⁰ Bombin, H., Martin-Delgado, M. A. "Topological quantum distillation". Physical review letters, 97(18), 180501, (2006). [1](#)
- ¹¹ Nussinov, Z. and Ortiz, G. "A symmetry principle for Topological Quantum Order", Cond-mat/0702377; "Symmetry and Topological Order". Cond-mat/0605316. [1](#)
- ¹² Nussinov, Z. and G. Ortiz, G. "Autocorrelations and thermal fragility of anyonic loops in topologically quantum ordered systems". Phys. Rev. B 77, 064302 (2008). [1](#)

- ¹³ Castelnovo, C. and Chamon, C. "The Toric-Boson model and quantum memory at finite temperature". Phys. Rev. B 79, 245122 (2009). [1](#)
- ¹⁴ Alicki, R., Fannes, M. and Horodecki, M. "On thermal stability of topological qubit in Kitaev's 4D model". J. Phys. A 40, 6451 (2007). [1](#)
- ¹⁵ Bravyi, S. and Kitaev, A. "Universal quantum computation with ideal clifford gates and noisy ancillas". Phys. Rev. A, 71(2):022316, 2005. [1](#), [11](#)
- ¹⁶ Campbell, E. T., Anwar, H., Browne, D. E. "Magic-state distillation in all prime dimensions using quantum reed-muller codes". Physical Review X, 2(4), 041021, (2012). [1](#), [11](#)
- ¹⁷ Lloyd, S. "Quantum computation with Abelian anyons. Quantum Information Processing", 1(1-2), 13-18, (2002) [1](#)
- ¹⁸ Pachos, J. K. "Quantum computation with Abelian anyons on the honeycomb lattice". International Journal of Quantum Information, 4(06), 947-954, (2006).
- ¹⁹ Raussendorf, R., Harrington, J., Goyal, K. "Topological fault-tolerance in cluster state quantum computation". New Journal of Physics, 9(6), 199, (2007).
- ²⁰ Wootton, J. R., Pachos, J. K. "Universal Quantum Computation with Abelian Anyon Models". Electronic Notes in Theoretical Computer Science, 270(2), 209-218, (2011). [1](#), [11](#)
- ²¹ Karimipour, V., Memarzadeh, L., Zarkeshian, P. "Kitaev-Ising model and the transition between topological and ferromagnetic order". Physical Review A, 87(3), 032322, (2013). [1](#), [2](#), [4](#)
- ²² Zarei, M. H. "On the robustness of topological quantum codes: Ising perturbation". arXiv preprint arXiv:1501.07619,(2015).
- ²³ Trebst, S., Werner, P., Troyer, M., Shtengel, K., Nayak, C. "Breakdown of a topological phase: Quantum phase transition in a loop gas model with tension". Physical review letters, 98(7), 070602, (2007).

- ²⁴ Hamma, A., Cincio, L., Santra, S., Zanardi, P., Amico, L. "Local response of topological order to an external perturbation". *Physical review letters*, 110(21), 210602, (2013).
- ²⁵ Halsz, G. B., Hamma, A. "Probing topological order with Rnyi entropy". *Physical Review A*, 86(6), 062330, (2012).
- ²⁶ Hamma, A., Lidar, D. A. "Adiabatic preparation of topological order". *Physical review letters*, 100(3), 030502, (2008).
- ²⁷ Vidal, J., Dusuel, S., Schmidt, K. P. "Low-energy effective theory of the toric code model in a parallel magnetic field". *Physical Review B*, 79(3), 033109, (2009). [1](#)
- ²⁸ Tupitsyn, I. S., Kitaev, A., Prokofev, N. V., Stamp, P. C. E. "Topological multicritical point in the phase diagram of the toric code model and three-dimensional lattice gauge Higgs model". *Physical Review B*, 82(8), 085114, (2010).
- ²⁹ Castelnovo, C., Trebst, S., Troyer, M. "Topological Order and Quantum Criticality". *arXiv preprint arXiv:0912.3272*. (2009)
- ³⁰ Viyuela, O. Rivas, A., Martin-Delgado, M. A. "Generalized toric codes coupled to thermal baths". *New Journal of Physics*, 14(3), 033044, (2012). [2](#)
- ³¹ Schulz, M. D., Dusuel, S., Ors, R., Vidal, J., Schmidt, K. P. "Breakdown of a perturbed Z_N topological phase". *New Journal of Physics*, 14(2), 025005, (2012). [1](#), [2](#), [9](#)
- ³² Franz Wegner, *Ann. Physik* **3** 77-91 (1994). [2](#), [6](#)
- ³³ C. Knetter and G.S. Uhrig, *Eur. Phys. J. B* **13** 209-225 (2000). [6](#), [7](#)
- ³⁴ C. Knetter, K.P. Schmidt, and G.S. Uhrig, *J. Phys. A* **36**, 7889 (2003). [6](#), [7](#)
- ³⁵ S. Dusuel, M. Kamfor, K. P. Schmidt, R. Thomale and J. Vidal, *Phys. Rev. B* **81** 064412 (2010). [2](#), [6](#), [7](#)
- ³⁶ Bullock, S. S., Brennen, G. K. "Qudit surface codes and gauge theory with finite cyclic groups". *Journal of Physics A: Mathematical and Theoretical*, 40(13), 3481,(2007). [2](#)
- ³⁷ Lieb, E. , Shultz, T. and Mattis, D. "Two soluble models of an antiferromagnetic chain", *Annals of Physics* Volume 16, Issue 3, December 1961, Pages 407-466. [2](#), [3](#)
- ³⁸ S. S. Jahromi, M. Kargarian, S. F. Masoudi, and K. P. Schmidt, *Phys. Rev. B*. **87**, 094413 (2013). [8](#)
- ³⁹ S. S. Jahromi, S. F. Masoudi, M. Kargarian, K.P. Schmidt, *Phys. Rev. B*. 88, 214411 (2013) [8](#)
- ⁴⁰ S. Sachdev, "Quantum Phase Transitions", Cambridge University Press, Cambridge, (2011). [8](#), [9](#)
- ⁴¹ Wei, T. C., Goldbart, P. M. "Geometric measure of entanglement and applications to bipartite and multipartite quantum states". *Physical Review A*, 68(4), 042307, (2003). [9](#)
- ⁴² Christian Knetter, "Perturbative continuous unitary transformations: spectral properties of low dimensional spin systems", phd thesis, Universität zu Köln, (2003). [9](#)
- ⁴³ Hbener, R., Kleinmann, M., Wei, T. C., Gonzalez-Guilln, C., Ghne, O. "Geometric measure of entanglement for symmetric states". *Physical Review A*, 80(3), 032324, (2009). [10](#)
- ⁴⁴ Huang, C. Y., Lin, F. L. "Multipartite entanglement measures and quantum criticality from matrix and tensor product states". *Physical Review A*, 81(3), 032304, (2010). [10](#)