

Maximal entanglement concentration for $(n + 1)$ -qubit states

Anindita Banerjee ^{a *}

Chitra Shukla ^b

Anirban Pathak ^b

November 5, 2018

^aDepartment of Physics and Center for Astroparticle Physics and Space Science,
Bose Institute, Block EN, Sector V, Kolkata 700091, India.

^bDepartment of Physics and Materials Science and Engineering,
Jaypee Institute of Information Technology, A-10, Sector-62, Noida, UP-201307, India.

Abstract

We propose two schemes for concentration of $(n + 1)$ -qubit entangled states that can be written in the form of $(\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$ where $|\varphi_0\rangle$ and $|\varphi_1\rangle$ are mutually orthogonal n -qubit states. The importance of this general form is that the entangled states like Bell, cat, GHZ, GHZ-like, $|\Omega\rangle$, $|Q_5\rangle$, 4-qubit cluster states and specific states from the 9 SLOCC-nonequivalent families of 4-qubit entangled states can be expressed in this form. The proposed entanglement concentration protocol is based on the local operations and classical communications (LOCC). It is shown that the maximum success probability for ECP using quantum nondemolition (QND) technique is $2\beta^2$ for $(n + 1)$ -qubit states of the prescribed form. It is shown that the proposed schemes can be implemented optically. Further it is also noted that the proposed schemes can be implemented using quantum dot and microcavity systems.

1 Introduction

Entanglement is one of the most important resources of quantum information processing. It plays a key role in quantum computation and communication. Specifically, maximally entangled states are often a prerequisite for quantum information processing. As entanglement between two or more distant parties cannot be created by local operations and classical communication (LOCC), it is required to be distributed. Usually a maximally entangled pure state is distributed among different parties by quantum channels and due to the presence of noise it transforms to either a mixed state or a less entangled pure state. Subsequently, maximally entangled state (MES) is extracted from an ensemble of mixed states or an ensemble of less entangled pure states. Entanglement concentration is a process in which one can extract maximally entangled state from nonmaximally (less entangled state) using LOCC with fidelity equal to 1. Similarly, the process of distilling a set of mixed states into a maximally entangled state is referred to as the entanglement purification.

Entanglement concentration protocol (ECP) was proposed by Bennett et al. [1] in 1996. This pioneering protocol was based on the Schmidt projection method. Several ECPs were proposed thereafter using different methods. For example, entanglement-swapping [2], qubit-assisted [3], POVM [4] and Bell-measurement [5] based ECPs were proposed. Initial proposals for ECPs were restricted to the concentration of nonmaximally entangled Bell states. However, with the recent advances in applications of multipartite entangled states in quantum communication and quantum computation, it has become extremely interesting to design ECPs for multipartite entangled states. For example, in the recent past, ECPs for various multipartite entangled states like GHZ and cat state [6], GHZ-like state [7], cluster state [8, 9, 10], arbitrary W state [11], 4-qubit entangled state [7], etc. have been proposed. Nevertheless, apart from using different methods for designing ECPs, the possibility of implementation of ECPs is also studied for different technologies, like linear optics [12, 13, 14, 15, 16], quantum dot and microcavity systems [17, 18, 19, 20] and electronic technologies [21, 22]. Yamamoto et al. [14] experimentally realized ECP from two copies of the state $|\Phi\rangle = \alpha|00\rangle + \beta|11\rangle$, where $|\alpha|^2 + |\beta|^2 = 1$ using polarising beam splitters (PBSs), wave plates and photon detectors. The maximum probability achieved in these cases for single pair concentration is $2\alpha^2\beta^2$. Another system which is used in ECP to obtain higher success probability is the cross-Kerr-nonlinearity [11, 12].

A qubit-assisted ECP for Bell state was initially proposed in [3] using CNOT gate and Von Neumann measurement. There, it was shown that using iterative process maximal success probability up to $2\beta^2$ can be obtained for Bell state. Recently, Sheng et al. [12] have proposed a qubit-assisted concentration protocol for Bell state using PBS. They have shown that it is possible to obtain a higher success probability by iteratively using cross-Kerr-nonlinearity. Various qubit-assisted entanglement concentration protocols [23, 24, 25, 26, 27, 28, 29, 30] for different entangled states using different systems have been reported thereafter, referring to the higher success probability mentioned in [12]. However, upper bound on the success

*email: aninditabanerjee.physics@gmail.com

Table 1: Entangled states and their corresponding less entangled counter parts expressed in the general form described by Eq. (1).

$(n + 1)$ -qubit state	Maximally entangled state	Non-maximally entangled state
2-qubit Bell	$\frac{ 00\rangle+ 11\rangle}{\sqrt{2}}$	$(\alpha 0\rangle 0\rangle + \beta 1\rangle 1\rangle)_{(1+1)}$
$(2 + 1)$ -qubit GHZ	$\frac{ 000\rangle+ 111\rangle}{\sqrt{2}}$	$(\alpha 00\rangle 0\rangle + \beta 11\rangle 1\rangle)_{(2+1)}$
$(n + 1)$ -qubit cat	$\frac{ 0\dots 00\rangle+ 1\dots 11\rangle}{\sqrt{2}}$	$(\alpha 0\dots 0\rangle 0\rangle + \beta 1\dots 1\rangle 1\rangle)_{(n+1)}$
$(2 + 1)$ -qubit GHZ-like	$\frac{ \psi^+0\rangle+ \phi^+1\rangle}{\sqrt{2}}$	$(\alpha \psi^+\rangle 0\rangle + \beta \phi^+\rangle 1\rangle)_{(2+1)}$
Q_5 state	$\frac{ 0000\rangle+ 1011\rangle+ 1101\rangle+ 1110\rangle}{2}$	$(\alpha(000\rangle + 111\rangle) 0\rangle + \beta(101\rangle + 110\rangle) 1\rangle)_{(3+1)}$
Cluster state	$\frac{ 0000\rangle+ 0011\rangle+ 1100\rangle- 1111\rangle}{2}$	$(\alpha(000\rangle + 110\rangle) 0\rangle + \beta(001\rangle - 111\rangle) 1\rangle)_{(3+1)}$

probability was not discussed in these works. In what follows, we propose two ECPs for all $(n + 1)$ -qubit entangled states that can be written in the form

$$(\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}, \quad (1)$$

where $|\varphi_0\rangle$ and $|\varphi_1\rangle$ are mutually orthogonal n -qubit states and $\alpha^2 + \beta^2 = 1 : \alpha \neq \frac{1}{\sqrt{2}}$. Both the ECPs proposed here may be viewed as qubit-assisted ECPs as each of them is assisted by an ancillary qubit having the same Schmidt coefficients as that of the less entangled state to be concentrated. Further, we show that one of the proposed ECP which utilizes QND techniques achieves the maximum possible success probability (i.e., $2\beta^2$). To justify the importance of the states of the form (1), in Table 1, we have listed different entangled states which can be written in the general form described in Eq. (1). Further, it may be noted that the states of this particular form have vast applications in secure quantum communication, quantum secret sharing (QSS), bidirectional quantum teleportation and hierarchical quantum communication schemes [31, 32] e.g., hierarchical quantum information splitting (HQIS), probabilistic HQIS and hierarchical quantum secret sharing (HQSS). The general nature and applicability of quantum states of the form $(\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$ have motivated us to construct ECPs for states of this form. Recently, we [7] have proposed a Bell state-assisted ECP for the states of this particular form. Here we aim to propose two more ECPs for the same states and to improve the efficiency. The qubit-assisted ECP provides higher efficiency as shown in [7] where efficiency of Bell state-assisted are compared with qubit-assisted ECP. Moreover, the latter requires less resource compared to the former. Further, if we use nonlinear resources (i.e., if we use QND technique or equivalently if we use cross-Kerr nonlinearity) then the efficiency approaches allowed upper bound.

Rest of the paper is organized as follows: In Section 2, we have presented our scheme. In Section 3, we have discussed that the entanglement is conserved during ECP and finally the paper is concluded in Section 4.

2 Maximal entanglement concentration

We have proposed the ECP for $(n + 1)$ -qubit state of the form

$$|\psi\rangle = (\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}.$$

The protocol can be implemented on all states shown in Table 1. The protocol is described in Subsection 2.1 and the corresponding quantum circuit is shown in Fig. 1. The purpose of presenting the quantum circuit is that the protocol can be realized in any technology. We propose concentration protocols using two independent methods: (i) post selection principle using linear optics and (ii) quantum nondemolition using cross-Kerr-nonlinearity as described in Subsection 2.2 and 2.3, respectively. Many concentration protocols have been realized using linear optics as already stated in Section 1. Linear optics has two important features first is post selection and second is requirement of sophisticated single photon detectors. Post selection destroys the photons and they cannot be further used. The cross-Kerr-nonlinearity is widely studied in the context of the CNOT gate [33], Bell state analysis [34], etc. It is a tool to construct quantum nondemolition detectors (QND) having the capability for conditioning the evolution of the system without necessarily destroying the state of the photon. It acts as parity check and single photon detector. Moreover, it is worthwhile to mention that the protocol here is implemented by linear optics, but can be used in other technologies as well for example in electronic systems like electronic polarization beam splitter and the charge detection [10], quantum dot spins [35], etc. The quantum nondemolition technique implemented using these electronic technologies have higher efficiency but the optimal value of success probability has not been attained in their works. In this work we show that by employing QND technique we can attain an optimal value concentration which is showed in Section 2.3.

2.1 Quantum circuit

Let Alice and Bob be two spatially separated parties. They share N number of $n + 1$ pure entangled states such that Bob possesses first n qubits and Alice possesses $(n + 1)^{th}$ -qubit. The entangled state is of the form $|\psi\rangle = (\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$




Figure 1: Proposed qubit-assisted maximal ECP for $(n+1)$ -qubit state $\alpha|\psi_0\rangle|0\rangle + \beta|\psi_1\rangle|1\rangle$.

where $|\varphi_0\rangle$ and $|\varphi_1\rangle$ are mutually orthogonal n -qubit states. The values of α and β are real such that $\alpha^2 + \beta^2 = 1$ and $\alpha \neq \beta$. Alice also possess one more qubit $|\psi_s\rangle = \alpha|0\rangle + \beta|1\rangle$ which has the same Schmidt coefficients as that of the non-maximally entangled state shared by Alice and Bob. Therefore, Alice possesses two qubits which are $(n+1)^{th}$ qubit and $(n+2)^{th}$ qubit.

Input state of the circuit shown in Fig. 1 is

$$\begin{aligned}
|\psi_1\rangle &= |\psi\rangle \otimes |\psi_s\rangle \\
&= (\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{1,2,\dots,n+1} \otimes (\alpha|0\rangle + \beta|1\rangle)_{n+2} \\
&= (\alpha^2|\varphi_0\rangle|00\rangle + \alpha\beta|\varphi_1\rangle|10\rangle \\
&\quad + \alpha\beta|\varphi_0\rangle|01\rangle + \beta^2|\varphi_1\rangle|11\rangle)_{1,2,\dots,n+1,n+2}.
\end{aligned} \tag{2}$$

Alice applies a CNOT gate on her qubits by using $(n+1)^{th}$ qubit as the control qubit and the $(n+2)^{th}$ qubit as the target qubit. As a consequence of this operation input state $|\psi_1\rangle$ transforms to

$$\begin{aligned}
|\psi_2\rangle &= (\alpha^2|\varphi_0\rangle|00\rangle + \alpha\beta|\varphi_1\rangle|11\rangle \\
&\quad + \alpha\beta|\varphi_0\rangle|01\rangle + \beta^2|\varphi_1\rangle|10\rangle)_{1,2,\dots,n+1,n+2} \\
&= [(\alpha^2|\varphi_0\rangle|0\rangle + \beta^2|\varphi_1\rangle|1\rangle)|0\rangle \\
&\quad + \alpha\beta(|\varphi_0\rangle|0\rangle + |\varphi_1\rangle|1\rangle)|1\rangle]_{1,2,\dots,n+1,n+2}.
\end{aligned} \tag{3}$$

Alice measures $(n+2)^{th}$ qubit of $|\psi_2\rangle$ in the computational $\{|0\rangle, |1\rangle\}$ basis. From (3), we can easily observe that if Alice's measurement yields $|1\rangle$ then the quantum state shared by Alice and Bob collapses to a normalized maximally entangled state $\frac{|\varphi_0\rangle|0\rangle + |\varphi_1\rangle|1\rangle}{\sqrt{2}}$, but the circuit (equivalently an ECP represented by the circuit) fails when Alice's measurement yields $|0\rangle$. Clearly, the success probability of the ECP is $2\alpha^2\beta^2$ and we are now left with a less entangled pure state with probability $1 - 2\alpha^2\beta^2$ i.e., $\alpha^4 + \beta^4$. We can repeat the ECP using the remaining less entangled pure states to maximize the number of maximally entangled states. The maximum probability that can be achieved by repeating this procedure is $2\beta^2$ which is provided later in Subsection 2.3.

2.2 Entanglement concentration using linear optics

In optical implementation of an ECP, we consider that qubits are realized using polarization states of the photon where horizontal (H) and vertical (V) photon represent the logical bits 0 and 1. Thus, the quantum state to be concentrated can be expressed as $\alpha|\varphi_0 H_a\rangle + \beta|\varphi_1 V_a\rangle$. A linear optics based scheme for entanglement concentration of this state is shown in Fig. 2 and the action of the same is described below. Initial state of the system can be expressed as

$$\begin{aligned}
|\psi_3\rangle = |\psi\rangle \otimes |\psi_s\rangle &= [\alpha|\varphi_0 H_a\rangle + \beta|\varphi_1 V_a\rangle \\
&\quad \otimes (\alpha|H_b\rangle + \beta|V_b\rangle)]_{n+2} \\
&= \alpha^2|\varphi_0 H_a H_b\rangle + \beta^2|\varphi_1 V_a V_b\rangle \\
&\quad + \alpha\beta|\varphi_0 H_a V_b\rangle + \alpha\beta|\varphi_1 V_a H_b\rangle.
\end{aligned} \tag{4}$$

Here, subscripts a and b represent the modes of Alice's photon. To be precise, a corresponds to the $(n+1)^{th}$ qubit of the shared entangled state and b corresponds to the additional single photon state ($(n+2)^{th}$ qubit) prepared by Alice to implement the ECP. Now, Alice uses a half wave plate HWP_{90} to rotate the polarization of photon in mode b by 90 degrees to transform $|\psi_3\rangle$ to

$$\begin{aligned}
|\psi_4\rangle &= \alpha^2|\varphi_0 H_a V_b\rangle + \beta^2|\varphi_1 V_a H_b\rangle + \\
&\quad \alpha\beta|\varphi_0 H_a H_b\rangle + \alpha\beta|\varphi_1 V_a V_b\rangle.
\end{aligned} \tag{5}$$

Photons in mode a and b enter polarized beam splitter (PBS). As the PBS transmits the horizontal polarization component and reflects the vertical polarization component after the PBS the state of the system transforms to




Figure 2: Proposed linear optics circuit for ECP for non-maximally entangled $(n + 1)$ -qubit state $\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle$.

$$|\psi_5\rangle = \alpha^2|\varphi_0 H_{b'} V_{b'}\rangle + \beta^2|\varphi_1 V_{a'} H_{a'}\rangle + \alpha\beta|\varphi_0 H_{b'} H_{a'}\rangle + \alpha\beta|\varphi_1 V_{a'} V_{b'}\rangle. \quad (6)$$

Note that in the first two terms in (6), the last two photons are in the same spatial mode, whereas in the last two terms one photon is present in each output mode. Alice selects the case where each mode contains one photon. Therefore, after this choice Alice and Bob obtain a shared state $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H_{a'} H_{b'}\rangle + |\varphi_1\rangle|V_{a'} V_{b'}\rangle)_{n+2}$ with probability $2\alpha^2\beta^2$. Now, Alice allows the photon in mode b' to pass through R_{45} which rotates the polarization by 45 degrees. As a consequence the state of the system transforms to

$$\begin{aligned} |\psi_6\rangle &= \frac{1}{2}(|\varphi_0\rangle|H_{a'} H_{b'}\rangle + |\varphi_0\rangle|H_{a' a'} V_{b'}\rangle \\ &+ |\varphi_1\rangle|V_{a'} H_{b'}\rangle - |\varphi_1\rangle|V_{a'} V_{b'}\rangle)_{n+2} \\ &= \frac{1}{2}((|\varphi_0\rangle|H_{a'}\rangle + |\varphi_1\rangle|V_{a'}\rangle)|H_{b'}\rangle \\ &+ (|\varphi_0\rangle|H_{a'}\rangle - |\varphi_1\rangle|V_{a'}\rangle)|V_{b'}\rangle)_{n+2}. \end{aligned}$$

Finally, if the detector detects $(n + 2)^{th}$ photon in the horizontal mode (i.e., if D_H clicks) then Alice and Bob obtain the state

$$\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle + |\varphi_1\rangle|V\rangle)_{n+1}$$

and similarly if the detector D_V clicks, then they obtain the state

$$\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle - |\varphi_1\rangle|V\rangle)_{n+1}.$$

It is easy to observe that on application of the unitary operation in both the cases we obtain a maximally entangled state $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|0\rangle + |\varphi_1\rangle|1\rangle)_{n+1}$. Therefore, the probability P_1 to obtain the maximally entangled state $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle + |\varphi_1\rangle|V\rangle)_{n+1}$ or $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle - |\varphi_1\rangle|V\rangle)_{n+1}$ is $2(\alpha\beta)^2$.

2.3 Entanglement concentration using QND detectors (cross-Kerr-nonlinearity)

Quantum nondemolition detector (QND) is generally based on cross-Kerr-nonlinearity. After their introduction by Nemoto and Munro [33] many works precisely in entanglement concentration protocols of pure entangled states were reported. The Hamiltonian of the nonlinear medium or interaction between the signal beam and probe beam can be written as

$$\hat{H} = \hbar\chi\hat{n}_s\hat{n}_p \quad (7)$$

where $\hbar\chi$ is the coupling strength of the cross-Kerr material, \hat{n}_s and \hat{n}_p are the number operators of signal mode s and probe mode p , respectively. Let the signal mode be in input state $|\varphi\rangle = a|H\rangle + b|V\rangle$ and the probe mode be in a coherent state $|\alpha\rangle$. The cross-Kerr nonlinear medium causes the combined input state of $|\varphi\rangle$ and coherent state $|\alpha\rangle$ to evolve as

$$\begin{aligned} U_{cross-Kerr}|\varphi\rangle|\alpha\rangle &= e^{iHt/\hbar}(a|0\rangle + b|1\rangle)|\alpha\rangle \\ &= a|0\rangle|\alpha\rangle + b|1\rangle|\alpha e^{i\theta}\rangle, \end{aligned} \quad (8)$$

where $\theta = \chi t$ and t is the interaction time. In Eq. (8), $|n\rangle : n \in 0, 1$ represents the Fock state contains n photons. Therefore, we observe that $|\varphi\rangle$ remains same and the probe state picks up a phase shift and it can be measured by homodyne detection. The phase shift is directly proportional to the number of photons. In this way, we can nondestructively determinate the state of the signal mode by measuring the probe mode. Thus, QND can condition the evolution of the system by not destroying




Figure 3: Schematic of the quantum nondemolition detector (QND) constructed by cross-Kerr-nonlinearity to distinguish between $|HH\rangle$ and $|VV\rangle$ from $|HV\rangle$ and $|VH\rangle$.

the single photon. This is non-destructive measurement. QND measurement is used for parity checking i.e., to differentiate between a pair of identical polarizations $|HH\rangle$ or $|VV\rangle$ from a pair of orthogonal polarizations $|HV\rangle$ or $|VH\rangle$ depending on the phase shift θ of the coherent state. QND based scheme for parity checking is shown in Fig. 3. This scheme is frequently used in quantum information and detail description of this scheme can be found in [15, 33]. Let $|\phi_1\rangle$ and $|\phi_2\rangle$ be two polarization qubits.

$$\begin{aligned} |\phi_1\rangle &= a_1|H\rangle + b_1|V\rangle \\ |\phi_2\rangle &= a_2|H\rangle + b_2|V\rangle \end{aligned}$$

The composite quantum system is $|\phi_1\rangle \otimes |\phi_2\rangle \otimes |\alpha\rangle$. This system will evolve to

$$\begin{aligned} [a_1a_2|HH\rangle + b_1b_2|VV\rangle] |\alpha e^{i\theta}\rangle + a_1b_2|HV\rangle |\alpha e^{i2\theta}\rangle \\ + b_1a_2|VH\rangle |\alpha\rangle \end{aligned}$$

The qubit is passed through PBS and enters the spatial mode which interacts with nonlinear medium. The coherent beam picks up a phase shift θ if there is $|HH\rangle$ or $|VV\rangle$ and 2θ or no phase shift if there are $|HV\rangle$ or $|VH\rangle$ in a mode, respectively. Our proposed scheme is shown in Fig. 4, where we place a QND measurement based parity checking box in place of the first PBS used in the linear optics based scheme described earlier (cf. Fig. 2). The input state in Fig. 4 is

$$\begin{aligned} |\psi_7\rangle &= |\psi_4\rangle |\alpha\rangle \\ &= (\alpha^2 |\varphi_0 H_a V_b\rangle + \beta^2 |\varphi_1 V_a H_b\rangle \\ &\quad + \alpha\beta |\varphi_0 H_a H_b\rangle + \alpha\beta |\varphi_1 V_a V_b\rangle) |\alpha\rangle. \end{aligned}$$

After operation of the QND measurement based parity checking box the state transforms to

$$\begin{aligned} |\psi_8\rangle &= \alpha^2 |\varphi_0 H_a V_b\rangle |\alpha e^{i2\theta}\rangle + \beta^2 |\varphi_1 V_a H_b\rangle |\alpha\rangle \\ &\quad + \alpha\beta (|\varphi_0 H_a H_b\rangle + |\varphi_1 V_a V_b\rangle) |\alpha e^{i\theta}\rangle. \end{aligned}$$

QND based scheme for ECP of the same state is shown in the Fig. 3. Therefore, if there is a phase shift of θ then Alice will communicate to Bob to keep the state. After the measurement if D_H clicks, then Alice and Bob obtain maximally entangled state $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle + |\varphi_1\rangle|V\rangle)_{n+1}$ as desired. Similarly, if D_V clicks, then the combined state of Alice and Bob, collapses to $\frac{1}{\sqrt{2}}(|\varphi_0\rangle|H\rangle - |\varphi_1\rangle|V\rangle)_{n+1}$ which can be transformed to the desired maximally entangled state by a phase flip operation. Thus, if D_V clicks, then Bob applies a phase flip operation to obtain the desired state. Now the states for which phase shift is either 2θ or no phase shift form the less entangled states and are given by

$$\alpha' |\varphi_0 H_{a'} V_{b'}\rangle + \beta' |\varphi_1 V_{a'} H_{b'}\rangle,$$

where $\alpha' = \frac{\alpha^2}{\sqrt{\alpha^4 + \beta^4}}$ and $\beta' = \frac{\beta^2}{\sqrt{\alpha^4 + \beta^4}}$. The qubit in mode b' passes through R_{45} and as a consequence, we obtain

$$(\alpha' |\varphi_0 H_{a'}\rangle + \beta' |\varphi_1 V_{a'}\rangle) |H_{b'}\rangle - (\alpha' |\varphi_0 H_{a'}\rangle - \beta' |\varphi_1 V_{a'}\rangle) |V_{b'}\rangle.$$

Therefore, if detector D_H clicks we get the state $\alpha' |\varphi_0 H_{a'}\rangle + \beta' |\varphi_1 V_{a'}\rangle$ and Alice needs to prepare another single photon having the same Schmidt coefficients i.e., $\alpha' |H_{a_2}\rangle + \beta' |V_{a_2}\rangle$. Alice can reconstruct the maximally entangled state following the same process. The probability of success of second iteration is given by

$$(1 - P_1)P_2 = (1 - 2\alpha^2\beta^2) \frac{2(\alpha'\beta')^2}{((\alpha')^2 + (\beta')^2)^2} = \frac{2\alpha^4\beta^4}{\alpha^4 + \beta^4}.$$

Thus, the probability of success after N iterations would be

$$P = P_1 + (1 - P_1)[P_2 + (1 - P_2)[P_3 \cdots + (1 - P_{N-1})[P_N]].$$




Figure 4: Schematic of ECP with QND.

This can be written as

$$\begin{aligned}
P &= 2\alpha^{2^1}\beta^{2^1} + \frac{2\alpha^{2^2} + \beta^{2^2}}{(\alpha^{2^2} + \beta^{2^2})} + \frac{2\alpha^{2^3} + \beta^{2^3}}{(\alpha^{2^2} + |\beta|^{2^2})(\alpha^{2^3} + \beta^{2^3})} + \dots \\
&+ \frac{2\alpha^{2^N}\beta^{2^N}}{(\alpha^{2^2} + \beta^{2^2})(\alpha^{2^3} + \beta^{2^3})(\alpha^{2^4} + \beta^{2^4}) \dots (\alpha^{2^N} + \beta^{2^N})} \\
&= 2\alpha^2\beta^2 + 2\beta^4 \left[\frac{x^{2^2-4}}{(1+x^{2^2})} + \frac{x^{2^3-4}}{(1+x^{2^2})(1+x^{2^3})} + \right. \\
&+ \left. \dots \frac{x^{2^N-4}}{(1+x^{2^2}) \dots (|\alpha|^{2^N} + |\beta|^{2^N})} \right],
\end{aligned}$$

where $x = \frac{\beta}{\alpha}$ and $0 < x < 1$. For large N , i.e., $N \rightarrow \infty$, above series (the series present within square bracket of the expression of P) becomes an infinite series and that converges to 1 for $x \in (0, 1)$. Therefore, $P = 2\beta^2 (\alpha^2 + \beta^2) = 2\beta^2$. Thus, the maximum probability with which we can obtain a MES from the corresponding less entangled state for $(n+1)$ -qubit state of the form $|\psi\rangle = (\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$ is $2\beta^2$. It is a straightforward exercise to show that the same bound on the maximum success probability is also applicable to the recent proposals of Sheng et. al. [11, 12].

3 Conservation of entanglement

It is well known that there exists a large number of measures of entanglement, each having its merit and demerit. In the investigations related to entanglement concentration and purification, a specific measure known as *entanglement of single pair purification* (ESPP) is often used [2, 3]. This measure was introduced by Bose et al. [2] as the maximum probability with which two parties sharing a non-maximally entangled pure Bell-type state can transform it to a Bell state by using LOCC (local operation and classical communication). Historically, Bose et al., defined ESPP in context of Bell states as in their original work they described an ECP for Bell state and showed that ESPP is such a measure of entanglement which remains conserved in the ECP described by them. Later on Bandyopadhyay [3] showed that ESPP remains conserved in a qubit-assisted ECP for Bell state. Motivated by this observation, here we plan to show that ESPP is also conserved in the QND-based ECP proposed here. To do so, first we need to generalize the definition of ESPP for states other than Bell state as our ECP is valid for all states of the form described in Eq. (1). The generalized definition of ESPP is as follows: ESPP is the maximum probability with which two parties sharing a non-maximally entangled pure entangled state can transform it to the corresponding maximally entangled state by using LOCC. Following Lo and Popescu [36] we may note that the maximum probability with which a nonmaximally entangled pure state of the form provided in Eq. (1) can be concentrated is twice the modulus square of the Schmidt coefficient of the smaller magnitude. Therefore, before concentration the average entanglement shared between Alice and Bob is $\langle E_{\text{before}} \rangle_{AB} = 2\beta^2$ where $\alpha > \beta$. In the proposed QND based protocol, the maximum probability for $(n+1)$ -qubit state after the concentration is $\langle E_{\text{after}} \rangle_{AB} = 2\alpha^2\beta^2 + 2\beta^4 = 2\beta^2$. This implies that the average ESPP is conserved, the maximum probability to concentrate $(n+1)$ -qubit state is $2\beta^2$ and that the remaining will be totally disentangled if the protocol is continued for sufficiently long time. Here we have followed the works of Bose et al. [2] and Bandyopadhyay [3] to establish the conservation of entanglement in the similar sense. However, it may be noted that this conclusion (i.e., conservation of entanglement) is entanglement measure specific and it may and may not hold for another measure of entanglement.

4 Conclusion

We have proposed two practical schemes for entanglement concentration for all entangled states that can be expressed in the form $(\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$. Specifically, we have proposed an ECP using linear optics i.e., using PBSs and single photon detectors. Subsequently, we have also proposed an ECP using cross-Kerr-nonlinearity and have obtained the maximum possible success probability $2\beta^2$. The proposed ECP uses less resources compared to the earlier proposals. We have presented a quantum circuit for the proposed ECP so that the protocol can be implemented using other technologies like atomic system, NMR, etc. The proposed ECP can be applied to many entangled states like Bell and cat states, GHZ, GHZ-like, $|\Omega\rangle$, $|Q_5\rangle$, 4-qubit cluster states and specific states from the 9 SLOCC-nonequivalent families of 4-qubit entangled states and other states which can be expressed as $(\alpha|\varphi_0\rangle|0\rangle + \beta|\varphi_1\rangle|1\rangle)_{n+1}$.

Acknowledgement

AB thanks DST-SERB project SR/S2/LOP-18/2012. AB also acknowledges S. Bandyopadhyay for some technical discussion and his interest in the present work. AP thanks Department of Science and Technology (DST), India for support provided through the DST project No. SR/S2/LOP-0012/2010

References

- [1] C. H. Bennett, H. J. Bernstein, S. Popescu and B. Schumacher, Concentrating partial entanglement by local operations, *Phys. Rev. A*, 53, 2046 (1996)
- [2] S. Bose, V. Vedral and P. L. Knight, Purification via entanglement swapping and conserved entanglement, *Phys. Rev. A*, 60, 194 (1999)
- [3] S. Bandyopadhyay, Qubit- and entanglement-assisted optimal entanglement concentration, *Phys. Rev. A*, 62, 032308 (2000)
- [4] Y. J. Gu, W. D. Li and G. C. Guo, Protocol and quantum circuits for realizing deterministic entanglement concentration, *Phys. Rev. A*, 73, 022321 (2006)
- [5] S. W. Lee and H. Jeong, Near-deterministic quantum teleportation and resource-efficient quantum computation using linear optics and hybrid qubits, *Phys. Rev. A*, 87, 022326 (2013)
- [6] B. S. Choudhury and A. Dhara, A three-qubit state entanglement concentration protocol assisted by two-qubit systems, *Int. J. Theor. Phys.*, 52, 3965 (2013)
- [7] C. Shukla, A. Banerjee and A. Pathak, Protocols and quantum circuits for implementing entanglement concentration in cat state, GHZ-like state and 9 families of 4-qubit entangled states, *Quantum Inf. Process.*, DOI 10.1007/s11128-015-0948-6, 2015
- [8] B. S. Choudhury and A. Dhara, An entanglement concentration protocol for cluster states, *Quantum Inf. Process.*, 12, 2577 (2013)
- [9] Z. Lan, Consequent entanglement concentration of a less-entangled electronic cluster state with controlled-not gates, *Chin. Phys. B*, 23, 050308 (2014)
- [10] S. Y. Zhao, J. Liu, L. Zhou, Y. B. Sheng, Two-step entanglement concentration for arbitrary electronic cluster state, *Quantum Inf. Process.*, 12, 3633 (2013)
- [11] Y. B. Sheng, L. Zhou and S. M. Zhao, Efficient two-step entanglement concentration for arbitrary W states, *Phys. Rev. A*, 85, 042302 (2012)
- [12] Y. B. Sheng, L. Zhou, S. M. Zhao and B. Y. Zheng, Efficient single-photon-assisted entanglement concentration for partially entangled photon pairs, *Phys. Rev. A*, 85, 012307 (2012)
- [13] Z. Zhao, T. Yang, Y. A. Chen, A. N. Zhang and J. W. Pan, Experimental realization of entanglement concentration and a quantum repeater, *Phys. Rev. Lett.*, 90, 207901 (2003)
- [14] T. Yamamoto, M. Koashi, S. K. Ozdemir and N. Imoto, Experimental extraction of an entangled photon pair from two identically decohered pairs, *Nature*, 421, 343 (2003)
- [15] Y. B. Sheng, F. G. Deng and H. Y. Zhou, Nonlocal entanglement concentration scheme for partially entangled multipartite systems with nonlinear optics. *Phys. Rev. A*, 77, 062325 (2008)
- [16] B. C. Ren, F. F. Du and F. G. Deng, Hyperentanglement concentration for two-photon four-qubit systems with linear optics, *Phys. Rev. A*, 88, 012302 (2013)

- [17] C. Wang, Efficient entanglement concentration for partially entangled electrons using a quantum-dot and microcavity coupled system. *Phys. Rev. A*, 86, 012323 (2012)
- [18] C. Wang, Y. Zhang and G. S. Jin, Entanglement purification and concentration of electron-spin entangled states using quantum-dot spins in optical microcavities, *Phys. Rev. A*, 84, 032307 (2011)
- [19] Y. B. Sheng, L. Zhou, L. Wang and S. M. Zhao, Efficient entanglement concentration for quantum dot and optical microcavities systems, *Quantum Inf. Process.*, 12, 1885 (2013)
- [20] Y. B. Sheng, J. Liu, S. Y. Zhao and L. Zhou, Multipartite entanglement concentration for nitrogen-vacancy center and microtoroidal resonator system, *Chin. Sci. Bull.*, 58, 3507 (2013)
- [21] B. C. Ren, H. R. Wei, T. Li, M. Hua and F. G. Deng, Optimal multipartite entanglement concentration of electron-spin states based on charge detection and projection measurements, *Quantum Inf. Process.*, 13, 825 (2014)
- [22] L. Zhou, Y. B. Sheng, W. W. Cheng, L. Y. Gong, S. M. Zhao, Efficient entanglement concentration for arbitrary single-photon multimode W state, *Journal of the Optical Society of America B*, 30, 71 (2013)
- [23] C. Wang, Efficient entanglement concentration for partially entangled electrons using a quantum-dot and microcavity coupled system, *Phys. Rev. A*, 86, 012323 (2012)
- [24] C. Cao, C. Wang, L. Y. He and R. Zhang, Atomic entanglement purification and concentration using coherent state input-output process in low-Q cavity QED regime, *Optics Express*, 21, Issue 4, 4093 (2013)
- [25] L. Zhou, Efficient entanglement concentration for electron-spin W state with the charge detection, *Quantum Inf. Process.*, 12, Issue 6, 2087 (2013)
- [26] L. Zhou, Efficient entanglement concentration for arbitrary less-entangled N-atom state, *Int. J. Theor. Phys.* DOI 10.1007/s10773-013-1974-8 (2014)
- [27] L. Zhou, Y. B. Sheng, W. W. Cheng, L. Y. Gong and S.-M. Zhao, Efficient entanglement concentration for arbitrary less-entangled NOON states, *Quant. Inf. Process.*, 12, 1307 (2013)
- [28] Y. B. Sheng and L. Zhou, Efficient W-state entanglement concentration using quantum-dot and optical microcavities, *JOSA B*, 30, Issue 3, 678 (2013)
- [29] L. Zhou, Y. B. Sheng, W. W. Cheng, L. Y. Gong and S. M. Zhao, Efficient entanglement concentration for arbitrary less-entangled NOON states, *Quantum Inf. Process.*, 12, 1307 (2013)
- [30] L. Y. Hea, C. Cao and C. Wang, *Optics Communications*, Entanglement concentration for multi-particle partially entangled W state using nitrogen vacancy center and microtoroidal resonator system, Volumes 298, 260 (2013)
- [31] C. Shukla and A. Pathak, Hierarchical quantum communication, *Phys. Lett. A*, 377, 1337 (2013)
- [32] S. Mishra, C. Shukla, A. Pathak, R. Srikanth and A. Venugopalan, An Integrated Hierarchical Dynamic Quantum Secret Sharing Protocol, *Int. J. Theor. Phys.*, DOI 10.1007/s10773-015-2552-z, (2015)
- [33] K. Nemoto and W. J. Munro, Nearly deterministic linear optical controlled-NOT gate, *Phys. Rev. Lett.*, 93, 250502 (2004)
- [34] S. D. Barrett, P. Kok, K. Nemoto, R. G. Beausoleil, W. J. Munro and T. P. Spiller, Symmetry analyzer for nondestructive Bell-state detection using weak nonlinearities, *Phys. Rev. A*, 71, 060302 (2005)
- [35] C. Wang, Y. Zhang and G. S. Jin, Entanglement purification and concentration of electron-spin entangled states using quantum-dot spins in optical microcavities, *Phys. Rev. A*, 84, 032307 (2011)
- [36] H. K. Lo and S. Popescu, Concentrating entanglement by local actions: Beyond mean values, *Phys. Rev. A*, 63, 022301 (2001)