

Refuting Unique Game Conjecture

Peng Cui

Key Laboratory of Data Engineering and Knowledge Engineering, MOE, School of Information Resource Management, Renmin University of China, Beijing 100872, P. R. China.

cuipeng@ruc.edu.cn

Abstract. In this short note, the author shows that the gap problem of some k -CSPs with the support of its predicate the ground of a balanced pairwise independent distribution can be solved by a modified version of Hast's Algorithm BiLin that calls Charikar&Wirth's SDP algorithm for two rounds in polynomial time, when k is sufficiently large, the support of its predicate is combined by the grounds of three biased homogeneous distributions and the three biases satisfy certain conditions. To conclude, the author refutes Unique Game Conjecture, assuming $P \neq NP$.

1 Introduction

Max k -CSP is the task of satisfying the maximum fraction of constraints when each constraint involves k variables, and each constraint accepts the same collection $C \subseteq G^k$ of local assignments. A challenging question is to identify constraint satisfaction problems (CSPs) that are extremely hard to approximate, so much so that they are NP-hard to approximate better than just outputting a random assignment. Such CSPs are called approximation resistant; famous examples include Max 3-SAT and Max 3-XOR[5]. A lot is known about such CSPs of arity at most four[6], but for CSPs of higher arity, results have been scattered.

To make progress, conditional results are obtained assuming the Unique Game Conjecture (UGC) of [7]. Under UGC, [2] shows that a CSP is approximation resistant if the support of its predicate is the ground of a balanced pairwise independent distribution. Traditionally, negations are allowed free in definition of k -CSP. In [1], the authors investigate k -CSP with no negations of variables and prove such k -CSP with the support of its predicate the ground of a biased pairwise independent distribution or uniformly positively correlated distribution or is approximation resistant in biased sense under Unique Game Conjecture.

The following is Theorem 3.1 in [2].

Theorem 1. *Let $k \geq 3$ be an integer, and C be a subset of G^k and the ground of a balanced pairwise independent distribution. For arbitrarily small constant ε , it is UG-hard to distinguish the following two cases given a instance P of Max C :*

- *Completeness:* $\text{val}(P) \geq 1 - \varepsilon$.

– *Soundness*: $\text{val}(P) \leq \frac{|C|}{2^k} + \varepsilon$.

In this short note, the author shows that the gap problem of this type of k -CSP can be solved by a modified version of Hast's Algorithm BiLin in polynomial time that calls Charikar&Wirth's SDP algorithm[3] for two rounds, when k is sufficiently large, the support of its predicate is combined by the grounds of three biased homogeneous distributions and the three biases satisfy certain conditions. To conclude, the author refutes Unique Game Conjecture, assuming $P \neq NP$.

Theorem 2. *Unique Game Conjecture does not hold true, assuming $P \neq NP$.*

This work has an origin that conditionally strengthens the previous known hardness for approximating Min 2-Lin-2 and Min Bisection, assuming a claim that refuting Unbalanced Max 3-XOR under biased assignments is hard on average[4]. In this paper, the author defines "bias" to be a parameter of pairwise independent subset (distribution), while he defines "bias" to be the fraction of variables assigned to value 1 in [4]. The author notices that biased pairwise independent distribution is defined in [2,1] and uniformly positively correlated distribution is defined in [1].

2 Definitions

As usual, let $[q] = \{1, 2, \dots, q\}$, and $-[q] = \{-q, -q+1, \dots, -1\}$.

Let $G = \{1, -1\}$, here 1 represent "0/false" and -1 represent "1/true" in standard Boolean algebra.

Random variables are denoted by italic boldface letters, such as \mathbf{z} . Suppose φ is a distribution over G^k , the ground of φ is defined as

$$G_\varphi = \{\varphi(\mathbf{z}) > 0 | \mathbf{z} \in G^k\}.$$

Definition 1. *For some $0 < \gamma < 1$, a distribution φ over G^k is biased pairwise independent if for every coordinate $i \in [k]$,*

$$\mathbb{P}[\mathbf{z}_i = 1] = \gamma$$

and for every two distinct coordinates $i, j \in [k]$,

$$\mathbb{P}[\mathbf{z}_i = 1, \mathbf{z}_j = 1] = \gamma^2,$$

where \mathbf{z} is a random element drawn from G^k by φ . γ is called bias of φ . If $\gamma = \frac{1}{2}$, we say φ is balanced pairwise independent.

The author notices the fact that a distribution over G^k can be thought as a linear superposition of several distributions over G^k .

Definition 2. Given m distributions φ_l over G^k with disjoint grounds G_{φ_l} , let ψ is a distribution over $[m]$ with $\psi_l > 0$ for each $l \in [m]$, and φ be the distribution over G^k such that

$$\varphi(\mathbf{z}) = \sum_{l=1}^m \psi_l \varphi_l(\mathbf{z}),$$

for each $\mathbf{z} \in G^k$. We say φ_l 's are disguised by ψ to φ .

3 Proof of Theorem 2

We construct the instances by three homogeneous distributions that are uniformly negatively correlated.

For each $l \in [3]$, suppose γ_l is a constant satisfying $0 \leq \gamma_l \leq 1$ and $\gamma_l k$ is an integer and ρ_l 's are constants such that

$$\gamma_1 = \frac{1}{2} + \rho_1 \frac{1}{\sqrt{k}}, \gamma_2 = \frac{1}{2} - \rho_2 \frac{1}{\sqrt{k}}, \gamma_3 = \frac{1}{2} + \rho_3 \frac{1}{\sqrt{k}}.$$

Let G_m denote the subset of G^k including all k -tuples with exactly m 1. For each $l \in [3]$, let φ_l be the uniform distribution over $G_{\gamma_l k}$, called γ_l -biased homogeneous distribution, which satisfies the following two properties:

For every coordinate $i \in [k]$,

$$\mathbb{P}[\mathbf{z}_i = 1] = \frac{1}{2},$$

and for every two distinct coordinates $i, j \in [k]$,

$$\mathbb{P}[\mathbf{z}_i = 1, \mathbf{z}_j = 1] = \frac{k}{k-1} \gamma_l^2 - \frac{1}{k-1} \gamma_l,$$

where \mathbf{z} is a random element drawn by φ_l .

We can prove the following lemma. (Section 4)

Lemma 1. Suppose $C = G_{\varphi_1} \cup G_{\varphi_2} \cup G_{\varphi_3}$, and $P^{(3)}(y)$ is the tri-linear term of the Fourier spectra of C . When k is sufficiently large, there are three absolute constants ρ_1, ρ_2 and ρ_3 satisfying $0 < \rho_1 < \rho_2 \approx \rho_3$, $\gamma_l k$'s are integers, and a distribution ψ over $[3]$ such that:

1. φ_l 's are disguised by ψ to a balanced pairwise independent distribution φ .
2. $P^{(3)}(y) \geq \iota$ for any $y \in C$, where $\iota = \Omega(\frac{1}{\sqrt{k}})$.

In the dictatorship test, for every three distinct coordinates $i_1, i_2, i_3 \in [k]$, let

$$\mathbb{P}[\mathbf{z}_{i_1} = 1, \mathbf{z}_{i_2} = 1, \mathbf{z}_{i_3} = 1] \triangleq a,$$

$$\begin{aligned} \mathbb{P}[\mathbf{z}_{i_1} = 1, \mathbf{z}_{i_2} = 1, \mathbf{z}_{i_3} = -1] &= \mathbb{P}[\mathbf{z}_{i_1} = 1, \mathbf{z}_{i_2} = -1, \mathbf{z}_{i_3} = 1] \\ &= \mathbb{P}[\mathbf{z}_{i_1} = 1, \mathbf{z}_{i_2} = 1, \mathbf{z}_{i_3} = -1] \triangleq b, \end{aligned}$$

$$\begin{aligned}\mathbb{P}[\mathbf{z}_{i_1} = 1, \mathbf{z}_{i_2} = -1, \mathbf{z}_{i_3} = -1] &= \mathbb{P}[\mathbf{z}_{i_1} = -1, \mathbf{z}_{i_2} = 1, \mathbf{z}_{i_3} = -1] \\ &= \mathbb{P}[\mathbf{z}_{i_1} = -1, \mathbf{z}_{i_2} = -1, \mathbf{z}_{i_3} = 1] \triangleq c,\end{aligned}$$

$$\mathbb{P}[\mathbf{z}_{i_1} = -1, \mathbf{z}_{i_2} = -1, \mathbf{z}_{i_3} = -1] \triangleq d,$$

where \mathbf{z} is a random element drawn by φ .

Then

$$\begin{pmatrix} 1 & 3 & 3 & 1 \\ 1 & -3 & 3 & -1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} = \begin{pmatrix} 1 \\ \alpha \\ \frac{1}{4} \\ \frac{1}{4} \end{pmatrix},$$

where

$$\alpha \approx 8\rho_1(\frac{1}{4} - \rho_2^2)\frac{1}{k\sqrt{k}},$$

which reduces to

$$\begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix} = \begin{pmatrix} \frac{1}{8} + \frac{\alpha}{8} \\ \frac{1}{8} - \frac{\alpha}{8} \\ \frac{1}{8} + \frac{\alpha}{8} \\ \frac{1}{8} - \frac{\alpha}{8} \end{pmatrix}.$$

By Theorem 1, given an instance P as in the statement of Lemma 1, for arbitrarily small constant ε , it is UG-hard to distinguish the following two cases: $\text{val}(P) \geq 1 - \varepsilon$; $\text{val}(P) \leq \frac{|C|}{2^k} + \varepsilon$.

On the other hand, consider the Fourier spectra of P , since C is folded and \mathbf{z} is balanced pairwise independent, there is no linear or bi-linear term. Let $I_{123}^{(3)}$ be the sum of tri-linear terms containing three variables with coordinate 1, 2 and 3 respectively in the Fourier spectra of P . Suppose $\text{val}(P) \geq 1 - \varepsilon$ for some ε , there is an assignment f^* under which $I_{123}^{(3)} \geq \Omega(\frac{1}{k^2\sqrt{k}})$ (cf. Lemma 4 in [6]).

Let $I_{123}^{(2)}$ be the sum of bi-linear terms defined as: For each tri-linear term $a_{i_1 i_2 i_3} x_{i_1}^{(1)} x_{i_2}^{(2)} x_{i_3}^{(3)}$ in $I_{123}^{(3)}$, introduce a bi-linear term $a_{i_1 i_2 i_3} x_{i_1}^{(1)} x_{i_2 i_3}^{(2)}$, where $x_{i_2 i_3}^{(2)}$'s are new variables in G , where $i_1 \in [M]$ and $i_2, i_3 \in [N]$.

We modify Hast's Algorithm BiLin as follows:

- Step 1, run Charikar&Wirth's SDP algorithm for the first round on $I_{123}^{(2)}$ to get an assignment $f^{(1)}$ on $x_{i_1}^{(1)}$'s and $x_{i_2 i_3}^{(2)}$'s.
- Step 2, run Charikar&Wirth's SDP algorithm for the second round on $I^{(3)}$ subject to $f^{(1)}$ to get an assignment $f^{(2)}$ to $x_{i_2}^{(2)}$'s and $x_{i_3}^{(3)}$'s.
- Step 3, let $f := f^{(1)}$ for $x_{i_1}^{(1)}$'s and let $f := f^{(2)}$ for $x_{i_2}^{(2)}$'s and $x_{i_3}^{(3)}$'s.
- Step 4, same as Step 3 in the original algorithm.
- Step 5, same as Step 4 in the original algorithm.

The first round returns $f^{(1)}$ under which $I_{123}^{(2)}$ is at least $\Omega(\frac{1}{k^2\sqrt{k}\log k})$ (cf. Lemma 5 in [3]). By enumeration arguments, there is an assignment f' to $x_{i_2}^{(2)}$'s and $x_{i_3}^{(3)}$'s under which $I^{(3)}$ subject to $f^{(1)}$ is at least $\Omega(\frac{1}{k^2\sqrt{k}\log k})$. Hence the second round returns $f^{(2)}$ under which $I^{(3)}$ subject to $f^{(1)}$ is at least $\Omega(\frac{1}{k^2\sqrt{k}\log^2 k})$ (cf. Lemma 5 in [3]).

Therefore, the modified version of BiLin returns a solution of P with value at least $\frac{|C|}{2^k} + \kappa$, where

$$\kappa = \Omega(\frac{1}{k^2\sqrt{k}\log^2 kk})^3 = \Omega(\frac{1}{\text{poly}(k)}),$$

(cf. Theorem 3 in [6]).

Therefore, Unique Game Conjecture does not hold true, assuming $P \neq NP$. The proof of Theorem 2 is accomplished.

4 Proof of Lemma 1

Let $m = 3$, consider the linear equations with ψ_l for $l \in [m]$,

$$\begin{pmatrix} 1 & 1 & 1 \\ \gamma_1 & \gamma_2 & \gamma_3 \\ \frac{k}{k-1}\gamma_1^2 - \frac{1}{k-1}\gamma_1 & \frac{k}{k-1}\gamma_2^2 - \frac{1}{k-1}\gamma_2 & \frac{k}{k-1}\gamma_3^2 - \frac{1}{k-1}\gamma_3 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{1}{2} \\ \frac{1}{4} \end{pmatrix},$$

which reduces to

$$\begin{pmatrix} 1 & 1 & 1 \\ \rho_1 & -\rho_2 & \rho_3 \\ \rho_1^2 & \rho_2^2 & \rho_3^2 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ \frac{1}{4} \end{pmatrix},$$

or

$$\begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} = \begin{pmatrix} \frac{-\rho_2\rho_3 + \frac{1}{4}}{(\rho_1 + \rho_2)(\rho_1 - \rho_3)} \\ \frac{\rho_3\rho_1 + \frac{1}{4}}{(\rho_2 + \rho_3)(\rho_2 + \rho_1)} \\ \frac{-\rho_1\rho_2 + \frac{1}{4}}{(\rho_3 - \rho_1)(\rho_3 + \rho_2)} \end{pmatrix}.$$

For $0 < \rho_1 < \rho_3$ and $0 < \rho_2$, $\psi_l > 0$ if and only if $\rho_1\rho_2 < \frac{1}{4}$ and $\rho_2\rho_3 > \frac{1}{4}$.

On the other hand, $P^{(3)}(y) = \sum_{\{i_1, i_2, i_3\} \subseteq [k]} a_{i_1 i_2 i_3} y_{i_1} y_{i_2} y_{i_3}$, and $a_{i_1 i_2 i_3} = 2^{-k} \sum_{y \in C} y_{i_1} y_{i_2} y_{i_3}$.

Suppose $\rho_2 \approx \rho_3$. For $y \in G_{\phi_l}$, we have for each $\{i_1, i_2, i_3\} \subseteq [k]$,

$$\begin{aligned} & a_{i_1 i_2 i_3} \\ &= \frac{1}{k\sqrt{k}2^k} \left(\binom{k}{\gamma_1 k} (6\rho_1(\frac{4}{3}\rho_1^2 - 1) + o_k(1)) - \binom{k}{\gamma_2 k} (6\rho_2(\frac{4}{3}\rho_2^2 - 1) + o_k(1)) \right. \\ & \quad \left. + \binom{k}{\gamma_3 k} (6\rho_3(\frac{4}{3}\rho_3^2 - 1) + o_k(1)) \right) \\ & \approx 6c(\rho_1(\frac{4}{3}\rho_1^2 - 1) + o_k(1)) \frac{1}{k^2}. \end{aligned}$$

and

$$\sum_{\{i_1, i_2, i_3\} \subseteq [k]} y_{i_1} y_{i_2} y_{i_3} = (\rho_l(\frac{4}{3}\rho_l^2 - 1) + o_k(1))k\sqrt{k},$$

hence

$$P^{(3)}(y) \approx 6c(\rho_l(\frac{4}{3}\rho_l^2 - 1)\rho_1(\frac{4}{3}\rho_1^2 - 1) + o_k(1))\frac{1}{\sqrt{k}},$$

c is an absolute positive.

When k is sufficiently large, we can determine ρ_l 's satisfying $\rho_1\rho_2 < \frac{1}{4}$, $\sqrt{\frac{1}{2}} < \rho_2 \approx \rho_3 < \frac{\sqrt{3}}{2}$, $\gamma_l k$'s are integers such that φ_l 's can be disguised by ψ to a balanced pairwise independent distribution, and the tri-linear term of Fourier spectra of C is at least $\iota = \Omega(\frac{1}{\sqrt{k}})$.

References

1. Austrin, P., & Håstad, J. (2013). On the usefulness of predicates. *ACM Transactions on Computation Theory (TOCT)*, 5(1), 1.
2. Austrin, P., & Mossel, E. (2009). Approximation resistant predicates from pairwise independence. *Computational Complexity*, 18(2), 249-271.
3. Charikar M. & Wirth A. (2004). Maximizing quadratic programs: Extending Grothendieck's inequality. In *Proceedings of the 45th annual IEEE Symposium on Foundations of Computer Science* (pp. 54-60). IEEE.
4. Cui, P. (2013, July). Strengthened hardness for minimum unique game and small set expansion. *Satellite Workshop of ICALP 2013: International Workshop on Approximation, Parameterized and EXact algorithms*. arXiv:1204.2026.
5. Håstad, J. (2001). Some optimal inapproximability results. *Journal of the ACM (JACM)*, 48(4), 798-859.
6. Hast, G. (2005). Beating a random assignment. In *Approximation, Randomization and Combinatorial Optimization. Algorithms and Techniques* (pp. 134-145). Springer Berlin Heidelberg.
7. Khot, S. (2002, May). On the power of unique 2-prover 1-round games. In *Proceedings of the 34th annual ACM symposium on Theory of computing* (pp. 767-775). ACM.