

Watch and Learn: Optimizing from Revealed Preferences Feedback

Aaron Roth*

Jonathan Ullman†

Zhiwei Steven Wu‡

December 19, 2021

Abstract

A *Stackelberg game* is played between a *leader* and a *follower*. The leader first chooses an action, then the follower plays his best response. The goal of the leader is to pick the action that will maximize his payoff given the follower’s best response. In this paper we present an approach to solving for the leader’s optimal strategy in certain Stackelberg games where the follower’s utility function (and thus the subsequent best response of the follower) is *unknown*.

Stackelberg games capture, for example, the following interaction between a producer and a consumer. The producer chooses the *prices* of the goods he produces, and then a consumer chooses to buy a utility maximizing bundle of goods. The goal of the seller here is to set prices to maximize his profit—his revenue, minus the production cost of the purchased bundle. It is quite natural that the seller in this example should not know the buyer’s utility function. However, he does have access to *revealed preference* feedback—he can set prices, and then observe the purchased bundle and his own profit. We give algorithms for efficiently solving, in terms of both computational and query complexity, a broad class of Stackelberg games in which the follower’s utility function is unknown, using only “revealed preference” access to it. This class includes in particular the profit maximization problem, as well as the optimal tolling problem in nonatomic congestion games, when the latency functions are unknown. Surprisingly, we are able to solve these problems even though the optimization problems are non-convex in the leader’s actions.

*University of Pennsylvania Department of Computer and Information Sciences. Partially supported by an NSF CAREER award, NSF Grants CCF-1101389 and CNS-1065060, and a Google Focused Research Award. aaroth@cis.upenn.edu.

†Northeastern University College of Computer and Information Science. Most of this work was done while the author was supported by a Junior Fellowship from the Simons Society of Fellows. jullman@ccs.neu.edu.

‡University of Pennsylvania Department of Computer and Information Sciences. wuzhiwei@cis.upenn.edu.

Contents

1	Introduction	1
1.1	Our Results and Techniques	2
1.2	Related Work	3
2	Preliminaries	4
2.1	Projected Subgradient Descent	5
2.2	Strong Convexity	5
2.3	Tools for Zeroth-Order Optimization	6
3	Profit Maximization From Revealed Preferences	7
3.1	The Model and Problem Setup	7
3.2	An Overview of Our Solution	8
3.3	Expressing Profit as a Function of the Bundle	8
3.4	Converting Bundles to Prices	10
3.5	Profit Maximization	14
4	General Framework of Stackelberg Games	18
4.1	Inducing a Target Action of the Follower	20
4.2	Optimizing Leader's Utility	20
5	Optimal Traffic Routing from Revealed Behavior	21
6	The Principal-Agent Problem	25
6.1	Inducing the Agent's Contribution Using Noisy Observations	26
6.2	Optimizing the Principal's Utility	27
7	Conclusion	30
A	A Routing Game Where Social Cost is Not Convex in The Tolls	32
B	Missing Proofs in Section 3	34
B.1	Properties of CES and Cobb-Douglas Utilities	34
B.1.1	Constant Elasticity of Substitution (CES)	34
B.1.2	Cobb-Douglas	36
C	Detailed Analysis of Section 4	38
D	Improvement with Ellipsoid in Noiseless Settings	41
D.1	The Ellipsoid Algorithm	41
D.2	Learning Prices with Ellipsoid	42
D.3	Learning Tolls with Ellipsoid	44

1 Introduction

Consider the following two natural problems:

1. **Profit Maximization via Revealed Preferences:** A retailer, who sells d goods, repeatedly interacts with a buyer. In each interaction, the retailer decides how to price the d goods by choosing $p \in \mathbb{R}_+^d$, and in response, the buyer purchases the bundle $x \in \mathbb{R}_+^d$ that maximizes her utility $v(x) - \langle x, p \rangle$, where v is an unknown concave valuation function. The retailer observes the bundle purchased, and therefore his profit, which is $\langle x, p \rangle - c(x)$, where c is an unknown convex cost function. The retailer would like to set prices that maximize his profit after only a polynomial number of interactions with the buyer.
2. **Optimal Tolling via Revealed Behavior:** A municipal authority administers m roads that form a network $G = (V, E)$. Each road $e \in E$ of the network has an unknown latency function $\ell_e : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which determines the time it takes to traverse the road given a level of congestion. The authority has the power to set constant tolls $\tau_e \in \mathbb{R}_+$ on the roads in an attempt to manipulate traffic flow. In rounds, the authority sets tolls, and then observes the *Nash equilibrium flow* induced by the non-atomic network congestion game defined by the unknown latency functions and the tolls, together with the social cost (average total latency) of the flow. The authority would like to set tolls that minimize the social cost after only a polynomial number of rounds.

Although these problems are quite different, they share at least one important feature—the retailer and the municipal authority each wish to optimize an *unknown* objective function given only query access to it. That is, they have the power to choose some set of prices or tolls, and then observe the value of their objective function that results from that choice. This kind of problem (alternately called *bandit* or *zeroth order* optimization) is well-studied, and is well understood in cases in which the unknown objective being maximized (resp. minimized) is concave (resp. convex). Unfortunately, the two problems posed above share another important feature—when posed as bandit optimization problems, the objective function being maximized (resp. minimized) is generally not concave (resp. convex). For the profit maximization problem, even simple instances lead to a non concave objective function.

Example 1. Consider a setting with one good ($d = 1$). The buyer’s valuation function $v(x) = \sqrt{x}$, and the retailer’s cost function is $c(x) = x$. The buyer’s utility for buying x units at price p is $\sqrt{x} - x \cdot p$. Thus, if the price is p , a utility-maximizing buyer will purchase $x^*(p) = \frac{1}{4p^2}$ units. The profit of the retailer is then

$$\text{Profit}(p) = p \cdot x^*(p) - c(x^*(p)) = \frac{1}{4p} - \frac{1}{4p^2}.$$

Unfortunately, this profit function is not concave.

Since the retailer’s profit function is not concave in the prices, it cannot be optimized efficiently using generic methods for concave maximization. This phenomenon persists into higher dimensions, where it is not clear how to efficiently maximize the non-concave objective. The welfare objective in the tolling problem is also non-convex in the tolls. We give an example in Appendix A.

Surprisingly, despite this non-convexity, we show that both of these problems can be solved efficiently subject to certain mild conditions. More generally, we show how to solve a large family of *Stackelberg games* in which the utility function of the “follower” is unknown. A *Stackelberg game* is played by a *leader* and a *follower*. The leader moves first and commits to an action (e.g., setting prices or tolls as in

our examples), and then the follower *best responds*, playing the action that maximizes her utility given the leader’s action. The leader’s problem is to find the action that will optimize his objective (e.g., maximizing profit, or minimizing social cost as in our examples) after the follower best responds to this action.

Traditionally, Stackelberg games are solved assuming that the leader knows the follower’s utility function, and thus his own utility function. But this assumption is very strong, and in many realistic settings the follower’s utility function will be unknown. Our results give general conditions—and several natural examples—under which the problem of computing an optimal Stackelberg equilibrium can be solved efficiently with only revealed preferences feedback to the follower’s utility function.

For clarity of exposition, we first work out our solution in detail for the special case of profit maximization from revealed preferences. We then derive and state our general theorem for optimally solving a class of Stackelberg games where the follower’s utility is unknown. Finally, we show how to apply the general theorem to other problems, including the optimal tolling problem mentioned above and a natural principal-agent problem.

1.1 Our Results and Techniques

The main challenge in solving our class of Stackelberg games is that for many natural examples, the leader’s objective function is not concave when written as a function of his own action. For instance, in our example, the retailer’s profit is not concave as a function of the price he sets. Our first key ingredient is to show that in many natural settings, *the leader’s objective is concave when written as a function of the follower’s action*.

Consider again the retailer’s profit maximization problem. Recall that if the buyer’s valuation function $v(x) = \sqrt{x}$, then when she faces a price p , she will buy the bundle $x^*(p) = 1/4p^2$. In this simple case, we can see that setting a price of $p^*(x) = 1/2\sqrt{x}$ will induce the buyer to purchase x units. In principle, we can now write the retailer’s profit function as a function of the bundle x . In our example, the retailer’s cost function is simply $c(x) = x$. So,

$$\text{Profit}(x) = p^*(x) \cdot x - c(x) = \frac{\sqrt{x}}{2} - x.$$

Written in terms of x , the profit function is concave! As we show, this phenomenon continues in higher dimensions, for arbitrary convex cost functions c and for a wide class of concave valuation functions satisfying certain technical conditions, including the well studied families of CES and Cobb-Douglas utility functions.

Thus, if the retailer had access to an oracle for the concave function $\text{Profit}(x)$, we could use an algorithm for bandit concave optimization to maximize the retailer’s profit. Unfortunately, the retailer does not directly get to choose the bundle purchased by the buyer and observe the profit for that bundle: he can only set *prices* and observe the buyer’s chosen bundle $x^*(p)$ at those prices, and the resulting profit $\text{Profit}(x^*(p))$.

Nevertheless, we have reduced the retailer’s problem to a possibly simpler one. In order to find the profit maximizing prices, it suffices to give an algorithm which simulates access to an oracle for $\text{Profit}(x)$ given only the retailer’s query access to $x^*(p)$ and $\text{Profit}(x^*(p))$. Specifically, if for a given bundle x , the retailer could find prices p such that the buyer’s chosen bundle $x^*(p) = x$, then he could simulate access to $\text{Profit}(x)$ by setting prices p and receiving $\text{Profit}(x^*(p)) = \text{Profit}(x)$.

Our next key ingredient is a “tâtonnement-like” procedure that efficiently finds prices that approximately induce a target bundle x given only access to $x^*(p)$, provided that the buyer’s valuation function is Hölder continuous and strongly concave on the set of feasible bundles. Specifically, given a target bundle x , our procedure finds prices p such that $|\text{Profit}(x^*(p)) - \text{Profit}(x)| \leq \varepsilon$. Thus, we can use our procedure to simulate approximate access to the function $\text{Profit}(x)$. Our procedure requires only $\text{poly}(d, 1/\varepsilon)$ queries

to $x^*(p)$. Using recent algorithms for bandit optimization due to Belloni et al. [BLNR15], we can maximize the retailer’s profits efficiently even with only approximate access to $\text{Profit}(x)$. When our algorithms receive *noiseless* feedback, we can improve the dependence on the approximation parameter ε to be only $\text{poly}(\log 1/\varepsilon)$.

A similar approach can be used to solve the optimal tolling problem assuming the unknown latency functions are convex and strictly increasing. As in the preceding example, the municipal authority’s objective function (social cost) is not convex in the tolls, but *is* convex in the induced flow. Whenever the latency function are strictly increasing, the *potential function* of the routing game is strongly convex, and so we can use our tâtonnement procedure to find tolls that induce target flows at equilibrium.

Our results for maximizing profits and optimizing tolls follow from a more general method that allows the leader in a large class of continuous action Stackelberg game to iteratively and efficiently maximize his objective function while only observing the follower’s response. The class requires the following conditions:

1. The follower’s utility function is strongly concave in her own actions and linear in the leader’s actions.
2. The leader’s objective function is concave when written as a function of the follower’s actions.¹

Finally, we show that our techniques are tolerant to two different kinds of noise. Our techniques work even if the follower only approximately maximizes his utility function, which corresponds to bounded, but adversarially chosen noise – and also if unbounded, but well behaved (i.e. zero mean and bounded variance) noise is introduced into the system. To illustrate this noise tolerance, we show how to solve a simple d -dimensional principal-agent problem, in which the principal contracts for the production of d types of goods that are produced as a stochastic function of the agent’s actions.

1.2 Related Work

There is a very large literature in operations research on solving so-called “bilevel programming” problems, which are closely related to Stackelberg games. Similar to a Stackelberg game, the variables in a bilevel programming problem are partitioned into two “levels.” The second-level variables are constrained to be the optimal solution to some problem defined by the first-level variables. See [CMS05] for a survey of the bilevel programming literature. Unlike our work, this literature does not focus substantially on computational issues (many of the algorithms are not polynomial time). [KCP10] show that optimally solving certain discrete Stackelberg games is NP-hard. Even ignoring computational efficiency, this literature assumes knowledge of the objective function of the “follower.” Our work departs significantly from this literature by assuming that the leader has no knowledge of the follower’s utility function.

There are two other works that we are aware of that consider solving Stackelberg games when the follower’s utility function is unknown. Letchford, Conitzer, and Munagala [LCM09] give algorithms for learning optimal leader strategies with a number of queries that is polynomial in the number of pure strategies of the leader. In our setting, the leader has a continuous and high dimensional action space, and so the results of [LCM09] do not apply. Blum, Haghtalab, and Procaccia [BHP14] consider the problem of learning optimal strategies for the leader in a class of *security games*. They exploit the structure of security games to learn optimal strategies for the leader in a number of queries that is polynomial in the representation size of the game (despite the fact that the number of pure strategies is exponential). The algorithm of [BHP14] is not computationally efficient – indeed, the problem they are solving is NP-hard. Neither of these techniques apply to our setting – and despite the fact that in our setting the leader has a continuous

¹When the leader and follower are instead trying to minimize a cost function, replace “concave” with “convex” in the above.

action space (which is exponentially large even under discretization), we are able to give an algorithm with both polynomial query complexity and polynomial running time.

There is also a body of related work related to our main example of profit maximization. Specifically, there is a recent line of work on *learning to predict from revealed preferences* ([BV06, ZR12, BDM⁺14]). In this line, the goal is to *predict* buyer behavior, rather than to optimize seller prices. Following these works, Amin et al. [ACD⁺15] considered how to find profit maximizing pricing from revealed preferences in the special case in which the buyer has a linear utility function and a fixed budget. The technique of [ACD⁺15] is quite specialized to linear utility functions, and does not easily extend to more general utility functions in the profit maximization problem, and not to Stackelberg games in general. “Revealed preferences” queries are quite similar to *demand queries* (see e.g. [BN09]). Demand queries are known to be sufficient to find welfare optimal allocations, and more generally, to be able to solve separable convex programs whose objective is social welfare. In contrast, our optimization problem is non-convex (and so the typical methodology by which demand queries are used does not apply), and our objective is not welfare.

The profit maximization application can be viewed as a dynamic pricing problem in which the seller has no knowledge of the buyers utilities. Babaioff et al. [BDKS15] study a version of this problem that is incomparable to our setting. On the one hand, [BDKS15] allow for *distributions* over buyers. On the other hand, [BDKS15] is limited to selling a single type of good, whereas our algorithms apply to selling bundles of many types of goods. There is also work related to our optimal tolling problem. In an elegant paper, Bhaskar et al. [BLSS14] study how one can iteratively find tolls such that a particular target flow is an equilibrium of a non-atomic routing game where the latency functions are unknown, which is a sub-problem we also need to solve in the routing application. Their technique is specialized to routing games, and requires that the unknown latency functions have a known simple functional form (linear or low-degree convex polynomial). In contrast, our technique works quite generally, and in the special case of routing games, does not require the latency functions to satisfy any known functional form (or even be convex). Our technique can also be implemented in a noise tolerant way, although at the expense of having a polynomial dependence on the approximation parameter, rather than a polylogarithmic dependence (in the absence of noise, our method can also be implemented to depend only polylogarithmically on the approximation parameter.)

Finally, our work is related in motivation to a recent line of work designed to study the *sample complexity of auctions* [BBHM08, CR14, HMR14, DHN14, CHN14, BMM15, MR15]. In this line of work, like in our work, the goal is to optimize an objective in a game theoretic setting when the designer has no direct knowledge of participant’s utility functions.

2 Preliminaries

We will denote the set of non-negative real numbers by $\mathbb{R}_+ = \{x \in \mathbb{R} \mid x \geq 0\}$ and the set of positive real numbers by $\mathbb{R}_{>0} = \{x \in \mathbb{R} \mid x > 0\}$. For a set $C \subseteq \mathbb{R}^d$ and a norm $\|\cdot\|$, we will use $\|C\| = \sup_{x \in C} \|x\|$ to denote the diameter of C with respect to the norm $\|\cdot\|$. When the norm is unspecified, $\|\cdot\|$ will denote the Euclidean norm $\|\cdot\|_2$.

An important concept we use is the *interior* of a set. In the following, we will use B_u to denote the unit ball centered at u for any $u \in \mathbb{R}^d$.

Definition 1. For any $\delta > 0$ and any set $C \subseteq \mathbb{R}^d$, the δ -interior $\text{Int}_{C,\delta}$ of C is a subset of C such that a point x is in the δ -interior $\text{Int}_{C,\delta}$ of C if the ball of radius δ centered at x is contained in C , that is:

$$x + \delta B_0 = \{x + \delta y \mid \|y\| \leq 1\} \subseteq C.$$

The interior Int_C of C is a subset of C such that a point x is in Int_C if there exists some $\delta' > 0$ such that x is in $\text{Int}_{C,\delta'}$.

We will also make use of the notions of Hölder continuity and Lipschitzness.

Definition 2. A function $f: C \rightarrow \mathbb{R}$ is (λ, β) -Hölder continuous for some $\lambda, \beta \geq 0$ if for any $x, y \in C$,

$$|f(x) - f(y)| \leq \lambda \|x - y\|^\beta.$$

A function f is λ -Lipschitz if it is $(\lambda, 1)$ -Hölder continuous.

2.1 Projected Subgradient Descent

A key ingredient in our algorithms is the ability to minimize a convex function (or maximize a concave function), given access only to the subgradients of the function (i.e. with a so-called “first-order” method). For concreteness, in this paper we do so using the projected sub gradient descent algorithm. This algorithm has the property that it is *noise-tolerant*, which is important in some of our applications. However, we note that any other noise-tolerant first-order method could be used in place of gradient descent to obtain qualitatively similar results. In fact, we show in the appendix that for applications that do not require noise tolerance, we can use the Ellipsoid algorithm, which obtains an exponentially better dependence on the approximation parameter. Because we strive for generality, in the body of the paper we restrict attention to gradient descent.

Let $C \subseteq \mathbb{R}^d$ be a compact and convex set that is contained in a Euclidean ball of radius R , centered at some point $x_1 \in \mathbb{R}^d$. Let $c: \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex “loss function.” Assume that c is also λ -Lipschitz—that is, $|c(x) - c(y)| \leq \lambda \|x - y\|_2$. Let Π_C denote the projection operator onto C ,

$$\Pi_C(x) = \underset{y \in C}{\operatorname{argmin}} \|x - y\|.$$

Projected subgradient descent is an iterative algorithm that starts at $x_1 \in C$ and iterates the following equations

$$\begin{aligned} y_{t+1} &= x_t - \eta g_t, \text{ where } g_t \in \partial c(x_t) \\ x_{t+1} &= \Pi_C(y_{t+1}) \end{aligned}$$

The algorithm has the following guarantee.

Theorem 3. The projected subgradient descent algorithm with $\eta = \frac{R}{\lambda\sqrt{T}}$ satisfies

$$c\left(\frac{1}{T} \sum_{t=1}^T x_t\right) \leq \min_{y \in C} c(y) + \frac{R\lambda}{\sqrt{T}}$$

Alternatively, the algorithm finds a solution within ε of optimal after $T = (R\lambda/\varepsilon)^2$ steps.

2.2 Strong Convexity

We will make essential use of *strong convexity/concavity* of certain functions.

Definition 4. Let $\phi: C \rightarrow \mathbb{R}$ be a function defined over a convex set $C \subseteq \mathbb{R}^d$. We say ϕ is σ -strongly convex if for every $x, y \in C$,

$$\phi(y) \geq \phi(x) + \langle \nabla \phi(x), y - x \rangle + \frac{\sigma}{2} \cdot \|y - x\|_2^2.$$

We say ϕ is σ -strongly concave if $(-\phi)$ is σ -strongly convex.

An extremely useful property of strongly convex functions is that any point in the domain that is close to the minimum in objective value is also close to the minimum in Euclidean distance.

Lemma 5. Let $\phi: C \rightarrow \mathbb{R}$ be a σ -strongly convex function, and let $x^* = \operatorname{argmin}_{x \in C} \phi(x)$ be the minimizer of ϕ . Then, for any $x \in C$,

$$\|x - x^*\|_2^2 \leq \frac{2}{\sigma} \cdot (\phi(x) - \phi(x^*)).$$

Similarly, if ϕ is σ -strongly concave, and $x^* = \operatorname{argmax}_{x \in C} \phi(x)$, then for any $x \in C$,

$$\|x - x^*\|_2^2 \leq \frac{2}{\sigma} \cdot (\phi(x^*) - \phi(x)).$$

2.3 Tools for Zeroth-Order Optimization

We briefly discuss a useful tool for noisy zeroth-order optimization (also known as *bandit* optimization) by [BLNR15], which will be used as blackbox algorithm in our framework. The important feature we require, satisfied by the algorithm from [BLNR15] is that the optimization procedure be able to tolerate a small amount of adversarial noise.

Definition 6. Let C be a convex set in \mathbb{R}^d . We say that C is well-rounded if there exist $r, R > 0$ such that $\mathcal{B}_2^d(r) \subseteq C \subseteq \mathcal{B}_2^d(R)$ and $R/r \leq O(\sqrt{d})$, where $\mathcal{B}_2^d(\gamma)$ denotes an ℓ_2 ball of radius γ in \mathbb{R}^d .

Let C be a well-rounded convex set in \mathbb{R}^d and $F, f: \mathbb{R}^d \rightarrow \mathbb{R}$ be functions such that f is convex and F satisfies

$$\sup_{x \in C} |F(x) - f(x)| \leq \varepsilon/d, \tag{1}$$

for some $\varepsilon > 0$. The function F can be seen as an oracle that gives a noisy evaluation of f at any point in C . Belloni et al. [BLNR15] give an algorithm that finds a point $x \in C$ that approximately optimizes the convex function f and only uses function evaluations of F at points in $x \in C$. The set C only needs to be specified via a membership oracle that decides if a point x is in C or not.

Lemma 7 ([BLNR15], Corollary 1). Let C be a well-rounded set in \mathbb{R}^d and f and F be functions that satisfy Equation (1). There is an algorithm **ZOO**(ε, C) (short for zeroth-order optimization) that makes $\tilde{O}(d^{4.5})$ calls² to F and returns a point $x \in C$ such that

$$\mathbb{E}[f(x)] \leq \min_{y \in C} f(y) + \varepsilon.$$

Naturally, the algorithm can also be used to approximately maximize a concave function.

²The notation $\tilde{O}(\cdot)$ hides the logarithmic dependence on d and $1/\varepsilon$.

3 Profit Maximization From Revealed Preferences

3.1 The Model and Problem Setup

Consider the problem of maximizing profit from revealed preferences. In this problem, there is a *producer*, who wants to sell a bundle x of d divisible goods to a *consumer*. The bundles are vectors $x \in C$ where $C \subseteq \mathbb{R}_+^d$ is some set of *feasible bundles* that we assume is *known* to both the producer and consumer.

- The producer has an *unknown cost function* $c : \mathbb{R}_+^d \rightarrow \mathbb{R}_+$. He is allowed to set prices $p \in \mathbb{R}_+^d$ for each good, and receives profit

$$r(p) = \langle p, x^*(p) \rangle - c(x^*(p)),$$

where $x^*(p)$ is the bundle of goods the consumer purchases at prices p . His goal is to find the profit maximizing prices

$$p^* = \operatorname{argmax}_{p \in \mathbb{R}_+^d} r(p).$$

- The consumer has a *valuation function* $v : \mathbb{R}_+^d \rightarrow \mathbb{R}_+$. The valuation function is *unknown to the producer*. The consumer has a *quasi-linear utility function* $u(x, p) = v(x) - \langle p, x \rangle$. Given prices p , the consumer will buy the bundle $x^*(p) \in C$ that maximizes her utility. Thus,

$$x^*(p) = \operatorname{argmax}_{x \in C} u(x, p) = \operatorname{argmax}_{x \in C} (v(x) - \langle x, p \rangle).$$

We call $x^*(p)$ the *induced bundle at prices* p .

In our model, in each time period t the producer will choose prices p^t and can observe the resulting induced bundle $x^*(p^t)$ and profit $r(p^t)$. We would like to design an algorithm so that after a polynomial number of observations T , the profit $r(p^T)$ is nearly as large as the optimal profit $r(p^*)$.

We will make several assumptions about the functions c and v and the set C . We view these assumptions as comparatively mild:

Assumption 3.1 (Set of Feasible Bundles). *The set of feasible bundles $C \subseteq \mathbb{R}_+^d$ is convex and well-rounded. It also contains the set $(0, 1]^d \subseteq C$ (the consumer can simultaneously buy at least one unit of each good). Also, $\|C\|_2 \leq \gamma$ (e.g. when $C = (0, 1]^d$, we have $\gamma = \sqrt{d}$). Lastly, C is downward closed, in the sense that for any $x \in C$, there exists some $\delta \in (0, 1)$ such that $\delta x \in C$ (the consumer can always choose buy less of each good).*

Assumption 3.2 (Producer's Cost Function). *The producer's cost function $c : \mathbb{R}_+^d \rightarrow \mathbb{R}$ is convex and Lipschitz-continuous.*

Assumption 3.3 (Consumer's Valuation Function). *The consumer's valuation function $v : \mathbb{R}_+^d \rightarrow \mathbb{R}$ is non-decreasing, Hölder-continuous, differentiable and strongly concave over C . For any price vector $p \in \mathbb{R}_+^d$, the induced bundle $x^*(p) = \operatorname{argmax}_{x \in C} u(x, p)$ is defined.*

Note that without the assumption that the consumer's valuation function is concave and that the producer's cost function is convex, even with full information, their corresponding optimization problems would not be polynomial time solvable. Our fourth assumption of *homogeneity* is more restrictive, but as we observe, is satisfied by a wide range of economically meaningful valuation functions including CES and Cobb-Douglas utilities. Informally, homogeneity is a scale-invariance condition — changing the *units* by which quantities of goods are measured should have a predictable multiplicative effect on the buyer valuation functions:

Definition 8. For $k \geq 0$, a function $v : \mathbb{R}_+^d \rightarrow \mathbb{R}_+$ is homogeneous of degree k if for every $x \in \mathbb{R}^d$ and for every $\sigma > 0$,

$$v(\sigma x) = \sigma^k v(x).$$

The function v is simply homogeneous if it is homogeneous of degree k for some $k \geq 0$.

Our fourth assumption is simply that the buyer valuation function is homogeneous of *some* degree:

Assumption 3.4. *The consumer's valuation function v is homogeneous.*

3.2 An Overview of Our Solution

We present our solution in three main steps:

1. First, we show that the profit function can be expressed as a concave function $r(x)$ of the consumer's induced bundle x , rather than as a (non-concave) function of the prices.
2. Next, we show that for a given candidate bundle x , we can iteratively find prices p such that $x \approx x^*(p)$. That is, in each time period s we can set prices p^s and observe the purchased bundle $x^*(p^s)$, and after a polynomial number of time periods S , we are guaranteed to find prices $p = p^S$ such that $x^*(p) \approx x$. Once we have found such prices, we can observe the profit $r(x^*(p)) \approx r(x)$, which allows us to simulate query access to $r(x)$.
3. Finally, we use our simulated query access to $r(x)$ as feedback to a bandit concave optimization algorithm, which iteratively queries *bundles* x , and quickly converges to the profit maximizing bundle.

3.3 Expressing Profit as a Function of the Bundle

First, we carry out Step 1 above and demonstrate how to rewrite the profit function as a function of the bundle x , rather than as a function of the prices p . Note that for any given bundle $x \in C$, there might be multiple price vectors that induce x . We denote the set of price vectors that induce x by:

$$P^*(x) = \{p \in \mathbb{R}^d \mid x^*(p) = x\}.$$

We then define the profit of a bundle x to be

$$r(x) = \max_{p \in P^*(x)} r(p) = \max_{p \in P^*(x)} \langle p, x \rangle - c(x).$$

Observe that the profit maximizing price vector $p \in P^*(x)$ is the price vector that maximizes *revenue* $\langle p, x \rangle$, since the *cost* $c(x)$ depends only on x , and so is the same for every $p \in P^*(x)$. The following lemma characterizes the revenue maximizing price vector that induces any fixed bundle $x \in C$.

Lemma 9. *Let $\hat{x} \in C$ be a bundle, and $P^*(\hat{x})$ be the set of price vectors that induce bundle \hat{x} . Then the price vector $p = \nabla v(\hat{x})$ is the revenue maximizing price vector that induces \hat{x} . That is, $\nabla v(\hat{x}) \in P^*(\hat{x})$ and for any price vector $p' \in P^*(\hat{x})$, $\langle p', \hat{x} \rangle \leq \langle \nabla v(\hat{x}), \hat{x} \rangle$.*

Proof. Observe that for any $x \in C$ the gradient of the consumer's utility $u(x, p) = v(x) - \langle p, x \rangle$ with respect to x is $(\nabla v - p)$. If the prices are $p = \nabla v(\hat{x})$, then since v is concave and $\nabla v(\hat{x}) - p = \mathbf{0}$, \hat{x} is a maximizer of the consumer's utility function. Thus, we have $x^*(\nabla v(\hat{x})) = \hat{x}$, and so $\nabla v(\hat{x}) \in P^*(\hat{x})$.

Suppose that there exists another price vector $p' \in P^*(\hat{x})$ such that $p' \neq \nabla v(\hat{x})$. Since the function $u(\cdot, p')$ is concave in x and $\hat{x} \in \arg \max_{x \in C} u(x, p')$, we know that for any $x' \in C$

$$\langle \nabla v(\hat{x}) - p', x' - \hat{x} \rangle \leq 0,$$

otherwise there is a feasible ascent direction, which contradicts the assumption that \hat{x} maximizes $u(x, p')$. By 3.1, we know there exists some $\delta < 1$ such that $\delta \hat{x} \in C$. Now consider $x' = \delta \hat{x}$, then it follows that

$$\langle \nabla v(\hat{x}) - p', (1 - \delta) \hat{x} \rangle = (1 - \delta) (\langle \nabla v(\hat{x}), \hat{x} \rangle - \langle p', \hat{x} \rangle) \geq 0.$$

Therefore, $\langle p', x \rangle \leq \langle \nabla v(x), x \rangle$, as desired. This completes the proof. \square

With this characterization of the revenue maximizing price vector, we can then rewrite the profit as a function of x in closed form for any $x \in C$:

$$r(x) = \langle \nabla v(x), x \rangle - c(x). \quad (2)$$

Next, we show that $r(x)$ is a concave function of x whenever the valuation v satisfies 3.3 (concavity and differentiability) and 3.4 (homogeneity).

Theorem 10. *If the consumer's valuation function v is differentiable, homogeneous, and concave over C , the producer's profit function $r(x) = \langle \nabla v(x), x \rangle - c(x)$ is concave over the domain C .*

To prove this result, we invoke Euler's theorem for homogeneous functions:

Theorem 11 (Euler's Theorem for Homogeneous Functions). *Let $v : C \rightarrow \mathbb{R}_+$ be continuous and differentiable. Then v is homogeneous of degree k if and only if*

$$\langle \nabla v(x), x \rangle = k \cdot v(x).$$

Proof of Theorem 10. Recall that:

$$r(x) = \langle \nabla v(x), x \rangle - c(x)$$

By the assumption that v is continuous, differentiable, and homogeneous of some degree $k \geq 0$, we have by Euler's theorem that

$$r(x) = kv(x) - c(x)$$

Because by assumption, $v(x)$ is concave, and $c(x)$ is convex, we conclude that $r(x)$ is concave. \square

Finally, we note that many important and well studied classes of valuation functions satisfy our assumptions – namely differentiability, strong concavity and homogeneity. Two classes of interest include

- **Constant Elasticity of Substitution (CES).** Valuation functions of the form:

$$v(x) = \left(\sum_{i=1}^d \alpha_i x_i^\rho \right)^\beta,$$

where $\alpha_i > 0$ for every $i \in [d]$ and $\rho, \beta > 0$ such that $\rho < 1$ and $\beta\rho < 1$. These functions are known to be differentiable, Hölder continuous and strongly concave over the set $(0, H]^d$ (see Appendix B.1 for a proof). Observe that $v(\sigma x) = (\sum_{i=1}^d \alpha_i (\sigma x_i)^\rho)^\beta = \sigma^{\rho\beta} (\sum_{i=1}^d \alpha_i x_i^\rho)^\beta = \sigma^{\rho\beta} v(x)$, so these functions are homogeneous of degree $k = \rho\beta$.

- **Cobb-Douglas.** These are valuation functions of the form

$$v(x) = \prod_{i=1}^d x_i^{\alpha_i},$$

where $\alpha_i > 0$ for every $i \in [d]$ and $\sum_{i=1}^d \alpha_i < 1$. These functions are known to be differentiable, Hölder continuous and strongly concave over the set $(0, H]^d$ (see Appendix B.1 for a proof). Observe that $v(\sigma x) = \prod_{i=1}^d (\sigma x_i)^{\alpha_i} = (\prod_{i=1}^d \sigma^{\alpha_i}) (\prod_{i=1}^d x_i^{\alpha_i}) = \sigma^{\sum_{i=1}^d \alpha_i} \cdot v(x)$, so these functions are homogeneous of degree $k = \sum_{i=1}^d \alpha_i$.

3.4 Converting Bundles to Prices

Next, we carry out Step 2 and show how to find prices \hat{p} to induce a given bundle \hat{x} . Specifically, the producer has a target bundle $\hat{x} \in C$ in mind, and would like to learn a price vector $\hat{p} \in \mathbb{R}_+^d$ such that the induced bundle $x^*(\hat{p})$ is “close” to \hat{x} . That is,

$$\|\hat{x} - x^*(\hat{p})\|_2 \leq \varepsilon,$$

for some $\varepsilon > 0$.

Our solution will actually only allow us to produce a price vector \hat{p} such that \hat{x} and $x^*(\hat{p})$ are “close in value.” That is

$$|u(\hat{x}, \hat{p}) - u(x^*(\hat{p}), \hat{p})| \leq \delta.$$

However, by strong concavity of the valuation function, this will be enough to guarantee that the actual bundle is close to the target bundle. The following is just an elaboration of assumption 3.3:

Assumption 3.5 (Quantitative version of 3.3). *The valuation function v is both*

1. $(\lambda_{\text{val}}, \beta)$ -Hölder continuous over the domain C with respect to the ℓ_2 norm—for all $x, x' \in C$,

$$|v(x) - v(x')| \leq \lambda_{\text{val}} \cdot \|x - x'\|_2^\beta,$$

for some constants $\lambda_{\text{val}} \geq 1$ and $\beta \in (0, 1]$, and

2. σ -strongly concave over the interior of C —for all $x, x' \in C$,

$$v(x') \leq v(x) + \langle \nabla v(x), x' - x \rangle - (\sigma/2) \cdot \|x - x'\|_2^2.$$

Our algorithm **LearnPrice** (\hat{x}, ε) is given as Algorithm 1. We will prove:

Theorem 12. *Let $\hat{x} \in C$ be a target bundle and $\varepsilon > 0$. Then **LearnPrice** (\hat{x}, ε) outputs a price vector \hat{p} such that the induced bundle satisfies $\|\hat{x} - x^*(\hat{p})\| \leq \varepsilon$ and the number of observations it needs is no more than*

$$T = d \cdot \text{poly} \left(\frac{1}{\varepsilon}, \frac{1}{\sigma}, \gamma, \lambda_{\text{val}} \right).$$

Algorithm 1 Learning the price vector to induce a target bundle: **LearnPrice**(\hat{x}, ε)

Input: A target bundle $\hat{x} \in C$, and target accuracy ε

Initialize: restricted price space $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}L\}$ where

$$L = (\lambda_{\text{val}})^{1/\beta} \left(\frac{4}{\varepsilon^2 \sigma} \right)^{(1-\beta)/\beta} \quad p_j^1 = 0 \text{ for all good } j \in [d] \quad T = \frac{32dL^2\gamma^2}{\varepsilon^4\sigma^2} \quad \eta = \frac{\sqrt{2}\gamma}{L\sqrt{dT}}$$

For $t = 1, \dots, T$:

Observe the purchased bundle by the consumer $x^*(p^t)$

Update price vector with projected subgradient descent:

$$\hat{p}_j^{t+1} = p_j^t - \eta (\hat{x}_j - x^*(p^t)_j) \text{ for each } j \in [d], \quad p^{t+1} = \Pi_{\mathcal{P}} [\hat{p}^{t+1}]$$

Output: $\hat{p} = 1/T \sum_{t=1}^T p^t$.

To analyze **LearnPrice**(\hat{x}, ε), we will start by defining the following convex program whose solution is the target bundle \hat{x} .

$$\max_{x \in C} v(x) \tag{3}$$

$$\text{such that } x_j \leq \hat{x}_j \text{ for every good } j \in [d] \tag{4}$$

Since v is non-decreasing, it is not hard to see that \hat{x} is the optimal solution. The *partial Lagrangian* of this program is defined as follows,

$$\mathcal{L}(x, p) = v(x) - \sum_{j=1}^d p_j (x_j - \hat{x}_j),$$

where p_j is the dual variable for each constraint (4) and is interpreted as the price of good j . By strong duality, we know that there is a *value* OPT such that

$$\max_{x \in C} \min_{p \in \mathbb{R}_+^d} \mathcal{L}(x, p) = \min_{p \in \mathbb{R}_+^d} \max_{x \in C} \mathcal{L}(x, p) = \text{OPT} = v(\hat{x}). \tag{5}$$

We know that $\text{OPT} = v(\hat{x})$ because \hat{x} is the optimal solution to (3)-(4).

We can also define the *Lagrange dual function* $g: \mathbb{R}^d \rightarrow \mathbb{R}$ to be

$$g(p) = \max_{x \in C} \mathcal{L}(x, p).$$

We will show that an approximately optimal price vector for g approximately induces the target bundle \hat{x} , and that **LearnPrice**(\hat{x}, ε) is using projected subgradient descent to find such a solution to g . In order to reason about the convergence rate of the algorithm, we restrict the space of the prices to the following bounded set:

$$\mathcal{P} = \left\{ p \in \mathbb{R}_+^d \mid \|p\|_2 \leq \sqrt{d} (\lambda_{\text{val}})^{1/\beta} \left(\frac{4}{\varepsilon^2 \sigma} \right)^{(1-\beta)/\beta} \right\}. \tag{6}$$

First, we can show that the minimax value of the Lagrangian remains closed to OPT even if we restrict the prices to the set \mathcal{P} .

Lemma 13. *There exists a value R-OPT such that*

$$\max_{x \in C} \min_{p \in \mathcal{P}} \mathcal{L}(x, p) = \min_{p \in \mathcal{P}} \max_{x \in C} \mathcal{L}(x, p) = R\text{-OPT}.$$

Moreover, $v(\hat{x}) \leq R\text{-OPT} \leq v(\hat{x}) + \frac{\varepsilon^2 \sigma}{4}$.

Proof. Since C and \mathcal{P} are both convex and \mathcal{P} is also compact, the minimax theorem [Sio58] shows that there is a value R-OPT such that

$$\max_{x \in C} \min_{p \in \mathcal{P}} \mathcal{L}(x, p) = \min_{p \in \mathcal{P}} \max_{x \in C} \mathcal{L}(x, p) = R\text{-OPT}. \quad (7)$$

Since $\mathcal{P} \subseteq \mathbb{R}_+^d$, by (5), we have $R\text{-OPT} \geq v(\hat{x})$. Thus, we only need to show that $R\text{-OPT} \leq v(\hat{x}) + \alpha$, where $\alpha = \varepsilon^2 \sigma / 4$. Let (x^\bullet, p^\bullet) be a pair of minimax strategies for (7). That is

$$x^\bullet \in \operatorname{argmax}_{x \in C} \min_{p \in \mathcal{P}} \mathcal{L}(x, p) \quad \text{and} \quad p^\bullet \in \operatorname{argmin}_{p \in \mathcal{P}} \max_{x \in C} \mathcal{L}(x, p)$$

It suffices to show that $\mathcal{L}(x^\bullet, p^\bullet) \leq v(\hat{x}) + \alpha$. Suppose not, then we have

$$v(\hat{x}) + \alpha < \mathcal{L}(x^\bullet, p^\bullet) = \min_{p \in \mathcal{P}} \mathcal{L}(x^\bullet, p) = v(x^\bullet) - \max_{p \in \mathcal{P}} \langle p, x^\bullet - \hat{x} \rangle \leq v(x^\bullet).$$

Now consider the bundle y such that $y_j = \max\{x_j^\bullet, \hat{x}_j\}$ for each $j \in [d]$. It is clear that $v(y) \geq v(x^\bullet) > v(\hat{x})$. Let $L = (\lambda_{\text{val}})^{1/\beta} \left(\frac{4}{\varepsilon^2 \sigma}\right)^{(1-\beta)/\beta}$, then we can construct the following price vector $p' \in \mathcal{P}$ such that $p'_j = L$ for each good j with $x_j^\bullet > \hat{x}_j$, and $p'_j = 0$ for all other goods. Since we assume that v is $(\lambda_{\text{val}}, \beta)$ -Hölder continuous with respect to ℓ_2 norm, we have

$$v(x^\bullet) - v(\hat{x}) \leq v(y) - v(\hat{x}) \leq \lambda_{\text{val}} \|y - \hat{x}\|_2^\beta \leq \lambda_{\text{val}} \|y - \hat{x}\|_1^\beta$$

It follows that

$$\begin{aligned} v(\hat{x}) + \alpha &< \mathcal{L}(x^\bullet, p^\bullet) \leq \mathcal{L}(x^\bullet, p') \\ &= v(x^\bullet) - \langle p', x^\bullet - \hat{x} \rangle \\ &= v(x^\bullet) - \sum_{j: x_j^\bullet > \hat{x}_j} L (y - \hat{x}_j) \\ &= v(x^\bullet) - L \|y - \hat{x}\|_1 \leq v(y) - L \|y - \hat{x}\|_2 \end{aligned}$$

Suppose that $\|y - \hat{x}\|_2 \geq 1$ or $\beta = 1$, we know that $\|y - \hat{x}\|_2^\beta \leq \|y - \hat{x}\|_2$. This means $v(\hat{x}) + \alpha < v(y) - L \|y - \hat{x}\|_2^\beta \leq v(y) - \lambda_{\text{val}} \|y - \hat{x}\|_2^\beta \leq v(\hat{x})$, a contradiction.

Next suppose that $\|y - \hat{x}\|_2 < 1$ and $\beta \in (0, 1)$. We also have that

$$\begin{aligned} \alpha &< v(y) - v(\hat{x}) - L \|y - \hat{x}\|_2 \\ &\leq \lambda_{\text{val}} \|y - \hat{x}\|_2^\beta - L \|y - \hat{x}\|_2 \\ &\leq \lambda_{\text{val}} \|y - \hat{x}\|_2^\beta \left(1 - \frac{L}{\lambda_{\text{val}}} \|y - \hat{x}\|_2^{1-\beta}\right) \end{aligned}$$

Since $\alpha > 0$, it must be that $\left(1 - \frac{L}{\lambda_{\text{val}}} \|y - \hat{x}\|_2^{1-\beta}\right)$ is also positive, and so $\|y - \hat{x}\|_2 < \left(\frac{\lambda_{\text{val}}}{L}\right)^{1/(1-\beta)}$. By the choice of our L ,

$$\alpha < \lambda_{\text{val}} \left(\frac{\lambda_{\text{val}}}{L}\right)^{\beta/(1-\beta)} = \frac{\varepsilon^2 \sigma}{4} = \alpha$$

which is a contradiction. Therefore, the minimax value of (7) is no more than $v(\hat{x}) + \alpha$. \square

The preceding lemma shows that \hat{x} is a primal optimal solution (even when prices are restricted). Therefore, if $\hat{p} = \operatorname{argmin}_{p \in \mathcal{P}} g(p)$ are the prices that minimize the Lagrangian dual, we must have that $\hat{x} = x^*(\hat{p})$ is the induced bundle at prices \hat{p} . The next lemma shows that if p' are prices that approximately minimize the Lagrangian dual, then the induced bundle $x^*(p')$ is close to \hat{x} .

Lemma 14. *Let $p' \in \mathcal{P}$ be a price vector such that $g(p') \leq \min_{p \in \mathcal{P}} g(p) + \alpha$. Let $x' = x^*(p')$ be the induced bundle at prices p' . Then x' satisfies*

$$\|x' - \hat{x}\| \leq 2\sqrt{\alpha/\sigma}.$$

Proof. Let R-OPT denote the Lagrangian value when we restrict the price space to \mathcal{P} . From Lemma 13, we have that $\text{R-OPT} = \min_{p \in \mathcal{P}} g(p) \in [v(\hat{x}), v(\hat{x}) + \alpha]$. By assumption, we also have

$$g(p') = \mathcal{L}(x', p') \leq \text{R-OPT} + \alpha \leq v(\hat{x}) + 2\alpha.$$

Note that $\mathcal{L}(\hat{x}, p') = v(\hat{x}) - \langle p', \hat{x} - \hat{x} \rangle = v(\hat{x})$ and x' is the maximizer for $\mathcal{L}(\cdot, p')$, so it follows that

$$0 \leq \mathcal{L}(x', p') - \mathcal{L}(\hat{x}, p') \leq 2\alpha.$$

Since we know that v is a σ -strongly concave function over C , the utility function $u(\cdot, p') = v(\cdot) - \langle p', \cdot \rangle$ is also σ -strongly concave over C .³ Then we have the following by Lemma 5 and the above argument,

$$2\alpha \geq \mathcal{L}(x', p') - \mathcal{L}(\hat{x}, p') = u(x', p') - u(\hat{x}, p') \geq \frac{\sigma}{2} \|x' - \hat{x}\|^2 \quad (8)$$

This means $\|x' - \hat{x}\| \leq 2\sqrt{\alpha/\sigma}$. \square

Based on Lemma 14, we can reduce the problem of finding the appropriate prices to induce the target bundle to finding the approximate optimal solution to $\operatorname{argmin}_{p \in \mathcal{P}} g(p)$. Even though the function g is unknown to the producer (because v is unknown), we can still approximately optimize the function using projected subgradient descent if we are provided access to subgradients of g . The next lemma shows that the bundle $x^*(p)$ purchased by the consumer gives a subgradient of the Lagrange dual objective function at p .

Lemma 15. *Let p be any price vector, and $x^*(p)$ be the induced bundle. Then*

$$(\hat{x} - x^*(p)) \in \partial g(p).$$

³If $f(\cdot)$ is a σ -strongly concave function over C and $g(\cdot)$ is a concave function over C , then $(f + g)(\cdot)$ is a σ -strongly concave function over C .

Proof. Given $x' = \arg \max_{x \in [0,1]^d} \mathcal{L}(x, p)$, we know by the envelope theorem that a subgradient of g can be obtained as follows

$$\frac{\partial g}{\partial p_j} = \hat{x}_j - x'_j \quad \text{for each } j \in [d].$$

Note that x' corresponds to the induced bundle of p because

$$\begin{aligned} x' &= \operatorname{argmax}_{x \in C} \mathcal{L}(x, p) \\ &= \operatorname{argmax}_{x \in C} [v(x) - \langle p, x - \hat{x} \rangle] \\ &= \operatorname{argmax}_{x \in C} [v(x) - \langle p, x \rangle] \\ &= \operatorname{argmax}_{x \in C} u(x, p) = x^*(p) \end{aligned}$$

Therefore, the vector $(\hat{x} - x^*(p))$ is a subgradient of g at the price vector p . \square

Now that we know the subgradients of the function g at p can be easily obtained from the induced bundle purchased by the consumer, it remains to observe that Algorithm **LearnPrice** (\hat{x}, ε) is performing projected gradient descent on the Lagrange dual objective, and to analyze its convergence.

Proof of Theorem 12. By Lemma 14, it suffices to show that the price vector \hat{p} returned by projected gradient descent satisfies

$$g(\hat{p}) \leq \min_{p \in \mathcal{P}} g(p) + \frac{\varepsilon^2 \sigma}{4}.$$

Note that the set \mathcal{P} is contained in the ℓ_2 ball centered at $\mathbf{0}$ with radius L . Also, for each p^t , the subgradient we obtain is bounded: $\|\hat{x} - x^*(p^t)\| \leq \sqrt{\|\hat{x}\|^2 + \|x^*(p^t)\|^2} \leq \sqrt{2}\gamma$ since $\|C\| \leq \gamma$. Since we set

$$T = \frac{32dL^2\gamma^2}{\varepsilon^4\sigma^2} \quad \eta = \frac{\sqrt{2}\gamma}{L\sqrt{dT}}$$

we can apply the guarantee of projected gradient descent from Theorem 3, which gives:

$$g(\hat{p}) - \min_{p \in \mathcal{P}} g(p) \leq \frac{\sqrt{2}L\gamma}{\sqrt{T}} = \frac{\varepsilon^2\sigma}{4}$$

By Lemma 14, we know that the resulting bundle $x^*(\hat{p})$ satisfies that $\|\hat{x} - x^*(\hat{p})\| \leq \varepsilon$. \square

Remark 16. *Since noise tolerance is not required in this setting, it is possible to approximately induce the target bundle only using poly-logarithmically in $(1/\varepsilon)$ number of observations. We will give an ellipsoid-based variant of **LearnPrice** in Appendix D that achieves this guarantee.*

3.5 Profit Maximization

Finally, we will show how to combine the algorithm **LearnPrice** with the zeroth order optimization algorithm **ZOO** to find the approximate profit-maximizing price vector. At a high level, we will use **ZOO** to (approximately) optimize the profit function r over the bundle space and use **LearnPrice** to (approximately) induce the optimal bundle.

Before we show how to use **ZOO**, we will verify that if we run the algorithm **LearnPrice** to obtain prices \hat{p} that approximately induce the desired bundle x , and observe the revenue generated from prices \hat{p} , we will indeed obtain an approximation to the revenue function $r(x)$.

Recall from Lemma 9 that the profit function can be written as a function of the bundle

$$r(x) = \langle \nabla v(x), x \rangle - c(x)$$

as long as the producer uses the profit maximizing price vector $\nabla v(x)$ to induce the bundle x . However, the price vector returned by **LearnPrice** might not be the optimal price vector for the induced bundle. In order to have an estimate of the optimal profit for each bundle, we need to guarantee that prices returned by **LearnPrice** are the profit maximizing ones. To do that, we will restrict the bundle space that **ZOO** is optimizing over to be the interior of C . Now we show that for every bundle in the interior of C , there is a unique price vector that induces that bundle. Thus, these prices are the profit-maximizing prices inducing that bundle.

Lemma 17. *Let x' be a bundle in Int_C . Then $\nabla v(x')$ is the unique price vector that induces x' .*

Proof. Let p' be a price vector such that $x^*(p') = x$. Since $\text{Int}_C \subseteq C$, we must have

$$x' = \operatorname{argmax}_{x \in \text{Int}_C} [v(x) - \langle p', x \rangle].$$

By the definition of Int_C , we know that there exists some $\delta > 0$ such that the ball $\delta B_{x'}$ is contained in C . Now consider the function $f: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $f(x) = u(x, p')$. It follows that x' is a local optimum of f neighborhood $\delta B_{x'}$. Since f is continuously differentiable, we must have $\nabla f(x') = \mathbf{0}$ by first-order conditions. Therefore, we must have

$$\nabla f(x') = \nabla v(x') - p' = \mathbf{0},$$

which implies that $p' = \nabla v(x')$. □

Instead of using the interior itself, we will use a simple and efficiently computable proxy for the interior obtained by slightly shifting and contracting C .

Claim 18. *For any $0 < \delta < 1/2$, let the set*

$$C_\delta = (1 - 2\delta)C + \delta \mathbf{1},$$

where $\mathbf{1}$ denotes the d -dimensional vector with 1 in each coordinate. Given 3.1, C_δ is contained in the $(\delta/2)$ -interior of C . That is, $C_\delta \subseteq \text{Int}_{C, \delta/2}$.

Proof. Our goal is to show that $C_\delta + \delta B_0 \subseteq C$, where B_0 denote the unit ball centered at $\mathbf{0}$. Any point in $C_\delta + (\delta/2)B_0$ can be written as $x' + (\delta/2)y'$ for $x' \in C_\delta$ and $y' \in B_0$. We will show that $x' + (\delta/2)y' \in C$. Since $x' \in C_\delta$, there exists $x \in C$ such that

$$x' = (1 - 2\delta)x + \delta \mathbf{1}.$$

Since $y' \in B_0$, there exists $y \in (0, 1]^d$ such that

$$\frac{1}{2}y' = 2y - \mathbf{1}.$$

To see this, note that $(0, 1]^d$ contains a ball of radius $1/4$ whose center is $(1/2) \cdot \mathbf{1}$. By Assumption 3.1, C contains $(0, 1]^d$, so $y \in C$. Therefore for some $x, y \in C$,

$$\begin{aligned} x' + (\delta/2)y' &= (1 - 2\delta)x + \delta\mathbf{1} + 2\delta y - \delta\mathbf{1} \\ &= \underbrace{(1 - 2\delta)x + 2\delta y}_{\in C}, \end{aligned}$$

where we used convexity of C . Hence, $x' + (\delta/2)y' \in C$, as desired. \square

We will let **ZOO** operate on the set C_δ instead of C , and we first want to show that there is little loss in profit if we restrict the induced bundle to C_δ . The following is just a formal, quantitative version of of Assumption 3.2:

Assumption 3.6 (Quantitative version of Assumption 3.2). *The producer's cost function $c: \mathbb{R}_+^d \rightarrow \mathbb{R}$ is λ_{cost} -Lipschitz over the domain C with respect to the ℓ_2 norm: for $x, x' \in C$,*

$$|c(x) - c(x')| \leq \lambda_{\text{cost}} \|x - x'\|.$$

Given this assumption, the profit function is also Hölder continuous.

Lemma 19. *For any $x, y \in C$ such that $\|x - y\| \leq 1$, the following holds*

$$|r(x) - r(y)| \leq (\lambda_{\text{val}} + \lambda_{\text{cost}}) \|x - y\|^\beta.$$

Proof. Recall the revenue component of the profit function is $\langle \nabla v(x), x \rangle$. Since v is a concave and homogeneous function, we know that the homogeneity degree satisfies $k \leq 1$. (See Appendix B for a proof). By Euler's theorem (Theorem 11),

$$\langle \nabla v(x), x \rangle = k \cdot v(x). \quad (9)$$

Since v is $(\lambda_{\text{val}}, \beta)$ -Hölder continuous C , by Equation 9 we know that the revenue $\langle \nabla v(x), x \rangle$ is also λ_{val} -Hölder continuous over C . Furthermore, since the cost function c is λ_{cost} -Lipschitz over C , the profit function satisfies the following: for any $x, y \in C$ such that $\|x - y\| \leq 1$, we have

$$|r(x) - r(y)| \leq |\langle \nabla v(x), x \rangle - \langle \nabla v(y), y \rangle| + |c(x) - c(y)| \leq \lambda_{\text{val}} \|x - y\|^\beta + \lambda_{\text{cost}} \|x - y\|$$

Since $\|x - y\| \leq 1$, we know that $\|x - y\|^\beta \geq \|x - y\|$, so $|r(x) - r(y)| \leq (\lambda_{\text{val}} + \lambda_{\text{cost}}) \|x - y\|^\beta$. \square

We can bound the difference between the optimal profits in C_δ and C .

Lemma 20. *For any $0 < \delta \leq 1/3\gamma$,*

$$\max_{x \in C} r(x) - \max_{x \in C_\delta} r(x) \leq (3\delta\gamma)^\beta (\lambda_{\text{val}} + \lambda_{\text{cost}}).$$

Proof. Let $x^* \in \arg \max_{x \in C} r(x)$. We know that $(1 - 2\delta)x^* + \delta\mathbf{1} \in C_\delta$, and

$$\|x^* - (1 - 2\delta)x^* - \delta\mathbf{1}\| \leq \delta \|2x^* - \mathbf{1}\| \leq 3\delta\gamma.$$

By Lemma 19, we then have

$$r(x^*) - r((1 - \delta)x^* + \delta\mathbf{1}) \leq (3\delta\gamma)^\beta (\lambda_{\text{val}} + \lambda_{\text{cost}}).$$

Furthermore, we also know $\max_{x \in C_\delta} r(x) \geq r((1 - \delta)x^* + \delta\mathbf{1})$, so we have shown the bound above. \square

Now we focus on how to optimize the profit function r over the set C_δ . Recall the algorithm **ZOO** requires approximate evaluations for the profit function r . Such evaluations can be implemented using our algorithm **LearnPrice**: for each bundle $x \in C_\delta$, run **LearnPrice**(x, ε) to obtain a price vector p such that $\|x - x^*(p)\| \leq \varepsilon$, and then the resulting profit $r(x^*(p))$ serves as an approximate evaluation for $r(x)$:

$$|r(x) - r(x^*(p))| \leq (\lambda_{\text{val}} + \lambda_{\text{cost}})\varepsilon^\beta.$$

Algorithm 2 Learning the price vector to optimize profit: **Opro**(C, α)

Input: Feasible bundle space C , and target accuracy α

Initialize:

$$\varepsilon = \min \left\{ \left(\frac{\alpha}{\lambda(d+1 + (12\gamma)^\beta)} \right)^{1/\beta}, \frac{1}{12\gamma} \right\} \quad \delta = 4\varepsilon \quad \alpha' = d\varepsilon^\beta(\lambda_{\text{val}} + \lambda_{\text{cost}})$$

restricted bundle space $C_\delta = (1 - 2\delta)C + \delta\mathbf{1}$ and number of iterations $T = \tilde{O}(d^{4.5})$

For $t = 1, \dots, T$:

ZOO(α', C_δ) queries the profit for bundle x^t

Let $p^t = \mathbf{LearnPrice}(x^t, \varepsilon)$ and observe the induced bundle $x^*(p^t)$

Send $r(x^*(p^t))$ to **ZOO**(α', C_δ) as an approximate evaluation of $r(x^t)$

$\hat{x} = \mathbf{ZOO}(\alpha', C_\delta)$

$\hat{p} = \mathbf{LearnPrice}(\hat{x}, \varepsilon)$

Output: the last price vector \hat{p}

Theorem 21. Let $\alpha > 0$ be the target accuracy. The instantiation **Opro**(C, α) computes a price vector \hat{p} such that the expected profit

$$\mathbb{E}[r(\hat{p})] \geq \max_{p \in \mathbb{R}_+^d} r(p) - \alpha,$$

the number of times it calls the algorithm **LearnPrice** is bounded by $\tilde{O}(d^{4.5})$, and the total observations it requires from the consumer is $\text{poly}(d, 1/\alpha)$.⁴

Proof. First we show that each induced bundle $x^*(p^t)$ is in the interior Int_C . Note that in the algorithm, we have $\varepsilon = \delta/4$. By the guarantee of **LearnPrice** in Theorem 12, we have that

$$\|x^t - x^*(p^t)\| \leq \varepsilon = \delta/4.$$

By 18, we know that $x^t \in \text{Int}_{C, \delta/2}$, so the ball of radius ε centered at x^t is contained in C , and hence $x^*(p^t)$ is in the interior of C . By Lemma 17 and Lemma 9, each vector $p^t = \nabla v(x^*(p^t))$ is the profit-maximizing prices for the induced bundle $x^*(p^t)$, so the profit the algorithm observes is indeed $r(x^*(p^t))$.

Next, to establish the accuracy guarantee, we need to bound two sources of error. First, we need to bound the error from **ZOO**. To simplify notation, let $\lambda = (\lambda_{\text{val}} + \lambda_{\text{cost}})$. Recall from Lemma 19 that the approximate profit evaluation $r(x^*(p^t))$ satisfies

$$|r(x^t) - r(x^*(p^t))| \leq \lambda\varepsilon^\beta.$$

⁴ In Appendix D, we give a variant of the algorithm with query complexity scaling poly-logarithmically in $1/\alpha$.

By the accuracy guarantee in Lemma 7, the final queried bundle \hat{x} satisfies

$$\mathbb{E}[r(\hat{x})] \geq \max_{x \in C_\delta} r(x) - d\lambda\varepsilon^\beta.$$

Since we know that $|r(\hat{x}) - r(x^*(\hat{p}))| \leq \lambda\varepsilon$, we also have

$$\mathbb{E}[r(x^*(\hat{p}))] \geq \max_{x \in C_\delta} r(x) - (d+1)\lambda\varepsilon^\beta.$$

Next, as we are restricting the bundle space to C_δ , there might be further loss of profit. Note that $\delta = 4\varepsilon \leq 1/3\gamma$, so we can bound it with Lemma 20:

$$\mathbb{E}[r(x^*(\hat{p}))] \geq \max_{x \in C} r(x) - \lambda \left[(d+1)\varepsilon^\beta + (3\delta\gamma)^\beta \right] = \max_{x \in C} r(x) - \lambda \left[(d+1)\varepsilon^\beta + (12\varepsilon\gamma)^\beta \right].$$

If we plug in our setting for parameter ε , we recover the desired bound since $r(x^*(\hat{p})) = r(\hat{p})$ and $\max_{x \in C} r(x) = \max_{p \in \mathbb{R}_+^d} r(p)$.

Finally, we need to bound the total number of observations the algorithm needs from the consumer. In each iteration, the instantiation **LearnPrice**(x^t, ε) requires number of observations bounded by according to Theorem 12

$$T' = d \cdot \text{poly} \left(\frac{1}{\varepsilon}, \frac{1}{\sigma}, \gamma, \lambda_{\text{val}} \right)$$

Therefore, after plugging in ε , we have that the total number of observations **Opro** needs is bounded by

$$O(T' \times T) = \text{poly}(d, 1/\alpha)$$

(hiding constants $\lambda_{\text{cost}}, \lambda_{\text{val}}, \sigma, \gamma$). □

4 General Framework of Stackelberg Games

Now that we have worked out a concrete application of our method in the context of learning to maximize revenue from revealed preferences, we will abstract our techniques and show how they can be used to solve a general family of Stackelberg games in which the objective of the follower is unknown to the leader. Along the way, we will also generalize our technique to operate in a setting in which the follower responds to the leader's actions by only *approximately* maximizing her utility function. In addition to generalizing the settings in which our approach applies, this avoids a technical concern that might otherwise arise – that bundles maximizing strongly concave utility functions might be non-rational. In addition to being able to handle approximations to optimal bundles that would be induced by taking a rational approximation, we show our method is robust to much larger errors.

In our general framework, we consider a *Stackelberg game* that consists of a *leader* with action set \mathcal{A}_L and a *follower* with action set \mathcal{A}_F . Each player has a utility function $U_L, U_F: \mathcal{A}_L \times \mathcal{A}_F \rightarrow \mathbb{R}$. In the corresponding Stackelberg game, the leader chooses an action $p \in \mathcal{A}_L$, and then the follower chooses a ζ -best response $x'(p)$ such that

$$U_F(p, x'(p)) \geq U_F(p, x^*(p)) - \zeta,$$

where $x^*(p) = \arg\max_{x \in \mathcal{A}_F} U_F(p, x)$ is the follower's *exact* best-response. Note that when $\zeta = 0$, $x'(p) = x^*(p)$.

The example of maximizing revenue from revealed preferences is a special case of this framework. The producer is the leader and his action space consists of prices p and the follower is the consumer and her

action space is the bundle x she purchases. The producer's utility for a pair (p, x) is his revenue minus the cost of producing x and the consumer's utility is her value for x minus the price she pays.

In general, we consider solving the leader's optimization problem—find $p \in \mathcal{A}_L$ such that $U_L(p, x^*(p))$ is (approximately) maximized. Formally, we consider a sub-class of Stackelberg games that have the following structure.

Definition 22. *An instance is a Stackelberg game $\mathcal{S}(\mathcal{A}_L, \mathcal{A}_F, \phi)$ which consists of two players—the leader and the follower such that:*

- *the leader has action set $\mathcal{A}_L \subseteq \mathbb{R}^d$, the follower has action set $\mathcal{A}_F \subseteq \mathbb{R}^d$, both of which are convex and compact;*
- *the follower's utility function $U_F: \mathcal{A}_L \times \mathcal{A}_F \rightarrow \mathbb{R}$ takes the form*

$$U_F(p, x) = \phi(x) - \langle p, x \rangle,$$

where $\phi: \mathbb{R}^d \rightarrow \mathbb{R}$ is a strongly concave, differentiable function unknown to the leader;

- *the leader's utility function $U_L: \mathcal{A}_L \times \mathcal{A}_F \rightarrow \mathbb{R}$ is an unknown function.*

The optimization problem associated with the game instance is $\max_{p \in \mathcal{A}_L} \psi(p, x^(p))$.*

Our first step to solve the problem is to rewrite the leader's utility function so that it can be expressed as a function only in the follower's action. For each action of the follower $x \in \mathcal{A}_F$, the set of leader's actions that induce x is

$$P^*(x) = \{p \in \mathcal{A}_L \mid x^*(p) = x\}.$$

Among all of the leader's actions that induce x , the optimal one is:

$$p^*(x) = \operatorname{argmax}_{p \in P^*(x)} U_L(p, x),$$

where ties are broken arbitrarily. We can then rewrite the leader's objective as a function of only x :

$$\psi(x) = U_L(p^*(x), x). \tag{10}$$

Note that to approximately solve the leader's optimization problem, it is sufficient to find the follower's action $\hat{x} \in \mathcal{A}_F$ which approximately optimizes $\psi_F(\cdot)$, together with the action $\hat{p} \in \mathcal{A}_L$ that approximately induces \hat{x} . Before we present the algorithm, we state the assumptions on the utility functions of the two players that we will need.

Assumption 4.1. *The game $\mathcal{S}(\mathcal{A}_L, \mathcal{A}_F, \phi)$ satisfies the following properties.*

1. *The function $\psi: \mathcal{A}_L \rightarrow \mathbb{R}$ defined in (10) is concave and λ_L -Lipschitz;*
2. *The function $\phi: \mathcal{A}_F \rightarrow \mathbb{R}$ is non-decreasing, σ -strongly concave and λ_F -Lipschitz;*
3. *The action space of the leader \mathcal{A}_L contains the following set*

$$\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}\lambda_F\}; \tag{11}$$

4. *The action space of the follower \mathcal{A}_F has bounded diameter, $\|\mathcal{A}_F\| \leq \gamma$.*

4.1 Inducing a Target Action of the Follower

We first consider the following sub-problem. Given a target action \hat{x} of the follower we want to learn an action \hat{p} for the leader such that the induced action satisfies

$$\|x'(\hat{p}) - \hat{x}\| \leq \varepsilon.$$

We now give an algorithm to learn \hat{p} that requires only polynomially many observations of the follower's ζ -approximate best responses.

Algorithm 3 Learning the leader's action to induce a target follower's action: **LearnLead**(\hat{x}, ε)

Input: A target follower action $\hat{x} \in \mathcal{A}_F$, and target accuracy ε
Initialize: restricted action space $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}\lambda_F\}$

$$p_j^1 = 0 \text{ for all } j \in [d] \quad T = \left(\frac{16\sqrt{2d}\lambda_F\gamma}{\varepsilon^2\sigma - 4\zeta} \right)^2 \quad \eta = \frac{\sqrt{2}\gamma}{\sqrt{d}\lambda_F\sqrt{T}}$$

For $t = 1, \dots, T$:

Observe the induced action by the follower $x^*(p^t)$

Update leader's action:

$$\tilde{p}_j^{t+1} = p_j^t - \eta (\hat{x}_j - x^*(p^t)_j) \text{ for each } j \in [d], \quad p^{t+1} = \Pi_{\mathcal{P}} [\tilde{p}^{t+1}]$$

Output: $\hat{p} = 1/T \sum_{t=1}^T p^t$.

Theorem 23. Let $\hat{x} \in \mathcal{A}_F$ be a target follower action and $\varepsilon > 0$. Then **LearnLead**(\hat{x}, ε) outputs a leader action \hat{p} such that the induced follower action satisfies $\|\hat{x} - x'(\hat{p})\| \leq \varepsilon$ and the number of observations it needs is no more than

$$T = O\left(\frac{d\lambda_F^2\gamma^2}{\varepsilon^4\sigma^2}\right)$$

as long as $\varepsilon > 2\sqrt{2\zeta}/\sigma$.

4.2 Optimizing Leader's Utility

Now that we know how to approximately induce any action of the follower using **LearnLead**, we are ready to give an algorithm to optimize the leader's utility function U_L . Recall that we can write the U_L as a function ψ that depends only of the follower's action. In order to obtain the approximately optimal utility $\psi(x)$, the leader must play the *optimal* action p that induces the follower to play approximately x .

Assumption 4.2. For any $\hat{x} \in \mathcal{A}_F$ and $\varepsilon > 0$, the instantiation **LearnLead**(\hat{x}, ε) returns \hat{p} such that

$$\hat{p} = p^*(x^*(\hat{p})).$$

Whenever this assumption holds, we can use **LearnLead** to allow the leader to obtain utility $U_L(\hat{p}, x^*(\hat{p})) = \psi(x^*(\hat{p}))$.

While 4.2 appears to be quite strong, we can often achieve it. Recall that we were able to satisfy 4.2 in our revealed preferences application by operating in the interior of the feasible region of the follower’s action space, and we can similarly do this in our principal-agent example. Moreover, it is trivially satisfied whenever the leader’s objective function *depends only on the follower’s action*, since in this case, every leader-action p which induces a particular follower-action x is optimal. This is the case, for example, in our routing games application in Section 5.

Now we will show how to use the algorithm **ZOO** to find an approximate optimal point for the function ψ . First, we will use **LearnLead** to provide approximate function evaluation for ψ at each $\hat{x} \in \mathcal{A}_F$: our algorithm first runs **LearnLead** (\hat{x}, ε) to learn a price vector \hat{p} , and we will use the observed function value on the induced follower’s approximate best response $\psi(x'(\hat{p}))$ as an approximation for $\psi(\hat{x})$. Since **LearnLead** guarantees that $\|x'(\hat{p}) - \hat{x}\| \leq \varepsilon$, by the Lipschitz property of ψ we have

$$|\psi(\hat{x}) - \psi(x'(\hat{p}))| \leq \lambda_L \varepsilon.$$

With these approximate evaluations, **ZOO** can then find a $(d\lambda_L\varepsilon)$ -approximate optimizer of ψ with only $\tilde{O}(d^{4.5})$ iterations by Lemma 7. The full algorithm is presented in Algorithm 4.

Algorithm 4 Leader learn to optimize: **LearnOpt** (\mathcal{A}_F, α)

Input: Follower action space C , and target accuracy α

Initialize: number of iterations $T = \tilde{O}(n^{4.5})$ and $\varepsilon = \frac{\alpha}{\lambda_L(d+1)}$

For $t = 1, \dots, T$:

ZOO $(d\varepsilon\lambda_L, \mathcal{A}_F)$ queries the objective value for action $x^t \in \mathcal{A}_F$

 Let $p^t = \mathbf{LearnLead}(x^t, \varepsilon)$ and observe the induced action $x'(p^t)$

 Send $\psi(x'(p^t))$ to **ZOO** $(d\varepsilon\lambda_L, C_\delta)$ as an approximate evaluation of $\psi(x^t)$

$\hat{x} = \mathbf{ZOO}(d\varepsilon\lambda_L, \mathcal{A}_F)$

$\hat{p} = \mathbf{LearnLead}(\hat{x}, \varepsilon)$

Output: the leader action \hat{p}

Theorem 24. *Let $\alpha > 0$ be the target accuracy. The instantiation **LearnOpt** (\mathcal{A}_F, α) computes a leader action \hat{p} along with its induced follower action $x^*(\hat{p})$ that satisfies*

$$\mathbb{E}[U_L(\hat{p}, x^*(\hat{p}))] \geq \max_{p \in \mathcal{A}_L} U_L(p, x^*(p)) - \alpha,$$

and the number of observations the algorithm requires of the follower is bounded by

$$\tilde{O}\left(\frac{d^{9.5}}{\alpha^4}\right),$$

as long as $\alpha \geq \Omega(d\lambda_L\sqrt{\zeta/\sigma})$.

5 Optimal Traffic Routing from Revealed Behavior

In this section, we give the second main application of our technique discussed in the introduction: how to find tolls to induce an approximately optimal flow in a non-atomic traffic routing game when the latency functions are unknown.

A nonatomic routing game $\mathcal{G}(G, \ell, \mathcal{D})$ is defined by a graph $G = (V, E)$, latency function ℓ_e on each edge $e \in E$, and the source, destination and demands for n commodities: $\mathcal{D} = \{(s_i, t_i, k_i)\}_{i \in [n]}$. The latency function $\ell_e: \mathbb{R}_+ \rightarrow [0, 1]$ represents the delay on each edge e as a function of the total flow on that edge. For simplicity, we assume $\sum_{i=1}^n k_i = 1$, and we let m denote the number of edges $|E|$.

For each commodity i , the demand k_i specifies the volume of flow from s_i to t_i routed by (self-interested) agents. The game is nonatomic: infinitely many agents each control only an infinitesimal amount of flow and each agent of type i selects an action (an s_i - t_i path) so as to minimize her total latency. The aggregate decisions of the agents induce a multicommodity flow $(f^i)_{i \in [n]}$, with each vector $f^i = (f_e^i)_{e \in E} \in \mathcal{F}_i$, where \mathcal{F}_i is the flow polytope for the i 'th commodity:

$$\mathcal{F}_i = \left\{ f^i \in \mathbb{R}_+^m \mid \sum_{(v,w) \in E} f_{vw}^i = \sum_{(u,v) \in E} f_{uv}^i, \forall v \in V \setminus \{s_i, t_i\}, \sum_{(s_i,w) \in E} f_{s_i w}^i - \sum_{(u,s_i) \in E} f_{u s_i}^i = k_i \right\}$$

Let $\mathcal{F} = \{f = \sum_{i=1}^n f^i \mid f^i \in \mathcal{F}_i \text{ for each } i\}$ denote the set of feasible flows. A flow f defines a latency $\ell_e(f_e)$ on each edge e . Given a path P , we write $\ell_P(f) = \sum_{e \in P} \ell_e(f_e)$ to denote the sum latency on all edges in the path. A Nash or Wardrop equilibrium is defined as follows:

Definition 25 (Wardrop equilibrium). *A multicommodity flow \hat{f} is a Wardrop equilibrium of a routing game if it is feasible and for every commodity i , and for all s_i - t_i paths P, Q with $f_P^i > 0$, we have $\ell_P(\hat{f}) \leq \ell_Q(\hat{f})$.*

Crucial to our application is the following well known lemma, which states that a Wardrop equilibrium can be found as the solution to a optimization problem (convex whenever the latencies are non-decreasing), which minimizes a potential function associated with the routing game

Lemma 26 ([MS96]). *A Wardrop equilibrium can be computed by solving the following optimization problem:*

$$\min_{f \in \mathcal{F}} \Phi(f) := \sum_e \int_0^{f_e} \ell_e(x) dx$$

Whenever the latency functions ℓ_e are each non-decreasing, this is a convex program. We call Φ the potential function of the routing game.

Now suppose there is a municipal authority which administers the network and wishes to minimize the social cost of the equilibrium flow:

$$\Psi(f) = \sum_{e \in E} f_e \cdot \ell_e(f_e).$$

The authority has the power to impose constant tolls on the edges. A toll vector $\tau = (\tau_e)_{e \in E} \in \mathbb{R}_+^m$ induces a new latency function on each edge: $\ell_e^\tau(f_e) = \ell_e(f_e) + \tau_e$, which gives rise to a different routing game $\mathcal{G}(G, \ell^\tau, \mathcal{D})$ with a new potential function Φ^τ . In particular, the equilibrium flow $f^*(\tau)$ induced by the toll vector is the Wardrop equilibrium of the tolled routing game:

$$f^*(\tau) = \operatorname{argmin}_{f \in \mathcal{F}} \Phi^\tau(f) = \operatorname{argmin}_{f \in \mathcal{F}} \left[\sum_{e \in E} \int_0^{f_e} (\ell_e(x) + \tau_e) dx \right] = \operatorname{argmin}_{f \in \mathcal{F}} \left[\Phi(f) + \sum_{e \in E} \tau_e \cdot f_e \right].$$

While the latency functions are unknown to the authority, his goal is to find a toll vector $\hat{\tau}$ such that the induced flow $f^*(\hat{\tau})$ approximately minimizes the total congestion function Ψ .

We can formulate this problem as an instance of the type of Stackelberg game we defined in Definition 22, where the authority is the leader, and there is a single “flow” player minimizing the game’s potential function, serving the role of the follower. We will refer to them as the toll player and the flow player respectively. In our setting:

1. The toll player has action set $\tau \in \mathbb{R}_+^m$ and the flow player has action set \mathcal{F} ;
2. The flow player has a utility function $U_F: \mathbb{R}_+^m \times \mathcal{F} \rightarrow \mathbb{R}$ of the form

$$U_F(\tau, f) = -\Phi(f) - \langle \tau, f \rangle;$$

3. The toll player has a utility function $U_L: \mathbb{R}_+^m \times \mathcal{F} \rightarrow \mathbb{R}$ of the form

$$U_L(\tau, f) = -\Psi(f).$$

Now we will apply the tools in Section 4 to solve this problem. Before we begin, we will impose the following assumptions on the latency functions to match with 4.1. We need two types of assumptions: one set to let us find tolls to induce a target flow, and another to guarantee that once we can induce such flows (and hence implement a “flow cost oracle”), we can optimize over flows.

To find tolls to induce a target flow, we require that the potential function Φ be strongly convex in the flow variables. The following conditions are sufficient to guarantee this:

Assumption 5.1. *For each edge $e \in E$, ℓ_e is differentiable and has derivative bounded away from zero: there exists some $\sigma > 0$ such that for all $x \in [0, 1]$, $\ell'_e(x) \geq \sigma$.*

Recall that the potential function $\Phi(x)$ is a function on m variables $(f_e)_{e \in E}$, and it’s Hessian $\nabla^2 \Phi$ at each $f \in \mathcal{F}$ is a diagonal matrix with entries $\ell'_e(f_e) \geq \sigma$. Therefore, we know that $\nabla^2 \Phi(f) \succeq \sigma I$ for any $f \in \mathcal{F}$, and so under Assumption 5.1, Φ is a σ -strongly convex function over \mathcal{F} . Note that the only condition we really require is that the potential function be strongly convex, and there are weaker conditions that imply this, but we state Assumption 5.1 because of its simplicity.

Once we can implement a flow oracle, we need to be able to use a bandit convex optimization algorithm to optimize social cost over flows. Hence, we require that the social cost function be convex and Lipschitz. The following assumptions are sufficient to guarantee this:

Assumption 5.2. *For each edge $e \in E$, ℓ_e is convex and (λ/m) -Lipschitz continuous over $[0, 1]$.*

Note that this guarantees that Ψ is λ -Lipschitz over \mathcal{F} .

We first show that we can use the algorithm **LearnLead** to learn a toll vector to induce any flow as a Wardrop equilibrium.

Lemma 27. *Fix any non-atomic routing game satisfying Assumption 5.1. Let $\hat{f} \in \mathcal{F}$ in a target flow and $\varepsilon > 0$. Then the instantiation **LearnLead** (\hat{f}, ε) outputs a toll vector $\hat{\tau}$ such that the induced Wardrop equilibrium flow $f^*(\hat{\tau})$ satisfies $\|\hat{f} - f^*(\hat{\tau})\| \leq \varepsilon$, and the number of observations on the flow behavior it needs is no more than*

$$O\left(\frac{m^3}{\varepsilon^4 \sigma^2}\right).$$

Proof. Before we apply Theorem 23, we still need to show that the potential function Φ of the original routing game (without tolls) is Lipschitz over \mathcal{F} . Note that this does not require any assumptions on the latency functions ℓ_e other than that they are bounded in $[0, 1]$. Let $f, g \in \mathcal{F}$, then we can write

$$\begin{aligned} |\Phi(f) - \Phi(g)| &= \left| \sum_e \left(\int_0^{f_e} \ell_e(x) dx - \int_0^{g_e} \ell_e(x) dx \right) \right| \\ &= \left| \sum_{e \in E} \int_{g_e}^{f_e} \ell_e(x) dx \right| \\ &\leq \sum_{e \in E} \max\{\ell_e(f_e), \ell_e(g_e)\} |f_e - g_e| \\ &\leq \sum_e |f_e - g_e| \leq \sqrt{m} \|f - g\|, \end{aligned}$$

where the last inequality follows from the fact that $\|x\|_1 \leq \sqrt{m} \|x\|_2$ for any $x \in \mathbb{R}^m$. Also, observe that each flow vector in \mathcal{F} has norm bounded by \sqrt{m} . Therefore, we know that Φ is a \sqrt{m} -Lipschitz function. Then we can instantiate Theorem 23 and obtain the result above. \square

Now we can instantiate Theorem 24 and show that **LearnOpt** can find a toll vector that induces the approximately optimal flow.

Pre-processing Step The set \mathcal{F} is not a well-rounded convex body in \mathbb{R}^m (it has zero volume), so we will have to apply the following standard pre-processing step to transform it into a well-rounded body. First, we find a maximal set \mathcal{I} of linearly independent points in \mathcal{F} . We will then embed the polytope \mathcal{F} into this lower-dimensional subspace spanned by \mathcal{I} , so that \mathcal{F} becomes full-dimensional. In this subspace, \mathcal{F} is a convex body with a relative interior. Next, we apply the transformation of [LV06] to transform \mathcal{F} into a well-rounded body within $\text{Span}(\mathcal{I})$.⁵ We will run **ZOO** over the transformed body.

Lemma 28. *Let $\alpha > 0$ be the target accuracy. The instantiation $\text{LearnOpt}(\mathcal{A}_F, \alpha)$ computes a toll vector $\hat{\tau}$ such that the induced flow $\hat{f} = f^*(\hat{\tau})$ is α -approximately optimal in expectation:*

$$\mathbb{E} \left[\Psi(\hat{f}) \right] \leq \min_{f \in \mathcal{F}} \Psi(f) + \alpha.$$

The total number of observations we need on the flow behavior is bounded by

$$\tilde{O} \left(\frac{m^{11.5}}{\alpha^4} \right).$$

Remark 29. *Just as with the profit maximization example, if we do not require noise tolerance, then we can improve the dependence on the approximation parameter α to be polylogarithmic. We show how to do this in the appendix.*

⁵See Section 5 of [LV06] for details of the rounding algorithm.

6 The Principal-Agent Problem

Our general framework applies even when the leader observes only the *noisy feedback* that arises when the follower only approximately maximizes her utility function. This corresponds to *adversarially chosen* noise of bounded magnitude. In this section, we show how to handle the natural setting in which the noise being added need not be bounded, but is well behaved – specifically has mean 0, and bounded variance. This can be used to model actual noise in an interaction, rather than a failure to exactly maximize a utility function. As a running example as we work out the details, we will discuss a simple *principal-agent* problem related to our profit-maximization example.

In a *principal-agent* problem, the principal (the leader) defines a contract by which the agent (the follower) will be paid, as a function of work produced by the agent. The key property of principal agent problems is that the agent is not able to deterministically produce work of a given quality. Instead, the agent chooses (and experiences cost as a function of) a level of *effort*, which stochastically maps to the quality of his work. However, the effort chosen by the agent is unobservable to the principal – only the quality of the finished product.

We consider a simple d -dimensional principal-agent problem, in which the result of the agent can be evaluated along d dimensions, each of which might require a different amount of effort. Since the agent knows how effort is stochastically mapped to realizations, we abstract away the agent’s choice of an “effort” vector, and instead (without loss of generality) view the agent as choosing a “target contribution” $x \in C \subseteq \mathbb{R}_+^d$ – the expected value of the agent’s ultimate contribution. The agent experiences some strongly convex cost $c(x)$ for producing a target contribution of x , but might nevertheless be incentivized to produce high quality contributions by the contract offered by the principal. However, the contribution that is actually realized (and that the principal observes) is a stochastically perturbed version of x : $\tilde{x} = x + \theta$, where $\theta \in \mathbb{R}^d$ is a noise vector sampled from the mean-zero Gaussian distribution $\mathcal{N}(\mathbf{0}, I)$.

The principal wants to optimize over the set of linear contracts: he will choose a price vector $p \in \mathbb{R}_+^d$, such that in response to the agent’s realized contribution \tilde{x} , the agent collects reward $\langle p, \tilde{x} \rangle$. His goal is to choose a price vector to optimize his expected value for the agent’s contribution, minus his own costs.

The agent’s strongly convex cost function $c: C \rightarrow \mathbb{R}_+$ is unknown to the principal. If the principal’s contract vector is p and the agent attempts to contribute x , then his utility is

$$U_a(p, x) = \langle p, (x + \theta) \rangle - c(x),$$

and his expected utility is just $u_a(p, x) = \mathbb{E}[U_a(p, x)] = \langle p, x \rangle - c(x)$. Fixing any price p , the agent will attempt to play the *induced* contribution vector: $x^*(p) = \operatorname{argmax}_{x \in C} (\langle p, x \rangle - c(x))$ in order to optimize his expected utility.

The principal has value v_i for each unit of contribution in the i -th dimension, and upon observing the realized contribution \tilde{x} , his utility is

$$u_p(p, \tilde{x}) = \langle v, \tilde{x} \rangle - \langle p, \tilde{x} \rangle = \langle v - p, \tilde{x} \rangle.$$

The principal’s goal is to find a price vector \hat{p} to (approximately) maximize his expected utility:

$$\mathbb{E}[u_p(p, x^*(p) + \theta)] = \mathbb{E}[\langle v - p, x^*(p) + \theta \rangle] = \langle v - p, x^*(p) \rangle.$$

This is an instantiation of our class of Stackelberg games in which the principal is the leader with action set \mathbb{R}_+^d and utility function $\psi(p, x) = \langle v - p, x \rangle$, and the agent is the follower with action set C and utility function $\phi(p, x) = \langle p, x \rangle - c(x)$. Indeed, in expectation, it is merely a “procurement” version of

our profit-maximization example. However, the crucial difference in this application (causing it to deviate from the general setting defined in Definition 22) is that the leader only gets to observe a noisy version of the follower’s best response at each round: $\tilde{x} = x^*(p) + \theta$. We will adapt the analysis from Section 3 and Section 4 to show that our algorithm is robust to noisy observations. We make the following assumptions, which correspond to the set of assumptions we made in our previous applications.

Assumption 6.1. *The following assumptions parallel 4.1 and 3.1.*

1. *The set of feasible contributions $C \subseteq \mathbb{R}_+^d$ is convex, closed, and bounded. It also contains the unit hypercube, $[0, 1]^d \subseteq C$ (the agent can simultaneously attempt to contribute at least one unit in each dimension) and in particular contains $\mathbf{0} \in \mathbb{R}^d$ (the agent can contribute nothing). Lastly, $\|C\|_2 \leq \gamma$;*
2. *the agent’s cost function c is homogeneous, 1-Lipschitz and σ -strongly convex;*
3. *the principal’s valuation vector has norm $\|v\| \leq 1$.*

6.1 Inducing the Agent’s Contribution Using Noisy Observations

We will first show that in general, **LearnLead** can learn the leader’s action which approximately induces any target follower action \hat{x} even if the algorithm only observes noisy perturbed best responses from the follower. This result holds in full generality, but we illustrate it by using the principal-agent problem.

First, given any target contribution \hat{x} , consider the following convex program similar to Section 3.4:

$$\min_{x \in C} c(x) \tag{12}$$

$$\text{such that } x_j \geq \hat{x}_j \text{ for every } j \in [d] \tag{13}$$

The Lagrangian of the program is

$$\mathcal{L}(x, p) = c(x) + \langle p, x - \hat{x} \rangle,$$

and the Lagrangian dual objective function is

$$g(p) = \min_{x \in C} \mathcal{L}(x, p).$$

By the same analysis used in the proof of Lemma 14, if we find a price vector $\hat{p} \in \mathcal{P}$ such that $g(\hat{p}) \geq \max_{p \in \mathcal{P}} g(p) - \alpha$, then we know that the induced contribution vector $x^*(\hat{p})$ satisfies $\|x^*(\hat{p}) - \hat{x}\| \leq \sqrt{2\alpha/\sigma}$. Now we show how to (approximately) optimize the function g based on the realized contributions of the agent, which correspond to mean-zero perturbations of the agent’s best response.

As shown in Lemma 15, a subgradient of g at price p is $(x^*(p) - \hat{x})$, but now since the principal only observes the realized contribution vector \tilde{x} , our algorithm does not have access to subgradients. However, we can still obtain an unbiased estimate of the subgradient: the vector $(\tilde{x} - \hat{x})$ satisfies $\mathbb{E}[\tilde{x} - \hat{x}] = (x^*(p) - \hat{x})$ because the noise vector is drawn from $\mathcal{N}(\mathbf{0}, I)$. This is sufficient to allow us to analyze **LearnLead** as stochastic gradient descent. The principal does the following: initialize $p^1 = \mathbf{0}$ and at each round $t \in [T]$, observes a realized contribution vector $\tilde{x}^t = x^*(p^t) + \theta^t$ and updates the contract prices as follows:

$$p^{t+1} = \Pi_{\mathcal{P}} [p^t + \eta(\tilde{x}^t - \hat{x})],$$

where each $\theta^t \sim \mathcal{N}(\mathbf{0}, I)$, η is a learning rate and $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}\}$; Finally, the algorithm outputs the average price vector $\hat{p} = 1/T \sum_{t=1}^T p^t$. We use the following standard theorem about the convergence guarantee for stochastic gradient descent (a more general result can be found in [NJLS09]).

Lemma 30. *With probability at least $1 - \beta$, the average vector \hat{p} output by stochastic gradient descent satisfies*

$$\max_{p \in \mathcal{P}} g(p) - g(\hat{p}) \leq O \left(\frac{\sqrt{d}}{\sqrt{T}} \left(\gamma + \sqrt{d} \log \left(\frac{Td}{\beta} \right) \right) \right).$$

Algorithm 5 Learning the price vector from noisy observations: **LearnPriceN**($\hat{x}, \varepsilon, \beta$)

Input: A target contribution $\hat{x} \in C$, target accuracy ε , and confidence parameter β

Initialize: restricted price space $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}\}$

$$p_j^1 = 0 \text{ for all } j \in [d] \quad T = \tilde{O} \left(\frac{d\gamma^2}{\varepsilon^4\sigma^2} \right) \quad \eta = \frac{\sqrt{2}\gamma}{\sqrt{d}\sqrt{T}}$$

For $t = 1, \dots, T$:

Observe the realized contribution by the agent $\tilde{x}^t = x^*(p^t) + \theta$, where $\theta \sim \mathcal{N}(\mathbf{0}, I)$

Update price vector:

$$\hat{p}_j^{t+1} = p_j^t + \eta (\hat{x}_j - \tilde{x}_j^t) \text{ for each } j \in [d], \quad p^{t+1} = \Pi_{\mathcal{P}} [\hat{p}^{t+1}]$$

Output: $\hat{p} = 1/T \sum_{t=1}^T p^t$.

Lemma 31. *Let $\hat{x} \in C$ be any target contribution vector. Then, with probability at least $1 - \beta$, the algorithm **LearnPriceN**($\hat{x}, \varepsilon, \beta$) outputs a contract price vector \hat{p} for the principal such that the induced contribution vector $x^*(\hat{p})$ satisfies*

$$\|\hat{x} - x^*(\hat{p})\| \leq \varepsilon,$$

and the number observations on the realized contributions of the agent it needs is no more than

$$T = \tilde{O} \left(\frac{d\gamma^2}{\varepsilon^4\sigma^2} \right).$$

6.2 Optimizing the Principal's Utility

Finally, we show how to optimize the principal's utility by combining **LearnPriceN** and **ZOO**.

Following from the same analysis of Lemma 9, we know that the principal's utility-maximizing price vector to induce expected contribution \hat{x} is $\nabla c(\hat{x})$. We can then rewrite the expected utility of the principal as a function of the attempted contribution of the agent:

$$u_p(x) = \langle v - \nabla c(x), x \rangle.$$

Since c is a homogeneous and convex function, by Theorem 10, u_p is a concave function.

Similar to Section 3.5, we will run **ZOO** to optimize over the interior subset:

$$C_\delta = (1 - 2\delta)C + \delta\mathbf{1},$$

so any price vector \hat{p} given by **LearnPriceN** is the unique price that induces the agent's attempted contribution vector $x^*(\hat{p})$ (Lemma 17). By the same analysis of Lemma 20, we know that there is little loss in principal's utility by restricting the contribution vectors to C_δ .

Lemma 32. *The function $u_p: C \rightarrow \mathbb{R}$ is 2-Lipschitz, and for any $0 < \delta < 1$,*

$$\max_{x \in C} u_p(x) - \max_{x \in C_\delta} u_p(x) \leq 6\delta\gamma.$$

Now we show how to use **LearnPriceN** to provide a noisy evaluation for u_p at each point of C_δ (scale of δ determined in the analysis). For each \hat{p} the **LearnPriceN** returns, the realized contribution vector we observe is $\tilde{x} = x^*(\hat{p}) + \theta$, so the utility experienced by the principal is

$$u_p(\hat{p}, \tilde{x}) = \langle v - \hat{p}, \tilde{x} \rangle.$$

We first demonstrate that $u_p(\hat{p}, \tilde{x})$ gives an unbiased estimate for u_p , and we can obtain an accurate estimate by taking the average of a small number realized utilities. In the following, let constant $a = \ln 2/(2\pi)$.

Lemma 33. *Let $x' \in C$ be the contribution vector such that $p' = \nabla c(x')$ is the unique price vector that induces x' . Let noise vectors $\theta^1, \dots, \theta^s \sim \mathcal{N}(\mathbf{0}, I)$ and $\tilde{x}^j = x' + \theta^j$ for each $j \in [s]$. Then with probability at least $1 - \beta$,*

$$\left| \frac{1}{s} \sum_{j=1}^s u_p(\hat{p}, \tilde{x}^j) - u_p(x') \right| \leq \sqrt{\frac{d}{s}} \sqrt{\frac{2}{a} \ln \frac{2}{\beta}}.$$

Proof. Let $b = v - p'$, then we can write

$$\begin{aligned} \frac{1}{s} \sum_{j=1}^s u_p(\hat{p}, \tilde{x}^j) - u_p(x') &= \frac{1}{s} \sum_{j=1}^s (\langle b, \tilde{x}^j \rangle - \langle b, x' \rangle) \\ &= \frac{1}{s} \sum_{j=1}^s \langle b, \theta^j \rangle \\ &= \frac{1}{s} \sum_{j=1}^s \sum_{i=1}^d b_i \theta_i^j \end{aligned}$$

Note that each θ_i^j is sampled from the Gaussian distribution $\mathcal{N}(0, 1)$, and we use the fact that if $X \sim \mathcal{N}(0, \sigma_1^2)$ and $Y \sim \mathcal{N}(0, \sigma_2^2)$ then $(bX + cY) \sim \mathcal{N}(0, b^2\sigma_1^2 + c^2\sigma_2^2)$. We can further derive that $\frac{1}{s} \sum_{j=1}^s \sum_{i=1}^d b_i \theta_i^j$ is a random variable with distribution $\mathcal{N}(0, \|b\|^2/s)$. Then we will use the following fact about Gaussian tails: let Y be a random variable sampled from distribution $\mathcal{N}(0, \iota^2)$ and $a = \ln 2/(2\pi)$, then for all $\zeta > 0$

$$\Pr[|Y| > \zeta] \leq 2 \exp(-a\zeta^2/\iota^2)$$

It follows that with probability at least $1 - \beta$, we have

$$\left| \frac{1}{s} \sum_{j=1}^s \sum_{i=1}^d b_i \theta_i^j \right| \leq \sqrt{\frac{\ln \frac{2}{\beta}}{as}} \|b\|.$$

Finally, note that we can bound $\|b\| = \|v - p'\| \leq \sqrt{2d}$, so replacing $\|b\|$ by $\sqrt{2d}$ recovers our bound. \square

Now we are ready to give the algorithm to optimize the principal's utility in Algorithm 6.

Algorithm 6 Learning the price vector to optimize under noisy observations: **OproN**(C, α, β)

Input: Feasible bundle space C , target accuracy α , and confidence parameter β

Initialize:

$$\varepsilon = \frac{\alpha}{12\gamma + 3d} \quad \delta = 2\varepsilon \quad \alpha' = 3d\varepsilon \quad \beta' = \beta/2T \quad s = \frac{2d \ln \frac{2}{\beta'}}{a\varepsilon^2}$$

restricted bundle space $C_\delta = (1 - 2\delta)C + \delta\mathbf{1}$ and number of iterations $T = \tilde{O}(d^{4.5})$

For $t = 1, \dots, T$:

ZOO(α', C_δ) queries the profit for bundle x^t

Let $p^t = \mathbf{LearnPriceN}(x^t, \varepsilon, \beta')$

For $j = 1, \dots, s$:

Principal post price p^t

Let $\tilde{x}^j(p^t)$ be the realized contribution and experiences utility $u(p^t, \tilde{x}^j(p^t))$

Send $\frac{1}{s} \sum_{j=1}^s u(p^t, \tilde{x}^j(p^t))$ to **ZOO**(α', C_δ) as an approximate evaluation of $u_p(x^t)$

$\hat{x} = \mathbf{ZOO}(\alpha', C_\delta)$

$\hat{p} = \mathbf{LearnPrice}(\hat{x}, \varepsilon)$

Output: the last price vector \hat{p}

Theorem 34. Let $\alpha > 0$ and $0 < \beta < 1/2$. With probability at least $1 - \beta$, the price vector \hat{p} output by **OproN**(C, α, β) satisfies

$$\mathbb{E}[u_p(\hat{p}, x^*(\hat{p}))] \geq \max_{p \in \mathcal{P}} u_p(p, x^*(p)) - \alpha,$$

and the number of observations on realized contributions is bounded by

$$\tilde{O}\left(\frac{d^{9.5}}{\alpha^4}\right).$$

Proof. First, by Lemma 31 and union bound, with probability at least $1 - \beta/2$, we have $\|x^t - x^*(p^t)\| \leq \varepsilon$ for all $t \in [T]$. We condition on this level of accuracy for the rest of the proof. By the same analysis of Footnote 4, we know that each target contribution $x^*(p^t)$ is in the interior Int_C , so we have that $u_p(x^*(p^t)) = u_p(p^t, x^*(p^t))$.

To establish the accuracy guarantee, we need to bound two sources of error. First, we need to bound the error from **ZOO**. Note that the target contribution $x^*(p^t)$ satisfies

$$|u_p(x^t) - u_p(x^*(p^t))| \leq 2\varepsilon.$$

By Lemma 33 and our setting of s , we have with probability at least $1 - \beta'$ that

$$\left| \frac{1}{s} \sum_{j=1}^s u_p(p^t, \tilde{x}^j(p^t)) - u_p(x^*(p^t)) \right| \leq \varepsilon.$$

By union bound, we know such accuracy holds for all $t \in [T]$ with probability at least $1 - \beta/2$. We condition on this level of accuracy, then the average utility provides an accurate evaluation for $u_p(x^t)$ at each queried point x^t

$$\left| \frac{1}{s} \sum_{j=1}^s u_p(p^t, \tilde{x}^j(p^t)) - u_p(x^t) \right| \leq 3\varepsilon.$$

By Lemma 7, we know that the vector \hat{x} output by **ZOO** satisfies

$$\mathbb{E}[u_p(\hat{x})] \geq \max_{x \in C_\delta} u_p(x) - 3d\varepsilon.$$

Finally, by Lemma 32 and the value of ε , we also have

$$\mathbb{E}[u_p(\hat{x})] \geq \max_{x \in C} u_p(x) - (12\varepsilon\gamma + 3d\varepsilon) = \max_{x \in C} u_p(x) - \alpha.$$

Note $\max_{x \in C} u_p(x) = \max_{p \in \mathcal{P}} u_p(p, x^*(p))$, so we have shown the accuracy guarantee. In each iteration, the algorithm requires $\tilde{O}\left(\frac{d\gamma^2}{\varepsilon^4\sigma^2}\right)$ noisy observations for running **LearnPriceN** and s observations for estimating $u_p(x^*(p^t))$, so the total number of observations is bounded by

$$\tilde{O}\left(d^{4.5} \times \left(\frac{\gamma^2 d(\gamma + d)^4}{\sigma^2 \alpha^4} + \frac{d(\gamma + d)^2}{\alpha^2}\right)\right) = \tilde{O}\left(\frac{d^{9.5}}{\alpha^4}\right)$$

where we hide constants σ, γ in the last equality. □

7 Conclusion

In this paper, we have given algorithms for optimally solving a large class of Stackelberg games in which the leader has only “revealed preferences” feedback about the follower’s utility function, with applications both to profit maximization from revealed preferences data, and optimal tolling in congestion games. We believe this is a very natural model in which to have access to agent utility functions, and that pursuing this line of work will be fruitful. There are many interesting directions, but let us highlight one in particular. In our profit maximization application, it would be very natural to consider a “Bayesian” version of our problem. At each round, the producer sets prices, at which point a new consumer, with valuation function drawn from an unknown prior, purchases her utility maximizing bundle. The producer’s goal is to find the prices that maximize her *expected* profit, over draws from the unknown prior. Under what conditions can we solve this problem efficiently? The main challenge (and the reason why it likely requires new techniques) is that the *expected value* of the purchased bundle need not maximize any well-behaved utility function, even if each individual consumer is maximizing a concave utility function.

Acknowledgements

We would like to thank Michael Kearns and Mallesh Pai for stimulating discussions about this work. In particular, we thank Mallesh for helpful discussions about principal-agent problems and conditions under which the profit function of the producer in the revealed preferences problem might be concave. We would also like to thank Tengyuan Liang and Alexander Rakhlin for very helpful discussions about [BLNR15].

References

- [ACD⁺15] Kareem Amin, Rachel Cummings, Lili Dworkin, Michael Kearns, and Aaron Roth. Online learning and profit maximization from revealed preferences. In *Twenty-Ninth AAAI Conference on Artificial Intelligence (AAAI-15)*, 2015.

- [BBHM08] Maria-Florina Balcan, Avrim Blum, Jason D Hartline, and Yishay Mansour. Reducing mechanism design to algorithm design via machine learning. *Journal of Computer and System Sciences*, 74(8):1245–1270, 2008.
- [BDKS15] Moshe Babaioff, Shaddin Dughmi, Robert Kleinberg, and Aleksandrs Slivkins. Dynamic pricing with limited supply. *ACM Transactions on Economics and Computation*, 3(1):4, 2015.
- [BDM⁺14] Maria-Florina Balcan, Amit Daniely, Ruta Mehta, Ruth Urner, and Vijay V Vazirani. Learning economic parameters from revealed preferences. In *Web and Internet Economics*, pages 338–353. Springer, 2014.
- [BHP14] Avrim Blum, Nika Haghtalab, and Ariel D Procaccia. Learning optimal commitment to overcome insecurity. In *Advances in Neural Information Processing Systems*, pages 1826–1834, 2014.
- [BLNR15] Alexandre Belloni, Tengyuan Liang, Hariharan Narayanan, and Alexander Rakhlin. Escaping the local minima via simulated annealing: Optimization of approximately convex functions. In *Proceedings of The 28th Conference on Learning Theory, COLT 2015, Paris, France, July 3-6, 2015*, pages 240–265, 2015.
- [BLSS14] Umang Bhaskar, Katrina Ligett, Leonard J. Schulman, and Chaitanya Swamy. Achieving target equilibria in network routing games without knowing the latency functions. In *55th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2014, Philadelphia, PA, USA, October 18-21, 2014*, pages 31–40. IEEE Computer Society, 2014.
- [BMM15] Avrim Blum, Yishay Mansour, and Jamie Morgenstern. Learning valuation distributions from partial observation. In *Twenty-Ninth AAAI Conference on Artificial Intelligence (AAAI-15)*, 2015.
- [BN09] Liad Blumrosen and Noam Nisan. On the computational power of demand queries. *SIAM Journal on Computing*, 39(4):1372–1391, 2009.
- [Bub14] Sébastien Bubeck. Theory of convex optimization for machine learning. *CoRR*, abs/1405.4980, 2014.
- [BV06] Eyal Beigman and Rakesh Vohra. Learning from revealed preference. In *Proceedings of the 7th ACM Conference on Electronic Commerce*, pages 36–42. ACM, 2006.
- [CHN14] Shuchi Chawla, Jason D. Hartline, and Denis Nekipelov. Mechanism design for data science. In Moshe Babaioff, Vincent Conitzer, and David Easley, editors, *ACM Conference on Economics and Computation, EC '14, Stanford, CA, USA, June 8-12, 2014*, pages 711–712. ACM, 2014.
- [CMS05] Benoît Colson, Patrice Marcotte, and Gilles Savard. Bilevel programming: A survey. *4OR*, 3(2):87–107, 2005.
- [CR14] Richard Cole and Tim Roughgarden. The sample complexity of revenue maximization. In *Proceedings of the 46th Annual ACM Symposium on Theory of Computing*, pages 243–252. ACM, 2014.
- [DHN14] Shaddin Dughmi, Li Han, and Noam Nisan. Sampling and representation complexity of revenue maximization. In *Web and Internet Economics*, pages 277–291. Springer, 2014.

- [FS96] Yoav Freund and Robert E. Schapire. Game theory, on-line prediction and boosting. In *Proceedings of the Ninth Annual Conference on Computational Learning Theory, COLT 1996, Desenzano del Garda, Italy, June 28-July 1, 1996.*, pages 325–332, 1996.
- [HMR14] Zhiyi Huang, Yishay Mansour, and Tim Roughgarden. Making the most of your samples. *arXiv preprint arXiv:1407.2479*, 2014.
- [KCP10] Dmytro Korzhyk, Vincent Conitzer, and Ronald Parr. Complexity of computing optimal stackelberg strategies in security resource allocation games. In *Twenty-Fourth AAAI Conference on Artificial Intelligence (AAAI-10)*, 2010.
- [LCM09] Joshua Letchford, Vincent Conitzer, and Kamesh Munagala. Learning and approximating the optimal strategy to commit to. In *Algorithmic Game Theory*, pages 250–262. Springer, 2009.
- [LV06] László Lovász and Santosh Vempala. Simulated annealing in convex bodies and an $O^*(n^4)$ volume algorithm. *J. Comput. Syst. Sci.*, 72(2):392–417, 2006.
- [MR15] Jamie Morgenstern and Tim Roughgarden. The pseudo-dimension of near-optimal auctions. *arXiv preprint arXiv:1506.03684*, 2015.
- [MS96] D Monderer and LS Shapley. Potential games. *Games and Economic Behavior*, 14(1):124–143, 1996.
- [NJLS09] Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.
- [Sio58] Maurice Sion. On general minimax theorems. *Pacific J. Math.*, 8(1):171–176, 1958.
- [Zin03] Martin Zinkevich. Online convex programming and generalized infinitesimal gradient ascent. In *Machine Learning, Proceedings of the Twentieth International Conference (ICML 2003), August 21-24, 2003, Washington, DC, USA*, pages 928–936, 2003.
- [ZR12] Morteza Zadimoghaddam and Aaron Roth. Efficiently learning from revealed preference. In Paul W. Goldberg, editor, *Internet and Network Economics - 8th International Workshop, WINE 2012, Liverpool, UK, December 10-12, 2012. Proceedings*, volume 7695 of *Lecture Notes in Computer Science*, pages 114–127. Springer, 2012.

A A Routing Game Where Social Cost is Not Convex in The Tolls

As we stated in the introduction, we can give a simple example of a routing game in which the function mapping a set of tolls on each of the edges to the social cost of the equilibrium routing in the routing game induced by those tolls is not a convex function of the tolls. The example is related to the canonical examples of *Braess’ Paradox* in routing games.

Let $SC(\tau_1, \tau_2)$ be the function that maps a pair of tolls for the two $A \rightarrow B$ edges to the social cost (excluding the tolls) of the equilibrium routing. For each of the inputs we consider, the equilibrium will be unique, so multiplicity of equilibria is irrelevant.




Figure 1: A routing game in which the function mapping tolls to social cost of the unique equilibrium routing is not convex. In this example, there are n players each trying to route $1/n$ units of flow from S to T . There are two edges from A to B with tolls τ_1 and τ_2 , and we assume without loss of generality that all other tolls are fixed to 0. Each edge is labeled with the latency function indicating the cost of using that edge when the congestion on that edge is $x \in [0, 1]$. Note that the latencies (excluding the tolls) on every edge are bounded in $[0, 1]$.

First, consider the set of tolls $\tau = \tau_1 = \tau_2 = 0$. It is not hard to verify that the unique equilibrium is for every player to use the route $S \rightarrow A \rightarrow B \rightarrow T$ using the $A \rightarrow B$ edge on the right (with latency 0).⁶ Each player will experience a total latency of 1 along their route. Thus $SC(\tau) = 8n/10$.

Now consider the tolls τ' in which $\tau_1 = 1, \tau_2 = 2$. At these tolls, it is not hard to verify that the unique equilibrium is for $n/2$ players to use $S \rightarrow A \rightarrow T$ and half to use $S \rightarrow B \rightarrow T$.⁷ Every player experiences a total latency of $2/10 + 1/2 = 7/10$. Thus $SC(\tau') = 7n/10$.

Finally, consider the convex combination $99\tau/100 + \tau'/100$ in which $\tau_1 = 1/100$ and $\tau_2 = 1/50$. In this case, the unique equilibrium routing will have every player use the route $S \rightarrow A \rightarrow B \rightarrow T$ but using the $A \rightarrow B$ edge on the left (with latency $1/200$ and latency-plus-toll $3/200$). To see why, observe that if a player were at A , then no matter what the other players are doing, the cheapest path to T is to go $A \rightarrow B \rightarrow T$ using the left edge (note that the right edge has latency-plus-toll $1/50$ whereas the left edge has latency-plus-toll $3/200$). Thus, the cost of going $B \rightarrow T$ is exactly $1/2$ and the cost of going $A \rightarrow B \rightarrow T$ is exactly $1/2 + 3/200$. Now, if the player is at S , going $S \rightarrow B \rightarrow T$ costs exactly 1, whereas going $S \rightarrow A \rightarrow B \rightarrow T$ costs *at most* $4/10 + 1/2 + 3/200 = 183/200 < 1/2$. Thus, every player will choose the path $S \rightarrow A \rightarrow B \rightarrow T$, using the left $A \rightarrow B$ edge. Every player experiences a total latency of exactly $183/200$. Thus, $SC(\tau'') = 183n/200$.

⁶Since the graph is a DAG, we can use backwards induction. From A , it can never cost more to go $A \rightarrow B \rightarrow T$ than to go $A \rightarrow T$. Since one can go from A to B for a cost of 0, players are indifferent about ending up at node A and node B . Since $S \rightarrow A$ can never cost more than $S \rightarrow B$, and players are indifferent between A and B , every player would choose the path $S \rightarrow A \rightarrow B \rightarrow T$ (using the 0 latency path from A to B).

⁷At these tolls, no player will ever use either $A \rightarrow B$ edge. Thus, they will balance the traffic so that $S \rightarrow A \rightarrow T$ and $S \rightarrow B \rightarrow T$ have equal cost. By symmetry, half will go through A and half through B .

But, since

$$\begin{aligned}
SC(99\tau/100 + \tau'/100) &= \frac{183n}{200} \\
&> \frac{99}{100} \cdot \frac{8n}{10} + \frac{1}{100} \cdot \frac{7n}{10} \\
&= \frac{99}{100} \cdot SC(\tau) + \frac{1}{100} \cdot SC(\tau'),
\end{aligned}$$

we can see that the function $SC(\tau)$ is not convex in τ .

B Missing Proofs in Section 3

Lemma 35. *Suppose function $v: \mathbb{R}_+^d \rightarrow \mathbb{R}_+$ is a concave and homogeneous of some degree $k \leq 0$. Then $k \leq 1$.*

Proof. First, we show that $v(\mathbf{0}) = 0$. To see this, observe that for any $b > 1$, we can write $v(\mathbf{0}) = v(b, \mathbf{0}) = b^k v(\mathbf{0})$. For any $x \in \mathbb{R}_+^d$ such that $x \neq 0$, we have the following due to the concavity of v :

$$v(x)/2 = \frac{1}{2} [v(\mathbf{0}) + v(x)] \leq v(x/2) = \left(\frac{1}{2}\right)^k v(x).$$

This means that $k \leq 1$. □

B.1 Properties of CES and Cobb-Douglas Utilities

In this sub-section, we give proofs showing that both CES and Cobb-Douglas utility functions are strongly concave and Hölder continuous in the convex region $(0, H]^d$.

B.1.1 Constant Elasticity of Substitution (CES)

Consider valuation functions of the form:

$$v(x) = \left(\sum_{i=1}^d \alpha_i x_i^\rho \right)^\beta,$$

where $\alpha_i > 0$ for every $i \in [d]$ and $\rho, \beta > 0$ such that $\rho < 1$ and $\beta\rho < 1$.

Theorem 36. *Let C be a convex set such that there exists some constant $H > 0$ such that $C \subseteq (0, H]^d$. Then v is R -strongly concave over the C for some constant R and is $((\max_i \alpha_i d)^\beta, \rho\beta)$ -Hölder continuous.*

Proof. We will derive the Hessian matrix $\nabla^2 v$ of the function v , and show that there exists some fixed $R > 0$ such that for every $x \in C$ and every $y \in \mathbb{R}^d$, we have $y^\top \nabla^2 v(x) y \leq -R \|y\|^2$. First, we have the first partial derivatives

$$\frac{\partial v}{\partial x_i} = \beta \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{(\beta-1)} \left(\rho \alpha_i x_i^{\rho-1} \right)$$

Now we take the second partial derivatives. For any $i \neq j$,

$$\frac{\partial^2 v}{\partial x_i \partial x_j} = \beta(\beta - 1) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left(\rho \alpha_j x_j^{\rho-1} \right) \left(\rho \alpha_i x_i^{\rho-1} \right)$$

and for any i ,

$$\frac{\partial^2 v}{\partial x_i^2} = \beta(\beta - 1) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left(\rho \alpha_i x_i^{\rho-1} \right)^2 + \beta \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \left(\rho(\rho - 1) \alpha_i x_i^{\rho-2} \right).$$

Recall that the ij -th entry of the Hessian matrix is $(\nabla^2 v)_{i,j} = \partial^2 v / \partial x_i \partial x_j$, and we could write

$$\begin{aligned} y^\top (\nabla^2 v(x)) y &= \sum_{i=1}^d \sum_{j=1}^d \frac{\partial^2 v}{\partial x_i \partial x_j} y_i y_j \\ &= 2 \sum_{i \neq j} \frac{\partial^2 v}{\partial x_i \partial x_j} y_i y_j + \sum_{i=1}^d \frac{\partial^2 v}{\partial x_i^2} y_i^2 \\ &= 2\beta(\beta - 1)\rho^2 \sum_{i \neq j} \alpha_i \alpha_j \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} (x_i^{\rho-1} y_i)(x_j^{\rho-1} y_j) \\ &\quad + \sum_{i=1}^d \left[\beta(\beta - 1) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left(\rho \alpha_i x_i^{\rho-1} y_i \right)^2 + \beta \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \left(\rho(\rho - 1) \alpha_i x_i^{\rho-2} y_i^2 \right) \right] \\ &= \beta(\beta - 1)\rho^2 \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left(\sum_{k=1}^d \alpha_k x_k^{\rho-1} y_k \right)^2 + \beta\rho(\rho - 1) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \left(\sum_{k=1}^d \alpha_k x_k^{\rho-2} y_k^2 \right) \end{aligned}$$

We will first consider the case where $\beta \leq 1$. Then both of the terms above are non-positive, and

$$\begin{aligned} y^\top (\nabla^2 v(x)) y &\leq -\beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \left(\sum_{k=1}^d \alpha_k x_k^{\rho-2} y_k^2 \right) \\ &\leq -\beta\rho(1 - \rho) \inf_{x \in C} \left\{ \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \right\} \inf_{k \in [d], x \in C} \{ \alpha_k x_k^{\rho-2} \} \left(\sum_{k=1}^d y_k^2 \right) \\ &\leq -\beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k H^\rho \right)^{\beta-1} \min_k \{ \alpha_k \} H^{\rho-2} \|y\|^2 \\ &= -\beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k \right)^{\beta-1} \min_k \{ \alpha_k \} H^{\rho\beta-2} \|y\|^2 \end{aligned}$$

Thus, $y^\top (\nabla^2 v(x)) y \leq -R \|y\|^2$ for $R = \beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k \right)^{\beta-1} \min_k \{ \alpha_k \} H^{\rho\beta-2}$.

Now we consider the case where $\beta > 1$. Since we have assumed that $\rho\beta < 1$, we also know that $(\beta - 1)\rho < 1 - \rho$. Let $\kappa = \frac{(\beta-1)\rho}{(1-\rho)}$ and we know that $0 < \kappa < 1$. It follows that

$$\begin{aligned}
y^\top (\nabla^2 v(x)) y &= \beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left[\kappa \left(\sum_{k=1}^d \alpha_k x_k^{\rho-1} y_k \right)^2 - \kappa \left(\sum_{k=1}^d \alpha_k x_k^\rho \right) \left(\sum_{k=1}^d \alpha_k x_k^{\rho-2} y_k^2 \right) \right] \\
&\quad - \beta\rho(1 - \rho) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-2} \left[(1 - \kappa) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right) \left(\sum_{k=1}^d \alpha_k x_k^{\rho-2} y_k^2 \right) \right] \\
(\text{Cauchy-Schwarz}) \quad &\leq -\beta\rho(1 - \rho)(1 - \kappa) \left(\sum_{k=1}^d \alpha_k x_k^\rho \right)^{\beta-1} \left(\sum_{k=1}^d \alpha_k x_k^{\rho-2} y_k^2 \right) \\
&\leq -\beta\rho(1 - \rho)(1 - \kappa) \left(\sum_{k=1}^d \alpha_k^\beta x_k^{\rho\beta-2} y_k^2 \right) \\
&\leq -\beta\rho(1 - \rho)(1 - \kappa) \inf_{k \in [d], x \in C} \left\{ \alpha_k^\beta x_k^{\rho\beta-2} \right\} \|y\|^2 \\
&\leq -\beta\rho(1 - \rho)(1 - \kappa) \min_k \left\{ \alpha_k^\beta \right\} H^{\rho\beta-2} \|y\|^2
\end{aligned}$$

This means, $y^\top (\nabla^2 v(x)) y \leq R \|y\|^2$ for $R = \beta\rho(1 - \rho)(1 - \kappa) \min_{k=1} \left\{ \alpha_k^\beta \right\} H^{\rho\beta-2}$. Therefore, we have shown that v is R strongly concave in C for some positive constant R .

Next, we will show that the function is Hölder continuous over C . Let $x, y \in C$ such that $x \neq y$. Without loss of generality, assume that $v(x) \geq v(y)$, and let $\varepsilon_i = |x_i - y_i|$ for each $i \in [d]$. Then we have

$$\begin{aligned}
\left(\sum_i \alpha_i x_i^\rho \right)^\beta - \left(\sum_i \alpha_i y_i^\rho \right)^\beta &\leq \left(\sum_i \alpha_i |x_i - y_i|^\rho \right)^\beta \\
&\leq \left(\max_i \alpha_i \right)^\beta \cdot \left(\sum_i \varepsilon_i^\rho \right)^\beta \\
&\leq \left(\max_i \alpha_i \right)^\beta \cdot (d \|x - y\|_2^\rho)^\beta \\
&\leq \left(\max_i \alpha_i \right)^\beta \cdot d^\beta \|x - y\|_2^{\rho\beta}
\end{aligned}$$

where the first step follows from the sub-additivity. This shows that the function is $((\max_i \alpha_i) d)^\beta, \rho\beta$ -Hölder continuous over C , which completes the proof. \square

B.1.2 Cobb-Douglas

Consider valuation functions of the form

$$v(x) = \prod_{i=1}^d x_i^{\alpha_i},$$

where $\alpha_i > 0$ for every $i \in [d]$ and $\sum_{i=1}^d \alpha_i < 1$.

Theorem 37. Let C be a convex set such that there exists some constant that $H > 0$ such that $C \subseteq (0, H]^d$. Then v is $(1, \sum_i \alpha_i)$ -Hölder continuous and R -strongly concave over C for some constant R .

Proof. Similar to the previous proof, we will show that there exists some constant $R > 0$ such that for every $x \in C$ and every $y \in \mathbb{R}^d$, we have $y^\top \nabla^2 v(x) y \leq -R \|y\|^2$. First, we could write down the following first and second partial derivatives of the function:

$$\frac{\partial v}{\partial x_i} = \alpha_i x_i^{\alpha_i - 1} \prod_{j \neq i} x_j^{\alpha_j}$$

$$\frac{\partial^2 v}{\partial^2 x_i} = \alpha_i (\alpha_i - 1) x_i^{\alpha_i - 2} \prod_{j \neq i} x_j^{\alpha_j}$$

and for any $i \neq j$,

$$\frac{\partial^2 v}{\partial x_i \partial x_j} = \alpha_i x_i^{\alpha_i - 1} \alpha_j x_j^{\alpha_j - 1} \prod_{k \neq i, j} x_k^{\alpha_k}$$

Let $y \in \mathbb{R}^d$, and let $\kappa = \sum_{i=1}^d \alpha_i \in (0, 1)$.

$$\begin{aligned} y^\top (\nabla^2 v(x)) y &= \sum_{i=1}^d \sum_{j=1}^d y_i \frac{\partial^2 v}{\partial x_i \partial x_j} y_j \\ &= 2 \sum_{i \neq j} \prod_{k \neq i, j} x_k^{\alpha_k} (\alpha_i x_i^{\alpha_i - 1} y_i) (\alpha_j x_j^{\alpha_j - 1} y_j) + \sum_{i=1}^d \alpha_i (\alpha_i - 1) x_i^{\alpha_i - 2} y_i^2 \prod_{j \neq i} x_j^{\alpha_j} \\ &= \left(\prod_{k=1}^d x_k^{\alpha_k} \right) \left[2 \sum_{i \neq j} (\alpha_i x_i^{-1} y_i) (\alpha_j x_j^{-1} y_j) + \sum_{i=1}^d \alpha_i (\alpha_i - 1) x_i^{-2} y_i^2 \right] \\ &= \left(\prod_{k=1}^d x_k^{\alpha_k} \right) \left[\left(\sum_{i=1}^d \alpha_i x_i^{-1} y_i \right)^2 - \sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right] \\ \text{(Cauchy-Schwarz)} \quad &\leq \left(\prod_{k=1}^d x_k^{\alpha_k} \right) \left[\left(\sum_{i=1}^d \alpha_i \right) \left(\sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right) - \sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right] \\ &\leq \left(\prod_{k=1}^d x_k^{\alpha_k} \right) \left[\kappa \left(\sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right) - \sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right] \\ &\leq - \left(\prod_{k=1}^d x_k^{\alpha_k} \right) (1 - \kappa) \left(\sum_{i=1}^d \alpha_i x_i^{-2} y_i^2 \right) \\ &\leq -H^{(\sum_k \alpha_k - 2)} (1 - \kappa) \min_k \{\alpha_k\} \|y\|^2 \end{aligned}$$

Therefore, v is R -strongly concave in C for $R = H^{(\sum_k \alpha_k - 2)} (1 - \kappa) \min_k \{\alpha_k\}$.

Next, we will show that the function is $(1, \sum_i \alpha_i)$ -Hölder continuous. Let $x, y \in C$ such that $x \neq y$. Without loss generality, assume that $v(x) \geq v(y)$. We could write

$$\begin{aligned} \prod_{i=1}^d x_i^{\alpha_i} - \prod_{i=1}^d y_i^{\alpha_i} &\leq \prod_{i=1}^d |x_i - y_i|^{\alpha_i} \\ &\leq \prod_{i=1}^d \|x - y\|_2^{\alpha_i} = \|x - y\|_2^{\sum_i \alpha_i} \end{aligned}$$

where the first step follows from the sub-additivity of v . This shows that the function is $(1, \sum_i \alpha_i)$ -Hölder continuous, which completes the proof. \square

C Detailed Analysis of Section 4

Just as in Section 3.4, we start by considering the following convex program associated with \hat{x}

$$\max_{x \in \mathcal{A}_F} \phi(x) \tag{14}$$

$$\text{such that } x_j \leq \hat{x}_j \text{ for every } j \in [d] \tag{15}$$

The Lagrangian of the program is

$$\mathcal{L}(x, p) = \phi(x) - \sum_{j=1}^d p_j (x_j - \hat{x}_j),$$

where p_j is the dual variable for each constraint (15) and the vector p is also interpreted as the action of the leader. We can interpret the Lagrangian as the payoff function of a zero-sum game, in which the leader is the minimization player and the follower is the maximization player. To guarantee the fast convergence of our later use of gradient descent, we will restrict the leader to play actions in \mathcal{P}_m as defined in (11). It is known that such a zero-sum game has value V : the leader has a minimax strategy p^* such that $\mathcal{L}(x, p^*) \leq V$ for all $x \in \mathcal{A}_F$, and the follower has a maxmin strategy x^* such that $\mathcal{L}(x^*, p) \geq V$ for all $p \in \mathcal{P}$. We first state the following lemma about the *value* of this zero-sum game.

Lemma 38. *The value of the Lagrangian zero-sum game is $V = \phi(\hat{x})$, that is*

$$\max_{x \in \mathcal{A}_F} \min_{p \in \mathcal{P}} \mathcal{L}(x, p) = \min_{p \in \mathcal{P}} \max_{x \in \mathcal{A}_F} \mathcal{L}(x, p) = \phi(\hat{x})$$

Proof. The proof is identical to the proof of Lemma 13. \square

An approximate minimax equilibrium of a zero-sum game is defined as follows.

Definition 39. *Let $\alpha \geq 0$. A pair of strategies $p \in \mathcal{P}$ and $x \in \mathcal{A}_F$ form an α -approximate minimax equilibrium if*

$$\mathcal{L}(x, p) \geq \max_{x' \in \mathcal{A}_F} \mathcal{L}(x', p) - \alpha \geq V - \alpha \quad \text{and} \quad \mathcal{L}(x, p) \leq \min_{p' \in \mathcal{P}} \mathcal{L}(x, p') + \alpha \leq V + \alpha.$$

First observe that fixing a strategy x , the maximization player's best response in this zero-sum game is the same as the follower's best response in the Stackelberg game.

Claim 40. Let $p' \in \mathcal{P}$ be any action of the leader, then

$$\arg \max_{x \in \mathcal{A}_F} \mathcal{L}(x, p') = x^*(p')$$

Proof. We can write

$$\begin{aligned} \arg \max_{x \in \mathcal{A}_F} \mathcal{L}(x, p') &= \arg \max_{x \in \mathcal{A}_F} [\phi(x) - \langle p', x - \hat{x} \rangle] \\ &= \arg \max_{x \in \mathcal{A}_F} [\phi(x) - \langle p', x \rangle] \\ &= \arg \max_{x \in \mathcal{A}_F} U_F(p', x) \end{aligned}$$

The second equality follows from the fact that $\langle p', \hat{x} \rangle$ is independent of the choice of x . \square

Now we show that if a strategy pair (p', x') is an approximate minimax equilibrium in the zero-sum game, then p' approximately induces the target action \hat{x} of the follower. Hence, our task reduces to finding an approximate minimax equilibrium of the Lagrangian game.

Lemma 41. Suppose that a pair of strategies $(p', x') \in \mathcal{P} \times \mathcal{A}_F$ forms an α -approximate minimax equilibrium. Then the induced follower action $x^*(p')$ satisfies $\|\hat{x} - x^*(p')\| \leq \sqrt{4\alpha/\sigma}$.

Proof. By the definition of approximate minimax equilibrium, we have

$$\phi(\hat{x}) - \alpha \leq \mathcal{L}(x', p') \leq \phi(\hat{x}) + \alpha$$

and also by 40,

$$\max_{x \in \mathcal{A}_F} \mathcal{L}(x, p') - \alpha = \mathcal{L}(x^*(p'), p') - \alpha \leq \mathcal{L}(x', p') \leq \phi(\hat{x}) + \alpha$$

Note that

$$\mathcal{L}(\hat{x}, p') = \phi(\hat{x}) - \langle p', \hat{x} - \hat{x} \rangle = \phi(\hat{x}).$$

It follows that $\mathcal{L}(x^*(p'), p') \leq \mathcal{L}(\hat{x}, p') + 2\alpha$. Since ϕ is a σ -strongly concave function, we know that fixing any leader's action p , \mathcal{L} is also a σ -strongly concave function in x . By Lemma 5 and the above argument, we have

$$2\alpha \geq \mathcal{L}(x^*(p'), p') - \mathcal{L}(\hat{x}, p') \geq \frac{\sigma}{2} \|x^*(p') - \hat{x}\|^2$$

Hence, we must have $\|x^*(p') - \hat{x}\| \leq \sqrt{4\alpha/\sigma}$. \square

To compute an approximate minimax equilibrium, we will use the following T -round no-regret dynamics: the leader plays online gradient descent (a “no-regret” algorithm), while the follower selects an ζ -approximate best response every round. In particular, the leader will produce a sequence of actions $\{p^1, \dots, p^T\}$ against the follower's best responses $\{x^1, \dots, x^T\}$ such that for each round $t \in [T]$:

$$p^{t+1} = \Pi_{\mathcal{P}} [p^t - \eta \cdot \nabla_p \mathcal{L}(x^t, p^t)] \quad \text{and} \quad x^t = x'(p^t).$$

At the end of the dynamics, the leader has *regret* defined as

$$\mathcal{R}_L \equiv \frac{1}{T} \sum_{t=1}^T \mathcal{L}(x^t, p^t) - \frac{1}{T} \min_{p \in \mathcal{P}} \sum_{t=1}^T \mathcal{L}(x^t, p).$$

Now take the average actions for both players for this dynamics: $\bar{x} = \frac{1}{T} \sum_{t=1}^T x^t$ and $\bar{p} = \frac{1}{T} \sum_{t=1}^T p^t$. A well-known result by [FS96] shows that the average plays form an approximate minimax equilibrium.

Theorem 42 ([FS96]). *The average action pair (\bar{p}, \bar{x}) forms a $(\mathcal{R}_L + \zeta)$ -approximate minimax equilibrium of the Lagrangian zero-sum game.*

To simulate the no-regret dynamics, we will have the following T -round dynamics between the leader and the follower: in each round t , the leader plays action p^t based the gradient descent update and observes the induced action $x^*(p^t)$. The gradient of the Lagrangian $\nabla_p \mathcal{L}$ can be easily computed based on the observations. Recall from 40 that the follower's best responses to the leader's actions are the same in both the Stackelberg game and the Lagrangian zero-sum game. This means that the gradient of the Lagrangian can be obtained as follows

$$\nabla_p \mathcal{L}(x^*(p'), p') = (\hat{x} - x^*(p')).$$

In the end, the algorithm will output the average play of the leader $1/T \sum_{t=1}^T p^t$. The full description of the algorithm **LearnLead** is presented in Algorithm 3, and before we present the proof of Theorem 23, we first include the no-regret guarantee of gradient descent.

Lemma 43 ([Zin03]). *Let \mathcal{D} be a closed convex set such that $\|\mathcal{D}\| \leq D$, and let c^1, \dots, c^T be a sequence of differentiable, convex functions with bounded gradients, that is for every $x \in \mathcal{D}, \|\nabla c^t(x)\|_2 \leq G$. Let $\eta = \frac{D}{G\sqrt{T}}$ and $\omega^1 = \Pi_{\mathcal{D}}[\mathbf{0}]$ be arbitrary. Then if we compute $\omega^1, \dots, \omega^T \in \mathcal{D}$ based on gradient descent $\omega^{t+1} = \Pi_{\mathcal{D}}[\omega^t - \eta \nabla c(\omega^t)]$, the regret satisfies*

$$\mathcal{R} \equiv \frac{1}{T} \sum_{t=1}^T c^t(\omega^t) - \min_{\omega \in \mathcal{D}} \frac{1}{T} \sum_{t=1}^T c^t(\omega) \leq \frac{GD}{\sqrt{T}} \quad (16)$$

Proof of Theorem 23. We will first bound the regret of the leader in the no-regret dynamics. Each vector of \mathcal{P} has norm bounded by $\sqrt{d}\lambda_F$. Also, since the gradient of the Lagrangian at point p' is

$$\nabla_p \mathcal{L}(x^*(p'), p') = (\hat{x} - x^*(p')),$$

we then know that the gradient is bounded the norm of \mathcal{A}_F , which is γ . The regret is bounded by

$$\mathcal{R}_L = \frac{\sqrt{2d}\lambda_F\gamma}{\sqrt{T}}.$$

Let $\bar{x} = 1/T \sum_{t=1}^T x^*(p^t)$ denote the average play of the follower. It follows from Theorem 42 that (\hat{p}, \bar{x}) forms an $(\mathcal{R}_L + \zeta)$ -approximate minimax equilibrium, and so by Lemma 41, we have

$$\|\hat{x} - x^*(\hat{p})\| \leq \sqrt{\frac{4(\mathcal{R}_L + \zeta)}{\sigma}} = \sqrt{\frac{4\sqrt{2d}\lambda_F\gamma}{\sigma\sqrt{T}} + \frac{4\zeta}{\sigma}}.$$

Plugging in our choice of T , we get $\|x^*(\hat{p}) - \hat{x}\| \leq \varepsilon/2$, and also the total number of observations on the follower is also T . Finally, by strong concavity of the function ϕ and Lemma 5, we have that

$$\|x'(\hat{p}) - x^*(\hat{p})\| \leq \sqrt{\frac{2\zeta}{\sigma}} \leq \varepsilon/2.$$

By triangle inequality, we could show that the approximate best-response satisfies $\|x'(\hat{p}) - \hat{x}\| \leq \varepsilon$. \square

Finally, we give the proof of Theorem 24.

Proof of Theorem 24. Since for each x^t , the approximate evaluation based on $x'(p^t)$ satisfies $|\psi(x^t) - \psi(x'(p^t))| \leq \lambda_L \varepsilon$, by Lemma 7 we know that the action \hat{x} output by **ZOO**($d\varepsilon\lambda_L, \mathcal{A}_F$) satisfies

$$\mathbb{E}[\psi(\hat{x})] \geq \max_{x \in \mathcal{A}_F} \psi(x) - d\varepsilon\lambda_L = \max_{p \in \mathcal{A}_L} U_L(p, x^*(p)) - d\varepsilon\lambda_L.$$

Finally, we will use **LearnLead**(\hat{x}, ε) to output a leader action \hat{p} such that $\|x^*(\hat{p}) - \hat{x}\| \leq \varepsilon$, and so $\psi(x^*(\hat{p})) \geq \psi(\hat{x}) - \lambda_L \varepsilon$. Therefore, in the end we guarantee that

$$\mathbb{E}[U_L(\hat{p}, x^*(\hat{p}))] = \mathbb{E}[\psi(\hat{p})] \geq \mathbb{E}[\psi(\hat{x})] - \lambda_L \varepsilon \geq \max_{p \in \mathcal{A}_L} U_L(p, x^*(p)) - (d+1)\varepsilon\lambda_L.$$

Plugging in our choice of ε , we recover the α accuracy guarantee. Note that in each iteration, the number of observations on the follower required by the call of **LearnLead** is

$$T' = O\left(\frac{d\lambda_F^2\gamma^2}{\varepsilon^4\sigma^2}\right)$$

Therefore the total number of observations our algorithm needs is bounded by

$$O(T' \times T) = \tilde{O}\left(\frac{d^{5.5}\lambda_F^2\gamma^2}{\varepsilon^4\sigma^2}\right)$$

Again plugging in our value of ε , the bound on the number of observations needed is

$$\tilde{O}\left(\frac{d^{9.5}\lambda_F^2\lambda_L^4\gamma^2}{\alpha^4\sigma^2}\right).$$

Hiding the constants ($\lambda_F, \lambda_L, \gamma$ and σ), we recover the bound above. □

D Improvement with Ellipsoid in Noiseless Settings

In this section, we present variants of the **LearnPrice** and **LearnLead** algorithms that uses the Ellipsoid algorithm as a first-order optimization method. In particular, this will allow us to improve the dependence of the query complexity on the target accuracy parameter α . For the technique we give in this section, the number of observations of the follower's actions will have a poly-logarithmic dependence on $1/\alpha$ instead of a polynomial one. We also improve the polynomial dependence on the dimension.

D.1 The Ellipsoid Algorithm

We will briefly describe the ellipsoid method without going into full detail. Let $\mathcal{P} \subset \mathbb{R}^d$ be a convex body, and $f: \mathcal{P} \rightarrow [-B, B]$ be a continuous and convex function. Let $r, R > 0$ be such that the set \mathcal{P} is contained in an Euclidean ball of radius R and it contains a ball of radius r . The *ellipsoid* algorithm solves the following problem: $\min_{p \in \mathcal{P}} f(p)$.

The algorithm requires access to a *separation oracle* for \mathcal{P} : given any $p \in \mathbb{R}^d$, it either outputs that x is a member of \mathcal{P} , or if $p \notin \mathcal{P}$ then it outputs a separating hyperplane between p and \mathcal{P} . It also requires access to a *first-order oracle*: given any $p \in \mathbb{R}^d$, it outputs a subgradient $w \in \partial f(p)$. The algorithm maintains an ellipsoid E^t with center c^t in \mathbb{R}^d over rounds, and in each round t does the following:

1. If the center of the ellipsoid $c^t \notin \mathcal{P}$, it calls the separation oracle to obtain a separating hyperplane $w_t \in \mathbb{R}^d$ such that $\mathcal{P} \subset \{p: (p - c_t)^\top w_t \leq 0\}$; otherwise it calls the first-order oracle at c_t to obtain $w_t \in \partial f(c_t)$.
2. Obtain a new ellipsoid E^{t+1} with center c_{t+1} based on the ellipsoid E^t and vector w_t . (See e.g. [Bub14] for details.) We will treat this ellipsoid update as a black-box step and write it as a function **ellip**(E, c, w) that takes an ellipsoid E along with its center c and also a vector w as input and returns a new ellipsoid E' with its center c' as output.

The sequence of ellipsoid centers $\{c_t\}$ produced by the algorithm has the following guarantee.

Theorem 44 (see e.g. [Bub14]). *For $T \geq 2d^2 \log(R/r)$, the ellipsoid algorithm satisfies $\{c_1, \dots, c_T\} \cap \mathcal{P} \neq \emptyset$ and*

$$\min_{c \in \{c_1, \dots, c_T\} \cap \mathcal{P}} f(c) - \min_{p \in \mathcal{P}} f(p) \leq \frac{2BR}{r} \exp\left(-\frac{T}{2d^2}\right).$$

In other words, with at most $T = O\left(d^2 \log\left(\frac{BR}{r\varepsilon}\right)\right)$ calls to the first-order oracles, the ellipsoid algorithm finds a point $p \in (\{c_t\} \cap \mathcal{P})$ that is ε -optimal for the function f .

D.2 Learning Prices with Ellipsoid

We will first revisit the problem in Section 3.4 and give an ellipsoid-based variant of **LearnPrice** that (when there is no noise) obtains a better query complexity for **LearnPrice**. Recall that we are interested in computing a price vector $\hat{p} \in \mathcal{P}$ such that the induced bundle $x^*(\hat{p})$ is close to the target bundle \hat{x} .

Recall from Lemma 14 that it is sufficient to find an approximately optimal solution to the Lagrangian dual function $g(p) = \operatorname{argmax}_{x \in C} \mathcal{L}(x, p)$. We will use the ellipsoid algorithm to find such a solution. Note that the feasible region for prices is given explicitly: $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}L\}$, so a separation oracle for \mathcal{P} is easy to implement. Furthermore, we know from Lemma 15 that for any p , we can obtain a gradient based on the buyer's purchased bundle: $(\hat{x} - x^*(p)) \in \partial g(p)$. This means we have both the separation oracle and first-order oracle necessary for running ellipsoid. The algorithm **LearnPE** is presented in Algorithm 7.

Algorithm 7 Learning the price vector to induce a target bundle: **LearnPE**(\hat{x}, ε)

Input: A target bundle $\hat{x} \in \operatorname{Int}_C$, and target accuracy ε

Initialize: restricted price space $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq \sqrt{d}L\}$

$$c^1 = \mathbf{0} \quad E^1 = \{p \in \mathbb{R}^d \mid \|p\| \leq \sqrt{d}L\} \quad T = 100d^2 \ln\left(\frac{d\lambda_{\text{val}}\gamma}{\varepsilon\sigma}\right) \quad L = (\lambda_{\text{val}})^{1/\beta} \left(\frac{4}{\varepsilon^2\sigma}\right)^{(1-\beta)/\beta}$$

For $t = 1, \dots, T$:

while $c^t \notin \mathcal{P}$ then let obtain a separating hyperplane w and let $(E^t, c^t) \leftarrow \mathbf{ellip}(E^t, c^t, w)$

Let $p^t = c^t$

Observe the purchased bundle by the consumer $x^*(p^t)$

Update the ellipsoid $(E^{t+1}, c^{t+1}) \leftarrow \mathbf{ellip}(E^t, c^t, (\hat{x} - x^*(p^t)))$:

Output: $\hat{p} = \operatorname{argmin}_{p \in \{p^1, \dots, p^T\}} \|\hat{x} - x^*(p)\|$

Theorem 45. Let $\hat{x} \in \text{Int}_C$ be a target bundle and $\varepsilon > 0$. Then **LearnPE**(\hat{x}, ε) outputs a price vector \hat{p} such that the induced bundle satisfies $\|\hat{x} - x^*(\hat{p})\| \leq \varepsilon$ and the number of observations it needs is no more than

$$T = O\left(d^2 \ln\left(\frac{d\lambda_{\text{val}}\gamma}{\varepsilon\sigma}\right)\right)$$

Proof. By Lemma 14, it suffices to show that there exists a price vector $\hat{p} \in \{p^t\}_{t=1}^T$ such that

$$g(\hat{p}) \leq \min_{p \in \mathcal{P}} g(p) + \frac{\varepsilon^2\sigma}{4}.$$

We will show this through the accuracy guarantee of ellipsoid. Note that the set \mathcal{P} is contained in a ball of radius $2\sqrt{d}\lambda_{\text{val}}$ and contains a ball of radius $\sqrt{d}\lambda_{\text{val}}$. Furthermore, the Lagrangian dual objective value is also bounded: for any $p \in \mathcal{P}$:

$$\begin{aligned} |g(p)| &= \left| \max_{x \in C} v(x) - \langle p, x - \hat{x} \rangle \right| \\ &\leq \max_{x \in C} v(x) + |\langle p, x^*(p) - \hat{x} \rangle| \\ &\leq \lambda_{\text{val}}\gamma + \|p\| \cdot \|x^*(p) - \hat{x}\| \\ &\leq \lambda_{\text{val}}\gamma + \sqrt{2d}\lambda_{\text{val}}\gamma \\ &\leq 2\sqrt{d}\lambda_{\text{val}}\gamma \end{aligned}$$

By Theorem 44, the following holds

$$\min_{p \in \{p^1, \dots, p^T\}} g(p) - \min_{p' \in \mathcal{P}} g(p') \leq \frac{\varepsilon^2\sigma}{4}.$$

By Lemma 14, there exists some $p \in \{p^1, \dots, p^T\}$ such that the resulting bundle $x^*(p)$ satisfies that $\|\hat{x} - x^*(p)\| \leq \varepsilon$. Since we are selecting \hat{p} as $\hat{p} = \text{argmin}_{p \in \{p^1, \dots, p^T\}} \|\hat{x} - x^*(p)\|$, we must have $\|\hat{x} - x^*(\hat{p})\| \leq \varepsilon$. \square

Now we could use **LearnPE** to replace **LearnPrice** in the algorithm **Opro** as a sub-routine to induce target bundles. The following result follows from the same proof of Footnote 4.

Theorem 46. Let $\alpha > 0$ be the target accuracy. If we replace **LearnPrice** by **LearnPE** in our instantiation of **Opro**(C, α), then the output price vector \hat{p} has expected profit

$$\mathbb{E}[[r(\hat{p})]] \geq \max_{p \in \mathbb{R}_+^d} r(p) - \alpha,$$

the number of times it calls **LearnPE** is bounded by $(d^{4.5} \cdot \text{polylog}(d, 1/\alpha))$, and the observations the algorithm requires from the consumer is at most

$$d^{6.5} \text{polylog}\left(\lambda_{\text{val}}, \gamma, \frac{1}{\alpha}, \frac{1}{\sigma}\right).$$

Algorithm 8 Learning the toll vector to induce a target flow: **LearnTE**(\widehat{f}, ε)

Input: A target flow $\widehat{f} \in \mathcal{F}$, and target accuracy ε
Initialize: restricted toll space $\mathcal{P} = \{p \in \mathbb{R}_+^d \mid \|p\| \leq m\}$

$$c^1 = \mathbf{0} \quad E^1 = \{x \in \mathbb{R}^d \mid \|x\| \leq m\} \quad T = 100 \left(m^2 \ln \left(\frac{m}{\varepsilon \sigma} \right) \right)$$

For $t = 1, \dots, T$:

while $c^t \notin \mathcal{P}$ then let obtain a separating hyperplane w and let $(E^t, c^t) \leftarrow \mathbf{ellip}(E^t, c^t, w)$

Let $\tau^t = c^t$

Observe the induced flow $f^*(\tau^t)$

Update the ellipsoid $(E^{t+1}, c^{t+1}) \leftarrow \mathbf{ellip}(E^t, c^t, -(\widehat{f} - f^*(\tau^t)))$:

Output: $\widehat{\tau} = \operatorname{argmin}_{\tau \in \{\tau^1, \dots, \tau^T\}} \|\widehat{f} - f^*(\tau)\|$

D.3 Learning Tolls with Ellipsoid

We will also revisit the problem in Section 5. We give a similar ellipsoid-based algorithm to induce target flow. See Algorithm 8.

Theorem 47. Let $\widehat{f} \in \mathcal{F}$ be a target bundle and $\varepsilon > 0$. Then **LearnTE**(\widehat{f}, ε) outputs a toll vector $\widehat{\tau}$ such that the induced flow satisfies $\|\widehat{f} - f^*(\widehat{\tau})\| \leq \varepsilon$ and the number of observations it needs is no more than

$$T = O \left(m^2 \ln \left(\frac{m}{\varepsilon \sigma} \right) \right)$$

Proof. Let function g be defined as

$$g(\tau) = \min_{f \in \mathcal{F}} \Phi(f) + \langle \tau, f - \widehat{f} \rangle.$$

It suffices to show that there exists some $\tau' \in \{\tau^1, \dots, \tau^T\}$ such that $g(\tau') \geq \min_{\tau \in \mathcal{P}} g(\tau) - \frac{\varepsilon^2 \sigma}{2}$. Before we instantiate the accuracy theorem of ellipsoid, note that the set \mathcal{P} is contained in a ball of radius m and contains a ball of radius $m/2$, and also that the value of $g(\tau)$ is bounded for any $\tau \in \mathcal{P}$:

$$\begin{aligned} |g(\tau)| &= \left| \min_{f \in \mathcal{F}} \Phi(f) + \langle \tau, f - \widehat{f} \rangle \right| \\ &\leq \max_{f \in \mathcal{F}} \Phi(f) + \max_{f \in \mathcal{F}} \|f - \widehat{f}\| \|\tau\| \\ &\leq m + \sqrt{2mm} \leq 2\sqrt{m^3} \end{aligned}$$

Given that $T = 4m^2 \ln(m/\varepsilon\sigma)$, we know by Theorem 44 that

$$\max_{\tau' \in \{\tau^1, \dots, \tau^T\}} g(\tau') - \max_{\tau \in \mathcal{P}} g(\tau) \geq \frac{\varepsilon^2 \sigma}{2}$$

Therefore, the output toll vector satisfies

$$\|\widehat{f} - f^*(\widehat{\tau})\| \leq \varepsilon.$$

This completes our proof. □

Finally, with this convergence bound, we could also improve the result of Lemma 28.

Theorem 48. *Let $\alpha > 0$ be the target accuracy. If we replace **LearnLead** by **LearnTE** in the instantiation of **LearnOpt**(\mathcal{A}_F, α), then the output toll vector $\hat{\tau}$ and its the induced flow $\hat{f} = f^*(\hat{\tau})$ is α -approximately optimal in expectation:*

$$\mathbb{E} \left[\Psi(\hat{f}) \right] \leq \min_{f \in \mathcal{F}} \Psi(f) + \alpha.$$

*The number of times it calls **LearnTE** is bounded by $(m^{4.5} \cdot \text{polylog}(m, 1/\alpha))$, and so the total number of observations we need on the flow behavior is bounded by*

$$m^{6.5} \text{polylog} \left(\lambda, \frac{1}{\alpha}, \frac{1}{\sigma} \right).$$