

SU(3) Clebsch-Gordan coefficients at large N_c

Fl. Stancu^a

University of Liège, Institute of Physics B5,

Sart Tilman, B-4000 Liège 1, Belgium

(Dated: October 1, 2020)

Abstract

It is argued that several papers where SU(3) Clebsch-Gordan coefficients were calculated in order to describe properties of hadronic systems are, up to a phase convention, particular cases of analytic formulae derived by Hecht in 1965 in the context of nuclear physics. This is valid for irreducible representations with multiplicity one in the corresponding Clebsch-Gordan series. For multiplicity two, Hecht has proposed an alternative which can provide correct $1/N_c$ sub-leading orders in large N_c studies.

arXiv:1504.01597v2 [hep-ph] 12 Oct 2015

^a E-mail address: fstancu@ulg.ac.be

I. INTRODUCTION

Since the 1963 classical paper of de Swart [1] where Clebsch-Gordan (CG) coefficients of $SU(3)$ were derived for the most important direct (or Kronecker) products of irreducible representations needed in particle physics at that time, namely $\mathbf{8} \times \mathbf{8}$, $\mathbf{8} \times \mathbf{10}$, $\mathbf{8} \times \mathbf{27}$, $\mathbf{10} \times \mathbf{10}$ and $\mathbf{10} \times \bar{\mathbf{10}}$, many authors devoted their papers or parts of them to the derivation of CG coefficients which were missing in de Swart's paper. As recalled in the next section this amounts to derive the corresponding isoscalar factor for each CG coefficient. In 1963 as well, numerical values for the $SU(3)$ isoscalar factors were published by Edmonds [2]. More tables were given in 1964 by McNamee and Chilton [3].

In recent years the $SU(3)$ flavor group was frequently used to study new hadronic properties and quark systems involving an arbitrary number of quarks as for example in large N_c QCD studies. The existing results seemed to be insufficient so that several authors derived their own tables. Here we show that some of them are particular cases of the analytic expressions obtained by Hecht in 1965 in the context of nuclear physics [4].

The purpose of this note is twofold: a) To draw attention to Hecht's work, which may not be known by particle physicists. Some analytic formulae obtained by Hecht for $SU(3)$ isoscalar factors can straightforwardly be used for particular cases. b) To show that Hecht's results for multiplicity two in the direct products are useful in large N_c studies, which give a qualitative insight into the structure of baryons. In the large $1/N_c$ expansion one has first to analyze formulae at arbitrary N_c and afterwards take $N_c = 3$ in applications.

II. REMINDER OF SOME $SU(3)$ CG PROPERTIES

In the chain $SU(3) \supset SU(2)_I \times U(1)_Y$ each $SU(3)$ CG coefficient factorizes into an $SU(2)$ -isospin CG coefficient and an $SU(3)$ isoscalar factor [1]

$$\left(\begin{array}{cc|c} (\lambda\mu) & (\lambda_a\mu_a) & (\lambda'\mu') \\ YII_3 & Y^a I^a I_3^a & Y' I' I_3' \end{array} \right)_\rho = \left(\begin{array}{cc|c} I & 1 & I' \\ I_3 & I_3^a & I_3' \end{array} \right) \left(\begin{array}{cc|c} (\lambda\mu) & (\lambda^a\mu^a) & (\lambda'\mu') \\ YI & Y^a I^a & Y' I' \end{array} \right)_\rho. \quad (1)$$

where $(\lambda\mu)$ labels an $SU(3)$ irreducible representation (irrep) and the index ρ distinguishes between identical representations occurring in the decomposition of a given direct product where the multiplicity of $(\lambda'\mu') = (\lambda\mu)$ is larger than one. The highest multiplicity considered here is two and in this case a typical example of direct product representations is when

one takes $(\lambda^a \mu^a) = (11)$, which is the adjoint representation of $SU(3)$, also denoted by its dimension **8**. The CG series reads

$$\begin{aligned} (\lambda\mu) \times (11) &= (\lambda + 1, \mu + 1) + (\lambda + 2, \mu - 1) + (\lambda\mu)_1 + (\lambda\mu)_2 \\ &+ (\lambda - 1, \mu + 2) + (\lambda - 2, \mu + 1) + (\lambda + 1, \mu - 2) + (\lambda - 1, \mu - 1). \end{aligned} \quad (2)$$

The isoscalar factors of $SU(3)$ satisfy an orthogonality relation resulting from the orthogonality relations of $SU(3)$ and $SU(2)$ CG coefficients. This is

$$\sum_{Y'' I'' Y^a I^a} \left(\begin{array}{c} (\lambda'' \mu'') \quad (\lambda^a \mu^a) \\ Y'' I'' \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda' \mu') \\ Y I \end{array} \right)_\rho \left(\begin{array}{c} (\lambda'' \mu'') \quad (\lambda^a \mu^a) \\ Y'' I'' \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda\mu) \\ Y' I' \end{array} \right)_\rho = \delta_{\lambda' \lambda} \delta_{\mu' \mu} \delta_{Y' Y} \delta_{I' I}, \quad (3)$$

and

$$\sum_{(\lambda\mu)\rho} \left(\begin{array}{c} (\lambda'' \mu'') \quad (\lambda^a \mu^a) \\ Y'' I'' \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda\mu) \\ Y I \end{array} \right)_\rho \left(\begin{array}{c} (\lambda'' \mu'') \quad (\lambda^a \mu^a) \\ Y'' I'' \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda\mu) \\ Y I \end{array} \right)_\rho = \delta_{Y'' Y''} \delta_{I'' I''} \delta_{Y^a Y^a} \delta_{I^a I^a}. \quad (4)$$

For completeness, we also recall that the isoscalar factors obey the following symmetry properties [4]

$$\begin{aligned} &\left(\begin{array}{c} (\lambda\mu) \quad (\lambda^a \mu^a) \\ Y I \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda' \mu') \\ Y' I' \end{array} \right) = \\ &(-)^{(\lambda - \mu + \lambda^a - \mu^a - \lambda' + \mu' + I + I^a - I')} \left(\begin{array}{c} (\lambda^a \mu^a) \quad (\lambda\mu) \\ Y^a I^a \quad Y I \end{array} \middle\| \begin{array}{c} (\lambda' \mu') \\ Y' I' \end{array} \right). \end{aligned} \quad (5)$$

and

$$\begin{aligned} &\left(\begin{array}{c} (\lambda\mu) \quad (\lambda^a \mu^a) \\ Y I \quad Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda' \mu') \\ Y' I' \end{array} \right) = \\ &(-)^{\frac{1}{3}(\mu' - \mu - \lambda' + \lambda + \frac{3}{2}Y^a) + I' - I} \sqrt{\frac{\dim(\lambda' \mu')(2I + 1)}{\dim(\lambda\mu)(2I' + 1)}} \left(\begin{array}{c} (\lambda' \mu') \quad (\lambda^a \mu^a) \\ Y' I' \quad -Y^a I^a \end{array} \middle\| \begin{array}{c} (\lambda\mu) \\ Y I \end{array} \right). \end{aligned} \quad (6)$$

where $\dim(\lambda\mu) = \frac{1}{2}(\lambda + 1)(\mu + 1)(\lambda + \mu + 2)$ is the dimension of the irrep $(\lambda\mu)$ of $SU(3)$. An alternative notation of the isoscalar factors is $\langle (\lambda\mu) Y I; (\lambda^a \mu^a) Y^a I^a || (\lambda' \mu') Y' I' \rangle$, see Hecht's paper.

III. CALCULATION OF SU(3) CLEBSCH-GORDAN COEFFICIENTS

The usual procedure to calculate CG coefficients is to start from the highest weight basis vector of a representation and use ladder operators, which are U_{\pm} , V_{\pm} and I_{\pm} in SU(3). Their matrix elements were first determined by Biedenharn [5]. Recursion relations among Clebsch-Gordan coefficients are obtained by coupling two states, as in the usual way, like for the rotation group. These recursion relations contain isoscalar factors.

To uniquely define the matrix elements of the ladder operators some phase conventions must be made. For the states in the same isomultiplet the standard Condon and Shortley has been chosen. Accordingly the non-vanishing matrix elements of I_{\pm} are positive. The relative phases between different isomultiplets were defined by the requirement that the non-vanishing matrix elements of V_{\pm} are real and positive [1] (for the phase convention of de Swart see [1], Section 10).

This procedure has been followed by Kaeding [6] who provided a large number of tables for $(\lambda^a \mu^a) = (10), (01), (20), (11), (30)$ and (21) or in dimensional notation **3**, $\bar{\mathbf{3}}$, **6**, **8**, **10** and **15'**.

More recently Hong [7] has derived the isoscalar factors of the direct product of **35** \times **8**, with the purpose of using them to the calculation of baryon magnetic moments and decuplet-to-octet transition magnetic moments. For multiplicity one, all the isoscalar factors are particular cases of the formulae derived by Hecht [4] in his Table 4, up to a phase convention (see next section).

In large N_c QCD Cohen and Lebed [8] derived N_c dependent SU(3) CG coefficients relevant for the coupling of large N_c baryons to mesons. They provided extended tables for the direct products for

$$(\lambda\mu) = \left(1, \frac{N_c - 1}{2}\right), \quad \left(3, \frac{N_c - 3}{2}\right) \quad (7)$$

denoted by "**8**" and "**10**" respectively and $(\lambda^a \mu^a) = (11)$ denoted by **8**. Their results, at multiplicity one, up to an overall phase, can directly be reproduced from Hecht's Table 4. For multiplicity two, for example, "**10**"_a \times **8** \rightarrow "**10**"_a they are different at arbitrary N_c , but identical at $N_c = 3$, as compared to those derived here using Hecht's analytic forms (see next section).

For the same direct products as those of Cohen and Lebed [8] partial tables were previously provided in Ref. [9].

The explicit algebraic expressions derived by Hecht [4] for SU(3) isoscalar factors were intended to nuclear physics applications, in particular to describing rotational states of deformed light nuclei from the $2s - 1d$ shell. The deformed nuclei possess collective states described by Elliott [10, 11] in a model where the SU(3) group is used. Thus the application of SU(3) in nuclear physics in 1958 predates the SU(3) classification of elementary particles of Gell-Mann [12] and Ne'eman [13] in 1961. The basic reason of using SU(3) in nuclear models is that intrinsic levels of nuclei can be described by the harmonic oscillator and SU(3) is the symmetry group of the harmonic oscillator in three dimensions (see, for example, Ref. [14] chapter 8). The physical states of a given angular momentum can be obtained by a projection technique [15].

In addition to the isoscalar factors needed for the $2s - 1d$ shell, Hecht had also derived explicit expressions for the direct product $(\lambda\mu) \times (11)$, considering such results as being of interest, not surprisingly, because (11) is the adjoint representation of SU(3). He used the standard technique of generating CG coefficients through recursion formulae containing matrix elements of the SU(3) generators, but introduced a phase convention different from that of de Swart. The difference is clearly explained in a footnote of Ref. [4]. In addition, when the irrep $(\lambda\mu)$ appears twice in the decomposition of the direct product $(\lambda\mu) \times (11)$, see Eq. (2), he introduced the quantum number ρ to label the independent modes of coupling, such as to have non-zero matrix elements of the SU(3) generators for only one state ρ . Then, according to the Wigner-Eckart theorem, the matrix elements of the generators T^a of SU(3) are

$$\begin{aligned} & \langle (\lambda'\mu') Y' I' I'_3; S' S'_3 | T^a | (\lambda\mu) Y I I_3; S S_3 \rangle = \\ & \delta_{SS'} \delta_{S_3 S'_3} \delta_{\lambda\lambda'} \delta_{\mu\mu'} \sum_{\rho=1,2} \langle (\lambda'\mu') || T^{(11)} || (\lambda\mu) \rangle_{\rho} \begin{pmatrix} (\lambda\mu) & (11) & (\lambda'\mu') \\ Y I I_3 & Y^a I^a I_3^a & Y' I' I'_3 \end{pmatrix}_{\rho}, \end{aligned} \quad (8)$$

where the reduced matrix elements are defined as [4]

$$\langle (\lambda\mu) || T^{(11)} || (\lambda\mu) \rangle_{\rho} = \begin{cases} \sqrt{C(\text{SU}(3))} & \text{for } \rho = 1 \\ 0 & \text{for } \rho = 2 \end{cases}, \quad (9)$$

in terms of the eigenvalue of the Casimir operator $C(\text{SU}(3)) = \frac{1}{3}g_{\lambda\mu}$ where

$$g_{\lambda\mu} = \lambda^2 + \mu^2 + \lambda\mu + 3\lambda + 3\mu. \quad (10)$$

Such a definition is useful for extending the method of calculation of isoscalar factors to other $SU(N)$ groups. It has been applied to the calculation of the matrix elements of $SU(6)$ generators, where one takes into account that $SU(3)$ is a subgroup of $SU(6)$ [17, 18].

The correspondence with other notations is

$$\begin{aligned}\rho = 1 &\iff (\lambda\mu)_2 \iff (\lambda\mu)_a, \\ \rho = 2 &\iff (\lambda\mu)_1 \iff (\lambda\mu)_s.\end{aligned}\tag{11}$$

where s and a stand for symmetric and antisymmetric respectively [19, 20]. Historically, following Gell-Mann, in Eq. (11), it is customary to call the symmetric combinations D coupling and the antisymmetric a combinations F coupling (the F and D notation is used in Ref. [9], for example).

Ambiguities in distinguishing the representations at multiplicity larger than one are typical for all groups, including the permutation group [21].

Another way to derive Clebsch-Gordan coefficients for $SU(3)$ is based on the tensor method (for an introduction see, for example, Ref. [14], Sec. 8.10). This method has been used for the Clebsch-Gordan series " $\mathbf{8}$ " \times " $\mathbf{8}$ " and " $\mathbf{10}$ " \times " $\mathbf{8}$ " in the systematic analysis of large N_c baryons [22].

IV. EXAMPLES

Here we wish to demonstrate the usefulness of Hecht's results, especially for multiplicity two, by using Table 4 of Ref. [4]. We use the same table format as that of de Swart because it helps in comparing with previous results found in the literature and moreover, it allows easy checking of the orthogonality relations (3) and (4). We consider two examples relevant for our purpose.

A. Example 1

The first example, shown in Table I, corresponds to one table obtained by Hong in Ref. [7]. It contains the isoscalar factors for all irreducible representations with $Y = 2$, $I = 2$ from the decomposition of the direct product $\mathbf{35} \times \mathbf{8}$. These are $\mathbf{81}$, $\mathbf{64}$, $\mathbf{35}_s$ and $\mathbf{35}_a$ in this case.

TABLE I. Isoscalar factors for the irreducible representations with $Y = 2$, $I = 2$ from the decomposition of the direct product $\mathbf{35} \times \mathbf{8}$. The first two columns indicate the hypercharge and isospin of $\mathbf{35}$ and $\mathbf{8}$ respectively. The phase convention is that of Hecht [4].

$Y_1 I_1;$	$Y_2 I_2$	$\mathbf{81}$	$\mathbf{64}$	$\mathbf{35}_s$	$\mathbf{35}_a$
$1, \frac{5}{2};$	$1, \frac{1}{2}$	$-\sqrt{\frac{1}{200}}$	$\sqrt{\frac{8}{25}}$	$\sqrt{\frac{5}{8}}$	$\sqrt{\frac{1}{20}}$
$1, \frac{3}{2};$	$1, \frac{1}{2}$	$\sqrt{\frac{144}{200}}$	$\sqrt{\frac{2}{25}}$	0	$-\sqrt{\frac{4}{20}}$
$2, 2;$	$0, 1$	$\sqrt{\frac{10}{200}}$	$\sqrt{\frac{5}{25}}$	$-\sqrt{\frac{2}{8}}$	$\sqrt{\frac{10}{20}}$
$2, 2;$	$0, 0$	$\sqrt{\frac{45}{200}}$	$-\sqrt{\frac{10}{25}}$	$\sqrt{\frac{1}{8}}$	$\sqrt{\frac{5}{20}}$

Note that one must use the symmetry property (5) to recover the phases for $\mathbf{8} \times \mathbf{35}$ as in Ref. [7], because here we consider $\mathbf{35} \times \mathbf{8}$. For the columns $\mathbf{81}$ and $\mathbf{64}$ the absolute values are the same as those of Hong. Incidentally column $\mathbf{81}$ also has the same phases as Hong and column $\mathbf{64}$ has an overall opposite phase. Our results for $\mathbf{35}_s$ and $\mathbf{35}_a$ are entirely different from those of [7] because the definition is different. In applications care must be taken in passing from one convention to another, especially for calculating transition matrix elements.

B. Example 2

The second example is exhibited in Table II and corresponds to a table of Cohen and Lebed [8], containing isoscalar factors with $Y = N_c/3$, $I = 3/2$ from the decomposition of the direct product $\mathbf{10} \times \mathbf{8}$. Cohen and Lebed obtained analytic expressions of the isoscalar factors as a function of N_c needed for large N_c baryon-meson coupling. Our table was obtained as a direct application of Hecht's Table 4, part of which is reproduced in Table III of the Appendix, referring to the irrep $\mathbf{10}$ with multiplicity 2, denoted here by $\mathbf{10}_a$ and $\mathbf{10}_s$ respectively. For completeness, to the three rows listed by Cohen and Lebed we have added a fourth one, corresponding to $Y_1 = N_c/3 - 1$, $I_1 = 2$ and $Y_2 = 1$, $I_2 = 1/2$, in order to check the orthogonality of columns, given by Eq. (3), valid at every N_c . Column

TABLE II. Isoscalar factors for the irreducible representations with $Y = N_c/3$, $I = 3/2$ from the decomposition of the direct product $\mathbf{10} \times \mathbf{8}$ obtained from Table III.

∞

$Y_1 I_1;$	$Y_2 I_2$	"35"	"27"	"10" _a	"10" _s
$\frac{N_c}{3}, \frac{3}{2};$	0, 1	$\sqrt{\frac{12}{16(N_c + 9)}}$	$\sqrt{\frac{5}{4(N_c + 1)}}$	$\sqrt{\frac{45}{N_c^2 + 6N_c + 45}}$	$-\sqrt{\frac{(N_c - 3)(N_c + 5)(N_c + 6)^2}{(N_c + 1)(N_c + 9)(N_c^2 + 6N_c + 45)}}$
$\frac{N_c}{3}, \frac{3}{2};$	0, 0	$\sqrt{\frac{60}{16(N_c + 9)}}$	$-\sqrt{\frac{9}{4(N_c + 1)}}$	$\sqrt{\frac{N_c^2}{N_c^2 + 6N_c + 45}}$	$\sqrt{\frac{45(N_c - 3)(N_c + 5)}{(N_c + 1)(N_c + 9)(N_c^2 + 6N_c + 45)}}$
$\frac{N_c}{3} - 1, 1;$	$1, \frac{1}{2}$	$\sqrt{\frac{15(N_c + 5)}{16(N_c + 9)}}$	$\sqrt{\frac{N_c + 5}{16(N_c + 1)}}$	$-\sqrt{\frac{9(N_c + 5)}{4(N_c^2 + 6N_c + 45)}}$	$\sqrt{\frac{5(N_c - 3)^3}{4(N_c + 1)(N_c + 9)(N_c^2 + 6N_c + 45)}}$
$\frac{N_c}{3} - 1, 2;$	$1, \frac{1}{2}$	$-\sqrt{\frac{(N_c - 3)}{16(N_c + 9)}}$	$\sqrt{\frac{15(N_c - 3)}{16(N_c + 1)}}$	$\sqrt{\frac{15(N_c - 3)}{4(N_c^2 + 6N_c + 45)}}$	$\sqrt{\frac{3(N_c + 5)(N_c + 21)^2}{4(N_c + 1)(N_c + 9)(N_c^2 + 6N_c + 45)}}$

"**35**" has the same phase for all entries as that of Cohen and Lebed and column "**27**" has opposite phase for all entries. It may happen that the phase conventions of de Swart and Hecht coincide sometimes. The column "**10**"_a \equiv "**10**"₂ is entirely different, inasmuch as we use the definition (9) of Hecht to define the representations with multiplicity 2. We have also added the column "**10**"_s \equiv "**10**"₁ where the first three entries vanish at $N_c = 3$, as observed in Ref. [8], but the last entry does not. Such a result may be important for large N_c baryon studies [23].

In large N_c studies the observables are described by operators expressed in terms of SU(6) generators when one considers three flavours, $N_f = 3$. The SU(6) generators are components of an irreducible SU(6) tensor operator which span the invariant subspace of the adjoint representation denoted here by the partition $[21^4]$, or otherwise by its dimensional notation **35**. Like for any other irreducible representation its matrix elements can be expressed in terms of a generalized Wigner-Eckart theorem [16, 17] which factorizes each matrix element into products of Clebsch-Gordan coefficients and a reduced matrix element, like in Eq. 8. The notation is as follows. The generic name for every generator is E^{ia} . An irrep of SU(6) is denoted by the partition $[f]$. Then one can write the matrix element of every SU(6) generator E^{ia} as

$$\begin{aligned} & \langle [f](\lambda'\mu')Y'I'I'_3S'S'_3 | E^{ia} | [f](\lambda\mu)YII_3SS_3 \rangle = \\ & \sqrt{C^{[f]}(\text{SU}(6))} \begin{pmatrix} S & S^i & S' \\ S_3 & S_3^i & S'_3 \end{pmatrix} \begin{pmatrix} I & I^a & I' \\ I_3 & I_3^a & I'_3 \end{pmatrix} \\ & \times \sum_{\rho=1,2} \left(\begin{array}{c} (\lambda\mu) \quad (\lambda^a\mu^a) \\ YI \quad Y^aI^a \end{array} \parallel \begin{array}{c} (\lambda'\mu') \\ Y'I' \end{array} \right)_{\rho} \left(\begin{array}{c} [f] \quad [21^4] \\ (\lambda\mu)S \quad (\lambda^a\mu^a)S^i \end{array} \parallel \begin{array}{c} [f] \\ (\lambda'\mu')S' \end{array} \right)_{\rho}, \end{aligned} \quad (12)$$

where $C^{[f]}(\text{SU}(6))$ is the SU(6) Casimir operator eigenvalue associated to the irreducible representation $[f]$, here being the reduced matrix element, followed by the familiar Clebsch-Gordan coefficients of SU(2)-spin and SU(2)-isospin. The sum over ρ contains products of isoscalar factors of SU(3) and SU(6) respectively. The label ρ is necessary whenever one has to distinguish between irreps $[f'] = [f]$ with multiplicities $m_{[f']}$ larger than one in the Clebsch-Gordan series [18]

$$[f] \times [21^4] = \sum_{[f']} m_{[f']} [f']. \quad (13)$$

The two values for ρ both in SU(6) and SU(3) reflects the multiplicity problem already appearing in the direct product of SU(3) irreducible representations, as discussed in Sec.

II. It is clear that one must make the sum over ρ in all cases. The large N_c behaviour is obtained from the analytic expressions of the isoscalar factors of SU(3) and SU(6). This behaviour is necessary for finding the most dominant contributions in the $1/N_c$ expansion. Examples of physical interest in baryon spectroscopy for the analytic expressions of SU(6) isoscalar factors can be found in Ref. [23] for $[f] = [N_c]$ and $[f] = [N_c - 1, 1]$. Here we discuss the large N_c behaviour resulting from SU(3) isoscalar factors.

For a comparison with Cohen and Lebed [8] let us consider the column "10"_a of Table II alone because the column "10"_s is missing in Ref. [8]. For the first three rows our isoscalar factors are of order $\mathcal{O}(N_c^{-1})$, $\mathcal{O}(N_c^0)$ and $\mathcal{O}(N_c^{-1/2})$ respectively while from Ref. [8] Table II at $Y = N_c/3$, $I = 1/2$, column "10"_a we obtain $\mathcal{O}(N_c^0)$, $\mathcal{O}(N_c^0)$ and $\mathcal{O}(N_c^{-1/2})$ respectively. Thus the large N_c behaviour is different from ours for $I_1 = 3/2, Y_1 = N_c/3, I_2 = 1, Y_2 = N_c/3$ and $I_1 = 3/2, Y_1 = N_c/3, I_2 = 0, Y_2 = 0$. For a proper analysis at large N_c the missing column $\rho = 2$ equivalent to "10"_s, is necessary as required by Eq. (12), even if some isoscalar factors vanish at $N_c = 3$. By summing up the contributions from "10"_a and "10"_s one would expect a similar answer in any convention, provided the SU(6) isoscalar factors are calculated consistently with those of SU(3). Moreover the case $I_1 = 2, Y_1 = N_c/3 - 1, I_2 = 1/2, Y_2 = 1$ is missing in Table II of Ref. [8], at $Y = N_c/3$ $I = 1/2$. Therefore, the results of Ref. [8] should be completed with extra rows and columns, whenever necessary, if one wishes to recover a proper large N_c behaviour. In the physical world of $N_c = 3$ they are sufficient for the exhibited I_1, Y_1, I_2, Y_2 cases.

It would be interesting to consider further applications of Hecht's SU(3) isoscalar factors either in large N_c QCD or in nuclear physics.

Appendix A

In Table III we reproduce part of Table 4 of Hecht's paper [4] which contains the analytic expressions of the isoscalar factors $\langle(\lambda\mu)Y_1I_1; (11)Y_2I_2||(\lambda\mu)YI\rangle$, often used in quark physics. Note that the entry in the column $\rho = 2$ for $Y_2 = 1, I_2 = \frac{1}{2}, I_1 = I + 1/2$ contains a misprint in Hecht's paper which has been here corrected. This means that in the numerator the bracket $(\lambda + \mu + 2 - q + 1)$ has been replaced by $(\lambda + \mu + 2 - q)$ and in the denominator the bracket $(\mu + p - q)$ has been replaced by $(\mu + p - q + 1)$. In Table III we have used $g_{\lambda\mu}$

TABLE III. Isoscalar factors $\langle (\lambda\mu)Y_1I_1; (11)Y_2I_2 || (\lambda\mu)YI \rangle$

 of Hecht's Table 4, p.31 [4] with corrections for the row $Y_2 = 1, I_2 = 1/2, I_1 = I + 1/2$.

	Y_2I_2	I_1	$(\lambda'\mu') = (\lambda\mu)$	
			$\rho = 1$	$\rho = 2$
11	$-1\frac{1}{2}$	$I + 1/2$	$\left[\frac{3(p+1)(\lambda-p)(\mu+2+p)}{2g_{\lambda\mu}(\mu+p-q+1)} \right]^{1/2}$	$\frac{[2g_{\lambda\mu}q - \mu(\lambda+\mu+1)(\lambda+2\mu+6)][(p+1)(\lambda-p)(\mu+2+p)]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)2g_{\lambda\mu}(\mu+p-q+1)]^{1/2}}$
	$-1\frac{1}{2}$	$I - 1/2$	$\left[\frac{3(q+1)(\mu-q)(\lambda+\mu+1-q)}{2g_{\lambda\mu}(\mu+p-q+1)} \right]^{1/2}$	$\frac{[2g_{\lambda\mu}p + \lambda(\mu+2)(\lambda-\mu+3)][(q+1)(\mu-q)(\lambda+\mu+1-q)]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)2g_{\lambda\mu}(\mu+p-q+1)]^{1/2}}$
	00	I	$-\frac{2\lambda+\mu-3p-3q}{[4g_{\lambda\mu}]^{1/2}}$	$\frac{\sqrt{3}}{2} \frac{\lambda\mu(\mu+2)(\lambda+\mu+1) - \mu(\lambda+\mu+1)(\lambda+2\mu+6)p + \lambda(\mu+2)(\lambda-\mu+3)q + 2g_{\lambda\mu}pq}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)g_{\lambda\mu}]^{1/2}}$
	01	$I + 1$	0	$\frac{[2(p+1)(\lambda-p)(\mu+2+p)q(\mu+1-q)(\lambda+\mu+2-q)g_{\lambda\mu}]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)(\mu+p-q+1)(\mu+p-q+2)]^{1/2}}$
	01	$I - 1$	0	$-\frac{[2p(\lambda+1-p)(\mu+1+p)(q+1)(\mu-q)(\lambda+\mu+1-q)g_{\lambda\mu}]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)(\mu+p-q+1)(\mu+p-q)]^{1/2}}$
	01	I	$\frac{[3(\mu+p-q)(\mu+p-q+2)]^{1/2}}{[4g_{\lambda\mu}]^{1/2}}$	$\frac{E}{2[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)g_{\lambda\mu}(\mu+p-q)(\mu+p-q+2)]^{1/2}}$
	$1\frac{1}{2}$	$I + 1/2$	$\left[\frac{3q(\mu+1-q)(\lambda+\mu+2-q)}{2g_{\lambda\mu}(\mu+p-q+1)} \right]^{1/2}$	$\frac{[2g_{\lambda\mu}p + \lambda(\mu+2)(\lambda-\mu+3)][q(\mu+1-q)(\lambda+\mu+2-q)]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)2g_{\lambda\mu}(\mu+p-q+1)]^{1/2}}$
	$1\frac{1}{2}$	$I - 1/2$	$-\left[\frac{3p(\lambda+1-p)(\mu+1+p)}{2g_{\lambda\mu}(\mu+p-q+1)} \right]^{1/2}$	$-\frac{[2g_{\lambda\mu}q - \mu(\lambda+\mu+1)(\lambda+2\mu+6)][p(\lambda+1-p)(\mu+1+p)]^{1/2}}{[\lambda(\lambda+2)\mu(\mu+2)(\lambda+\mu+1)(\lambda+\mu+3)2g_{\lambda\mu}(\mu+p-q+1)]^{1/2}}$

TABLE IV. Values of λ' , μ' , p and q needed for $Y = N_c/3$, $I = 3/2$ to calculate the isoscalar factors of "10" \times 8 using Table III. The label $(\lambda'\mu')$ identifies the irreps of the Clebsch-Gordan series (2) for a given $(\lambda\mu)$ in the left hand side. The isoscalar factors are presented in Table II.

	λ'	μ'	p	q	$(\lambda'\mu')$
"35"	4	$\frac{N_c - 1}{2}$	3	$\frac{N_c - 1}{2}$	$(\lambda + 1, \mu + 1)$
"27"	2	$\frac{N_c + 1}{2}$	2	$\frac{N_c - 1}{2}$	$(\lambda - 1, \mu + 2)$
"10"	3	$\frac{N_c - 3}{2}$	3	$\frac{N_c - 3}{2}$	$(\lambda\mu)$

defined by Eq. (10) and E defined by

$$\begin{aligned}
E = & \lambda(\lambda + \mu + 1)\mu(\mu + 2)(2\lambda + \mu + 6) + 2(\lambda + \mu + 1)\mu \\
& \times [\lambda(\lambda + 2) - (\mu + 2)(\mu + 3)]p - \mu(\lambda + \mu + 1)(\lambda + 2\mu + 6)p^2 \\
& - 2\lambda[(\mu + 1)(\lambda + \mu + 1)(2\lambda + \mu + 6) - \mu g_{\lambda\mu}]q + \lambda(\mu + 2)(\lambda - \mu + 3)q^2 \\
& - 2[\lambda(\lambda + \mu + 1)(2\lambda + \mu + 6) - g_{\lambda\mu}]pq + 2g_{\lambda\mu}(p^2q + pq^2). \quad (A1)
\end{aligned}$$

Table III and the rest Table 4 of Hecht can straightforwardly be applied to a given $(\lambda'\mu')$ with definite values of Y and I , from which one can obtain the integers p and q defined as

$$Y = p + q - \frac{2\lambda' + \mu'}{2}, \quad I = \frac{\mu' + p - q}{2} \quad (A2)$$

given by Hecht where Y is related to the a quantity called ϵ by

$$\epsilon = -3Y. \quad (A3)$$

For $\lambda = 3$ and $\mu = \frac{N_c - 3}{2}$ the values of λ' , μ' together with p and q defined by Eqs. (A2) are listed in Table IV.

We believe there is no reason to reproduce the full Table 4 of Hecht which contains four distinct tables.

Acknowledgments

This research was supported by the Fond de la Recherche Scientifique - FNRS, Belgium,

under the grant 4.4501.05.

-
- [1] J. J. de Swart, Rev. Mod. Phys. **35** (1963) 916 [Erratum-ibid. **37** (1965) 326].
 - [2] A. R. Edmonds, Proc. Roy. Soc. **A268** (1963) 436.
 - [3] P. S. J. McNamee and F. Chilton, Rev. Mod. Phys. **36** (1964) 1005.
 - [4] K. T. Hecht, Nucl. Phys. **62** (1965) 1 .
 - [5] L. C. Biedenharn, Phys. Lett. **3** (1962) 69; Phys. Lett. **3** (1962) 254. See also G. E. Baird and L. C. Biedenharn, J. Math. Phys. **4** (1963) 1449.
 - [6] T. A. Kaeding, Atom. Data Nucl. Data Tabl. **61** (1995) 233 [nucl-th/9502037].
 - [7] S. T. Hong, J. Korean Phys. Soc. **66** (2015) 2, 158.
 - [8] T. D. Cohen and R. F. Lebed, Phys. Rev. D **70** (2004) 096015.
 - [9] D. Diakonov, V. Petrov and A. A. Vladimirov, Phys. Rev. D **88** (2013) 7, 074030.
 - [10] J. P. Elliott, Proc. Roy. Soc. Lond. A **245** (1958) 128.
 - [11] J. P. Elliott, Proc. Roy. Soc. Lond. A **245** (1958) 562.
 - [12] M. Gell-Mann, Phys. Rev. **125** (1962) 1067.
 - [13] Y. Ne'eman, Nucl. Phys. **26** (1961) 222.
 - [14] F. Stancu, "Group theory in subnuclear physics," Oxford Stud. Nucl. Phys. **19** (1996) 1, (Oxford University Press, Oxford).
 - [15] K. T. Hecht, 'Collective models' in 'Selected topics in nuclear spectroscopy' (ed. B. J. Verhaar), North-Holland, Amsterdam, 1964.
 - [16] N. Matagne and F. Stancu, Phys. Rev. D **73** (2006) 114025.
 - [17] N. Matagne and Fl. Stancu, Nucl. Phys. A **826** (2009) 161.
 - [18] N. Matagne and Fl. Stancu, Phys. Rev. D **83** (2011) 056007.
 - [19] D. B. Lichtenberg, *Unitary symmetries and elementary particles*, Academic Press, New York and London (1970), Chapter 8.
 - [20] For details on the derivation of SU(3) Clebsch-Gordan coefficients see J. F. Cornwell, *Group Theory in Physics*, Academic Press, London, San Diego, New York (1984), vol. II, Ch. 11, Sec. 4 and Ch. 16, Secs. 5 and 6.
 - [21] Fl. Stancu and S. Pepin, Few-Body Systems, **26** (2004) 113.
 - [22] R. F. Dashen, E. E. Jenkins and A. V. Manohar, Phys. Rev. D **49** (1994) 4713 [Erratum-ibid.

D **51** (1995) 2489].

[23] N. Matagne and F. Stancu, Rev. Mod. Phys. **87** (2015) 211.