

Non-Hermitian heat engine with all-quantum-adiabatic-process cycle

S. Lin and Z. Song[†]

School of Physics, Nankai University, Tianjin 300071, China

Abstract. As a quantum device, a quantum heat engine (QHE) is described by a Hermitian Hamiltonian. However, since it is an open system, reservoirs have to be imposed phenomenologically without any description in the context of quantum mechanics. A non-Hermitian system is expected to describe an open system which exchanges energy and particles with external reservoirs. Correspondingly, such an exchange can be adiabatic in the context of quantum mechanics. We first propose a non-Hermitian QHE by a concrete simple two-level system, which is an $S = 1/2$ spin in a complex external magnetic field. The non-Hermitian \mathcal{PT} -symmetric Hamiltonian, as a self-contained one, describes both working medium and reservoirs. A heat-engine cycle is composed of completely quantum adiabatic processes. Surprisingly, the heat efficiency is obtained to be the same as that of Hermitian quantum Otto cycle. A classical analogue of this scheme is also presented. Our finding paves the way for revealing what the role of a non-Hermitian Hamiltonian plays in physics.

PACS numbers: 03.65.-w, 11.30.Er, 05.70.-a, 05.70.Ce

[†] songtc@nankai.edu.cn

1. Introduction

Non-Hermitian system has received intensive studies due to two reasons: It can possess full real eigenvalues [1] which is fundamental in the quantum world. A non-Hermitian potential breaks the probability conservation, which is supposed to characterize the source or drain of particles and energy. The connection between the imaginary potentials and the environment has been investigated in discrete system [2]. A non-Hermitian system could be employed to describe an open system which exchanges energy and particles with external reservoirs. An alternative way to deeply understand the exact meaning of a non-Hermitian Hamiltonian in physics is to find out its function which is similar to some phenomena in practice. An exemplified example is the invention that an imaginary potential can be the function of laser or anti-laser [3, 4] when spectral singularity is reached [5].

The topic of the quantum heat engine (QHE) [6, 7, 8, 9, 10, 11, 12, 13] has attracted a lot of interest since it was first proposed by Scovil and Schultz-Dubois [14]. A QHE converts heat into useful mechanical work from quantum working mediums, such as multilevel systems [15], harmonic oscillator systems [16, 17, 18, 19], spins or coupled spins [20], optomechanical systems [21], relativistic particles [22] and so on. Many investigations have been carried out to explore various possible improvement of the heat engine that the efficiency of a QHE can surpass the efficiency of a classical Carnot heat engine, and a QHE can extract work from a single heat bath [23]. Due to the quantum properties of the working medium, or the effects of the quantum heat bath, some unusual and exotic phenomena have been manifested by considering and using a squeezed reservoir [24, 25], quantum coherence [23], or coupled spins [26, 27, 28, 29]. Moreover, via constructing a QHE which is a two-level quantum system and undergoes quantum adiabatic process and energy exchanges with heat baths at different stages in a work cycle, some important aspects of the second law of thermodynamics have been clarified by Kieu [30]. Very recently, many other investigations have been carried out and demonstrated about the Carnot statement of the second law of thermodynamics and the quantum Jarzynski equality in quantum systems described by pseudo-Hermitian Hamiltonians [31].

While, in contrast to those quantum devices, external reservoirs are inevitable. QHEs considered in the literature are mainly described by a Hermitian Hamiltonian and imposed reservoirs. With the proposal and development of theoretical explorations [32, 33, 34], non-Hermitian quantum theories arise as effective descriptions of certain open quantum systems, and therefore induce to an effective non-Hermitian QHE in the presence of absorption and gain. It is natural to establish a non-Hermitian description for QHEs.

In this paper, we employ a non-Hermitian \mathcal{PT} -symmetric Hamiltonian to study a QHE. It is a two-level system, which describes an $S = 1/2$ spin in a complex external magnetic field. The non-Hermitian QHE behaves quite differently from a Hermitian one in all the approaches so far, providing an alternative description. We do not provide a Hamiltonian to describe the working medium, whereas a fully description of working medium and external reservoirs is included. The process of exchanging heat can be adiabatic in the context of quantum mechanics. We will show that such a non-Hermitian system can fully describe a QHE without imposed external reservoirs, which can operate at Otto efficiency for an optimal cycle. In order to get a clear physical picture of the non-Hermitian engine we construct a classical analogue of this scheme, which is also a variable-mass cycle.




Figure 1. (Color online) Schematic diagrams of a variable-mass Otto cycle for the non-Hermitian QHE. The cycle is shown as a closed loop in the (a) $J - \phi$ and (b) $\varepsilon_+ - P_+$ planes, respectively, consisting of four steps, two isospectrum processes $A \rightarrow B$ and $C \rightarrow D$; and two adiabatic processes $B \rightarrow C$ and $D \rightarrow A$. According to Eq. (17), the mechanical work done by the engine equals the surface integral of a nonzero function over the surface enclosed by the loop.

2. Model

We consider a non-Hermitian spin-1/2 system in an external magnetic field, which can be described by the following Hamiltonian

$$H = \vec{B} \cdot \vec{\sigma}, \quad (1)$$

where $\vec{B} = (B_x, B_y, i\gamma J)$ is a time-dependent complex magnetic field and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ is Pauli matrices. Taking

$$\begin{aligned} B_x &= J \cos \phi, B_y = -J \sin \phi, \\ \cos \theta &= i\gamma / \sqrt{1 - \gamma^2}, \end{aligned} \quad (2)$$

we rewrite the Hamiltonian as the form

$$H = J(t) \sqrt{1 - \gamma^2} \begin{pmatrix} \cos \theta & \sin \theta e^{i\phi(t)} \\ \sin \theta e^{-i\phi(t)} & -\cos \theta \end{pmatrix} \quad (3)$$

where $J(t)$ and $\phi(t)$ are real functions of time, and $\theta(\gamma)$ is a complex (real) constant. Obviously, the Hamiltonian H is \mathcal{PT} -symmetric, i.e., $[\mathcal{P}, H] \neq 0$ and $[\mathcal{T}, H] \neq 0$, but $[\mathcal{PT}, H] = 0$, in the sense of $\mathcal{T}i\mathcal{T} = -i$ and $\mathcal{P} = \sigma_x$. The diagonal elements of the matrix can be regarded as imaginary on-site potentials of a two-site tight-binding model, the existence of which violates the law of conservation of mass and energy. It is thought as an open system with the source and drain. The aim of this paper is trying to establish a full quantum mechanical description of the QHE. We will consider a non-Hermitian two-level quantum engine, in which the source and drain are thought as the channel to the heat bath. To this end, we will seek the solution of the Schrodinger equation of the system. Considering the adiabatic time-evolution of an initial eigenstate of $H(0)$ under the

quantum adiabatic condition, we have

$$|\Psi_\lambda(t)\rangle = e^{i\Lambda_\lambda} |\psi_\lambda(t)\rangle, \quad (4)$$

where $|\psi_\lambda(t)\rangle$ ($\lambda = \pm$) is the instantaneous eigenstate of $H(t)$,

$$|\psi_+\rangle = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{-i\phi} \end{pmatrix}, |\psi_-\rangle = \begin{pmatrix} -\sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} e^{-i\phi} \end{pmatrix}, \quad (5)$$

with eigenvalue $\varepsilon_\lambda = \lambda J(t) \sqrt{1 - \gamma^2}$ and $J(t) > 0$ (see Appendix). Phase Λ_λ is the key quantity of this work, which can be expressed as

$$\begin{aligned} \Lambda_\lambda = & -\lambda \sqrt{1 - \gamma^2} \int_0^t J(t) dt \\ & + \frac{1}{2} [\phi(t) - \phi(0)] \left(1 - i\lambda\gamma / \sqrt{1 - \gamma^2} \right). \end{aligned} \quad (6)$$

In the rest of the paper, our investigation does not involve a coherent superposition of $|\psi_\lambda(t)\rangle$. We neglect the real part of Λ_λ and take

$$\Lambda_\lambda = -\frac{i\lambda\gamma}{2} [\phi(t) - \phi(0)] / \sqrt{1 - \gamma^2}. \quad (7)$$

We note that Λ_λ is path independent and proportional to the difference of ϕ . And the nonzero imaginary adiabatic phase is quite essential to the thermal process of QHE, since it can vary the amplitude of instantaneous eigenstates [35], leading to the change of population distributions. The non-Hermitian Hamiltonian is rather different from a Hermitian one, treating the system as a whole in describing the working medium and reservoirs.

Before start dealing with a heat engine cycle, we would like to briefly remark on the time evolution of a non-Hermitian system, which involves both non-unitary evolution and complex geometrical phase of a state. The Schrödinger equation is the common basis of both conventional and non-Hermitian systems. A quantum state evolves along the solution of the Schrödinger equation. According to the solution, the Dirac probability is no longer conservative. In this sense, the non-unitary evolution and the complex geometric phase are nothing but the direct and natural results of the Schrödinger equation.

3. All-quantum-adiabatic cycle

Now we investigate a thermodynamic process for an adiabatic passage of the non-Hermitian system. For a traditional (Hermitian) QHE, the temperature of a quantum working medium is introduced based on the thermalization assumption [9, 10, 30, 36], that is, the working medium has the same distribution of occupation probability as the heat bath in thermodynamic equilibrium. Then for such a small quantum system, the probability distribution is related to a parameter called spectral temperature. In this paper, we still employ such a concept for the non-Hermitian two-level system, as an assumption. Consider an initially prepared mixed state, which has Dirac probability $P_\lambda(t=0)$ on the eigenstate $|\psi_\lambda(t=0)\rangle$ of $H(0)$ with eigenenergy $\lambda J_0 \sqrt{1 - \gamma^2}$. The system parameters are taken as (J_0, ϕ_0) initially where $J_0 = J(0)$ and $\phi_0 = \phi(0)$. And the initial populations for the upper and lower levels are $P_+(0) = p_0$ and $P_-(0) = 1 - p_0$, respectively. We express this mixed state by a density matrix

$$\rho(0) = \sum_{\lambda=\pm} P_\lambda(0) |\psi_\lambda(0)\rangle \langle \psi_\lambda(0)|, \quad (8)$$

Table 1. State parameters for a quantum variable-mass Otto cycle illustrated in Fig 1. Here T, ε_+ and U are in the unit of $\sqrt{1 - \gamma^2}$.

	J	ϕ	T	ε_+	P_+, P_-	U	S
A	J_1	ϕ_1	$\frac{2J_1}{k_B \ln(p_0^{-1} - 1)}$	J_1	$p_0, 1 - p_0$	$J_1 (2p_0 - 1)$	$\frac{-k_B p_0 \ln p_0}{-k_B (1 - p_0) \ln (1 - p_0)}$
B	J_1	ϕ_2	$\frac{2J_1}{k_B \ln[(p_0^{-1} - 1)\xi^{-2}]}$	J_1	$p_0 \xi, (1 - p_0) \xi^{-1}$	$p_0 \frac{(\xi + \xi^{-1})}{-\xi^{-1} J_1} J_1$	$\frac{-k_B p_0 \xi \ln(p_0 \xi)}{-k_B (1 - p_0) \xi^{-1} \ln[(1 - p_0) \xi^{-1}]}$
C	J_2	ϕ_2	$\frac{2J_2}{k_B \ln[(p_0^{-1} - 1)\xi^{-2}]}$	J_2	$p_0 \xi, (1 - p_0) \xi^{-1}$	$p_0 \frac{(\xi + \xi^{-1})}{-\xi^{-1} J_2} J_2$	$\frac{-k_B p_0 \xi \ln(p_0 \xi)}{-k_B (1 - p_0) \xi^{-1} \ln[(1 - p_0) \xi^{-1}]}$
D	J_2	ϕ_1	$\frac{2J_2}{k_B \ln(p_0^{-1} - 1)}$	J_2	$p_0, 1 - p_0$	$J_2 (2p_0 - 1)$	$\frac{-k_B p_0 \ln p_0}{-k_B (1 - p_0) \ln (1 - p_0)}$

Table 2. Variances of state parameters related to four processes in a quantum variable-mass Otto cycle illustrated in Fig 1. Here ΔT and ΔU are in the unit of $\sqrt{1 - \gamma^2}$.

	ΔT	ΔU	ΔQ	ΔW	ΔS
$A \rightarrow B$	$\frac{4J_1 \ln \xi}{k_B \ln[(p_0^{-1} - 1)\xi^{-2}] \ln(p_0^{-1} - 1)}$	$p_0 \frac{(\xi + \xi^{-1} - 2)}{+ (1 - \xi^{-1})} J_1$	ΔU	0	$\frac{-k_B p_0 \ln(p_0^{\xi^{-1}} \xi^\xi) - k_B (1 - p_0)}{\times \ln[(1 - p_0)^{\xi^{-1} - 1} (\xi^{-1})^{\xi^{-1}}]}$
$B \rightarrow C$	$\frac{2(J_2 - J_1)}{k_B \ln[(p_0^{-1} - 1)\xi^{-2}]}$	$p_0 \frac{(\xi + \xi^{-1})(J_2 - J_1)}{-\xi^{-1}(J_2 - J_1)}$	0	ΔU	0
$C \rightarrow D$	$-\frac{4J_2 \ln \xi}{k_B \ln[(p_0^{-1} - 1)\xi^{-2}] \ln(p_0^{-1} - 1)}$	$-p_0 \frac{(\xi + \xi^{-1} - 2)}{(1 - \xi^{-1})} J_2$	ΔU	0	$\frac{k_B p_0 \ln(p_0^{\xi^{-1}} \xi^\xi) + k_B (1 - p_0)}{\times \ln[(1 - p_0)^{\xi^{-1} - 1} (\xi^{-1})^{\xi^{-1}}]}$
$D \rightarrow A$	$\frac{2(J_1 - J_2)}{k_B \ln(p_0^{-1} - 1)}$	$(J_1 - J_2)(2p_0 - 1)$	0	ΔU	0

where eigenstate $|\psi_\lambda(0)\rangle$ is normalized in the context of Dirac inner product (see Appendix). Here we assume that the Boltzmann distribution still refers to Dirac probability even for a non-Hermitian system. When a two-level system couples to a heat bath, both systems obey the thermal equilibrium Boltzmann distributions and then the two-level mixed state has its temperature. A density matrix would represent a thermal equilibrium state with temperature

$$T_0 = \frac{2J_0 \sqrt{1 - \gamma^2}}{k_B \ln(p_0^{-1} - 1)}, \quad (9)$$

where k_B is the Boltzmann constant. In quantum statistics, the definition of temperature does not require the preservation of particle probability. It is only determined by the distribution of particle probability on each level. For a two-level system, one can always find a temperature which only depends on the ratio of the probabilities and the energy difference of two levels, i.e.,

$$\ln \frac{P_+}{P_-} = -\frac{\varepsilon_+ - \varepsilon_-}{k_B T} \quad (10)$$

where P_λ and ε_λ are the particle probability and the energy of two levels. At time t , state $|\psi_\lambda(0)\rangle$ evolves into state $|\Psi_\lambda(t)\rangle$ under the time-dependent Hamiltonian $H(t)$. We note that the initial state does not involve a coherent superposition of $|\psi_+(0)\rangle$ and $|\psi_-(0)\rangle$. Then the evolved mixed state does not involve a coherent superposition of $|\Psi_+(t)\rangle$ and $|\Psi_-(t)\rangle$. We only concern about the time evolution of the pure state $|\Psi_\lambda(t)\rangle$.

It is an acceptable consensus that the Schrödinger equation is the common basis for both Hermitian and non-Hermitian quantum mechanics. Whether the operator $H(t)$ is Hermitian or not, an evolved state, vector $|\Psi_\lambda(t)\rangle$ should be the solution of the Schrödinger equation

$$i \frac{\partial}{\partial t} |\Psi_\lambda(t)\rangle = H(t) |\Psi_\lambda(t)\rangle, \quad (11)$$

with the initial condition $|\Psi_\lambda(0)\rangle = |\psi_\lambda(0)\rangle$. Accordingly, the density matrix of the evolved mixed state becomes

$$\rho(t) = \sum_{\lambda=\pm} P_\lambda(0) |\Psi_\lambda(t)\rangle \langle \Psi_\lambda(t)|, \quad (12)$$

where $|\Psi_\lambda(t)\rangle$ is probably not Dirac normalized so as to contribute to the population $P_\lambda(t)$. However, it is tough to get an analytical solution for an arbitrary time-dependent operator $H(t)$. Fortunately, when we only consider the time evolution under the quantum adiabatic condition, i.e., there is no transition between two levels, an approximate solution can be obtained. Then the density matrix of the evolved mixed state can be expressed as

$$\rho(t) = \sum_{\lambda=\pm} P_\lambda(t) |\psi_\lambda(t)\rangle \langle \psi_\lambda(t)|, \quad (13)$$

where

$$P_\lambda(t) = P_\lambda(0) \exp[-2\text{Im}(\Lambda_\lambda)]. \quad (14)$$

And the corresponding temperature of the thermal state $\rho(t)$ is

$$T(t) = \frac{2J(t)\sqrt{1-\gamma^2}}{k_B \ln[(p_0^{-1}-1)\xi^{-2}]}, \quad (15)$$

where $\xi = \exp\{\gamma[\phi(t) - \phi_0]/\sqrt{1-\gamma^2}\}$ characterizes the amplification or attenuation of the whole probability in the working medium.

Under the quantum adiabatic condition, the solution of Eq. (4) tells us that any evolution along an arbitrary path in the $J - \phi$ plane is always adiabatic in the context of quantum mechanics, i.e., there is no transition between two levels. Nevertheless, it does not mean that the occupation probabilities for the levels and their distributions are always remaining invariant due to the contribution of the Λ_λ in Eq. (7), which are unusual and exotic properties in the non-Hermitian system in contrast to the Hermitian one. Moreover, we would demonstrate that the “adiabatic” in our paper for the non-Hermitian Hamiltonian may be different from the one in the Hermitian system

because the occupation probability for each level is not invariant due to the image part of Λ_λ . But what both adiabatic processes of the Hermitian and non-Hermitian system have in common is that there is no tunneling between each level. Therefore here we could call it “generalized adiabatic” in our non-Hermitian system. Since all the processes considered in our work are “generalized adiabatic”, we only write it as the “adiabatic” for simplification. Now we consider a loop in the $J - \phi$ plane as a thermal cycle.

According to the first law of thermodynamics, the total heat transferred in the cycle is

$$\begin{aligned}\Delta Q &= \oint_c dQ = \oint_c \sum_\lambda \varepsilon_\lambda dP_\lambda \\ &= \frac{2\gamma}{Z} \iint_A \cosh\left(\ln \sqrt{\frac{1}{p_0} - 1} - \ln \xi\right) dJ d\phi,\end{aligned}\quad (16)$$

where $Z = 2 \cosh(\ln \sqrt{\frac{1}{p_0} - 1})$ is the partition function. Here A denotes the area enclosed by the loop c in the $J - \phi$ plane. We can see that ΔQ is nonzero for a non-trivial cycle, since a hyperbolic cosine function is positive definite. Similarly, the net work done in the cycle is

$$\Delta W = \oint_c dW = \oint_c \sum_\lambda P_\lambda d\varepsilon_\lambda = -\Delta Q, \quad (17)$$

which leads to

$$\Delta U = \Delta Q + \Delta W = 0. \quad (18)$$

Furthermore, a straightforward derivation shows that the entropy variation in the loop is

$$\Delta S = \oint_c \frac{1}{T} dQ = \oint_c \frac{1}{T} \sum_\lambda \varepsilon_\lambda dP_\lambda = 0. \quad (19)$$

According with the traditional thermodynamics, these indicate that the internal energy U and entropy S are state functions, which are only determined by the system parameters (J, ϕ) . We would like to point out that the facts of $\Delta Q \neq 0$ and $\Delta W \neq 0$ in the loop are exclusive for a non-Hermitian system. Taking $\gamma = 0$, the system is reduced to a Hermitian one. We can see that $\Delta Q = \Delta W = 0$, which indicates that the Hermitian system is trivial as a QHE. More detailed, as mentioned above, the complex parameter represents the effect of reservoirs phenomenologically. Once $\gamma = 0$, the working medium is decoupled from reservoirs. There is no exchange between the working medium and the reservoirs. Then the heat engine becomes trivial, since we do not assume extra reservoirs imposed on the system. This is different from the conventional Hermitian QHE, which is always accompanied by the reservoirs. This reflects the key point of our work, replacing the imposed reservoirs by the imaginary parameter γ . In the context of quantum mechanics, processes for an arbitrary loop is spontaneously adiabatic without any extra condition. This may enhance the feasibility of experimental realizations of the present scheme.

4. Cycle with Otto efficiency

So far we have established an alternative description for QHEs via a non-Hermitian Hamiltonian. Remarkably, all the thermal processes are adiabatic in a quantum mechanics manner. By this we mean that irrespective of whether there is energy transfer between the heat engine and reservoirs or not all the processes are adiabatic evolutions driven by a time-dependent Hamiltonian. To understand such quantum processes, it is useful to contrast them to classical thermal processes.




Figure 2. (Color online) A classical analogue to the quantum variable-mass cycle consisting of four steps, two isothermal processes and two adiabatic processes. Here the isothermal process is associated with particle transfer. The cycle is shown as a closed loop in the (a) $P - V$ and (b) $T - M$ planes, respectively.

In the following, we try to connect such a description to a classical engine, showing that the non-Hermitian QHE is not a non-physics toy model.

To this end, we consider a specific cycle, which consists of four processes connecting four points, $A: (J_1, \phi_1)$, $B: (J_1, \phi_2)$, $C: (J_2, \phi_2)$, and $D: (J_2, \phi_1)$. There are two types of processes involved: (I) Isospectrum process, $A \rightarrow B$ and $C \rightarrow D$; (II) Adiabatic process, $B \rightarrow C$ and $D \rightarrow A$. A schematic illustration is listed in Figs. 1(a) and 1(b). In type I process, varying ϕ can change the populations on two levels due to the imaginary adiabatic phase Λ_λ , but leave the level structure unchanged. The level structure can be completely characterized by quantity ε_+ . While, in type II process, varying J can change the level structure ε_+ , whereas leave populations on two levels unchanged. Based on the adiabatic solution of Eq. (4), quantities, including the state parameters at each point and their variances in each process, are obtained and listed in Tables 1 and 2, respectively. According to the definition in the Refs. [30, 9, 10], our calculations indicate that such an engine can operate at Otto efficiency

$$\eta = \frac{Q_1 + Q_2}{Q_2} = 1 - \frac{J_2}{J_1}. \quad (20)$$

Such a non-Hermitian QHE does not violate the Carnot efficiency limit which is of great significance in thermodynamics. This indicates that the non-Hermitian Hamiltonian still obeys some underlying rules in nature.

We note that the probability is not conservative in the cycle, which differs from that in a Hermitian system. We refer to this cycle as a variable-mass Otto cycle. However, we would like to point out that the particle-number conservation still holds in such a cycle [37]. Here we use the term variable-mass in order to correspond with its classical analogue. For a particle-exchange heat engine, it has been investigated in Ref. [38]. Moreover, to understand such a non-Hermitian cycle and its related issues, we propose a variable-mass cycle for classical ideal gas in the following.

5. Classical analogue

In order to obtain a clear physical picture of the non-Hermitian QHE we construct a classical analogue of this scheme, which is a variable-mass cycle and consists of the following four steps, which is schematically illustrated in Fig. 2.

1. Reversible isothermal addition of gas at the high temperature T_1 . During this step a volume ΔV_1 of the gas with the same temperature T_1 and the same density as the working medium is added. It does no work on the surroundings. The temperature of the gas does not change during the process, and thus the addition is isothermal. The gas addition is associated by absorption of heat energy Q_1 and increase of entropy $\Delta S = Q_1/T_1$ from the high temperature reservoir.

2. Reversible adiabatic expansion of gas. For this step, the gas is allowed to expand and does work on the surroundings, whose expansion causes it to cool to the low temperature T_2 , with the entropy remaining unchanged.

3. Reversible isothermal deduction of gas at the low temperature T_2 . During this step a volume ΔV_2 of the gas with the same temperature T_2 and the same density as the working medium is removed. Such an amount of the gas contains heat energy Q_2 and entropy $\Delta S = Q_2/T_2$, which flows to the low temperature reservoir.

4. Reversible adiabatic compression of gas. During this step, the surroundings do work on the gas, increasing its internal energy and compressing it, causing the temperature to rise to T_1 . The gas goes back to the state at the start.

For a standard cycle, mass is conserved, i.e., the working medium does not lose or increase mass. The present cycle transfers mass from hot to cool reservoirs, associating with the heat transfer. The essence of this cycle is that there is a volume of gas transferred from hot to cool reservoirs, rather than transfer the heat only. Then the heat efficiency is always $1 - T_2/T_1$, which is independent of the mass transferred. Therefore, the proposed variable-mass cycle would help us to make a better understanding of the non-Hermitian cycle in some sense.

6. Summary and discussion

We have studied a time-dependent non-Hermitian \mathcal{PT} -symmetric two-level system, which could be regarded as a QHE. A non-Hermitian Hamiltonian turns out to be quite different from a Hermitian one. It does not solely describe the working medium, but offers a full description including the working medium and external reservoirs. We have shown that any cycle can be implemented by an adiabatic time-evolution along a quantum adiabatic passage. As an example, a specific cycle which operates at Otto efficiency is presented. Moreover, we have also discussed a classical analogue of this scheme, which corresponds to a variable-mass cycle, transferring not only heat but also mass from hot to cool reservoirs. Our main conclusion is that the proposed non-Hermitian Hamiltonian can provide an alternative description for the QHE, leading to a deeper and better understanding of the role that a non-Hermitian Hamiltonian takes in physics.

Finally, we would like to point that there is a long way to go for connecting the current non-Hermitian 2-level system to a Hermitian description of QHE by using a rigorous mathematical formulation. It is due to the subtle relation between Hermitian and non-Hermitian Hamiltonians: on the one hand, a non-Hermitian Hamiltonian is usually thought to be a better candidate for describing some natural processes in an open system, which do not follow the conservation of law of mass and energy. However, on the other hand, it is still a challenge to get a perfect connection between a non-Hermitian Hamiltonian and a Hermitian one with a clear physical picture, although

many efforts have been dedicated to this topic [2].

7. Appendix

The instantaneous eigenstates of $H(t)$ and $H^\dagger(t)$ with eigenvalue $\varepsilon_\lambda = \lambda J(t) \sqrt{1 - \gamma^2}$ are

$$|\psi_+\rangle = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{-i\phi} \end{pmatrix}, |\psi_-\rangle = \begin{pmatrix} -\sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} e^{-i\phi} \end{pmatrix}, \quad (21)$$

and

$$|\eta_+\rangle = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} e^{i\phi} \end{pmatrix}^*, |\eta_-\rangle = \begin{pmatrix} -\sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} e^{i\phi} \end{pmatrix}^*, \quad (22)$$

respectively, which can construct biorthonormal complete set $\{|\psi_\lambda\rangle, |\eta_\lambda\rangle\}$ ($\lambda = \pm$), i.e.,

$$\langle \eta_{\lambda'} | \psi_\lambda \rangle = \delta_{\lambda\lambda'}, \sum_\lambda |\psi_\lambda\rangle \langle \eta_\lambda| = 1. \quad (23)$$

Here we demonstrate that when computing the density matrix or detecting the probability in experiments, both eigenstates of $H(t)$ must be normalized in Dirac inner product, i.e., we need to take $|\psi_\lambda\rangle \rightarrow |\psi_\lambda\rangle / \sqrt{\Theta}$ with $\Theta = |\cos \frac{\theta}{2}|^2 + |\sin \frac{\theta}{2}|^2$. Now we extend the adiabatic theorem to the non-Hermitian Hamiltonian and we will show that such an extension is correct. This extension can be simply applied by taking the biorthonormal inner product to replace the Dirac inner product. Then under the quantum adiabatic condition

$$\left| \frac{\langle \eta_- | \frac{\partial H}{\partial t} | \psi_+ \rangle}{(\varepsilon_+ - \varepsilon_-)^2} \right| = \left| \frac{\dot{\phi}}{4J(t)(1 - \gamma^2)} \right| \ll 1, \quad (24)$$

phase Λ_λ can be expressed as

$$\begin{aligned} \Lambda_\lambda &= - \int_0^t \varepsilon_\lambda dt + i \int_0^t \langle \eta_\lambda | \frac{\partial}{\partial t} | \psi_\lambda \rangle dt \\ &= - \lambda \sqrt{1 - \gamma^2} \int_0^t J(t) dt \\ &\quad + \frac{1}{2} [\phi(t) - \phi_0] \left(1 - i\lambda\gamma / \sqrt{1 - \gamma^2} \right). \end{aligned} \quad (25)$$

Similar works have been done in many systems [39, 40, 41, 42, 43], and this conclusion can also be obtained by an exact solution of Schrödinger equation

$$i \frac{\partial}{\partial t} |\Psi\rangle = H |\Psi\rangle, \quad (26)$$

in this concrete example. Here our \mathcal{PT} -symmetric Hamiltonian H could be used to describe an open system, which is different from [44] that is used to describe a closed system. Actually, the propagator of the Hamiltonian H

$$U = \mathcal{T} \exp(-i \int_0^t H(t) dt) \quad (27)$$

can be obtained as the form

$$\begin{aligned}
U_{11} &= \left[\cos(\Omega\tau) - i \frac{\Delta}{2\Omega} \sin(\Omega\tau) \right] e^{i\omega\tau/2}, \\
U_{22} &= \left[\cos(\Omega\tau) + i \frac{\Delta}{2\Omega} \sin(\Omega\tau) \right] e^{-i\omega\tau/2}, \\
U_{12} &= -\frac{i}{\Omega} \sin(\Omega\tau) e^{i\omega\tau/2}, \\
U_{21} &= -\frac{i}{\Omega} \sin(\Omega\tau) e^{-i\omega\tau/2},
\end{aligned} \tag{28}$$

when we take $\phi(\tau) = \omega\tau$. Here \mathcal{T} is time-order operator and

$$\tau = \int_0^t J(t) dt, \Omega = \frac{1}{2} \sqrt{4 + \Delta^2}, \tag{29}$$

$$\Delta = 2i\gamma + \omega. \tag{30}$$

The adiabatic condition in Eq. (24) reduces to

$$\left| \frac{\omega}{1 - \gamma^2} \right| \ll 1, \tag{31}$$

under which the Taylor expansion gives

$$\begin{aligned}
-\Omega\tau &= -\sqrt{1 - \gamma^2} \sqrt{1 + \frac{i\gamma\omega}{1 - \gamma^2}} \tau \\
&\approx -\int_0^t \sqrt{1 - \gamma^2} J(t) dt - \frac{i\gamma\omega\tau}{2\sqrt{1 - \gamma^2}}.
\end{aligned} \tag{32}$$

This solution indicates that the adiabatic process exists under a certain condition, i.e., there is no tunnelling between instantaneous eigenstates during the time evolution. Then the corresponding geometrical phase can be obtained naturally. While, in the article [45], dynamics near exceptional point is considered and it is natural that the adiabatic theorem cannot hold in that situation. The process cannot hold when the system closes to the exceptional point (or quantum phase transition point for breaking \mathcal{PT} symmetry). Actually, similar behavior can occur in a Hermitian system, for example, an adiabatic process cannot be achieved near a quantum phase transition point. A simple demonstration is Landau-Zener formula.

Moreover, in the biorthonormal formalism, compared with right ket $|\Psi\rangle$ which obeys the Eq. (26), the left bra $\langle\Phi|$ follows the different evolution equation

$$i \frac{\partial}{\partial t} |\Phi\rangle = H^\dagger |\Phi\rangle, \tag{33}$$

and correspondingly the propagator

$$\tilde{U} = \mathcal{T} \exp(-i \int_0^t H^\dagger(t) dt) \tag{34}$$

can be obtained as the form

$$\tilde{U}_{11} = \left[\cos(\Omega^*\tau) - i \frac{\Delta^*}{2\Omega^*} \sin(\Omega^*\tau) \right] e^{i\omega\tau/2},$$

$$\begin{aligned}
\tilde{U}_{22} &= \left[\cos(\Omega^* \tau) + i \frac{\Delta^*}{2\Omega^*} \sin(\Omega^* \tau) \right] e^{-i\omega\tau/2}, \\
\tilde{U}_{12} &= -\frac{i}{\Omega^*} \sin(\Omega^* \tau) e^{i\omega\tau/2}, \\
\tilde{U}_{21} &= -\frac{i}{\Omega^*} \sin(\Omega^* \tau) e^{-i\omega\tau/2},
\end{aligned} \tag{35}$$

then we can prove

$$\tilde{U}^\dagger U = 1. \tag{36}$$

As a result, it is unitary according to the biorthonormal inner product. However, we would like to point out that the complex phase for a non-Hermitian system is a natural result (see references [39, 40, 41, 42, 43]). The Berry phase and the unitarity of time evolution are compatible with each other.

Next, explicit derivations of Eq. (16), (17), and (19) are given. Together with $P_+ = p_0 \xi$, $P_- = (1 - p_0) \xi^{-1}$ (here ξ is only the function of ϕ) and ε_λ , we get the heat exchange, the work done, and the variation of entropy of any arbitrary path in the $J - \phi$ plane as following,

$$\begin{aligned}
\Delta Q &= \oint_c dQ = \oint_c \sum_\lambda \varepsilon_\lambda dP_\lambda \\
&= \frac{\sqrt{1-\gamma^2}}{Z} \oint_c J d \left[e^{-(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi)} - e^{(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi)} \right] \\
&= -\frac{2\sqrt{1-\gamma^2}}{Z} \oint_c J d \left[\sinh \left(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi \right) \right] \\
&= \frac{2\gamma}{Z} \oint_c J \cosh \left(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi \right) d\phi \\
&= \frac{2\gamma}{Z} \iint_A \cosh \left(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi \right) dJ d\phi,
\end{aligned} \tag{37}$$

and

$$\begin{aligned}
\Delta W &= \oint_c dW = \oint_c \sum_\lambda P_\lambda d\varepsilon_\lambda \\
&= \frac{\sqrt{1-\gamma^2}}{Z} \oint_c \sum_\lambda \left[\lambda e^{-\lambda(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi)} \right] dJ \\
&= \frac{\sqrt{1-\gamma^2}}{Z} \oint_c \left[e^{-(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi)} - e^{(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi)} \right] dJ \\
&= -\frac{2\sqrt{1-\gamma^2}}{Z} \oint_c \sinh \left(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi \right) dJ \\
&= -\frac{2\gamma}{Z} \iint_A \cosh \left(\ln \sqrt{\frac{1}{p_0}-1} - \ln \xi \right) dJ d\phi.
\end{aligned} \tag{38}$$

Obviously we have

$$\Delta Q + \Delta W = 0. \tag{39}$$

Similarly, the variation of entropy is

$$\begin{aligned}
\Delta S &= \oint_c \frac{1}{T} dQ = \oint_c \frac{1}{T} \sum_{\lambda} \varepsilon_{\lambda} dP_{\lambda} \\
&= -\frac{2k_B}{Z} \oint_c \left(\ln \sqrt{\frac{1}{p_0} - 1} - \ln \xi \right) d \left[\sinh \left(\ln \sqrt{\frac{1}{p_0} - 1} - \ln \xi \right) \right] \\
&= \frac{2k_B \gamma}{Z \sqrt{1 - \gamma^2}} \oint_c \left(\ln \sqrt{\frac{1}{p_0} - 1} - \ln \xi \right) \cosh \left(\ln \sqrt{\frac{1}{p_0} - 1} - \ln \xi \right) d\phi \\
&= 0.
\end{aligned} \tag{40}$$

Acknowledgement

We acknowledge the support of the National Basic Research Program (973 Program) of China under Grant No. 2012CB921900 and the CNSF (Grant No. 11374163).

References

- [1] Bender C M and Boettcher S 1998 *Phys. Rev. Lett.* **80** 5243
- [2] Jin L and Song Z 2010 *Phys. Rev. A* **81** 032109; 2011 *Phys. Rev. A* **83** 052102; 2011 *J. Phys. A: Math. Theor.* **44** 375304
- [3] Longhi S 2010 *Phys. Rev. A* **82** 031801
- [4] Zhang X Z, Jin L and Song Z 2013 *Phys. Rev. A* **87** 042118
- [5] Mostafazadeh A 2009 *Phys. Rev. Lett.* **102** 220402
- [6] Alicki R 1979 *J. Phys. A* **12** L103
- [7] E Geva and Kosloff R 1992 *J. Chem. Phys.* **96** 3054
- [8] Bender C M, Brody D C and Meister B K 2000 *J. Phys. A: Math. Gen.* **33** 4427
- [9] Quan H T, Liu Y X, Sun C P and Nori F 2007 *Phys. Rev. E* **76** 031105
- [10] Henrich M J, Rempp F and Mahler G 2007 *Eur. Phys. J. Spec. Top.* **151** 157
- [11] Abah O, Roßnagel J, Jacob G, Deffner S, Schmidt-Kaler F, Singer K and Lutz E 2012 *Phys. Rev. Lett.* **109** 203006
- [12] Zhang X Y, Huang X L and Yi X X 2014 *J. Phys. A: Math. Theor.* **47** 455002 (10pp)
- [13] Altintas F, Hardal Ali Ü C and Müstecaplıoğlu Özgür E 2015 *Phys. Rev. A* **91** 023816
- [14] Scovil H and Schulz-Dubois E 1959 *Phys. Rev. Lett.* **2** 262
- [15] Quan H T, Zhang P and Sun C P 2005 *Phys. Rev. E* **72** 056110
- [16] Lin B and Chen J 2003 *Phys. Rev. E* **67** 046105
- [17] Kosloff R 1984 *J. Chem. Phys.* **80** 1625
- [18] Scully M O 2002 *Phys. Rev. Lett.* **88** 050602
- [19] Rezek Y and Kosloff R 2006 *New J. Phys.* **8** 83
- [20] He J Z, Chen J C and Hua B 2002 *Phys. Rev. E* **65** 036145
- [21] Zhang K Y, Bariani F and Meystre P 2014 *Phys. Rev. Lett.* **112** 150602
- [22] Muñoz E and Peña F J 2012 *Phys. Rev. E* **86** 061108
- [23] Scully M O, Zubairy M S, Agarwal G S and Walther H 2003 *Science* **299** 862
- [24] Huang X L, Wang Tao and Yi X X 2012 *Phys. Rev. E* **86** 051105
- [25] Roßnagel J, Abah O, Schmidt-Kaler F, Singer K and Lutz E 2014 *Phys. Rev. Lett.* **112** 030602
- [26] Zhang T, Liu W T, Chen P X and Li C Z 2007 *Phys. Rev. A* **75** 062102
- [27] Thomas G and Johal R S 2011 *Phys. Rev. E* **83** 031135
- [28] Huang X L, Wang L C and Yi X X 2013 *Phys. Rev. E* **87** 012144
- [29] Huang X L, Liu Y, Wang Z and Niu X Y 2014 *Eur. Phys. J. Plus* **129** 4
- [30] Kieu T D 2004 *Phys. Rev. Lett.* **93** 140403; 2006 *Eur. Phys. J. D* **39** 115
- [31] Gardas B, Deffner S and Saxena A 2016 *Sci. Rep.* **6** 23408
- [32] Muga J G, Palao J P, Navarro B and Egusquiza I L 2004 *Phys. Rep.* **395** 357
- [33] Graefe Eva-Maria, Korsch Hans Jürgen and Niederle Astrid Elisa 2010 *Phys. Rev. A* **82** 013629
- [34] Lee Tony E and Chan Ching-Kit 2014 *Phys. Rev. X* **4** 041001

- [35] Lin S, Zhang X Z and Song Z 2015 *Phys. Rev. A* **92** 012117
- [36] Gemmer J, Michel M and Mahler G 2004 *Quantum Thermodynamics* (Springer, Berlin, Heidelberg)
- [37] For a non-Hermitian Hamiltonian, conservations of probability and particle are two different concepts, which is investigated in Ref. Jin L and Song Z 2013 *Ann. Phys.* **330** 142
- [38] Humphrey T E and Linke H 2005 *Physica E* **29** 390
- [39] Gao X -C, Xu J -B and Qian T -Z 1992 *Phys. Rev. A* **46** 3626
- [40] Nesterov A I and Aceves de la Cruz F 2008 *J. Phys. A: Math. Theor.* **41** 485304 (25pp)
- [41] Nesterov A I and Ovchinnikov S G 2008 *Phys. Rev. E* **78** 015202(R)
- [42] Cui X -D and Zheng Y 2012 *Phys. Rev. A* **86** 064104
- [43] Liang Shi-Dong and Huang Guang-Yao 2013 *Phys. Rev. A* **87** 012118
- [44] Gong Jiangbin and Wang Qing-hai 2013 *J. Phys. A: Math. Theor.* **46** 485302 (20pp)
- [45] Milburn T J, Doppler J, Holmes C A, Portolan S, Rotter S and Rabl P arXiv:1410.1882.