

PREPARED FOR SUBMISSION TO JHEP

# Cartan's Supersymmetry and the Decay of a $H^0(0^+)$

**Sadataka Furui**

*Graduate School of Science and Engineering, Teikyo University  
2-17-12 Toyosatodai, Utsunomiya, 320-0003 Japan*

*E-mail:* [furui@umb.teikyo-u.ac.jp](mailto:furui@umb.teikyo-u.ac.jp)

**ABSTRACT:** We compare the decay of a Higgs boson  $H^0(0^+) \rightarrow \ell\bar{\ell}\ell\bar{\ell} \rightarrow \gamma\gamma$ ,  $H^0(0^+) \rightarrow W\bar{W} \rightarrow \ell\bar{\nu}\ell\nu$ , and  $H^0(0^+) \rightarrow Z\bar{Z} \rightarrow \ell\bar{\ell}\ell\bar{\ell}$  using Cartan's supersymmetry, that defines coupling of a vector particle  $x_i$  and Dirac spinors  $\psi$  and  $\phi$ .

Apparent discrepancy of  $H^0(0^+) \rightarrow \gamma\gamma/\ell\bar{\ell}\ell\bar{\ell}$  experimental data of ATLAS and CMS collaborations disappears, since the sum of transition to  $\ell\bar{\ell}\ell\bar{\ell}$  and  $\gamma\gamma$  have the meaning. The trilinearity of leptons vector fields coupling and the initial wave functions to be large components define Higgs-leptons couplings.

The ratio of Higgs boson decay branching ratios  $H \rightarrow WW = 21.6 \pm 0.9$ ,  $H \rightarrow \gamma\gamma = 0.228 \pm 0.011$  can be reproduced when we multiply coupling constant  $\alpha_s \simeq 1$  for  $WW$  and  $\alpha_e = 1/137$  for  $\gamma\gamma$  to the number of independent diagrams.

Comparison with Atiyah-Witten's quark-gluon dynamics on a manifold of  $G_2$  holonomy theory is added.

**KEYWORDS:** Supersymmetric Effective Theories, Higgs bosons,  $S$  mesons

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Analysis of Higgs boson decay into <math>\gamma\gamma</math> using Cartan's supersymmetry</b>	<b>2</b>
<b>3</b>	<b>Analysis of Higgs boson decay into vector bosons using Cartan's supersymmetry</b>	<b>4</b>
<b>4</b>	<b>Nature of the <math>S(750 \text{ GeV})</math> and an enhancement of <math>\gamma\gamma \rightarrow S</math> at 13GeV</b>	<b>5</b>
<b>5</b>	<b>Discussion and conclusion</b>	<b>5</b>

---

## 1 Introduction

The decay properties of Higgs boson  $H^0(0^+)$  of mass 125GeV is not well understood. It decays as  $H \rightarrow WW = 21.6 \pm 0.9\%$ ,  $H \rightarrow \gamma\gamma = 0.228 \pm 0.011\%$  and  $H \rightarrow ZZ = 2.67 \pm 0.11\%$ [1].

The Higgs boson decay to  $\gamma\gamma$  is not simple, since  $Z\bar{Z}$  decays into  $\ell\bar{\ell}\ell\bar{\ell}$  and  $\ell\bar{\ell}$  can make a  $\gamma$ . The measurements of the decay branching ratios of  $H^0 \rightarrow \gamma\gamma$  and  $H^0 \rightarrow \ell\bar{\ell}\ell\bar{\ell}$  by the ATLAS Collaboration and by the CMS collaboration [2–9] are not in agreement, but the sum of the branching ratios of  $\gamma\gamma$  and  $\ell\bar{\ell}\ell\bar{\ell}$  are in agreement.

Experimentally, the ratio of the branching ratio of  $\gamma\gamma$  and  $W\bar{W}$  or  $Z\bar{Z}$  are also measured. The boson  $W$  decays in the average 10.86% to  $\ell\bar{\nu}$ , and  $Z$  decays in the average 3.6% to  $\ell\bar{\ell}$  ( $\ell = e, \mu$  or  $\tau$ ). The signal strength of  $H^0 \rightarrow W\bar{W}$  of the ATLAS Collaboration and the CMS collaboration differ by about 20%[8]. It is therefore interesting to calculate the decay branching ratios of the Higgs boson to  $\gamma$  and vector bosons in a model beyond the standard model.

In the book of the theory of spinors[12], Cartan proposed two types of Dirac spinors  $\phi$  and  $\psi$  and their charge conjugates  $\mathcal{C}\phi$  and  $\mathcal{C}\psi$  and two types of vector particles  $\mathbf{E}$  and  $\mathbf{E}'$ . In [13] we studied decays of  $H^0(0^+)$  into photons, and in [14] we studied the decay of  $\chi_b(nP)$  into  $\Upsilon(mS)\gamma$  and possible decay of Higgs partner  $h^0(0^+)$  into  $\Upsilon(mS)\gamma$  ( $m = 1, 2$ ). Experimental results of  $h^0(0^+)$  are not clear, and in this paper, we study decays of  $H^0(0^+)$  into  $\ell\bar{\ell}\ell\bar{\ell}$  and compare with the decay of  $H^0(0^+) \rightarrow \ell\bar{\ell}\ell\bar{\ell} \rightarrow \gamma\gamma$ ,  $H^0(0^+) \rightarrow Z\bar{Z}$  and  $H^0(0^+) \rightarrow W\bar{W}$ .

The experimental decay branching ratio of  $H^0(0^+) \rightarrow Z\bar{Z}$  is obtained from  $H^0(0^+) \rightarrow \ell\bar{\ell}\ell\bar{\ell}$ , and the strength parameter in this channel seems to be very sensitive to the mass of  $H^0(0^+)$ , and whether the final  $\ell\bar{\ell}\ell\bar{\ell}$  originates from  $Z\bar{Z}$  is not clear. By inclusion of  $g\bar{g} \rightarrow \ell\bar{\ell}\ell\bar{\ell}$ , the signal strength of  $H^0(0^+) \rightarrow Z\bar{Z}$  was reduced to 0.99[15], and by inclusion of  $g\bar{g} \rightarrow Z\bar{Z}$ , the signal strength was reduced to 0.93[16].

In the supersymmetric theory, the coupling of the Higgs boson to two leptons is given by the right chiral leptons, defined as  $\mathcal{E}_i$  ( $i = 1, 2, 3$ )

$$\begin{aligned}\mathcal{E}_1 &= \tilde{\phi}_{\bar{e}} + \theta \cdot \chi_{\bar{e}} + \frac{1}{2}\theta \cdot \theta F_{\bar{e}} \\ \mathcal{E}_2 &= \tilde{\phi}_{\bar{\mu}} + \theta \cdot \chi_{\bar{\mu}} + \frac{1}{2}\theta \cdot \theta F_{\bar{\mu}} \\ \mathcal{E}_3 &= \tilde{\phi}_{\bar{\tau}} + \theta \cdot \chi_{\bar{\tau}} + \frac{1}{2}\theta \cdot \theta F_{\bar{\tau}}\end{aligned}$$

and left chiral leptons, defined as

$$\mathcal{L}_1 = \begin{pmatrix} \tilde{\phi}_{\nu_e} \\ \tilde{\phi}_e \end{pmatrix} + \theta \cdot \begin{pmatrix} \chi_{\nu_e} \\ \chi_e \end{pmatrix} + \frac{1}{2}\theta \cdot \theta \begin{pmatrix} F_{\nu_e} \\ F_e \end{pmatrix}$$

and  $\mathcal{L}_2$  and  $\mathcal{L}_3$  defined similarly [17].

The Higgs coupling to  $Z$  through leptons is given by using  $\chi_{\nu_{L_i}}$  part of  $\mathcal{L}_j$ ,

$$-y_\ell^{ij} \mathcal{E}_i(\mathcal{L}_j \circ \mathcal{H}_d) = -y_\ell^{ij} \nu_\ell (\bar{\ell}_{L_i} \ell_{L_j} + \bar{\ell}_{L_j} \ell_{L_i})$$

where  $L_i, L_j$  specify left handed  $SU(2)$  lepton degrees of freedom.

The Higgs coupling to  $W$  through leptons is given by

$$-y_\ell^{ij} \mathcal{E}_i(\mathcal{L}_j \circ \mathcal{H}_d) = -y_\ell^{ij} \nu_\ell (\bar{\ell}_{L_i} \ell_{\nu_j})$$

and that to  $\bar{W}$  is

$$-y_\ell^{ji} \mathcal{E}_j(\mathcal{L}_i \circ \mathcal{H}_d) = -y_\ell^{ji} \nu_\ell (\ell_{L_j} \bar{\ell}_{\nu_i}).$$

Couplings to quarks of  $u, c, t$  of Higgs bosons are defined by

$$y_u^{ij} \mathcal{U}_i(\mathcal{Q}_j \circ \mathcal{H}_u) = y_u^{ij} \nu_u \bar{u}_i u_j.$$

where  $\mathcal{Q}$  defines the colour and  $\mathcal{U}$  defines the flavour

In sect.2, we formulate Higgs boson decay into  $\gamma\gamma$  using Cartan's supersymmetric theory of spinors, and in sect.3 Higgs boson decay into vector bosons using Cartan's supersymmetry. Nature of the  $S(750 \text{ GeV})$  and a possible enhancement of  $\gamma\gamma \rightarrow S$  at 13GeV from Cartan's supersymmetry is discussed in sect.4. Discussion and conclusion are given in sect.5.

## 2 Analysis of Higgs boson decay into $\gamma\gamma$ using Cartan's supersymmetry

Following Cartan's supersymmetry, we express scalar particles by  $\Psi$  and  $\Phi$ , and leptons by  $\psi, \mathcal{C}\phi$  and antileptons by  $\mathcal{C}\psi, \phi$ [12]. The coupling of a vector field  $x_i$  ( $i = 1, 2, 3, 4$ ) to Dirac spinors,

$$\begin{aligned}\psi &= \xi_1 \mathbf{i} + \xi_2 \mathbf{j} + \xi_3 \mathbf{k} + \xi_4 \mathbf{I} = \begin{pmatrix} \xi_4 + i\xi_3 & i\xi_1 - \xi_2 \\ i\xi_1 + \xi_2 & \xi_4 - i\xi_3 \end{pmatrix} \\ \mathcal{C}\psi &= -\xi_{234} \mathbf{i} - \xi_{314} \mathbf{j} - \xi_{124} \mathbf{k} + \xi_{123} \mathbf{I} = \begin{pmatrix} \xi_{123} - i\xi_{124} & -i\xi_{234} + \xi_{314} \\ -i\xi_{234} - \xi_{314} & \xi_{123} + i\xi_{124} \end{pmatrix}\end{aligned}\quad (2.1)$$

and the spinor operator

$$\begin{aligned}\phi &= \xi_{14}\mathbf{i} + \xi_{24}\mathbf{j} + \xi_{34}\mathbf{k} + \xi_0\mathbf{I} = \begin{pmatrix} \xi_0 + i\xi_{34} & i\xi_{14} - \xi_{24} \\ i\xi_{14} + \xi_{24} & \xi_0 - i\xi_{34} \end{pmatrix} \\ C\phi &= -\xi_{23}\mathbf{i} - \xi_{31}\mathbf{j} - \xi_{12}\mathbf{k} + \xi_{1234}\mathbf{I} = \begin{pmatrix} \xi_{1234} - i\xi_{12} & -i\xi_{23} + \xi_{31} \\ -i\xi_{23} - \xi_{31} & \xi_{1234} + i\xi_{12} \end{pmatrix}\end{aligned}\quad (2.2)$$

are specified by trilinearity [12, 18, 19]

$$\begin{aligned}\mathcal{F}_H &= {}^t\phi CX\psi = {}^t(C\phi)\gamma_0 x_i \gamma_i \psi + {}^t\phi \gamma_0 x_{i'} \gamma_i (C\psi) \\ &= x_1(\xi_{12}\xi_{314} - \xi_{31}\xi_{124} - \xi_{14}\xi_{123} + \xi_{1234}\xi_1) \\ &\quad + x_2(\xi_{23}\xi_{124} - \xi_{12}\xi_{234} - \xi_{24}\xi_{123} + \xi_{1234}\xi_2) \\ &\quad + x_3(\xi_{31}\xi_{234} - \xi_{23}\xi_{314} - \xi_{34}\xi_{123} + \xi_{1234}\xi_3) \\ &\quad + x_4(-\xi_{14}\xi_{234} - \xi_{24}\xi_{314} - \xi_{34}\xi_{124} + \xi_{1234}\xi_4) \\ &\quad + x_{1'}(-\xi_0\xi_{234} + \xi_{23}\xi_4 - \xi_{24}\xi_3 + \xi_{34}\xi_2) \\ &\quad + x_{2'}(-\xi_0\xi_{314} + \xi_{31}\xi_4 - \xi_{34}\xi_1 + \xi_{14}\xi_3) \\ &\quad + x_{3'}(-\xi_0\xi_{124} + \xi_{12}\xi_4 - \xi_{14}\xi_2 + \xi_{24}\xi_1) \\ &\quad + x_{4'}(\xi_0\xi_{123} - \xi_{23}\xi_1 - \xi_{31}\xi_2 - \xi_{12}\xi_3)\end{aligned}\quad (2.3)$$

In the case of coupling of  $\gamma\gamma$  to a Higgs boson, I first fix the types of polarization of  $\ell\bar{\ell}$  that creates  $\gamma$  which are expressed by large components of Dirac wave functions, and couple quarks of different  $q\bar{q}$  pairs by  $\xi_{1234}$  and antiquarks to Higgs boson as shown in the left hand side of Fig.1. The right hand side of Fig.1 are diagrams in which roles of quarks and antiquarks are interchanged and  $\xi_{1234}$  is interchanged to  $\xi_{123}$ .

In these 12 diagrams, lepton pairs which move are components of  $(\psi, \mathcal{C}\psi)$  and  $(\phi, \mathcal{C}\phi)$ , which can be regarded as relatively anti-particles and produce a  $\gamma$ .

Couplings of quarks to electromagnetic field is defined as

$$\mathcal{F}_L = \bar{\psi}_L^k(\gamma_L(i\partial_\mu - eA_\mu) - m)\psi_L^k + \bar{\phi}_L^k(\gamma_L(i\partial_\mu - eA_\mu) - m)\phi_L^k$$

where

$$\psi_L^k = \begin{pmatrix} \psi^k \\ \mathcal{C}\psi^k \end{pmatrix} \text{ and } \phi_L^k = \begin{pmatrix} \mathcal{C}\phi^k \\ \phi^k \end{pmatrix}.$$

In electromagnetic interactions  $\psi$  v.s.  $\mathcal{C}\phi$  and  $\phi$  v.s.  $\mathcal{C}\psi$  have no differences, but we expect that our electronic detector is sensitive only to  $\psi_L^k$ .

When the momenta of the 4 leptons are parallel,  $H^0$  decay to  $\ell\bar{\ell}\ell\bar{\ell}$  and to  $\gamma\gamma$  through exchange of vector particle  $X$  is possible. However kinematically the configuration is suppressed, and we expect  $H^0$  decay to  $\gamma\gamma$  mainly occurs via  $\ell\bar{\ell}\ell\bar{\ell}$  channels.

### 3 Analysis of Higgs boson decay into vector bosons using Cartan's supersymmetry

In the standard model, the vector boson  $Z$  and the vector field  $A$  are expressed as

$$\begin{aligned} Z_\mu &= \cos \theta_W \cdot W_\mu^3 + \sin \theta_W \cdot B_\mu \\ A_\mu &= -\sin \theta_W \cdot W_\mu^3 + \cos \theta_W \cdot B_\mu \end{aligned}$$

and  $\sin^2 \theta_W \simeq 0.22$  is the Weinberg angle. Relative strength of  $B_\mu$  near the energy of the mass of Higgs boson is not so clear.

The Lagrangian of  $W^\pm$  and  $Z$  is given by

$$\mathcal{L} = m_W^2 W^{+\mu} W_\mu^- + \frac{1}{2} m_Z^2 Z_\mu Z^\mu$$

and we define  $W_\mu^\pm = \frac{1}{\sqrt{2}}(W_\mu^1 \pm i W_\mu^2)$  and

$$\frac{1}{2} m_Z^2 Z_\mu Z^\mu = \frac{\nu^2}{4} (g W^3 - g' B)^2$$

In contrast to the field of  $\gamma$ , I make a Cayley-Dickson doubling process and assign complex fields at the position of the Higgs potential. Couplings of quark and antiquarks to  $W$  which are expressed as  $X_1 \pm i X_2$  are assumed to satisfy the same trilinearity.

Couplings of  $W^\pm$  fields to the Higgs boson are described by  $(X_1 + i X_2)(X_1 - i X_2) = X_1^2 + X_2^2$ . Since  $X_1 + i X_2$  and  $X_1 - i X_2$  are independent fields there are 8 independent diagrams.

In these four diagrams  $X_3$  in the final state can be interpreted as  $Z(\ell\bar{\ell})$ .

In the world of hadron dynamics, photons couple to Higgs boson via  $Z$  bosons, and in addition to the coupling to leptons, there are couplings to quarks  $d, s, b$  via  $y_d^{ij}$  defined by [17]

$$-y_d^{ij} \mathcal{D}_i(\mathcal{Q}_j \circ \mathcal{H}_d) = -y_d^{ij} \nu_d \bar{d}_i d_j.$$

Here,  $\nu_d$  is defined by the minimum of the potential  $H_d = {}^t(\nu_d, 0)$ , and after diagonalization masses of three generations  $m_{d,s,b} = \nu_d y_{d,s,b}$  appear.

In the weak interactions there are differences in  $\phi_L^k$  and  $\psi_L^k$ , as the final states of the supersymmetric transformation [13].

In the expression of  $-y_d^{ij} \nu_\ell \bar{\ell}_{L_i} \ell_{L_j}$  coupling,

$$\ell_{L_j} = \begin{pmatrix} \psi^j \\ \mathcal{C}\psi^j \end{pmatrix} \text{ and } \bar{\ell}_{L_i} = (\mathcal{C}\phi^i, \phi^i).$$

In the coupling of  $y_\ell^{ij} \mathcal{E}_i(\mathcal{L}_j \circ \mathcal{H}_d)$ ,  $i, j$  specify generations of right handed  $\bar{\ell}$  and left-handed  $\ell_i$  or  $\nu_j$ , respectively. The Higgs boson couples to  $W$  by  $H^0 \ell \bar{\nu} \ell \nu$ .

The ratio of Higgs boson decay branching ratios  $H \rightarrow WW = 21.6 \pm 0.9\%$ ,  $H \rightarrow \gamma\gamma = 0.228 \pm 0.011\%$  [1] can be reproduced when we multiply coupling constant  $\alpha_s \simeq 1$  for  $WW$  and  $\alpha_e = 1/137$  for  $\gamma\gamma$  to the number of independent diagrams. Branching ratio

$H \rightarrow ZZ = 2.67 \pm 0.11\%$  is difficult to interpret, since  $Z$  are detected mainly as  $\ell\bar{\ell}s$ , which decay to  $\gamma s$ .

The main difference of Higgs decays to  $W$ ,  $Z$  and  $\gamma$  is that  $\gamma$  belongs to  $U(1)_{em}$  and vector field  $x$  and  $x'$  are treated as identical as shown in Fig.1, while  $W$  and  $Z$  belong to  $SU(2)_w$  and higgs field  $h_0$  and  $h'_0$  are treated as different as shown in Fig.2 and 3.

In the energy region of Higgs boson mass, the decay to  $\gamma\gamma$  can occur also through  $\ell\bar{\ell}\ell\bar{\ell}$  in the final state of  $Z\bar{Z}$ . The boson  $Z$  has a partner of a vector field  $A$  in this energy region. Labelle[17] assigned the field  $A$  as a photon, but we assign the field here as a gluon, and expect the difference of the  $H^0$  signal strength of  $Z\bar{Z}$  final state  $1.11 \pm 0.3$  and that of  $W\bar{W}$  final state  $0.87 \pm 0.2$  is due to the contamination of  $A = g\bar{g} \rightarrow \ell\bar{\ell}\ell\bar{\ell}$  in the final state of  $Z\bar{Z}$ .

#### 4 Nature of the $S(750 \text{ GeV})$ and an enhancement of $\gamma\gamma \rightarrow S$ at 13GeV

Since leptons and quarks in scalar particles  $\Psi$  and  $\Phi$  have charge conjugated states, we can extend the same argument starting from the charge conjugated states  $\mathcal{C}\Psi$  and  $\mathcal{C}\Phi$ .

The coherent decays of Higgs pairs on two planes defined by a unique time

$$(\Psi\Phi)^r, (\mathcal{C}\Psi\mathcal{C}\Phi)^r, (\Psi\Phi)^g, (\mathcal{C}\Psi\mathcal{C}\Phi)^g, \quad \text{and} \quad (\Psi\Phi)^b, (\mathcal{C}\Psi\mathcal{C}\Phi)^b$$

into  $\gamma\gamma$  can make a peak at  $125\text{GeV} \times 6 = 750\text{GeV}$  scalar meson  $S$ , which was observed in LHC experiments of run energy  $\sqrt{s} = 13\text{TeV}$  [21, 22].

Review of electroweak symmetry breaking in Higgs boson decay was given in [23]. In the standard theory, a large component of a lepton or a quark remains large component after coupling to photons. But Cartan's couplings of  $\gamma$  to spinors contain  $\gamma$  couplings to  $\ell\bar{\ell}$  or  $q\bar{q}$  vertices.

There are discussion on parent states of the  $S$  meson and decays of the  $S$  meson into dark matter [24, 25]. We think dark matters are weakly interacting particles but electromagnetic waves emitted from them are transformed by  $G_{12}, G_{13}, G_{123}$  and  $G_{132}$  and insensitive to our detectors.

There are arguments on the decay width of 750 GeV resonance  $S$  to  $\gamma\gamma$  and the  $W_R$  vector of 1.9TeV decay to  $WZ$  based on trinification model[26, 27]

$$\frac{\Gamma(W_R^\pm \rightarrow W^\pm Z)}{\Gamma(S \rightarrow \gamma\gamma)} = \frac{(1 \pm 13) \text{ fb}}{(8 \pm 2) \text{ fb}}.$$

The mass of parent state  $W_R^\pm \rightarrow W^\pm Z$  is about 3 times the mass of  $S$ , and which is about 1/6 of the energy of the parent state of  $S \rightarrow \gamma\gamma = 12\text{TeV}$ . The factor 1/2 can be understood as due to a choice of the charge + or - and 1/3 can be understood as due to a choice of the colour.

#### 5 Discussion and conclusion

We studied the decay of Higgs boson  $H^0(0^+)$  to  $\gamma\gamma$  via  $\ell\bar{\ell}\ell\bar{\ell}$  states. Cartan's supersymmetry allows presence of a scalar boson  $\Psi$  and a scalar boson  $\Phi$  which decay into  $\ell\bar{\ell}\ell\bar{\ell}$ , and reduce

to  $\gamma\gamma$ . It is important that the direct decay of  $\Psi$  or  $\Phi$  to  $\gamma\gamma$  is not allowed. Experimental discrepancy of the branching ratios of  $\gamma\gamma$  and  $\ell\bar{\ell}\ell\bar{\ell}$  of the ATLAS collaboration and the CMS collaboration [8] is not a problem since branching ratio of  $\gamma\gamma$  comes from  $\ell\bar{\ell}\ell\bar{\ell}$  in our model, and the sum of the branching ratios of  $\gamma\gamma$  and  $\ell\bar{\ell}\ell\bar{\ell}$  has the physical meaning.

The Higgs meson of 125 GeV expects  $S$  meson of mass 750 GeV whose hint was observed in  $\sqrt{s} = 13\text{TeV}$ , since coherent sum of 3 colour states times 2 charge conjugates makes  $6 \times 125 = 750\text{GeV}$ . If there is a parent state of  $S$  mesons of energy  $2^4 \times 750\text{GeV} = 12\text{TeV}$ , where the factor  $2^4$  comes from the selection of  $S\bar{S}$  or  $\bar{S}S$  in 4 dimensional space-time of  $S$  or  $\bar{S}$ , we can understand disappearance of  $S$  meson in  $\sqrt{s} = 7$  and 8 TeV experiment.

When the supersymmetric transformation is  $G_{13}$ , transformations  $\mathbf{E} \rightarrow \mathcal{C}\psi$  and  $\mathbf{E}' \rightarrow \psi$  in the lepton sector, and the spinor  $\phi_L^k \rightarrow \tilde{\phi}_L^k$  appears in the quark or lepton sector, where  $\tilde{\phi}_L^k$  means that particle-antiparticle transformation is done on  $\phi_L^k$  [20]. When the transformation is  $G_{12}$ , transformations  $\mathbf{E} \rightarrow \tilde{\phi}$  in the lepton sector and  $\psi_L^k \rightarrow \tilde{\psi}_L^k$  appears in the quark or lepton sector.

When the transformation is  $G_{132}$ , transformations  $\mathbf{E} \rightarrow \phi$  and  $\mathbf{E}' \rightarrow \mathcal{C}\phi$  occurs in the quark or lepton sector and the spinor  $\phi_L^k \rightarrow \tilde{\psi}_L^k$  appears in the quark or lepton sector. When the transformation is  $G_{123}$ , transformations of  $\mathbf{E} \rightarrow \tilde{\mathcal{C}}\psi$  and  $\mathbf{E}' \rightarrow \tilde{\psi}$  occurs in the quark or lepton sector and  $\psi_L^k \rightarrow \tilde{\phi}_L^k$  in the quark or lepton sector.

Possible appearance of  $\gamma\gamma$  resonance of 750GeV was proposed in trification models [26, 27], in which Higgs bosons  $H_1$  and  $H_2$  in some representations of  $SU(3)_L \otimes SU(3)_R \otimes SU(3)_c$  gauge symmetry were introduced. Higgs bosons derived from Cartan's supersymmetry contain trilinearity, and we can work in  $SU(2)_L \otimes U(1)_Y \otimes SU(3)_c$  gauge framework.

Corresponding to the triality of Cartan, one can consider three  $G_2$  manifolds  $X_i$  and a flat  $C$ -field of the symmetry class  $H^3(Y; U(1))$  and  $H^3(X_i; U(1))$  [28].

Observation of supersymmetry as a relation between fluctuations in the  $C$ -field and fluctuations in metric [28] agree with each other is interesting. The fact that the trilinear quark antiquark vector field couplings of Cartan yields Higgs  $\gamma\gamma$  decay branching ratio and Higgs  $WW$  decay branching ratio consistent with experiments indicates that five dimensional spinor  $\xi$  like Kaluza-Klein spinor can play important roles for understanding quark gluon dynamics.. Cartan's low energy supersymmetry is based on our number system containing octonions and rich in applicability at multiscale levels.


**Acknowledgement** I thank Stan Brodsky (SLAC) for calling my attention to new experimental results on  $H^0$  and  $S$  meson decays, Naoki Kondo (Teikyo University) for discussions on Clifford Algebra and Serge Dos Santos (INSA Centre Val de Loire) for helpful comments.

## References


- [1] The ATLAS and CMS Collaborations, ATLAS-CONF-2015-044 (2015); CMS PAS HIG-15-002 (2015), Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC  $pp$  collision data at  $\sqrt{s} = 7$  and 8 TeV, JHEP 08 (2016) 045.
- [2] ATLAS Collaboration, Measurement of Higgs boson and couplings in diboson final states with the ATLAS detector at the LHC, Physics Letters **B 726** (2013) 88-119. [*Erratum ibid* **B 734** (2014) 406 ].
- [3] Aad, G. et al (ATLAS collaboration), Measurement of the Higgs boson mass from the  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4\ell$  channels in  $pp$  collision at center-of-mass energies of 7 and 8 TeV with the ATLAS detector, Phys. Rev. **D90**,052004 (2014).
- [4] The CMS Collaboration, Measurement of Higgs boson production and properties in the WW decay channel with leptonic final states, JHEP 01(2014) 096, arXiv 1312.1129 v2.
- [5] The CMS Collaboration, Observation of the diphoton decay of the Higgs boson and measurement of its properties, Eur. Phys. J.C. (2014) 74: 3076.
- [6] The CMS Collaboration, Evidence for the direct decay of the 125 GeV Higgs boson to fermions, nature Physics, DOI:10.1038/NPHYS3005 (2014).
- [7] The CMS Collaboration, Performance of photon reconstruction and identification with the CMS detector in proton-proton collision at  $\sqrt{s}=8\text{TeV}$ , arXiv:1502.02702 v2.
- [8] The ATLAS and CMS Collaborations, ATLAS note, ATLAS-CONF-2015-081 (2015); CMS note, CMS PAS EXO-15-004 (2015), Search for resonances decaying to photon pairs in  $3.2\text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector, (2015).
- [9] The ATLAS Collaboration, Study of  $(W/Z)H$  production and Higgs boson couplings using  $H \rightarrow WW^*$  decays with the ATLAS detector, CERN-PH-EP-2015-104, submitted to JHEP (2015). S-cion: Where else should we look for it?, Phys. Lett. B 757 (2016) 39.
- [10] The CMS Collaboration, Search for resonant production of high mass photon pairs using  $12.9\text{ fb}^{-1}$  of proton-proton collisions at  $\sqrt{s} = 13$  TeV and combined interpretation of searches at 8 and 13 TeV. The CMS Physics Analysis Summary.
- [11] The ATLAS Collaboration, Search for scalar diphoton resonances with  $15.4\text{ fb}^{-1}$  of data collected at  $\sqrt{s} = 13$  TeV in 2015 and 2016 with the ATLAS detector (2016).
- [12] É. Cartan, *The theory of Spinors*, Dover Pub. (1966)
- [13] S. Furui, Triality selection rules of Octonion and Quantum Mechanics, [hep-ph/1409.3761].
- [14] S. Furui, Cartan's Supersymmetry and the Decay of a  $h^0$  with the mass  $m_{h^0} \simeq 11\text{GeV}$  to  $\Upsilon(1S)\gamma$  and to  $\Upsilon(2S)\gamma$ , [hep-ph/1502.07011].
- [15] J.S. Gainer, J. Lykken, K.T. Matchev, S. Mrenna and M. Park, Beyond Geolocating: Constraining Higher Dimensional Operators in  $H \rightarrow 4\ell$  with Off-Shell Production and More, Phys. Rev. **D 91** (2014) 035011
- [16] L. Finco, Recent Results of the Higgs Boson Properties in the  $H \rightarrow ZZ \rightarrow 4\ell$  decay channel at CMS, in *Proceedings of the Second Annual Conference on Large Hadron Collider Physics*, Columbia University, New York, U.S.A. June 2014, [hep-ex/1409.2646v2].
- [17] P. Labelle, *Supersymmetry Demystified*, McGraw Hill (2010).




- [18] S. Furui, Fermion Flavors in Quaternion Basis and Infrared QCD, *Few Body Syst.* **52**, (2012) 171-187, [hep-ph/1104.1225].
- [19] S. Furui, The flavor symmetry in the standard model and the triality symmetry, *Int. J. Mod. Phys. A* **27** (2012) 1250158, [hep-ph/1203.5213].
- [20] S. Furui, Cartan's Supersymmetry and Weak and Electromagnetic Interactions, *Few Body Syst.* **56**, (2015) 703.
- [21] ATLAS note, Search for resonances decaying to photon pairs in  $3.2\text{ fb}^{-1}$  of  $pp$  collision at  $\sqrt{s} = 13\text{ TeV}$  with ATLAS detector, ATLAS-CONF-2015-081 (2015).
- [22] CMS note, Search for new physics in high mass diphoton events in proton-proton collision at 13 TeV, CMS PAS EXO-15-004 (2015).
- [23] A. Djouadi, The anatomy of electro-weak symmetry breaking. @I: The Higgs boson in the standard model, *Phys. Rept.* **457**,1 (2008); arXiv:hep-ph/0503172.
- [24] R. Franceschini et al., What is the  $\gamma\gamma$  resonance at 750 GeV?, arXiv:1512.04933.
- [25] A. Salvio, F. Staub, A. Strumia and A. Urbano, On the maximal diphoton width, CERN-TH-2016-028, arXiv:1602.01460.
- [26] J. Hetzel and B. Stech, Low-energy phenomenology of trinification: An effective left-right-symmetric model, *Phys. Rev. D* **91**, 055026 (2015).
- [27] G.M. Pelaggi, A. Strumia and E. Vigiani, Trinification can explain the di-photon and di-boson LHC anomalies, CERN-PH-TH-2015-311, arXiv:1512.07225 v1[hep-ph].
- [28] M. Atiyah and F. Witten, M theory Dynamics on a Manifold of  $G_2$  Holonomy, hep-th/0107177 v3.



**Figure 1.** Higgs boson decay into  $\gamma(\ell\bar{\ell})\gamma(\ell\bar{\ell})$ .



**Figure 2.** Higgs boson decay into  $W(\ell\bar{\nu})W(\bar{\ell}\nu)$ .



**Figure 3.** Higgs boson decay into  $W_3(\ell\bar{\nu})B(\bar{\ell}\ell)$ .