

Hyperon Semileptonic decay constants with flavor SU(3) symmetry breaking

Ghil-Seok Yang,^a and Hyun-Chul Kim^{b,c}

^a*Department of Physics, Soongsil University,
Seoul 156-743, Republic of Korea*

^b*Department of Physics, Inha University,
Incheon 402-751, Republic of Korea*

^c*School of Physics, Korea Institute for Advanced Study (KIAS),
Seoul 130-722, Republic of Korea*

E-mail: ghsyang@ssu.ac.kr, hchkim@inha.ac.kr

ABSTRACT: We investigate the hyperon semileptonic decay constants, f_2/f_1 , and g_1/f_1 , within a general framework of a chiral soliton model. All relevant parameters for the SU(3) baryon wave functions were unambiguously determined by using the experimental data for the masses of the baryon octet and the decuplet. Using then the existing experimental data for the magnetic moments of the baryon octet and the decay constants of hyperon semileptonic decays, we are able to determine all the hyperon semileptonic decay constants f_2/f_1 and g_1/f_1 of the baryon octet unequivocally. In addition, we also present the results of the axial-vector transition constants from the baryon decuplet to the octet.

KEYWORDS: hyperon semileptonic decay constants, vector constants, axial-vector constants, chiral soliton model

Contents

1	Introduction	1
2	Baryon matrix elements of the vector and axial-vector currents	3
3	Analysis of the HSD constants for the baryon octet	6
4	Analysis for the transition from the baryon decuplet to the octet	13
5	Summary and outlook	16

1 Introduction

The Cabibbo-Kobayashi-Maskawa matrix elements $|V_{ud}|$ and $|V_{us}|$ characterize the quark mixings in the process $d \rightarrow ue^- \bar{\nu}_e$ and $s \rightarrow ue^- \bar{\nu}_e$ [1, 2] in the standard model. So far the most precise determination of $|V_{ud}|$ and $|V_{us}|$ are obtained respectively from super-allowed Fermi transition together with pion decays, and from leptonic and semileptonic kaon decays [3–5]. On the other hand, hyperon semileptonic decays (HSDs) can also provide independent constraints on $|V_{ud}|$ and $|V_{us}|$ [6, 7]. For last years, new experimental results for HSD have been reported: the KTeV collaboration first announced the measurement of the $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ decay [8] and determined the corresponding form factors [9] as $f_2(0)/f_1(0) = 2.0 \pm 1.2_{\text{stat}} \pm 0.5_{\text{syst}}$ and $g_1(0)/f_1(0) = 1.32^{+0.21}_{-0.17\text{stat}} \pm 0.05_{\text{syst}}$. The NA48/1 collaboration brought about the branching ratios for the same process with higher statistics in comparison with the KTeV experiments: $g_1(0)/f_1(0) = 1.21 \pm 0.05$ for $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ [10]. Now there are experimental data for the $g_1(0)/f_1(0)$ ratios in five different decay channels in HSDs, collected by the Particle Data Group (PDG) [11].

It is also of great importance to understand the HSD constants, since they provide essential information on properties of the nucleon and low-lying hyperons. The data for HSDs reveal experimentally the pattern of flavor SU(3) symmetry breaking. In exact flavor SU(3) symmetry the ratios of the axial-vector and vector constants g_1/f_1 are expressed only by the two constants F and D . Similarly, the ratios of the vector constants f_2/f_1 are written in terms of the anomalous magnetic moments of the proton and the neutron with flavor SU(3) symmetry assumed. However, the experimental data for HSDs show that flavor SU(3) symmetry is manifestly broken. For example, the ratio $g_1/f_1 = 1.21 \pm 0.05$ for the HSD process $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ measured by the KTeV collaboration [8, 9] is equal to that of the neutron β decay ($g_1/f_1 = 1.2695 \pm 0.0029$) in flavor SU(3) symmetry. Thus, complete information on HSDs will furnish the pattern of flavor SU(3) symmetry breaking in nature. It is also interesting to see that the experimental data for the HSD constants give a clue how isospin symmetry is broken. The above-mentioned ratio of the HSD constants $g_1/f_1 =$

1.20 ± 0.05 from the KTeV collaboration should be equal to that of its isospin partner process $\Xi^- \rightarrow \Sigma^0 e^- \bar{\nu}_e$ ($g_1(0)/f_1(0) = 1.25_{-0.16}^{+0.14}$) [12] even with flavor SU(3) symmetry breaking. If it is possible to measure more precisely the HSD constants for the decay $\Xi^- \rightarrow \Sigma^0 e^- \bar{\nu}_e$, HSDs will shed light on how isospin symmetry is broken.

For this reason, there has been a great deal of theoretical works. Chiral perturbation theory (χ PT) is used as a framework to analyze flavor SU(3) breaking effects on $f_1(0)$ beyond the first order [13–16], in which other form factors of HSD have been also studied. The results for V_{us} within HSDs were discussed in Ref. [17] and for the large N_c expansion in Ref. [18]. In lattice QCD [16] all form factors of the $\Sigma^- \rightarrow n l \nu$ decay were first time studied and the results were given as: $f_2(0)/f_1(0) = -1.52 \pm 0.81$ and $g_1(0)/f_1(0) = -0.287 \pm 0.052$. Flavor SU(3) symmetry breaking in HSDs was also investigated within the $1/N_c$ expansion [19]. In the chiral quark-soliton model (χ QSM) with a *model-independent approach* considered, the HSD constants were investigated many years ago [20]. Being distinguished from the self-consistent χ QSM [21, 22], almost all dynamical parameters were fixed by using the experimental data of the masses and the HSD constants for the baryon octet in this *model-independent* approach. In Ref. [20], however, the baryon wave functions were not completely determined and experimental information was then not enough to fix unambiguously the parameters for the HSD constants.

In the present work we will examine the HSD constants of the baryon octet and decuplet within a general framework of a chiral soliton model (χ SM), fixing *all* dynamical parameters unequivocally. To derive the collective wave functions for the baryon octet, one has to take into account both flavor SU(3) symmetry and isospin symmetry breakings such that the experimental data for the masses of the baryon octet can be employed as a whole. In particular, sources of isospin symmetry breaking arise from the electromagnetic (EM) interaction as well as from the mass difference of the up and down quarks. In Ref. [23], the EM mass differences between baryons in the same isospin multiplet were analyzed within the χ SM and the corresponding model parameters were fixed by the experimental data. Together with the isospin symmetry breaking from the mass difference of the up and down quarks, Ref. [24, 25] showed that the collective wave functions of the baryon octet and decuplet were uniquely determined. In addition, a more complete analysis of the HSD constants can be carried out, the KTeV data for the $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ process considered. Since now the five experimental data for the ratios g_1/f_1 , we are able to fix all dynamical parameters for the HSD axial-vector constants with the singlet axial-vector charge from the data of deep inelastic scattering employed. Similarly, all dynamical parameters for the HSD vector constants can be fixed by using the experimental data for the magnetic moments of the baryon octet. Using the fixed parameters, we compute the HSD axial-vector and vector constants for the baryon octet. In addition, we also present the results of the transition axial-vector constants for the baryon decuplet, which will be of great use in determining the meson-baryon strong coupling constants.

The present work is sketched as follows. In Section II, we explain how to fix all dynamical parameters, using the experimental data. We also recapitulate relevant formulae for the HSD coupling constants. In Section III, we show the results of the present work and discuss physical implications of them. In the final Section, we summarize the present

work and draw conclusions. Additional detailed formulae are given in Appendices.

2 Baryon matrix elements of the vector and axial-vector currents

The transition matrix elements of the vector and axial-vector currents for the baryon octet are expressed respectively in terms of the six real form factors:

$$\begin{aligned}\langle B_2 | V_\mu^\chi | B_1 \rangle &= \bar{u}_{B_2}(p_2, s_2) \left[f_1^{B_1 \rightarrow B_2}(q^2) \gamma_\mu - \frac{i f_2^{B_1 \rightarrow B_2}(q^2) \sigma_{\mu\nu} q^\nu}{M_{B_1}} \right. \\ &\quad \left. + \frac{f_3^{B_1 \rightarrow B_2}(q^2) q_\mu}{M_{B_1}} \right] u_{B_1}(p_1, s_1), \\ \langle B_2 | A_\mu^\chi | B_1 \rangle &= \bar{u}_{B_2}(p_2, s_2) \left[g_1^{B_1 \rightarrow B_2}(q^2) \gamma_\mu + \frac{i g_2^{B_1 \rightarrow B_2}(q^2) \sigma_{\mu\nu} q^\nu}{M_{B_1}} \right. \\ &\quad \left. + \frac{g_3^{B_1 \rightarrow B_2}(q^2) q_\mu}{M_{B_1}} \right] \gamma_5 u_{B_1}(p_1, s_1),\end{aligned}\quad (2.1)$$

where the vector and axial-vector currents are defined as

$$V_\mu^\chi(x) = \bar{\psi}(x) \gamma_\mu \frac{1}{2} \lambda^\chi \psi(x), \quad A_\mu^\chi(x) = \bar{\psi}(x) \gamma_\mu \gamma_5 \frac{1}{2} \lambda^\chi \psi(x). \quad (2.2)$$

The u_{B_1} (\bar{u}_{B_2}) denote the Dirac spinors corresponding to the initial and final baryon states, respectively. The λ^χ designate flavor Gell-Mann matrices for strangeness conserving $\Delta S = 0$ transitions ($\chi = 3, 8, 1 \pm i2$) and for $|\Delta S| = 1$ ones ($\chi = 4 \pm i5$), respectively. The $q^2 = -Q^2$ stands for the square of the momentum transfer $q = p_2 - p_1$. The form factors $f_i^{B_1 \rightarrow B_2}$ and g_i are real quantities due to CP -invariance, depending only on the square of the momentum transfer. We can neglect $f_3^{B_1 \rightarrow B_2}$ and $g_3^{B_1 \rightarrow B_2}$ for the reason that their contributions to the decay rate is proportional to the ratio $m_l^2/M_{B_1}^2 \ll 1$, where m_l represents a mass of the lepton (e or μ) in the final state and that of the baryon in the initial state, M_{B_1} , respectively. The $g_2^{B_1 \rightarrow B_2}$ are finite only with the effects of flavor SU(3) symmetry and isospin symmetry breakings because of its opposite G parity to the axial-vector current, so that they are very small for the baryon octet. Moreover, the Ademollo-Gatto theorem [26] does not allow $f_1^{B_2 \rightarrow B_1}$ to acquire the linear-order corrections of flavor SU(3) symmetry breaking, so that $f_1^{B_2 \rightarrow B_1}$ are merely expressed in terms of the SU(3) Clebsch-Gordan coefficients.

Concerning the HSD constants for the transition from the baryon decuplet to the octet, it is difficult to determine the vector HSD constants because of a lack of experimental data for the $M1$ and $E2$ transitions from the baryon decuplet to the octet. For this reason, we will consider only the axial-vector transitions in the present work. There are several different ways of decomposing the matrix elements of the axial-vector current between the baryon decuplet and the octet. We will follow here the formalism of the Alder form factors $C_i^A(q^2)$ [27–30] to describe the axial-vector transitions of the baryon decuplet. The matrix elements between the baryon decuplet and the octet can be written as follows

$$\langle B_8 | A_\mu^\chi | B_{10} \rangle = \bar{u}_{B_8}(p_2, s_2) [C_5^A(q^2) g_{\mu\nu} + C_6^A(q^2) q_\mu q_\nu$$

$$+ \{C_3^A(q^2)\gamma^\alpha + C_4^A(q^2)p'^\alpha\} (q_\alpha g_{\mu\nu} - q_\nu g_{\alpha\mu})] u_{B_{10}}^\nu(p_1, s_1), \quad (2.3)$$

where the $u_{B_{10}}^\nu(p_1, s_1)$ represents the Rarita-Schwinger spinor that describes the baryon decuplet with spin 3/2. p' is defined as $p' = (M_{B_{10}}, 0, 0, 0)$.

Since we want to determine the HSD constants within the framework of a χ SM with all dynamical parameters in the model fixed by the experimental data, we first explain the χ SM in brief. The detailed formalism can be found for example in Ref. [24] and references therein. The χ SM is characterized by the following collective Hamiltonian:

$$H = M_{\text{cl}} + H_{\text{rot}} + H_{\text{sb}}, \quad (2.4)$$

where M_{cl} stands for the classical soliton mass. The collective Hamiltonian H_{rot} comes from the collective quantization of the chiral soliton by considering its slow rotation. We often call it the $1/N_c$ rotational correction and express it as

$$H_{\text{rot}} = \frac{1}{2I_1} \sum_{i=1}^3 \hat{J}_i^2 + \frac{1}{2I_2} \sum_{p=4}^7 \hat{J}_p^2, \quad (2.5)$$

where $I_{1,2}$ denote the moments of inertia of the soliton, which depend on dynamics of specific formulations of the χ SM. \hat{J}_i stand for the usual spin operators and \hat{J}_p represent the group generators corresponding to the right rotation in flavor SU(3) space. The last term in eq.(2.4) comes from the flavor SU(3) symmetry breaking, which are expressed as

$$\begin{aligned} H_{\text{sb}} = & (m_d - m_u) \left(\frac{\sqrt{3}}{2} \alpha D_{38}^{(8)}(\mathcal{R}) + \beta \hat{T}_3 + \frac{1}{2} \gamma \sum_{i=1}^3 D_{3i}^{(8)}(\mathcal{R}) \hat{J}_i \right) \\ & + (m_s - \bar{m}) \left(\alpha D_{88}^{(8)}(\mathcal{R}) + \beta \hat{Y} + \frac{1}{\sqrt{3}} \gamma \sum_{i=1}^3 D_{8i}^{(8)}(\mathcal{R}) \hat{J}_i \right) \\ & + (m_u + m_d + m_s) \sigma, \end{aligned} \quad (2.6)$$

where m_u, m_d , and m_s denote the up, down, and strange current quark masses, respectively. \bar{m} designates the average of the up and down current quark mass. $D_{ab}^{(R)}(\mathcal{R})$ are the SU(3) Wigner D functions in a R irreducible representation. \hat{T}_3 and \hat{Y} stand for the operators corresponding to the third component of the isospin and the hypercharge, respectively. The dynamical parameters α, β , and γ encode a specific dynamics of a certain chiral soliton model. They are defined as

$$\alpha = - \left(\frac{2}{3} \frac{\Sigma_{\pi N}}{m_u + m_d} - \frac{K_2}{I_2} \right), \quad \beta = - \frac{K_2}{I_2}, \quad \gamma = 2 \left(\frac{K_1}{I_1} - \frac{K_2}{I_2} \right), \quad (2.7)$$

where $\Sigma_{\pi N}$ indicates the πN sigma term. The σ in eq. (2.6) is defined as

$$\sigma = -(\alpha + \beta) = \frac{1}{3} \frac{\Sigma_{\pi N}}{\bar{m}}. \quad (2.8)$$

In the χ SM, the value of the eighth component of the soliton angular velocity J_8 provides a very important constraint on the collective quantization. In the Skyrme model, it has its origin in the Wess-Zumino term [31–33], whereas it is related to the presence of

the discrete valence quark level in the Dirac-sea spectrum in the SU(3) chiral quark-soliton model [34, 35]. Its presence has no effects on the chiral soliton but allows one to consider only the flavor SU(3) irreducible representations with zero triality.

The wave functions for the baryon octet and decuplet are derived by diagonalizing the collective SU(3) Hamiltonian of the χ SM [24, 25, 34, 35] and are expressed in terms of the SU(3) Wigner D functions:

$$\langle A | \mathcal{R}, B(Y T T_3, Y' J J_3) \rangle = \sqrt{\dim(\mathcal{R})} (-)^{J_3+Y'/2} D_{(Y, T, T_3)(-Y', J, -J_3)}^{(\mathcal{R})*}(A), \quad (2.9)$$

where \mathcal{R} designates the allowed irreducible representations of the flavor SU(3) group, i.e. $\mathcal{R} = 8, 10, \dots$. $Y T$, and T_3 denote the corresponding hypercharge, isospin and its third component, respectively. The right hypercharge is constrained to be $Y' = 2J_8/\sqrt{3} = -N_c/3 = -1$ that selects a tower of allowed flavor SU(3) representations. The baryon octet and decuplet, which are the lowest representations among them, coincide with those of the quark model.

In the presence of flavor SU(3) symmetry breaking, a baryon state is no more pure state but a mixed one with higher representations. Taking into account the strange current quark mass m_s as a perturbation, we obtain the collective wave functions for the baryon octet and decuplet mixed with higher representations as

$$\begin{aligned} |B_8\rangle &= |8_{1/2}, B\rangle + c_{\overline{10}}^B |\overline{10}_{1/2}, B\rangle + c_{27}^B |27_{1/2}, B\rangle, \\ |B_{10}\rangle &= |10_{3/2}, B\rangle + a_{27}^B |27_{3/2}, B\rangle + a_{35}^B |35_{3/2}, B\rangle, \end{aligned} \quad (2.10)$$

where the spin indices J_3 have been suppressed. The m_s -dependent coefficients in eq. (2.10) are written as

$$c_{\overline{10}}^B = c_{\overline{10}} \begin{bmatrix} \sqrt{5} \\ 0 \\ \sqrt{5} \\ 0 \end{bmatrix}, \quad c_{27}^B = c_{27} \begin{bmatrix} \sqrt{6} \\ 3 \\ 2 \\ \sqrt{6} \end{bmatrix}, \quad a_{27}^B = a_{27} \begin{bmatrix} \sqrt{15/2} \\ 2 \\ \sqrt{3/2} \\ 0 \end{bmatrix}, \quad a_{35}^B = a_{35} \begin{bmatrix} 5/\sqrt{14} \\ 2\sqrt{5/7} \\ 3\sqrt{5/14} \\ 2\sqrt{5/7} \end{bmatrix}, \quad (2.11)$$

respectively in the basis $[N, \Lambda, \Sigma, \Xi]$ and $[\Delta, \Sigma^*, \Xi^*, \Omega]$. The mixing coefficients in eq. (2.11) that contain m_s and \overline{m} are written as

$$\begin{aligned} c_{\overline{10}} &= -\frac{I_2}{15} (m_s - \overline{m}) \left(\alpha + \frac{1}{2}\gamma \right), & c_{27} &= -\frac{I_2}{25} (m_s - \overline{m}) \left(\alpha - \frac{1}{6}\gamma \right), \\ a_{27} &= -\frac{I_2}{8} (m_s - \overline{m}) \left(\alpha + \frac{5}{6}\gamma \right), & a_{35} &= -\frac{I_2}{24} (m_s - \overline{m}) \left(\alpha - \frac{1}{2}\gamma \right). \end{aligned} \quad (2.12)$$

In order to determine the dynamical parameters α, β, γ and the mixing coefficients in eq.(2.12) by using the experimental data, we have to take into account the effects of isospin symmetry breaking, which consist of the electromagnetic interactions and the up and down quark mass differences. The corresponding formalisms and analyses in detail can be found in Ref. [23] and Ref. [24], respectively. The numerical values of the coefficients that were obtained from Ref. [24] are given as

$$c_{\overline{10}} = 0.0434 \pm 0.0006, \quad c_{27} = 0.0203 \pm 0.0003,$$

$$a_{27} = 0.0903 \pm 0.0013, \quad a_{35} = 0.0181 \pm 0.0003. \quad (2.13)$$

Note that it is of great importance to know these mixing coefficients, since they allow one to carry out an unambiguous analysis of the vector and axial-vector HSD constants.

In a χ SM, the collective operators for the HSD vector and axial-vector constants can be expressed in terms of the SU(3) Wigner D functions [20, 36, 37]:

$$\begin{aligned} \hat{f}_2^{B_1 \rightarrow B_2} &= w_1 D_{\chi^3}^{(8)} + w_2 d_{pq3} D_{\chi^p}^{(8)} \hat{J}_q + \frac{w_3}{\sqrt{3}} D_{\chi^8}^{(8)} \hat{J}_3 + \frac{w_4}{\sqrt{3}} d_{pq3} D_{\chi^p}^{(8)} D_{8q}^{(8)} \\ &\quad + w_5 \left(D_{\chi^3}^{(8)} D_{88}^{(8)} + D_{\chi^8}^{(8)} D_{83}^{(8)} \right) + w_6 \left(D_{\chi^3}^{(8)} D_{88}^{(8)} - D_{\chi^8}^{(8)} D_{83}^{(8)} \right), \\ \hat{g}_1^{B_1 \rightarrow B_2} &= a_1 D_{\chi^3}^{(8)} + a_2 d_{pq3} D_{\chi^p}^{(8)} \hat{J}_q + \frac{a_3}{\sqrt{3}} D_{\chi^8}^{(8)} \hat{J}_3 + \frac{a_4}{\sqrt{3}} d_{pq3} D_{\chi^p}^{(8)} D_{8q}^{(8)} \\ &\quad + a_5 \left(D_{\chi^3}^{(8)} D_{88}^{(8)} + D_{\chi^8}^{(8)} D_{83}^{(8)} \right) + a_6 \left(D_{\chi^3}^{(8)} D_{88}^{(8)} - D_{\chi^8}^{(8)} D_{83}^{(8)} \right), \end{aligned} \quad (2.14)$$

where w_i and a_i stand for the parameters encoding the specific dynamics of a chiral solitonic model as in the case of the α , β , and γ . For example, they can be explicitly computed by using the χ QSM [21, 22]. Each w_i has its own dynamical meaning: w_1 (a_1) corresponds to the leading-order contribution, w_2 (a_2) and w_3 (a_3) arise from the rotational $1/N_c$ corrections, and w_4 (a_4), w_5 (a_5) and w_6 (a_6) are originated from flavor SU(3) symmetry breaking, in which the strange current quark mass m_s is included. There are two different linear m_s corrections to the HSD constants $\hat{f}_2^{B_1 \rightarrow B_2}$ and $\hat{g}_1^{B_1 \rightarrow B_2}$: The m_s corrections from the collective operators given in eq.(2.14) and those from the baryon wave functions because of the mixing coefficients in eq.(2.12).

3 Analysis of the HSD constants for the baryon octet

We are now in a position to determine the parameters w_i and a_i in eq.(2.14). The experimental data for the magnetic moments of the baryon octet, which are listed in Table 1, can be used for the determination of w_i .

Experimental data [11]	
μ_p	$2.792847356 \pm 0.000000023$
μ_n	-1.9130427 ± 0.0000005
μ_Λ	-0.613 ± 0.004
μ_{Σ^+}	2.458 ± 0.010
μ_{Σ^-}	-1.160 ± 0.025
μ_{Ξ^0}	-1.250 ± 0.014
μ_{Ξ^+}	-0.6507 ± 0.0025

Table 1. The experimental data for the magnetic moments of the baryon octet [11] in units of the nuclear magneton μ_N , which are used as input in the present work.

In fact, the analyses of the magnetic moments in a similar scheme to the present one were carried out in Refs. [36–38]. However, it was then not possible to determine all relevant parameters uniquely, so that the results were shown as functions either of m_s or of the πN

sigma term. With the mixing coefficients in eq.(2.13) completely determined, however, we are able to proceed to fix the parameters w_i unambiguously by using the experimental data for the baryon magnetic moments listed in Table 1. Since we have the seven independent experimental data as shown in Table 1, whereas we need to find the six parameters (w_i), we use the χ^2 -fit method. Let us set the magnetic moments of the octet baryon and the parameters w_i as vectors

$$\boldsymbol{\mu} = (\mu_p, \mu_n, \mu_\Lambda, \mu_{\Sigma^+}, \mu_{\Sigma^-}, \mu_{\Xi^0}, \mu_{\Xi^-}), \quad \boldsymbol{w} = (w_1, w_2, w_3, w_4, w_5, w_6) \quad (3.1)$$

such that the relation between $\boldsymbol{\mu}$ and \boldsymbol{w} is cast into a matrix equation

$$\boldsymbol{\mu} = \boldsymbol{A} \cdot \boldsymbol{w}, \quad (3.2)$$

where

$$\boldsymbol{A} = \begin{bmatrix} -\frac{2}{15} - \frac{4}{45}c_{27} & \frac{1}{15} - \frac{8}{45}c_{27} & \frac{1}{30} + \frac{2}{15}c_{27} & -\frac{2}{135} - \frac{1}{18} - \frac{1}{30} \\ -\frac{1}{10} + \frac{1}{3}c_{10} - \frac{2}{45}c_{27} & -\frac{1}{20} + \frac{1}{3}c_{10} - \frac{4}{45}c_{27} & \frac{1}{60} + \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & \frac{7}{270} - \frac{1}{18} - \frac{1}{30} \\ \frac{1}{20} - \frac{1}{10}c_{27} & -\frac{1}{40} - \frac{1}{5}c_{27} & \frac{1}{120} + \frac{3}{20}c_{27} & -\frac{1}{60} - 0 - 0 \\ -\frac{2}{15} - \frac{2}{45}c_{27} & \frac{1}{15} - \frac{4}{45}c_{27} & \frac{1}{30} + \frac{1}{15}c_{27} & -\frac{1}{135} - \frac{1}{90} - \frac{1}{30} \\ \frac{1}{30} + \frac{1}{3}c_{10} - \frac{2}{45}c_{27} & -\frac{1}{60} + \frac{1}{3}c_{10} - \frac{4}{45}c_{27} & -\frac{1}{20} + \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & \frac{7}{270} - \frac{1}{18} - \frac{1}{30} \\ \frac{1}{10} - \frac{2}{45}c_{27} & -\frac{1}{20} - \frac{4}{45}c_{27} & \frac{1}{60} + \frac{1}{15}c_{27} & -\frac{1}{135} - \frac{1}{90} - \frac{1}{30} \\ \frac{1}{30} - \frac{4}{45}c_{27} & -\frac{1}{60} - \frac{8}{45}c_{27} & -\frac{1}{20} + \frac{2}{15}c_{27} & -\frac{2}{135} - \frac{1}{18} - \frac{1}{30} \end{bmatrix}. \quad (3.3)$$

Having solved eq.(3.2), we determine the parameters w_i as listed in Table 2. We want to

w_1	w_2	w_3	w_4	w_5	w_6
-13.51 ± 0.01	4.15 ± 0.93	8.54 ± 0.86	-3.79 ± 0.21	-4.93 ± 0.86	-2.01 ± 0.84

Table 2. The dynamical parameters w_i for the HSD vector constants f_2

point out that the results of w_i in Ref. [36] suffered from an uncertainty, so that the magnetic moments of the baryon decuplet were given as a function of an unknown parameter p . On the other hand, the results of w_i in the present work are uniquely determined, as shown in Table 2.

Using the results of w_i in Table 2, we obtain straightforwardly f_2/f_1 for various HSD modes:

$$\begin{bmatrix} f_2/f_1 (n \rightarrow p) \\ f_2/f_1 (\Sigma^+ \rightarrow \Lambda) \\ f_2/f_1 (\Sigma^- \rightarrow n) \\ f_2/f_1 (\Xi^- \rightarrow \Lambda) \\ f_2/f_1 (\Xi^- \rightarrow \Sigma^0) \\ f_2/f_1 (\Sigma^- \rightarrow \Lambda) \\ f_2/f_1 (\Lambda \rightarrow p) \\ f_2/f_1 (\Sigma^- \rightarrow \Sigma^0) \\ f_2/f_1 (\Xi^- \rightarrow \Xi^0) \\ f_2/f_1 (\Xi^0 \rightarrow \Sigma^+) \end{bmatrix} = \mathbf{M}_V \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \\ w_5 \\ w_6 \end{bmatrix} \quad (3.4)$$

with the matrix \mathbf{M}_V expressed as

$$\mathbf{M}_V = \begin{bmatrix} -\frac{7}{30} - \frac{1}{3}c_{10} - \frac{2}{45}c_{27} & \frac{7}{60} - \frac{1}{3}c_{10} - \frac{4}{45}c_{27} & \frac{1}{60} - \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & -\frac{11}{270} & -\frac{1}{9} & -\frac{1}{15} \\ -\frac{3}{20} - \frac{1}{4}c_{10} - \frac{1}{12}c_{27} & \frac{3}{40} - \frac{1}{4}c_{10} - \frac{1}{6}c_{27} & -\frac{1}{40} - \frac{1}{8}c_{10} + \frac{1}{8}c_{27} & -\frac{7}{180} & -\frac{1}{30} & 0 \\ \frac{1}{15} - \frac{1}{45}c_{27} & -\frac{1}{30} - \frac{2}{45}c_{27} & \frac{1}{15} + \frac{1}{30}c_{27} & -\frac{1}{270} & -\frac{1}{45} & \frac{1}{30} \\ -\frac{1}{30} - \frac{1}{30}c_{27} & \frac{1}{60} - \frac{1}{15}c_{27} & \frac{1}{20} + \frac{1}{20}c_{27} & -\frac{1}{180} & \frac{1}{30} & -\frac{1}{30} \\ -\frac{7}{30} + \frac{1}{6}c_{10} + \frac{1}{45}c_{27} & \frac{7}{60} + \frac{1}{6}c_{10} + \frac{2}{45}c_{27} & \frac{1}{60} + \frac{1}{12}c_{10} - \frac{1}{30}c_{27} & \frac{11}{540} & \frac{1}{18} & \frac{1}{30} \\ -\frac{3}{20} - \frac{1}{4}c_{10} - \frac{1}{12}c_{27} & \frac{3}{40} - \frac{1}{4}c_{10} - \frac{1}{6}c_{27} & -\frac{1}{40} - \frac{1}{8}c_{10} + \frac{1}{8}c_{27} & -\frac{7}{180} & -\frac{1}{30} & 0 \\ -\frac{2}{15} + \frac{1}{6}c_{10} + \frac{1}{30}c_{27} & \frac{1}{15} + \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & \frac{1}{30} + \frac{1}{12}c_{10} - \frac{1}{20}c_{27} & \frac{1}{45} & 0 & -\frac{1}{30} \\ -\frac{1}{12} - \frac{1}{6}c_{10} & \frac{1}{24} - \frac{1}{6}c_{10} & \frac{1}{24} - \frac{1}{12}c_{10} & -\frac{1}{60} & -\frac{1}{30} & \frac{1}{30} \\ \frac{1}{15} + \frac{2}{45}c_{27} & -\frac{1}{30} + \frac{4}{45}c_{27} & \frac{1}{15} - \frac{1}{15}c_{27} & \frac{1}{135} & \frac{2}{45} & -\frac{1}{15} \\ -\frac{7}{30} + \frac{1}{6}c_{10} + \frac{1}{45}c_{27} & \frac{7}{60} + \frac{1}{6}c_{10} + \frac{2}{45}c_{27} & \frac{1}{60} + \frac{1}{12}c_{10} - \frac{1}{30}c_{27} & \frac{11}{540} & \frac{1}{18} & \frac{1}{30} \end{bmatrix}. \quad (3.5)$$

It is well known that when the effects of SU(3) symmetry breaking are turned off, the vector HSD constants $f_2(0)/f_1(0)$ are expressed in terms of the anomalous magnetic moments of the proton and the neutron as listed in Table 3. Note that the constants $f_1(0)$ do not have any linear m_s corrections because of the Ademollo-Gatto theorem [26]. However, one has to use the values of κ_p and κ_n in the exact SU(3) symmetry to compute

Decay mode	$f_1(0)$	$f_2(0)$	$f_2(0)/f_1(0)$
$n \rightarrow p$	1	$\frac{1}{2}(\kappa_p - \kappa_n)$	$\frac{1}{2}(\kappa_p - \kappa_n)$
$\Sigma^- \rightarrow \Lambda$	0	$-\frac{1}{2}\sqrt{\frac{3}{2}}\kappa_n$	$\sqrt{\frac{3}{2}}f_2^{\Sigma^- \rightarrow \Lambda}(0)$
$\Sigma^0 \rightarrow \Sigma^+$	$\sqrt{2}$	$\frac{1}{\sqrt{2}}(\kappa_p + \frac{1}{2}\kappa_n)$	$\frac{1}{2}(\kappa_p + \frac{1}{2}\kappa_n)$
$\Sigma^- \rightarrow \Sigma^0$	$\sqrt{2}$	$\frac{1}{\sqrt{2}}(\kappa_p + \frac{1}{2}\kappa_n)$	$\frac{1}{2}(\kappa_p + \frac{1}{2}\kappa_n)$
$\Xi^- \rightarrow \Xi^0$	1	$\frac{1}{2}(\kappa_p + 2\kappa_n)$	$\frac{1}{2}(\kappa_p + 2\kappa_n)$
$\Lambda \rightarrow p$	$\sqrt{\frac{3}{2}}$	$\frac{1}{2}\sqrt{\frac{3}{2}}\kappa_p$	$\frac{1}{2}\kappa_p$
$\Sigma^0 \rightarrow p$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2\sqrt{2}}(\kappa_p + 2\kappa_n)$	$\frac{1}{2}(\kappa_p + 2\kappa_n)$
$\Sigma^- \rightarrow n$	1	$\frac{1}{2}(\kappa_p + 2\kappa_n)$	$\frac{1}{2}(\kappa_p + 2\kappa_n)$
$\Xi^0 \rightarrow \Sigma^+$	1	$\frac{1}{2}(\kappa_p - \kappa_n)$	$\frac{1}{2}(\kappa_p - \kappa_n)$
$\Xi^- \rightarrow \Lambda$	$\sqrt{\frac{3}{2}}$	$\frac{1}{2}\sqrt{\frac{3}{2}}(\kappa_p + \kappa_n)$	$\frac{1}{2}(\kappa_p + \kappa_n)$
$\Xi^- \rightarrow \Sigma^0$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2\sqrt{2}}(\kappa_p - \kappa_n)$	$\frac{1}{2}(\kappa_p - \kappa_n)$

Table 3. The expressions of $f_1(0)$, $f_2(0)$, and $g_1(0)$ in exact SU(3) symmetry. The κ_p and the κ_n denote the anomalous magnetic moments of the proton and the neutron, respectively.

$f_2(0)$ in SU(3) symmetry, since the experimental data for the nucleon anomalous magnetic moments already include the effects of SU(3) symmetry breaking. The numerical values of anomalous magnetic moments of proton and neutrons for the contributions of the order

$\mathcal{O}(m_s^0)$ and $\mathcal{O}(m_s^0) + \mathcal{O}(m_s^1)$ are obtained as follows:

$$\begin{aligned}\kappa_p^{\text{sym}} [\mathcal{O}(m_s^0)] &= 1.363 \pm 0.069, \\ \kappa_n^{\text{sym}} [\mathcal{O}(m_s^0)] &= -1.416 \pm 0.049, \\ \kappa_p [\mathcal{O}(m_s^0) + \mathcal{O}(m_s^1)] &= 1.793 \pm 0.087, \\ \kappa_n [\mathcal{O}(m_s^0) + \mathcal{O}(m_s^1)] &= -1.913 \pm 0.069,\end{aligned}\tag{3.6}$$

where κ_p^{sym} and κ_n^{sym} represent the anomalous magnetic moments of the proton and the neutron in the exact SU(3) symmetry. We can estimate the effect of SU(3) symmetry breaking from eq.(3.6), which is approximately 25 %.

Decay mode	$(f_2/f_1)^{\text{sym}}$	$(f_2/f_1)^{\text{br}}$	Experimental data [11]
$n \rightarrow p$	1.389 ± 0.042	1.853 ± 0.056	
$\Sigma^- \rightarrow \Lambda$	1.062 ± 0.037	1.435 ± 0.052	
$\Sigma^- \rightarrow \Sigma^0$	0.328 ± 0.037	0.418 ± 0.047	
$\Xi^- \rightarrow \Xi^0$	-0.734 ± 0.060	-1.017 ± 0.082	
$\Lambda \rightarrow p$	0.681 ± 0.035	0.896 ± 0.043	
$\Sigma^- \rightarrow n$	-0.734 ± 0.060	-1.017 ± 0.082	-0.97 ± 0.14
$\Xi^0 \rightarrow \Sigma^+$	1.389 ± 0.042	1.853 ± 0.056	$2.0 \pm 1.2 \pm 0.5$
$\Xi^- \rightarrow \Lambda$	-0.026 ± 0.042	-0.060 ± 0.056	
$\Xi^- \rightarrow \Sigma^0$	1.389 ± 0.042	1.853 ± 0.056	

Table 4. Numerical results for the ratios of the vector HSD constants f_2/f_1 of the baryon octet. The superscripts “sym” and “br” denote the contribution in exact SU(3) symmetry and the total results with flavor SU(3) symmetry breaking, respectively. Note that the values for the $\Sigma^- \rightarrow \Lambda$ mode are given for $\sqrt{3}/2 f_2$ instead of f_2/f_1 .

In Table 4 we list the present results for the vector HSD constants $f_2(0)/f_1(0)$. There are only two existing data for the decay processes $\Sigma^- \rightarrow ne^- \bar{\nu}_e$ and $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ from the PDG. The present results are in agreement with these experimental data, though the experimental uncertainties of $f_2(0)/f_1(0)$ for the $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ decay are rather large. Note that there are old data for the $\Lambda \rightarrow pe^- \bar{\nu}_e$ decay [12, 39, 40]. In Ref. [39], the vector HSD constant for $\Lambda \rightarrow pe^- \bar{\nu}_e$ decay is given as $f_2(0)/f_1(0)(\Lambda \rightarrow p) = 0.15 \pm 0.30$. However, this value needs to be scrutinized experimentally, because firstly its uncertainty is very large and secondly its value itself is quite different from most theoretical predictions. In fact, Cabbibo et al. [6] reviewed the old experimental data [41, 42] and extracted the following values: $f_1 = 1.238 \pm 0.024$ and $f_2 = 1.34 \pm 0.20$, which is in line with the old data: $f_1 = 1.229 \pm 0.035$ [42]. The prediction from the present work is $f_2 = 1.098 \pm 0.053$, which is closer to that from Ref. [6]. As shown in Table 4, the effects of SU(3) symmetry breaking on the HSD vector constants appear consistently to be more than 20 %.

In order to determine the dynamical parameters a_i in eq. (2.14), we use the experimental data for the HSD constants $g_1/f_1(B_1 \rightarrow B_2)$ listed in Table 5. Since there exist only five experimental data for them as listed in Table 5 but six unknown parameters a_i , we need at least one more data. There are three known diagonal axial-vector constants, that is, $g_A^{(0)}$, $g_A^{(3)}$, and $g_A^{(8)}$. However, the triplet axial-vector constant $g_A^{(3)}$ is the same as

that for the neutron β decay $g_1/f_1(n \rightarrow p)$, we have to avoid it. The octet one $g_A^{(8)}$ is usually determined by using the HSD constants with SU(3) symmetry assumed, it is also not suitable for the present purpose. The rest is the singlet axial-vector constant $g_A^{(0)}$, which implies the quark content of the nucleon spin and is determined by the data for polarized electron-proton deep inelastic scattering. Thus, in addition to the experimental data for the five known HSD constants, we utilize $g_A^{(0)}$ to fix a_i .

	Experimental data	References
$g_1/f_1 (n \rightarrow p)$	1.2701 ± 0.0025	PDG [11]
$g_1/f_1 (\Lambda \rightarrow p)$	0.718 ± 0.015	PDG [11]
$g_1/f_1 (\Sigma^- \rightarrow n)$	-0.340 ± 0.017	PDG [11]
$g_1/f_1 (\Xi^- \rightarrow \Lambda)$	0.25 ± 0.05	PDG [11]
$g_1/f_1 (\Xi^0 \rightarrow \Sigma^+)$	1.21 ± 0.05	PDG [11]
g_A^0	0.36 ± 0.03	Bass et al. [43]

Table 5. The experimental data for the HSD constants from the Particle Data Group [11].

With these six input data from experiments at hand, the dynamical parameters a_i can be determined by solving the matrix equation similar to eq.(3.2). The parameters a_i determined by eq. (3.7) are listed in Table 6. Having fixed a_i , we can straightforwardly

a_1	a_2	a_3	a_4	a_5	a_6
-3.51 ± 0.01	3.44 ± 0.03	0.60 ± 0.03	-1.21 ± 0.07	0.48 ± 0.02	-0.74 ± 0.04

Table 6. The dynamical parameters of axial-vector transitions

solve the following matrix equation to find other HSD constants and g_A^8 :

$$\begin{bmatrix} g_1/f_1 (n \rightarrow p) \\ g_1/f_1 (\Sigma^+ \rightarrow \Lambda) \\ g_1/f_1 (\Lambda \rightarrow p) \\ g_1/f_1 (\Sigma^- \rightarrow n) \\ g_1/f_1 (\Xi^- \rightarrow \Lambda) \\ g_1/f_1 (\Xi^- \rightarrow \Sigma^0) \\ g_1/f_1 (\Sigma^- \rightarrow \Lambda) \\ g_1/f_1 (\Sigma^- \rightarrow \Sigma^0) \\ g_1/f_1 (\Xi^- \rightarrow \Xi^0) \\ g_1/f_1 (\Xi^0 \rightarrow \Sigma^+) \\ g_A^0(p) \\ g_A^3(p) \\ g_A^8(p) \end{bmatrix} = \mathbf{M}_A \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \\ a_6 \end{bmatrix}, \quad (3.7)$$

where the matrix \mathbf{M}_A is expressed as

$$\mathbf{M}_A = \begin{bmatrix} -\frac{7}{30} - \frac{1}{3}c_{10} - \frac{2}{45}c_{27} & \frac{7}{60} - \frac{1}{3}c_{10} - \frac{4}{45}c_{27} & \frac{1}{60} - \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & -\frac{11}{270} - \frac{1}{9} - \frac{1}{15} \\ -\frac{3}{20} - \frac{1}{4}c_{10} - \frac{1}{12}c_{27} & \frac{3}{40} - \frac{1}{4}c_{10} - \frac{1}{6}c_{27} & -\frac{1}{40} - \frac{1}{8}c_{10} + \frac{1}{8}c_{27} & -\frac{7}{180} - \frac{1}{30} & 0 \\ -\frac{2}{15} + \frac{1}{6}c_{10} + \frac{1}{30}c_{27} & \frac{1}{15} + \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & \frac{1}{30} + \frac{1}{12}c_{10} - \frac{1}{20}c_{27} & \frac{1}{45} & 0 & -\frac{1}{30} \\ \frac{1}{15} - \frac{1}{45}c_{27} & -\frac{1}{30} - \frac{2}{45}c_{27} & \frac{1}{15} + \frac{1}{30}c_{27} & -\frac{1}{270} - \frac{1}{45} & \frac{1}{30} \\ -\frac{1}{30} - \frac{1}{30}c_{27} & \frac{1}{60} - \frac{1}{15}c_{27} & \frac{1}{20} + \frac{1}{20}c_{27} & -\frac{1}{180} & \frac{1}{30} & -\frac{1}{30} \\ -\frac{7}{30} + \frac{1}{6}c_{10} + \frac{1}{45}c_{27} & \frac{7}{60} + \frac{1}{6}c_{10} + \frac{2}{45}c_{27} & \frac{1}{60} + \frac{1}{12}c_{10} - \frac{1}{30}c_{27} & \frac{11}{540} & \frac{1}{18} & \frac{1}{30} \\ -\frac{3}{20} - \frac{1}{4}c_{10} - \frac{1}{12}c_{27} & \frac{3}{40} - \frac{1}{4}c_{10} - \frac{1}{6}c_{27} & -\frac{1}{40} - \frac{1}{8}c_{10} + \frac{1}{8}c_{27} & -\frac{7}{180} - \frac{1}{30} & 0 \\ -\frac{1}{12} - \frac{1}{6}c_{10} & \frac{1}{24} - \frac{1}{6}c_{10} & \frac{1}{24} - \frac{1}{12}c_{10} & -\frac{1}{60} - \frac{1}{30} & \frac{1}{30} \\ \frac{1}{15} + \frac{2}{45}c_{27} & -\frac{1}{30} + \frac{4}{45}c_{27} & \frac{1}{15} - \frac{1}{15}c_{27} & \frac{1}{135} & \frac{2}{45} & -\frac{1}{15} \\ -\frac{7}{30} + \frac{1}{6}c_{10} + \frac{1}{45}c_{27} & \frac{7}{60} + \frac{1}{6}c_{10} + \frac{2}{45}c_{27} & \frac{1}{60} + \frac{1}{12}c_{10} - \frac{1}{30}c_{27} & \frac{11}{540} & \frac{1}{18} & \frac{1}{30} \\ 0 & 0 & 1 & 0 & -\frac{1}{5} & \frac{1}{5} \\ -\frac{7}{30} - \frac{1}{3}c_{10} - \frac{2}{45}c_{27} & \frac{7}{60} - \frac{1}{3}c_{10} - \frac{4}{45}c_{27} & \frac{1}{60} - \frac{1}{6}c_{10} + \frac{1}{15}c_{27} & -\frac{11}{270} - \frac{1}{9} - \frac{1}{15} \\ -\frac{1}{10\sqrt{3}} + \frac{1}{\sqrt{3}}c_{10} - \frac{2}{5\sqrt{3}}c_{27} & \frac{1}{20\sqrt{3}} + \frac{1}{\sqrt{3}}c_{10} - \frac{4}{5\sqrt{3}}c_{27} & \frac{\sqrt{3}}{20} + \frac{1}{2\sqrt{3}}c_{10} + \frac{\sqrt{3}}{5}c_{27} & \frac{1}{30\sqrt{3}} & 0 & 0 \end{bmatrix}. \quad (3.8)$$

In the exact SU(3) symmetry, all axial-vector decay amplitudes are given in terms of two reduced matrix elements F and D as follows:

$$g_1^{(B_1 \rightarrow B_2)}(0) = C_F^{(B_1 \rightarrow B_2)} F + C_D^{(B_1 \rightarrow B_2)} D. \quad (3.9)$$

Here $C_F^{(B \rightarrow B')}$ and $C_D^{(B \rightarrow B')}$ are SU(3) Clebsch-Gordan coefficients that appear when the axial-vector collective operator $\hat{g}_1^{B_1 \rightarrow B_2}$ in eq.(2.14) is sandwiched between octet states. Thus, the F and D are determined in terms of a_1 , a_2 , and a_3

$$\begin{aligned} F &= -\frac{1}{12} \left(a_1 - \frac{1}{2}a_2 \right) + \frac{1}{24}a_3 = 0.461 \pm 0.002, \\ D &= -\frac{3}{20} \left(a_1 - \frac{1}{2}a_2 \right) - \frac{1}{40}a_3 = 0.769 \pm 0.003. \end{aligned} \quad (3.10)$$

The results for the F and D are in good agreement with the empirical ones extracted from the experiments [44], given as $F = 0.463 \pm 0.008$ and $D = 0.804 \pm 0.008$.

In Table 8, the results for the axial-vector HSD constants g_1/f_1 are listed. In contrast to the results for f_2/f_1 , the effects of flavor SU(3) symmetry breaking on the axial-vector HSD constants are rather small and stable. However, when it comes to the singlet axial charge, they are not negligible. Note that though $g_A^{(0)}$ is used as an input, we still can estimate the effects of SU(3) symmetry breaking, which turn out to be approximately 40 %. The reason why the contribution of SU(3) symmetry breaking is large in the case the singlet axial charge can be found in eqs.(3.7, 3.8). The expression for $g_A^{(0)}$ is written as

$$g_A^{(0)} = a_3 - \frac{1}{5}(a_5 - a_6). \quad (3.11)$$

The singlet axial charge does not have any contribution from the leading order in the $1/N_c$ expansion. Indeed, this explains why the Skyrme model predicts the smallness of $g_A^{(0)}$.

Decay mode	$f_1(0)$	$g_1(0)$	$g_1(0)/f_1(0)$
$n \rightarrow p$	1	$F + D$	$F + D$
$\Sigma^- \rightarrow \Lambda$	0	$\sqrt{\frac{2}{3}}D$	D
$\Sigma^0 \rightarrow \Sigma^+$	$\sqrt{2}$	$\sqrt{2}F$	F
$\Sigma^- \rightarrow \Sigma^0$	$\sqrt{2}$	$\sqrt{2}F$	F
$\Xi^- \rightarrow \Xi^0$	1	$F - D$	$F - D$
$\Lambda \rightarrow p$	$\sqrt{\frac{3}{2}}$	$\sqrt{\frac{3}{2}}(F + \frac{1}{3}D)$	$F + \frac{1}{3}D$
$\Sigma^0 \rightarrow p$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}(F - D)$	$F - D$
$\Sigma^- \rightarrow n$	1	$F - D$	$F - D$
$\Xi^0 \rightarrow \Sigma^+$	1	$F + D$	$F + D$
$\Xi^- \rightarrow \Lambda$	$\sqrt{\frac{3}{2}}$	$\sqrt{\frac{3}{2}}(F - \frac{1}{3}D)$	$F - \frac{1}{3}D$
$\Xi^- \rightarrow \Sigma^0$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}(F + D)$	$F + D$

Table 7. The expressions of $g_1(0)$ and $g_1(0)/f_1(0)$ in exact SU(3) symmetry.

Decay modes	$(g_1/f_1)^{\text{sym}}$	$(g_1/f_1)^{\text{br}}$	Exp. (Input)	Refs.
$n \rightarrow p$	1.230 ± 0.004	1.269 ± 0.006	1.2701 ± 0.0025	[11]
$\Sigma^- \rightarrow \Lambda$	0.769 ± 0.003	0.794 ± 0.004		
$\Sigma^- \rightarrow \Sigma^0$	0.461 ± 0.002	0.439 ± 0.003		
$\Xi^- \rightarrow \Xi^0$	-0.308 ± 0.002	-0.245 ± 0.004		
$\Lambda \rightarrow p$	0.717 ± 0.003	0.718 ± 0.003	0.718 ± 0.015	[11]
$\Sigma^- \rightarrow n$	-0.308 ± 0.002	-0.340 ± 0.003	-0.340 ± 0.017	[11]
$\Xi^0 \rightarrow \Sigma^+$	1.230 ± 0.004	1.210 ± 0.005	1.21 ± 0.05	[11]
$\Xi^- \rightarrow \Lambda$	0.204 ± 0.002	0.250 ± 0.002	0.25 ± 0.05	[11]
$\Xi^- \rightarrow \Sigma^0$	1.230 ± 0.004	1.210 ± 0.005		
$g_A^{(0)}$	0.604 ± 0.030	0.361 ± 0.031	0.36 ± 0.03	[43]
$g_A^{(8)}$	0.354 ± 0.003	0.325 ± 0.004		

Table 8. Numerical results for the ratios of the axial-vector HSD constants g_1/f_1 of the baryon octet. The superscripts “sym” and “br” denote the contribution in exact SU(3) symmetry and the total results with flavor SU(3) symmetry breaking, respectively. Note that the values for the $\Sigma^- \rightarrow \Lambda$ mode are given for $\sqrt{3/2}g_1$ instead of g_1/f_1 . The experimental data are taken from the Particle Data Group [11], which are used for the input.

Because of this fact, there is no wavefunction corrections to $g_A^{(0)}$. Moreover, the relative signs of a_5 and a_6 are different, so that the linear m_s corrections turn out to be rather large. Thus, the effects of SU(3) symmetry breaking play an essential role in describing the quark content of the nucleon spin.

The octet axial-vector constant $g_A^{(8)}$ was usually extracted from the HSD data with flavor SU(3) symmetry assumed and its value was obtained to be $g_A^{(8)} = 0.58 \pm 0.03$ [45].

On the other hand, the present result for $g_A^{(8)}$ in the SU(3) symmetric case is given as $g_A^{(8)} = 0.354 \pm 0.003$. With the linear m_s corrections taken into account, the value of $g_A^{(8)}$ is reduced to $g_A^{(8)} = 0.325 \pm 0.004$, which is not much different from that with SU(3) symmetry. However, compared with the extracted value from Ref. [45], the present result for $g_A^{(8)}$ is approximately 40 % smaller than that. Note that the octet axial charge is expressed in terms of F and D as follows:

$$g_A^{(8)} = \frac{1}{\sqrt{3}}(3F - D) = \frac{1}{\sqrt{3}} \left[-\frac{1}{10}(a_1 - \frac{1}{2}a_2) + \frac{1}{20}a_3 \right]. \quad (3.12)$$

Inserting the values of F and D given in Ref. [44], we find $g_A^{(8)} = 0.338 \pm 0.015$, which is quite different from that of Ref. [45] but is compatible with the present value.

4 Analysis for the transition from the baryon decuplet to the octet

In this Section, we want to present the results for the axial-vector transition constants of the baryon decuplet. Since the baryon decuplet except for Ω^- decays into the baryon octet strongly, there are no direct experimental data for the axial-vector HSD constants for the baryon decuplet to date. However, there are at least two evident reasons why the axial-vector transition constants for the baryon decuplet are interesting. Firstly, the baryon decuplet can be produced by exclusive neutrino-nucleon scattering such as $\nu + N \rightarrow \mu + \Delta$ and $\nu + N \rightarrow \mu + B_{10} + M_8$ processes, where M_8 stands for the pseudo-scalar meson octet. These reactions are also interesting, since they may bring out information on weak generalized parton distributions [46]. The axial-vector transition constants for the baryon decuplet might be extracted from these reactions. Secondly, using the Goldberger-Treiman relation, one can relate the axial-vector transition constants to the strong coupling constants for vertices of the baryon decuplet-octet and the meson octet. Because of these reasons, it is of great interest to understand the axial-vector transition constants for the baryon decuplet.

There are four different axial-vector form factors or Adler form factors $C_i^A(q^2)$ ($i = 3, 4, 5, 6$) as given in eq.(2.3). However, C_3^A and C_4^A are often neglected, since the partially conserved axial-vector current (PCAC) relation yields an expression, where the coefficients in front of C_3^A and C_4^A are proportional to the mass difference of a decuplet baryon and a octet one divided by the mass sum, and its squares, respectively [47]. Thus, their contribution is negligible in the transition amplitude of $\nu N \rightarrow \mu B_{10}$ scattering. In the chiral limit, C_6^A can be related to C_5^A , so that we will concentrate on C_5^A in this work.

Before we present the results, we want to discuss possible relations and sum rules for the axial-vector transition constants C_5^A . With the linear m_s corrections turned on, we

find the following eight relations for C_5^A :

$$\begin{aligned}
(\Delta^0 \rightarrow p) &= \frac{1}{\sqrt{3}} (\Delta^- \rightarrow n) = -\frac{1}{\sqrt{3}} (\Delta^{++} \rightarrow p) = -(\Delta^+ \rightarrow n), \\
(\Sigma^{*0} \rightarrow \Sigma^+) &= (\Sigma^{*-} \rightarrow \Sigma^0) = (\Sigma^{*+} \rightarrow \Sigma^0) = (\Sigma^{*0} \rightarrow \Sigma^-), \\
(\Sigma^{*-} \rightarrow \Lambda) &= (\Sigma^{*+} \rightarrow \Lambda), \\
(\Xi^{*-} \rightarrow \Xi^0) &= (\Xi^{*0} \rightarrow \Xi^-), \\
(\Delta^{++} \rightarrow \Sigma^+) &= \sqrt{\frac{3}{2}} (\Delta^+ \rightarrow \Sigma^0) = \sqrt{3} (\Delta^0 \rightarrow \Sigma^-), \\
(\Sigma^{*0} \rightarrow p) &= \frac{1}{\sqrt{2}} (\Sigma^{*-} \rightarrow n), \\
(\Sigma^{*+} \rightarrow \Xi^0) &= \sqrt{2} (\Sigma^{*0} \rightarrow \Xi^-), \\
(\Xi^{*0} \rightarrow \Sigma^+) &= \sqrt{2} (\Xi^{*-} \rightarrow \Sigma^0). \tag{4.1}
\end{aligned}$$

These relations come from isospin symmetry. We also obtain the following six sum rules for the C_5^A :

$$\begin{aligned}
(\Delta^0 \rightarrow p) &= -(\Xi^{*-} \rightarrow \Xi^0) - \frac{1}{\sqrt{2}} (\Xi^{*-} \rightarrow \Sigma^0) + \sqrt{\frac{3}{2}} (\Xi^{*-} \rightarrow \Lambda) + \frac{2}{\sqrt{3}} (\Omega^- \rightarrow \Xi^0), \\
(\Sigma^{*0} \rightarrow \Sigma^+) &= \frac{1}{\sqrt{2}} (\Xi^{*-} \rightarrow \Xi^0) + (\Xi^{*-} \rightarrow \Sigma^0) - \frac{1}{\sqrt{6}} (\Omega^- \rightarrow \Xi^0), \\
(\Sigma^{*-} \rightarrow \Lambda) &= -\sqrt{\frac{3}{2}} (\Xi^{*-} \rightarrow \Xi^0) + (\Xi^{*-} \rightarrow \Lambda) + \frac{1}{\sqrt{2}} (\Omega^- \rightarrow \Xi^0), \\
(\Xi^{*-} \rightarrow \Xi^0) &= \sqrt{\frac{2}{3}} (\Xi^{*-} \rightarrow \Lambda) - \sqrt{\frac{2}{3}} (\Sigma^{*-} \rightarrow \Lambda) + \frac{1}{\sqrt{3}} (\Omega^- \rightarrow \Xi^0), \\
(\Sigma^{*0} \rightarrow p) &= -\frac{1}{2} (\Xi^{*-} \rightarrow \Sigma^0) + \frac{\sqrt{3}}{2} (\Xi^{*-} \rightarrow \Lambda) + \frac{1}{\sqrt{6}} (\Omega^- \rightarrow \Xi^0), \\
(\Delta^{++} \rightarrow \Sigma^+) &= -\sqrt{6} (\Xi^{*-} \rightarrow \Sigma^0) + \sqrt{6} (\Sigma^{*0} \rightarrow \Xi^-) + (\Omega^- \rightarrow \Xi^0). \tag{4.2}
\end{aligned}$$

Though we do not expect that the sum rules in eq.(4.2) will be confirmed by the experimental data in near future, they provide a consistency check of the present results.

In Tables 9 and 10, we list the results for the axial-vector transition constants of the baryon decuplet decaying into the octet in the strangeness-conserving cases ($\Delta S = 0$). Table 9 presents those for the positive charge difference between the initial state and the final state ($\Delta Q = +1$), and Table 10 for the negative charge difference ($\Delta Q = -1$). Tables 11 and 12 list the results for C_5^A in the strangeness-changing cases ($\Delta S = 1$). Note that all the results presented in Tables 9, 10, 11 and 12 satisfy the isospin relations given in eq.(4.1) and also the sum rules in eq.(4.2). The effects of flavor SU(3) symmetry breaking turn out to be rather small in the case of C_5^A .

$B_{10} \xrightarrow{X=1+i2} B_8$	$C_5^A(\text{sym})$	$C_5^A(\text{br})$
$\Delta^0 \rightarrow p$	-0.954 ± 0.003	-1.040 ± 0.005
$\Delta^- \rightarrow n$	-1.653 ± 0.006	-1.801 ± 0.008
$\Sigma^{*0} \rightarrow \Sigma^+$	0.675 ± 0.002	0.614 ± 0.004
$\Sigma^{*-} \rightarrow \Sigma^0$	0.675 ± 0.002	0.614 ± 0.004
$\Sigma^{*-} \rightarrow \Lambda$	-1.169 ± 0.004	-1.231 ± 0.005
$\Xi^{*-} \rightarrow \Xi^0$	0.954 ± 0.003	0.903 ± 0.006

Table 9. Axial-vector transition constants of the baryon decuplet to the octet with $X = 1 + i2$, i.e. in the strangeness-conserving ($\Delta S = 0$) case. The charge difference between the initial state and the final state is $\Delta Q = +1$.

$B_{10} \xrightarrow{X=1-i2} B_8$	$C_5^A(\text{sym})$	$C_5^A(\text{br})$
$\Delta^{++} \rightarrow p$	1.653 ± 0.006	1.801 ± 0.008
$\Delta^+ \rightarrow n$	0.954 ± 0.003	1.040 ± 0.005
$\Sigma^{*+} \rightarrow \Sigma^0$	0.675 ± 0.002	0.614 ± 0.004
$\Sigma^{*+} \rightarrow \Lambda$	1.169 ± 0.004	1.231 ± 0.005
$\Sigma^{*0} \rightarrow \Sigma^-$	0.675 ± 0.002	0.614 ± 0.004
$\Xi^{*0} \rightarrow \Xi^-$	0.954 ± 0.003	0.903 ± 0.006

Table 10. Axial-vector transition constants of the baryon decuplet to octet with the flavor transition operator of $X = 1 - i2$, i.e. in the strangeness-conserving ($\Delta S = 0$) case. The charge difference between the initial state and the final state is $\Delta Q = -1$.

$B_{10} \xrightarrow{X=4+i5} B_8$	$C_5^A(\text{sym})$	$C_5^A(\text{br})$
$\Sigma^{*0} \rightarrow p$	-0.675 ± 0.002	-0.755 ± 0.004
$\Sigma^{*-} \rightarrow n$	-0.954 ± 0.003	-1.067 ± 0.005
$\Xi^{*0} \rightarrow \Sigma^+$	0.954 ± 0.003	0.896 ± 0.004
$\Xi^{*-} \rightarrow \Sigma^0$	0.675 ± 0.002	0.633 ± 0.003
$\Xi^{*-} \rightarrow \Lambda$	-1.169 ± 0.004	-1.266 ± 0.005
$\Omega^- \rightarrow \Xi^0$	1.653 ± 0.006	1.612 ± 0.007

Table 11. Axial-vector transition constants of the baryon decuplet to the octet with $X = 4 + i5$, i.e. in the strangeness-changing ($\Delta S = 1$) case. The charge difference between the initial state and the final state is $\Delta Q = +1$.

In the SU(6) quark model, the $C_5^A(n \rightarrow \Delta^+)$ can be related to the axial-vector HSD constant $g_1/f_1(n \rightarrow p)$ [27, 48] as

$$C_5^A(\Delta^+ \rightarrow n) = \frac{2\sqrt{3}}{5} g_1/f_1(n \rightarrow p). \quad (4.3)$$

Equation (4.3) has been often used to determine the $\pi N \Delta$ coupling constant. Thus, it is interesting to examine it based on the present result. Using the experimental data for $g_1/f_1(n \rightarrow p)$, we obtain $C_5^A(\Delta^+ \rightarrow n) = 0.88$, whereas the present result from Table 10 is

$B_{10} \xrightarrow{X=4-i5} B_8$	$C_5^A(\text{sym})$	$C_5^A(\text{br})$
$\Delta^{++} \rightarrow \Sigma^+$	-1.653 ± 0.006	-1.547 ± 0.007
$\Delta^+ \rightarrow \Sigma^0$	-1.350 ± 0.005	-1.263 ± 0.005
$\Delta^+ \rightarrow \Lambda$	0	0
$\Delta^0 \rightarrow \Sigma^-$	-0.954 ± 0.003	-0.893 ± 0.004
$\Sigma^{*+} \rightarrow \Xi^0$	-0.954 ± 0.003	-0.928 ± 0.004
$\Sigma^{*0} \rightarrow \Xi^-$	-0.675 ± 0.002	-0.656 ± 0.003

Table 12. Axial-vector transition constants of the baryon decuplet to the octet with $X = 4 - i5$, i.e. in the strangeness-changing ($\Delta S = 1$) case. The charge difference between the initial state and the final state is $\Delta Q = -1$.

$C_5^A(\Delta^+ \rightarrow n) = 0.95$. Thus, the value from the SU(6) relation is deviated from the present one approximately by 10 %.

5 Summary and outlook

In the present work, we aimed at investigating the hyperon semileptonic decay constants of the baryon octet within the framework of a chiral soliton model, determining all the dynamical parameters by the experimental data. We first reviewed the analysis of the mass splittings of the baryon octet and decuplet in brief. Then, we studied the vector hyperon semileptonic decay constants f_2/f_1 with the dynamical parameters w_i fixed in an unambiguous manner by the experimental data for the magnetic moments of the baryon octet. we found that the effects of flavor SU(3) symmetry breaking contributed to the f_2/f_1 approximately by 25 %.

The dynamical parameters a_i were fixed by using the experimental data for the axial-vector hyperon semileptonic decay constants as well as the singlet axial charge $g_A^{(0)}$ unequivocally. We first derived the F and D in exact SU(3) symmetry and the results were consistent with the empirical data. We also predicted other g_1/f_1 which are not measured yet. The octet $g_A^{(8)}$ was obtained to be $g_A^{(8)} = 0.325 \pm 0.004$. The contribution of flavor SU(3) symmetry breaking turned out to be rather small in the case of the axial-vector hyperon semileptonic decay constants. Finally, we predicted the axial-vector constants for the transitions from the baryon decuplet to the octet. In addition, we derived the isospin relations and their six different sum rules.

The present results for the axial-vector transition constants for the baryon decuplet can be used to determine the meson-baryon Yukawa coupling constants by the Goldberger-Treiman relations. The whole analysis including the baryon decuplet-decuplet-meson coupling constants is under way. Last but not least, the vector transition constants for the baryon decuplet are also interesting and important. However, experimental information on the E_2/M_1 ratio for the baryon decuplet is essential but only that of the Δ isobar is known. On the other hand, if one assumes SU(3) symmetry, it might be possible to study the vector transition constants for the baryon decuplet. The corresponding investigation will appear elsewhere.

Acknowledgments

The present work was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (Grant Numbers: NRF-2013R1A1A2063590 and NRF-2012R1A1A2001083).

References

- [1] N. Cabibbo, *Unitary symmetry and leptonic decays*, Phys. Rev. Lett. **10** (1963) 531.
- [2] M. Kobayashi and T. Maskawa, *CP Violation in the Renormalizable Theory of Weak Interaction*, Prog. Theor. Phys. **49** (1973) 652.
- [3] E. Blucher, E. De Lucia, G. Isidori, V. Lubicz, H. Abele, V. Cirigliano, R. Flores-Mendieta and J. Flynn *et al.*, *Status of the Cabibbo angle*, hep-ph/0512039.
- [4] J. C. Hardy and I. S. Towner, *The measurement and interpretation of superallowed $0^+ \rightarrow 0^+$ nuclear β decay*, J. Phys. G **41** (2014) 11, 114004 [arXiv:1312.3587 [nucl-ex]].
- [5] M. Antonelli, V. Cirigliano, G. Isidori, F. Mescia, M. Moulson, H. Neufeld, E. Passemar and M. Palutan *et al.*, *An Evaluation of $|V_{us}|$ and precise tests of the Standard Model from world data on leptonic and semileptonic kaon decays*, Eur. Phys. J. C **69** (2010) 399 [arXiv:1005.2323 [hep-ph]].
- [6] N. Cabibbo, E. C. Swallow and R. Winston, *Semileptonic hyperon decays*, Ann. Rev. Nucl. Part. Sci. **53** (2003) 39 [hep-ph/0307298].
- [7] N. Cabibbo, E. C. Swallow and R. Winston, *Semileptonic hyperon decays and CKM unitarity*, Phys. Rev. Lett. **92** (2004) 251803 [hep-ph/0307214].
- [8] A. A. Affolder *et al.* [KTeV E832/E799 Collaboration], *Observation of the decay $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$* , Phys. Rev. Lett. **82** (1999) 3751.
- [9] A. Alavi-Harati *et al.* [KTeV Collaboration], *First measurement of form-factors of the decay $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$* , Phys. Rev. Lett. **87** (2001) 132001 [hep-ex/0105016].
- [10] J. R. Batley *et al.* [NA48/I Collaboration], *Measurement of the branching ratios of the decays $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}_e$ and $\Xi^0 \rightarrow \bar{\Sigma}^+ e^+ \nu_e$* , Phys. Lett. B **645** (2007) 36 [hep-ex/0612043].
- [11] K. A. Olive *et al.* [Particle Data Group Collaboration], *Review of Particle Physics*, Chin. Phys. C **38** (2014) 090001.
- [12] M. Bourquin *et al.* [Bristol-Geneva-Heidelberg-Orsay-Rutherford-Strasbourg Collaboration], *Measurements of Hyperon Semileptonic Decays at the CERN Super Proton Synchrotron. II. The $\Lambda \rightarrow p e \bar{\nu}_e$, $\Xi^- \rightarrow \Lambda e \bar{\nu}_e$, and $\Xi^- \rightarrow \Sigma^0 e \bar{\nu}_e$ Decay Modes*, Z. Phys. C **21** (1983) 1.
- [13] G. Villadoro, *Chiral corrections to the hyperon vector form factors*, Phys. Rev. D **74** (2006) 014018 [hep-ph/0603226].
- [14] A. Lacour, B. Kubis and U. G. Meissner, *Hyperon decay form-factors in chiral perturbation theory*, JHEP **0710** (2007) 083 [arXiv:0708.3957 [hep-ph]].

- [15] A. Faessler, T. Gutsche, B. R. Holstein, M. A. Ivanov, J. G. Korner and V. E. Lyubovitskij, *Phys. Rev. D* **78**, 094005 (2008) [arXiv:0809.4159 [hep-ph]].
- [16] D. Guadagnoli, V. Lubicz, M. Papinutto and S. Simula, *First Lattice QCD Study of the $\Sigma^- \rightarrow n$ Axial and Vector Form Factors with $SU(3)$ Breaking Corrections*, *Nucl. Phys. B* **761** (2007) 63 [hep-ph/0606181].
- [17] V. Mateu and A. Pich, *V_{us} determination from hyperon semileptonic decays*, *JHEP* **0510** (2005) 041 [hep-ph/0509045].
- [18] R. Flores-Mendieta, *V_{us} from hyperon semileptonic decays*, *Phys. Rev. D* **70** (2004) 114036 [hep-ph/0410171].
- [19] R. Flores-Mendieta, E. E. Jenkins and A. V. Manohar, *$SU(3)$ symmetry breaking in hyperon semileptonic decays*, *Phys. Rev. D* **58**, 094028 (1998) [hep-ph/9805416].
- [20] H. -Ch. Kim, M. Praszalowicz and K. Goeke, *Hyperon semileptonic decays and quark spin content of the proton*, *Phys. Rev. D* **61**, 114006 (2000) [hep-ph/9910282].
- [21] H. -Ch. Kim, M. V. Polyakov, M. Praszalowicz and K. Goeke, *Semileptonic decay constants of octet baryons in the chiral quark-soliton model*, *Phys. Rev. D* **57**, 299 (1998) [hep-ph/9709221].
- [22] T. Ledwig, A. Silva, H. -Ch. Kim and K. Goeke, *Semileptonic hyperon decays in the self-consistent $SU(3)$ chiral quark-soliton model*, *JHEP* **0807**, 132 (2008) [arXiv:0806.4072 [hep-ph]].
- [23] G. -S. Yang, H. -Ch. Kim and M. V. Polyakov, *Electromagnetic mass differences of $SU(3)$ baryons within a chiral soliton model*, *Phys. Lett. B* **695**, 214 (2011) [arXiv:1009.5250 [hep-ph]].
- [24] G. -S. Yang and H. -Ch. Kim, *Mass splittings of $SU(3)$ baryons within a chiral soliton model*, *Prog. Theor. Phys.* **128**, 397 (2012) [arXiv:1010.3792 [hep-ph]].
- [25] G. -S. Yang and H. -Ch. Kim, *Mass splittings of the baryon decuplet and antidecuplet with the second-order flavor symmetry breakings within a chiral soliton model*, *J. Korean Phys. Soc.* **61**, 1956 (2012) [arXiv:1102.1786 [hep-ph]].
- [26] M. Ademollo and R. Gatto, *Nonrenormalization Theorem for the Strangeness Violating Vector Currents*, *Phys. Rev. Lett.* **13**, 264 (1964).
- [27] C. H. Llewellyn Smith, *Neutrino Reactions At Accelerator Energies*, *Phys. Rept.* **3**, 261 (1972).
- [28] T. R. Hemmert, B. R. Holstein and N. C. Mukhopadhyay, *NN , $N\Delta$ couplings and the quark model*, *Phys. Rev. D* **51**, 158 (1995) [arXiv:hep-ph/9409323].
- [29] B. Golli, S. Sirca, L. Amoreira and M. Fiolhais, *Axial amplitudes for Delta excitation in chiral quark models*, *Phys. Lett. B* **553**, 51 (2003) [arXiv:hep-ph/0210014].
- [30] C. Alexandrou, T. Leontiou, J. W. Negele and A. Tsapalis, *The axial $N \rightarrow \Delta$ transition form factors from lattice QCD*, *Phys. Rev. Lett.* **98**, 052003 (2007) [arXiv:hep-lat/0607030].
- [31] E. Witten, *Current Algebra, Baryons, and Quark Confinement*, *Nucl. Phys. B* **223** (1983) 433.

- [32] E. Guadagnini, *Baryons as Solitons and Mass Formulae*, Nucl. Phys. B **236** (1984) 35.
- [33] S. Jain and S. R. Wadia, *Large N Baryons: Collective Coordinates of the Topological Soliton in $SU(3)$ Chiral Model*, Nucl. Phys. B **258** (1985) 713.
- [34] A. Blotz, D. Diakonov, K. Goeke, N. W. Park, V. Petrov and P. V. Pobylitsa, *The $SU(3)$ Nambu-Jona-Lasinio soliton in the collective quantization formulation*, Nucl. Phys. A **555** (1993) 765.
- [35] C. V. Christov, A. Blotz, H.-Ch. Kim, P. Pobylitsa, T. Watabe, T. Meissner, E. Ruiz Arriola and K. Goeke, *Baryons as nontopological chiral solitons*, Prog. Part. Nucl. Phys. **37** (1996) 91 [hep-ph/9604441].
- [36] H. -Ch. Kim, M. Praszalowicz and K. Goeke, *Magnetic moments of the $SU(3)$ decuplet baryons in the chiral quark-soliton model*, Phys. Rev. D **57**, 2859 (1998) [hep-ph/9706531].
- [37] G. -S. Yang, H. -Ch. Kim, M. Praszalowicz and K. Goeke, *Octet, decuplet and antidecuplet magnetic moments in the chiral quark soliton model revisited*, Phys. Rev. D **70**, 114002 (2004) [hep-ph/0410042].
- [38] H. -Ch. Kim, M. Praszalowicz, M. V. Polyakov and K. Goeke, *Flavor structure of the octet magnetic moments*, Phys. Rev. D **58**, 114027 (1998) [hep-ph/9801295].
- [39] J. Dworkin, P. T. Cox, E. C. Dukes, O. E. Overseth, R. Handler, R. Grobel, B. Lundberg and L. Pondrom *et al.*, *High statistics measurement of g_A/g_V in $\Lambda \rightarrow p + e^- + \bar{\nu}$* , Phys. Rev. D **41**, 780 (1990).
- [40] M. Bourquin *et al.* [Bristol-Geneva-Heidelberg-Orsay-Rutherford-Strasbourg Collaboration], *Measurements of Hyperon Semileptonic Decays at the CERN Super Proton Synchrotron. I. The $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$ Decay Mode*, Z. Phys. C **12** (1982) 307.
- [41] J. Wise, D. A. Jensen, M. N. Kreisler, F. Lomanno, R. Poster, M. S. Z. Rabin, K. Raychaudhuri and M. Way *et al.*, *Precise Measurement of γ ($\Lambda^0 \rightarrow pe^- \bar{\nu}$) / γ ($\Lambda^0 \rightarrow p\pi^-$)*, Phys. Lett. B **91**, 165 (1980).
- [42] J. Wise, D. A. Jensen, M. N. Kreisler, F. Lomanno, R. Poster, M. S. Z. Rabin, K. Raychaudhuri and M. Way *et al.*, *Precise Measurement of the Ratio of the Axial Vector Coupling to Vector Coupling in $\Lambda^0 \rightarrow pe^- \bar{\nu}$* , Phys. Lett. B **98**, 123 (1981) [Erratum-ibid. B **100**, 519 (1981)].
- [43] S. D. Bass, *The Spin structure of the proton*, Rev. Mod. Phys. **77**, 1257 (2005) [hep-ph/0411005].
- [44] Y. Goto *et al.* [Asymmetry Analysis Collaboration], *Polarized parton distribution functions in the nucleon*, Phys. Rev. D **62**, 034017 (2000) [hep-ph/0001046].
- [45] F. E. Close and R. G. Roberts, *Consistent analysis of the spin content of the nucleon*, Phys. Lett. B **316**, 165 (1993) [hep-ph/9306289].
- [46] A. Psaker, W. Melnitchouk and A. V. Radyushkin, *Weak Deeply Virtual Compton Scattering*, Phys. Rev. D **75** (2007) 054001 [hep-ph/0612269].
- [47] S. M. Berman and M. J. G. Veltman, *Baryon Resonance Production By Neutrinos*, Nuovo Cim. **38** (1965) 993.

- [48] C. H. Albright and L. S. Liu, *Weak N^* Production and $SU(6)$* , Phys. Rev. **140** (1965) B748.
- [49] S. D. Bass and A. W. Thomas, *The Nucleon's octet axial-charge $g_A^{(8)}$ with chiral corrections*, Phys. Lett. B **684**, 216 (2010) [arXiv:0912.1765 [hep-ph]].
- [50] J. Lindquist, E. C. Swallow, R. L. Sumner, J. M. Watson, R. Winston, D. M. Wolfe, P. R. Phillips and K. Reibel *et al.*, *Measurement of the Angular Correlation Parameters in the beta Decay of Polarized Lambda Hyperons*, Phys. Rev. D **16**, 2104 (1977).