

Localized States and Dynamics in the Nonlinear Schrödinger / Gross-Pitaevskii Equation

M.I. Weinstein *

Department of Applied Physics and Applied Mathematics,
and Department of Mathematics
Columbia University
New York, NY 10027

July 24, 2021

1 Introduction

Nonlinear dispersive waves are wave phenomena resulting from the interacting effects of nonlinearity and dispersion. *Dispersion* refers to the property that waves of different wavelengths travel at different velocities. This property may, for example, be due to the material properties of the medium, *e.g.* chromatic dispersion [1], or to the geometric arrangement of material constituents, *e.g.* a periodic medium with Floquet-Bloch *band dispersion* [3]. Nonlinearity distorts the shape of a localized structure or creates amplitude dependent non-uniformities in phase. This may concentrate or localize energy in a region of space (attractive / focusing nonlinearity) or tend to expel energy from compact sets (repulsive / defocusing nonlinearity). In electromagnetics, nonlinearities may arise due to the intensity dependence of dielectric parameters (*e.g.* Kerr effect [1]). Physical phenomena in which the effects of both dispersion and nonlinearity play a role are ubiquitous. Some examples are: (a) long waves of small amplitude at a water-air interface [112], (b) a nearly mono-chromatic laser beam propagating through air, glass or water [36], (c) light-pulses propagating through optical fiber waveguides [1] and (d) the macroscopic dynamics of weakly correlated quantum particles in a Bose-Einstein condensate; see, for example, [86, 30, 49]. Interest in nonlinear dispersive waves and their interaction with nonhomogeneous media ranges from Fundamental to Applied Science with great promise in engineering / technological applications due to major advances in materials science and micro- and nano-structure fabrication techniques; see, for example, [107].

An important feature of many nonlinear dispersive systems is the prevalence of coherent structures. They range from phenomena we can only passively observe in nature, such as large-scale localized meteorological events, to those we can control, such as the flow of energy in computer

*This work is supported in part by U.S. NSF grant DMS-10-08855 and DMS-14-12560.

chips and other nano-patterned media. In general, a coherent structure is one which has a spatially localized core, *e.g.* a solitary wave concentrated in a compact region of space or a front-like state which transitions between different asymptotic states across a spatially compact transition region.

We focus on a key model in the mathematical theory of nonlinear dispersive waves, the *nonlinear Schrödinger / Gross Pitaevskii* (NLS/GP) equation:

$$i\partial_t\Phi = -\Delta\Phi + V(x)\Phi + g|\Phi|^2\Phi. \quad (1.1)$$

In particular, we consider its *nonlinear bound states*. Here, $\Phi(x, t)$ is a complex-valued function for $x \in \mathbb{R}^d$ and $t \in \mathbb{R}$. $V(x)$ is a real-valued “potential”, and g is a real parameter. The following table lists the physical significance of Φ , V and g in the settings of nonlinear optics and macroscopic quantum physics:

	Nonlinear Optics	Macroscopic Quantum Physics
$\Phi(x, t)$	Electric field complex envelope	Macroscopic quantum wave function
$V(x)$	Refractive index	Magnetic trap
g	Kerr nonlinearity coefficient, $g < 0$	Microscopic 2-body scattering length ($g > 0$ or $g < 0$)

1.1 Outline

Basic mathematical properties of NLS and NLS/GP are discussed in section 2. In section 3 nonlinear bound states and aspects of their stability theory are discussed from variational and bifurcation perspectives. Examples are also presented and the particular cases of $V(x)$ given by a single-well, double-well potential and periodic potential are discussed in detail. In section 4 we discuss particle-like dynamics of solitons interacting with a potential over a large, but finite, time interval. In sections 5 and 6 we consider the very long time behavior of solutions to NLS / GP. In particular, the focus is placed on the important resonant radiation damping mechanism that drives the relaxation of the system to a nonlinear ground state and underlies the phenomena of *Ground State Selection* and *Energy Equipartition*. Sections 5.1-5.2 discuss linear and nonlinear *toy minimal models*, which illustrate these mechanisms. Regarding the overall style of this article, we seek to emphasize the key ideas, and therefore do not present all detailed technical hypotheses, leaving that to the references.

Acknowledgements: The author would like to thank J. Marzuola and E. Shlizerman for stimulating discussions and helpful comments on this article. He also wishes to thank Bjorn Sandstede and the referees for their careful reading and suggested improvements.

2 NLS and NLS/GP

In this section we review basic properties of NLS/GP:

$$i\partial_t \Phi = -\Delta \Phi + V(x)\Phi + g|\Phi|^{2\sigma}\Phi. \quad (2.1)$$

Here, g is a real parameter and we assume that V is a bounded and smooth real-valued potential. The equation (2.1) is often called the nonlinear Schrödinger / Gross-Pitaevskii (NLS/GP).

We consider solutions to the initial value problem for (2.1) with initial conditions

$$\Phi(x, 0) = \Phi_0(x) \in H^1(\mathbb{R}^d). \quad (2.2)$$

The basic well-posedness theorem states that for $d = 1, 2$ and all $0 \leq \sigma < \infty$ and for $d \geq 3$ and $0 \leq \sigma < 2/(d-2)$, the initial value problem (2.1)-(2.2) has a unique local-in-time solution $\Phi \in C^0([0, T]; H^1(\mathbb{R}^d))$ [64, 100, 6, 101, 37].

NLS / GP is a Hamiltonian system, expressible in the form

$$i\partial_t \Phi = \frac{\delta \mathcal{H}[\Phi, \Phi^*]}{\delta \Phi^*}, \quad (2.3)$$

with Hamiltonian energy functional

$$\mathcal{H}[F, F^*] = \int_{\mathbb{R}^d} \nabla F(x) \cdot \nabla F^*(x) + V(x) F(x) F^*(x) + \frac{g}{\sigma+1} (F(x) F^*(x))^{\sigma+1} dx. \quad (2.4)$$

The solution of the initial value problem satisfies the conservation laws:

$$\mathcal{H}[\Phi(\cdot, t)] = \mathcal{H}[\Phi_0] \quad (2.5)$$

$$\mathcal{N}[\Phi(\cdot, t)] = \mathcal{N}[\Phi_0] \quad (2.6)$$

where $\mathcal{H}[\cdot]$ is defined in (2.4) and

$$\mathcal{N}[F, F^*] = \int_{\mathbb{R}^d} F(x) F^*(x) dx. \quad (2.7)$$

The conserved integrals \mathcal{H} and \mathcal{N} are associated, respectively, with the invariances $t \mapsto t + t_0$, $t_0 \in \mathbb{R}$ (time-translation) and $\Phi \mapsto \Phi e^{i\theta}$, $\theta \in \mathbb{R}$.

For the nonlinear Schrödinger equation (NLS), where $V \equiv 0$, we have the additional symmetries

1. $\Phi(x, t) \mapsto \Phi(x + x_0, t)$, $x_0 \in \mathbb{R}^d$ (Translation invariance)
2. $\Phi(x, t) \mapsto \mathcal{G}_\xi[\Phi](x, t) = e^{i\xi(x-\xi t)} \Phi(x - 2\xi t, t)$, $\xi \in \mathbb{R}^d$ (Galilean invariance)

For NLS with the homogeneous power nonlinearity in (2.1) the equation is also dilation invariant:

$$\Phi(x, t) \mapsto \lambda^{\frac{1}{\sigma}} \Phi(\lambda x, \lambda^2 t) \quad (2.8)$$

The case where $g < 0$ is called the *focusing case* and the case where $g > 0$ is called the *defocusing case*. This terminology refers to the nonlinear term's tendency to focus (localize) energy or defocus (spread) energy. Independently of the nonlinear term, the linear potential may be an attractive potential, which concentrates energy, or a repulsive potential which allows energy to spread.

Concerning global-in-time behavior of solutions, the initial value problem is globally well-posed for any H^1 initial condition if either $g > 0$ or if $g < 0$ and $\sigma < 2/d$ (subcritical case). For $g < 0$ and $\sigma \geq 2/d$ solutions with H^1 initial data may develop singularities in finite time (blow up). See, for example, [64, 100, 6, 101, 108, 37].

3 Bound States - Linear and Nonlinear

3.1 Linear bound states

We start with the linear Schrödinger equation:

$$i\partial_t \Psi = (-\Delta + V(x)) \Psi = H \Psi \quad (3.1)$$

Here V denotes a smooth and real-valued potential satisfying

$$V(x) \rightarrow 0 \text{ sufficiently rapidly as } |x| \rightarrow \infty. \quad (3.2)$$

Bound states are solutions of the form $\Psi = e^{-iEt} \psi(x)$, where ψ satisfies the eigenvalue problem

$$H\psi = (-\Delta + V)\psi = E\psi, \quad \psi \in H^1(\mathbb{R}^d). \quad (3.3)$$

We expect the eigenvalue problem (3.3) to have a non-trivial solutions if V is an attractive “potential well”; intuitively, the set where $V(x) \leq 0$ should be sufficiently wide and deep.

The following result gives a condition on the existence of a ground state, a nontrivial solution of (3.3) of minimal energy, E [15, 76].

Theorem 3.1 *Denote by*

$$E_{0\star} = \inf_{\int_{\mathbb{R}^d} |f|^2 = 1} \langle Hf, f \rangle \equiv \inf_{\int_{\mathbb{R}^d} |f|^2 = 1} \int_{\mathbb{R}^d} |\nabla f(x)|^2 + V(x)|f(x)|^2 dx.$$

If $-\infty < E_{0\star} < 0$ then the infimum is attained at a positive function $\psi_{0\star} \in H^1(\mathbb{R}^d)$, which is a solution of

$$H\psi_{0\star} = E_{0\star}\psi_{0\star}, \quad E_{0\star} = \langle H\psi_{0\star}, \psi_{0\star} \rangle$$

3.2 Nonlinear bound states

Here we consider spatially localized solutions of NLS/GP of the form

$$\Phi(x, t) = e^{-iEt} \psi_E(x), \quad (3.4)$$

where ψ_E satisfies the nonlinear elliptic equation

$$(-\Delta + V)\psi_E + g|\psi_E|^{2\sigma}\psi_E = E\psi_E, \quad \psi_E \in H^1(\mathbb{R}^d) \quad (3.5)$$

Solutions of the nonlinear eigenvalue problem (3.5) are often called *nonlinear bound states* or solitary waves. A positive and decaying solution is often called a nonlinear ground state. In the case where $V(x)$ is non-trivial, such states are often called nonlinear defect modes. In this case, the spatially varying potential, $V(x)$, is viewed as a localized defect in a homogeneous (translation-invariant) background.

The question of existence of nonlinear bound states of NLS / GP has been studied extensively by variational and bifurcation methods. See, for example, [99, 5, 108, 77, 88] as well as the discussion in sections 3.6 [72] and 3.7 [59].

The following result, which we shall refer to later in this article, concerns bifurcation of a nonlinear ground state from the ground eigenstate of the linear operator $-\Delta + V$; see, for example, [88, 85]

Theorem 3.2 *Assume V is real-valued, smooth and rapidly decreasing at infinity and that $-\Delta + V$ has at least one discrete eigenvalue. Let $(\psi_{0\star}, E_{0\star})$ denote the ground state eigenpair given by Theorem 3.1; here, $\psi_{0\star}$ may be chosen to be positive and normalized in L^2 . Consider the equation for nonlinear bound states of NLS / GP with $g = -1$ (attractive nonlinear potential):*

$$(-\Delta + V(x) - |\psi_E|^{2\sigma})\psi_E = E\psi_E, \quad E < 0, \quad \sigma \geq 1. \quad (3.6)$$

Then, there is a constant $\delta_0 > 0$ and a non-empty interval $\mathcal{I} = [E_{0\star} - \delta_0, E_{0\star})$ such that for any $E \in \mathcal{I}$, equation (3.6) has a positive solution, which for E tending to $E_{0\star}$ has the expansion:

$$\psi_E(x) = \rho(E) \left(\psi_{0\star}(x) + \mathcal{O}(\rho(E)^{2\sigma}) \right), \quad \rho(E) = |E_{0\star} - E|^{\frac{1}{2}} \left(\int \psi_{0\star}^{2\sigma+2} \right)^{-\frac{1}{2\sigma}} \quad (3.7)$$

3.3 Orbital stability of nonlinear bound states

In this section we discuss the *orbital Lyapunov stability* of nonlinear ground states. In particular, we prove nonlinear stability in $H^1(\mathbb{R}^n)$ modulo the natural group of symmetries. We shall in this section consider the case where $g = -1$, the case of focusing nonlinearity. Given a positive H^1 solution of (3.6) introduce its *orbit*:

$$\begin{aligned} \mathcal{O}_{\text{gs}} &= \{ \Psi_E(x) e^{i\gamma} : \gamma \in [0, 2\pi) \}, \quad V \neq 0 \\ \mathcal{O}_{\text{gs}} &= \{ \Psi_E(x + x_0) e^{i\gamma} : \gamma \in [0, 2\pi), x_0 \in \mathbb{R}^n \}, \quad V \equiv 0. \end{aligned} \quad (3.8)$$

We say that the ground state is *orbitally stable* in $H^1(\mathbb{R}^n)$ if the following holds. If $\Phi(x, t = 0)$ is close to element of \mathcal{O}_{gs} in H^1 then for all $t \neq 0$, $\Phi(x, t)$ is H^1 close to some (typically t dependent)

element of \mathcal{O}_{gs} . In order to make this precise, we introduce a metric which measures the distance from an arbitrary H^1 function to the ground state orbit:

$$\begin{aligned} \text{dist}(u, \mathcal{O}_{\text{gs}}) &= \inf_{\gamma} \|u - \Psi_0 e^{i\gamma}\|_{H^1}, \quad V \neq 0 \\ \text{dist}(u, \mathcal{O}_{\text{gs}}) &= \inf_{\gamma, x_0} \|u(\cdot + x_0) - \Psi_0 e^{i\gamma}\|_{H^1}, \quad V \equiv 0. \end{aligned} \quad (3.9)$$

A more precise statement of H^1 -Lyapunov stability is as follows. For any $\epsilon > 0$ there is a $\delta > 0$ such that if

$$\text{dist}(\Phi(\cdot, 0), \mathcal{O}_{\text{gs}}) < \delta \quad (3.10)$$

then for all $t \neq 0$

$$\text{dist}(\Phi(\cdot, t), \mathcal{O}_{\text{gs}}) < \epsilon.$$

Nonlinear bound states are critical points of the Hamiltonian \mathcal{H} subject to fixed L^2 norm, \mathcal{N} . In particular, a nonlinear bound state, ψ_E , of frequency E satisfies $\delta \mathcal{E}_E[f, f^*]/\delta f^* = 0$, where

$$\begin{aligned} \mathcal{E}_E[f] &= \mathcal{H}[f] - E \int |f|^2 \\ \mathcal{H}[f] &\equiv \int |\nabla f|^2 + V(x)|f|^2 - \frac{1}{\sigma+1} |f|^{2\sigma+2} \end{aligned}$$

For subcritical nonlinearities, $\sigma < 2/d$, stable nonlinear ground states may be realized as constrained global minimizers of the variational problem:

$$\inf_{f \in H^1} \mathcal{H}[f] \quad \text{subject to fixed } \mathcal{N}[f] > 0. \quad (3.11)$$

Orbital H^1 stability is a consequence of the compactness properties of arbitrary minimizing sequences [12, 77].

However in general, depending on the details of the nonlinear terms, stable solitary waves may arise as *local* minimizers of \mathcal{H} subject to fixed \mathcal{N} . We now discuss stability, in this more general setting. Introduce the linearized operators L_{\pm} , which are real and imaginary parts of the second variation of the energy \mathcal{E}_E :

$$L_+ \equiv -\Delta + V(x) - (2\sigma + 1)\psi_E^{2\sigma} - E. \quad (3.12)$$

$$L_- \equiv -\Delta + V(x) - \psi_E^{2\sigma} - E \quad (3.13)$$

We define the *index* of ψ_E by

$$\text{index}(\psi_E) = \text{number of strictly negative eigenvalues of } L_+ . \quad (3.14)$$

Theorem 3.3 (H^1 Orbital Stability) *1. Assume the conditions*

(S1) Spectral condition: $\psi_E > 0$ and $\text{index}(\psi_E) = 1$

(S2) Slope condition

$$\frac{d}{dE}\mathcal{N}[\psi_E] \equiv \frac{d}{dE} \int |\psi_E(x)|^2 dx < 0 \quad (3.15)$$

Then, ψ_E is H^1 orbitally stable.

2. Assume $\psi_E > 0$. If $\text{index}(\psi_E) \geq 2$ or $\frac{d}{dE}\mathcal{N}[\psi_E] > 0$ then ψ_E is linearly exponentially unstable. That is, the linearized evolution equation (see (3.19) below) has a spatially localized solution which grows exponentially with t .

See [109, 110, 51, 62, 50, 91]. In [106] it was shown that $\partial_E\mathcal{N}[\psi_E] > 0$ implies the existence of an exponentially growing mode of the linearized evolution equation.

We give the idea of the proof of stability. For simplicity, suppose $V(x)$ is non-trivial, In this case the ground state orbit consists of all phase translates of ψ_E ; see the first line of (3.8). Let ϵ be an arbitrary positive number. We have for $t \neq 0$, by choosing δ in (3.10) sufficiently small

$$\begin{aligned} \epsilon^2 &\sim \mathcal{E}_E[\Psi(\cdot, 0)] - \mathcal{E}_E[\psi_E] \\ &= \mathcal{E}_E[\Psi(\cdot, t)] - \mathcal{E}_E[\psi_E], \quad \text{by conservation laws,} \\ &= \mathcal{E}_E[\Psi(\cdot, t)e^{i\gamma}] - \mathcal{E}_E[\psi_E], \quad \text{by phase invariance} \\ &= \mathcal{E}_E[\psi_E + u(\cdot, t) + iv(\cdot, t)] - \mathcal{E}_E[\psi_E] \quad (\text{definition of the perturbation } u + iv, \ u, v \in \mathbb{R}) \\ &\sim \langle L_+ u(t), u(t) \rangle + \langle L_- v(t), v(t) \rangle \quad (\text{by Taylor expansion and } \delta\mathcal{E}_E[\psi_E] = 0) \end{aligned} \quad (3.16)$$

If L_+ and L_- were positive definite operators, implying the existence of positive constants C_+ and C_- such that

$$\langle L_+ u, u \rangle \geq C_+ \|u\|_{H^1}^2, \quad (3.17)$$

$$\langle L_- v, v \rangle \geq C_- \|v\|_{H^1}^2 \quad (3.18)$$

for all $u, v \in H^1$, then it would follow from (3.16) that the perturbation about the ground state, $u(x, t) + iv(x, t)$, would remain of order ϵ in H^1 for all time $t \neq 0$. The situation is however considerably more complicated. The relevant facts to note are as follows.

- (1) $L_- \psi_E = 0$, with $\psi_E > 0$. Hence, ψ_E is the ground state of L_- , $0 \in \sigma(L_-)$, and L_- is a non-negative with continuous spectrum $[|E|, \infty)$.
- (2) For small L^2 nonlinear ground states, L_+ has exactly one strictly negative eigenvalue and continuous spectrum $[|E|, \infty)$.

The zero eigenvalue of L_- and the negative eigenvalue of L_+ constitute two *bad* directions, which are treated as follows, noting that $u(\cdot, t)$ and $v(\cdot, t)$ are not arbitrary H^1 functions but are rather constrained by the dynamics of NLS.

To control L_- , we choose $\gamma(t)$ so as to minimize the distance of the solution to the ground state orbit, (3.9). This yields the codimension one constraint on v : $\langle v(\cdot, t), \psi_E \rangle = 0$, subject to which (3.18) holds with $C_- > 0$.

To control L_+ , we observe that since the L^2 norm is invariant for solutions, we have the codimension constraint on u : $\langle u(\cdot, t), \psi_E \rangle = 0$. Although ψ_E is not the ground state of L_+ , it can be shown by constrained variational analysis that if the slope condition (3.15) holds, then constraint on u places u in the positive cone of L_+ , *i.e.* $C_+ > 0$ in (3.17). Thus, positivity (coercivity) estimates (3.17) and (3.18) hold and \mathcal{E}_E serves as a Lyapunov functional which controls the distance of the solution to the ground state orbit. The detailed argument is presented in [110].

We also note that the role played by the second variation of \mathcal{E} in the linearized time-dynamics [109]. Let $\Psi(t) = (\psi_E + u + iv)e^{-iEt}$. The linearized perturbation, $(u(t), v(t))^t$, satisfies the linear Hamiltonian system:

$$\partial_t \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & L_- \\ -L_+ & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}, \quad (3.19)$$

with conserved (time-invariant) energy

$$\mathcal{Q}(u, v) \equiv \langle L_+ u, u \rangle + \langle L_- v, v \rangle \quad (3.20)$$

The above constrained variational analysis underlying nonlinear orbital theory corresponds to the H^1 boundedness of the linearized flow (3.19), restricted on a finite codimension subspace. This subspace is expressible in terms of symplectic orthogonality to first order generators of symmetries acting on the ground state. This structure is used centrally in many works on nonlinear asymptotic stability and nonlinear scattering theory of solitary waves; see section 6 and references cited.

In the following subsections, we give several examples: a) the free NLS soliton, b) the nonlinear defect mode of a simple potential well, c) nonlinear bound states and symmetry breaking for the double well, and d) *gap solitons* of NLS / GP with a periodic potential.

3.4 The free soliton of focusing NLS: $V \equiv 0$ and $g = -1$

In this case, there is a unique (up to spatial translation) positive and symmetric solution, a ground state, which is smooth and decays exponentially as $|x| \rightarrow \infty$. For results on existence, symmetry and uniqueness of the ground state see, for example, [99, 5, 44, 74].

For example, the focusing one-dimensional cubic ($\sigma = 1$) nonlinear Schrödinger equation

$$i\partial_t \psi = -\partial_x^2 \psi - |\psi|^2 \psi, \quad (3.21)$$

has the solitary standing wave:

$$\psi_{sol}(x, t) = e^{it} \sqrt{2} \operatorname{sech}(x), \quad (3.22)$$

By the symmetries of NLS (section 2), we have the extended family of solitary traveling waves of arbitrary negative frequency $E = -\lambda^2$, $\lambda > 0$:

$$\mathcal{G}_{\lambda, x_0, v, \theta} [\psi_{sol}] (x, t) = \lambda e^{i\lambda^2 t} \sqrt{2} \operatorname{sech}(\lambda (x - x_0 - 2vt)) e^{iv(x - x_0 - vt)} e^{i\theta}; \quad v, \theta, x_0 \in \mathbb{R}. \quad (3.23)$$

Theorem 3.3 implies that the positive solitary standing wave of translation invariant NLS is orbitally stable if $\sigma < 2/d$ and is unstable if $\sigma \geq 2/d$ [108, 110, 51]. The solid curve in Figure 3.5 shows the family of solitary waves, (3.23), of (3.21) bifurcating from the zero solution at the edge-energy of the continuous spectrum of $-\partial_x^2$.

3.5 $V(x)$, a simple potential well; model of a pinned nonlinear defect mode

In [88] the bifurcation of nonlinear bound states of NLS/GP from localized eigenstates of the linear Schrödinger operator $-\Delta + V$ was studied. The simplest case of such a *nonlinear defect mode* is for NLS / GP with $V(x)$ is taken to be a Dirac delta function potential well:

$$i\partial_t \psi = H\psi - |\psi|^2 \psi, \quad H = -\partial_x^2 - \gamma \delta(x), \quad \gamma > 0. \quad (3.24)$$

For the well-posedness theory of the initial-value-problem for (3.24) in $C^0([0, T]; H^1(\mathbb{R}))$ see [60] and, more specifically, section 8D of [27]. In this case, the nonlinear defect mode of frequency $E = -\lambda^2 < -(\gamma/2)^2$ is explicitly given by:

$$\psi_{sol}(x, t; \gamma) = \lambda e^{i\lambda^2 t} \sqrt{2} \operatorname{sech}\left(\lambda|x| + \tanh^{-1} \frac{\gamma}{\lambda}\right) e^{i\theta}$$

This family of nonlinear bound states, which is pinned to the “defect” at $x = 0$, bifurcates from the zero solution at $E = E_\star = -(\gamma/2)^2$, the unique negative eigenvalue of H ; see the dashed curve in Figure 3.5.




Figure 1: Squared L^2 norm, \mathcal{N} , vs. frequency for the free soliton (solid) and pinned soliton (dashed).

3.6 NLS / GP: Double well potential with separation, L

Consider now the NLS / GP with a double-well potential, $V_L(x)$:

$$i\partial_t\psi = -\Delta\psi - V_L(x)\psi - |\psi|^2\psi .$$

We take the double-well, $V_L(x)$ to be constructed by centering two identical single-bound state potentials, $V_0(x)$, of the sort considered in the example of section 3.5, at the positions $x = \pm L$:

$$V_L(x) = V_0(x + L) + V_0(x - L) \quad (3.25)$$

A sketch of a one-dimensional double well potential, $V_L(x)$, and the associated spectrum of $H = -\partial_x^2 + V_L(x)$ is displayed in the right panels of Figure 2.




Figure 2: Left: Single-well potential and its spectrum: one discrete eigenvalue, marked by “o”, and continuous spectrum, \mathbb{R}_+ . Right: Double-well potential and its spectrum: two nearby discrete eigenvalues (marked “*” and “x”) and continuous spectrum, \mathbb{R}_+

As in the example of section 3.5 (see [88]) there are branches of nonlinear bound states which bifurcate from the zero solution at the discrete eigenvalue energies. We focus on the branch emanating from the zero solution at the ground state eigenvalue of H_L . For small squared L^2 norm, \mathcal{N} , this is the unique (up to the symmetry $\psi \mapsto e^{i\theta}\psi$) nontrivial solution branch. This solution has the same symmetries as the ground state of the linear double well potential [52]. That is, for \mathcal{N} small, $\psi_{E(\mathcal{N})}$ is bi-modal with peaks centered at $x \pm L$. Increasing the L^2 norm (or its square, \mathcal{N}) we find that there is a critical value $\mathcal{N}_{\text{cr}} > 0$, such that for $\mathcal{N} > \mathcal{N}_{\text{cr}}$ there are multiple nonlinear bound states; see Figure 3. In particular, at $(E(\mathcal{N}_{\text{cr}}), \psi_{\mathcal{N}_{\text{cr}}})$ there is a symmetry breaking

bifurcation. Specifically, for $\mathcal{N} > \mathcal{N}_{\text{cr}}$, there are three families of solutions: the continuation of the symmetric branch (dashed curve continuation of the symmetric branch) and two branches of asymmetric states, corresponding to solutions whose mass is concentrated on the left or right side wells. The bifurcation diagram in Figure 3 show only two branches beyond the bifurcation point. The solid leftward branching curve represents both asymmetric branches, one set of states being obtained from the other via a reflection about $x = 0$.

Figure 3 also encodes stability properties. For $0 < \mathcal{N} < \mathcal{N}_{\text{cr}}$, the (symmetric) ground state is stable, while for $\mathcal{N} > \mathcal{N}_{\text{cr}}$ the symmetric state is unstable. at $\mathcal{N} = \mathcal{N}_{\text{cr}}$ stability is transferred to the asymmetric branches (solid curve for $\mathcal{N} > \mathcal{N}_{\text{cr}}$).




Figure 3: Bifurcations from the two discrete eigenvalues of the double-well: Bifurcation curve emanating from the zero state at the ground state (lowest) energy shows symmetry breaking at some positive \mathcal{N}_{cr} . Bifurcation curve emanating from the zero state at the excited state energy shows no symmetry breaking.

The above discussion is summarized in the following [65, 72]:

Theorem 3.4 (Symmetry Breaking Bifurcation) *Consider the nonlinear eigenvalue problem*

$$(-\Delta + V_L)\psi_E + g|\psi_E|^2\psi_E = E\psi_E, \quad \psi_E \in H^1(\mathbb{R}^d) \quad (3.26)$$

where $g < 0$ and V_L denotes a double well potential with separation parameter, L , as in (3.25). Denote the ground and excited state eigenvalues of $-\Delta + V_L$ by:

$$E_{0\star}(L) < E_{1\star}(L) < 0. \quad (3.27)$$

Let $\mathcal{N}[f] = \int_{\mathbb{R}^d} |f|^2$.

There exists a positive constant, L_0 , such that the following holds. For all $L \geq L_0$, there exists $\mathcal{N}_{\text{cr}} = \mathcal{N}_{\text{cr}}(L) > 0$ such that

- (a) For any $\mathcal{N} < \mathcal{N}_{\text{cr}}$ there is a unique non-trivial symmetric state.

- (b) The point $(E, \psi_E) = (E_{cr}, \psi_{E_{cr}})$ is a bifurcation point i.e. there are, for $\mathcal{N} > \mathcal{N}_{cr}$, two bifurcating branches asymmetric states, consisting respectively of states concentrated about $x = \pm L$.
- (c) Concerning the stability of these branches:
- (c1) $\mathcal{N} < \mathcal{N}_{cr}$: The symmetric branch is orbitally stable.
- (c2) $\mathcal{N} > \mathcal{N}_{cr}$: The symmetric branch is linearly exponentially unstable; these states have index = 2; see (3.14).
- (c3) $\mathcal{N} > \mathcal{N}_{cr}$: The asymmetric branch is orbitally stable.
- (d) Symmetry breaking threshold:

$$\mathcal{N}_{cr}(L) \sim E_{1\star}(L) - E_{0\star}(L) \quad (3.28)$$

(The gap, $E_{1\star}(L) - E_{0\star}(L)$ is exponentially small for large L [52, 60].)




Figure 4: Symmetric, asymmetric and anti-symmetric modes arising in the bifurcation diagram of Figure 3.

Theorem 3.4 is proved by methods of bifurcation theory [45, 82]. Specifically, for L sufficiently large we can use a Lyapunov-Schmidt reduction strategy to reduce the nonlinear eigenvalue problem to a weakly perturbed system of two nonlinear algebraic equations depending on \mathcal{N} . The unknowns of this system are essentially the projections of the sought nonlinear bound state onto the linear ground and excited states of H_L . The bifurcation structure of this pair of nonlinear algebraic equations is of the type displayed in Figure 3. This bifurcation structure is then shown to persist under the (infinite dimensional) perturbing terms to this finite algebraic reduction. The latter are controlled via PDE estimates and an implicit function theorem argument.

An illustrative study of the *global bifurcation structure* for the exactly solvable case where $V_L(x)$ is taken to be a sum of two attractive Dirac-delta wells centered at $x = \pm L$ is presented

in [60]. In [63] the local bifurcation methods described above are used to study the NLS / GP with a triple well potential. A recent study of global properties of the bifurcating branches and their stability properties for general nonlinearities, $|\psi|^{2\sigma}\psi$, with $\sigma \geq 1/2$ is presented in [65]. In this work the authors prove, for the case of symmetric double-well potentials, that as L varies a symmetry breaking bifurcation occurs once the potential develops a local maximum at the origin. The analysis in [65] allows for the bifurcation to occur at large \mathcal{N}_{cr} , outside the weakly nonlinear regime where the analysis in [72] applies.

Local bifurcation methods can be applied as well to study excited state branches. An example is the branch emanating from the linear excited state energy, along which it can be shown, for small \mathcal{N} , that there are no secondary bifurcations; see Figure 3 and the corresponding mode curve which changes sign in Figure 4.

An approach which complements the local bifurcation approach is variational. For example, in [2] symmetry breaking in a three-dimensional nonlinear Hartree model having an attractive non-local potential is proved for \mathcal{N} sufficiently large. Their arguments are straightforwardly adaptable to the variational formulation (3.11) and yield that for $\sigma < 2/d$ the ground state is asymmetric. The proof is based on a trial function argument and the intuition that for \mathcal{N} sufficiently large it is energetically preferable to concentrate mass in the left well or in the right well but not both.

3.7 NLS / GP: $V(x)$ periodic and the bifurcations from the spectral band edge

We now consider the nonlinear bound states of NLS / GP for the case where V is periodic on \mathbb{R}^d . In this case, the spectrum of $H = -\Delta + V(x)$ is absolutely continuous. Let ψ_μ denote a nonlinear bound state of frequency μ . Any bifurcation of nontrivial solutions of (3.5) from the zero solution must occur at an energy in the continuous spectrum. Indeed, the example of the free soliton of section 3.4 is an example; $V \equiv 0$ is periodic (!) and has continuous spectrum equal to the half-line $\mathbb{R}_+ = [0, \infty)$. The soliton bifurcates from the zero-amplitude solution (as measured say in L^∞ or L^2) from the continuous spectral edge (solid curve in Figure 3). Indeed, for NLS with $V \equiv 0$ and power nonlinearity $-|\psi|^{2\sigma}\psi$

$$\psi_\mu(x) = (-\mu)^{\frac{1}{2\sigma}} \psi_{-1}(\sqrt{-\mu}x) \quad (3.29)$$

and therefore

$$\mathcal{N}[\psi_\mu] = \|\psi_\mu\|_{L^2}^2 = (-\mu)^{\frac{1}{\sigma} - \frac{d}{2}} \|\psi_{-1}\|_{L^2}^2. \quad (3.30)$$

For $V(x)$ a general periodic potential the spectrum of $H = -\Delta + V$ is the union of spectral bands obtained as follows. Consider the d - parameter family of periodic elliptic eigenvalue problems:

$$\begin{aligned} \left(-(\nabla + ik)^2 + V(x) \right) u(x) &= \mu u(x) \\ u(x) &\text{ is periodic with the periodicity of } V(x), \end{aligned} \quad (3.31)$$

where k varies over a fundamental dual lattice cell, the Brillouin zone, $\mathcal{B} \subset \mathbb{R}^d$. For each fixed $k \in \mathcal{B}$, the spectrum of (3.31) is discrete and consists of eigenvalues denoted:

$$\mu_1(k) \leq \mu_2(k) \leq \cdots \leq \mu_b(k) \leq \cdots, \quad (3.32)$$

listed with multiplicities and tending to positive infinity. The corresponding eigenfunctions are denoted, $p_j(x; k)$. The states $\{e^{ikx} p_j(x; k)\}$ where $j \geq 1$ and $k \in \mathcal{B}$ are complete in $L^2(\mathbb{R}^d)$.

It can be shown that nonlinear bound states bifurcate from edges of the spectral bands: to the left, if the nonlinearity is attractive ($g < 0$) and to the right if the nonlinearity is repulsive ($g > 0$) [59, 84, 89]. We expect similar scaling of the L^2 norm of such more general edge-bifurcations:

$$\mathcal{N}[\psi_\mu] = \|\psi_\mu\|_{L^2}^2 \sim |\mu - \mu_\star|^{\frac{1}{\sigma} - \frac{d}{2}} \quad (3.33)$$

where $|\mu - \mu_\star|$ is the distance to the spectral band edge located at μ_\star . Now the details of the bifurcation at the edge depend on the periodic structure. We first give a heuristic picture and then state a precise theorem.

For $|\mu - \mu_\star|$ small, that is for μ near a spectral band edge, the nonlinear bound state, $\psi_\mu(x)$, should be exponentially localized with decay $\psi_\mu(x) \sim \exp(-|\mu - \mu_\star|^{\frac{1}{2}}|x|)$ for $|x| \rightarrow \infty$. Due to the separation of length scales: $|\mu - \mu_\star|^{-1} \gg \text{period of } V$, we expect $\psi_\mu(x)$ should oscillate like the Floquet-Bloch mode $p_1(x; 0) = p_\star(x)$, associated with the band edge energy, μ_\star , with a slowly varying and spatially localized amplitude, $F(x)$, :

$$\psi \sim \delta^{\frac{1}{\sigma}} F(\delta x) \times p_\star(x), \quad \delta = |\mu - \mu_\star|^{\frac{1}{2}}, \quad (3.34)$$

where $F(x)$ satisfies an effective medium (homogenized / constant coefficient) NLS equation.

To simplify the discussion we will focus on bifurcation from the lowest energy (bottom) of the continuous spectrum, $\mu_\star = \mu_1(0)$, for the case of an attractive nonlinearity ($g < 0$). Before stating a precise result, we introduce the inverse effective mass matrix associated with the bottom of the continuous spectrum defined by

$$\begin{aligned} A_{\text{eff}} \equiv \frac{1}{2} D^2 \mu_1(k=0) &= \frac{1}{2} \left(\frac{\partial^2 \mu_1(k=0)}{\partial k_i \partial k_j} \right)_{1 \leq i, j \leq d} \\ &= \delta_{ij} - \frac{4 \langle \partial_{x_j} p_\star, (-\Delta + V - \mu_\star)^{-1} \partial_{x_i} p_\star \rangle}{\langle p_\star, p_\star \rangle} \end{aligned} \quad (3.35)$$

and Γ_{eff} , the effective nonlinearity coefficient defined by

$$\Gamma_{\text{eff}} = \frac{\int p_\star(x)^{2\sigma+2} dx}{\int p_\star(x)^2 dx} > 0. \quad (3.36)$$

The matrix A_{eff} , is clearly symmetric and, for the lowest band edge, it is positive definite [73].

The following result, proved in [59], describes the bifurcation from the bottom of continuous spectrum (left end point of the first spectral band). See also [84, 98, 89]. A proof of the case of a general band edge is discussed in [59].

Theorem 3.5 (Edge bifurcations of nonlinear bound states for periodic potentials) *Let $V(x)$ denote a smooth and even periodic potential in dimension $d = 1, 2$ or $d = 3$. Let x_0 denote any local minimum or maximum of $V(x)$. Denote by $\mu_\star = \inf \sigma(-\Delta + V)$. Let $F_{A,\Gamma}(y)$ the unique, centered at $y = 0$, positive $H^1(\mathbb{R}^d)$ solution (ground state) of the effective / homogenized nonlinear Schrödinger equation with inverse effective mass matrix, A , and effective nonlinearity, Γ .*

$$-\sum_{i,j=1}^d \frac{\partial}{\partial y_i} A_{\text{eff},ij} \frac{\partial}{\partial y_j} F(y) - \Gamma_{\text{eff}} F^{2\sigma+1}(y) = -F(y). \quad (3.37)$$

Then, there exists $\delta_0 > 0$ such that for all $\mu \in (\mu_\star - \delta_0, \mu_\star)$ there is a family of nonlinear bound states $\mu \mapsto \psi_\mu(x)$

$$\psi_\mu(x) - (\mu_\star - \mu)^{\frac{1}{2\sigma}} F_{A_{\text{eff}}, \Gamma_{\text{eff}}} \left(\sqrt{\mu_\star - \mu} (x - x_0) \right) p_\star(x) \rightarrow 0, \text{ as } \mu \uparrow \mu_\star \text{ in } H^1(\mathbb{R}^d). \quad (3.38)$$

The proof of Theorem 3.5 [59], like the study of bifurcations from discrete eigenvalues, proceeds via Lyapunov-Schmidt reduction and application of the implicit function theorem. However, while the analysis of bifurcation from discrete spectrum leads to a finite dimensional (nonlinear algebraic) bifurcation equation, bifurcation from the continuous spectrum leads to an infinite dimensional bifurcation equation, a nonlinear homogenized partial differential equation (3.37); see also [25, 26]. Examples of recent applications of these and related ideas to other systems appears in [29, 28, 61, 34, 35, 31, 32, 33].

Since (3.37) is a constant coefficient equation, $F_{A,\Gamma}$ can be related, via scaling, to the nonlinear ground state of the translation invariant nonlinear Schrödinger equation. Thus, the shape of the bifurcation diagram $\mu \mapsto \mathcal{N}[\psi_\mu]$ (see Figure 3) can be deduced for μ near μ_\star ; see [59]. We also note that the stability / instability properties of ground states of (3.37) for the time-dependent effective nonlinear Schrödinger equation, are a consequence of Theorem 3.3 in the translation invariant case.

4 Soliton / Defect Interactions

In this section we turn to the detailed time dynamics of a soliton-like nonlinear bound state interacting with a potential. This question has fundamental and applications interest. From the fundamental perspective it is an important problem in the direction of developing a nonlinear scattering theory for non-integrable Hamiltonian PDEs. And, from an applications perspective, nonlinear waves often arise in systems with impurities, *e.g.* random defects in fabrication or those deliberately inserted into the medium to influence the propagation; see, for example, [48].

Referring to Figure 4 we sketch the different stages present in the dynamics of a soliton which is initially incident upon a potential well:

The results of formal asymptotic and numerical studies of soliton / defect interactions (see, for example, [48, 46, 11]) suggests a description of the time-dynamics in terms of overlapping time




Figure 5: $P = \mathcal{N}[\psi_\mu]$ vs. μ for the 1d - NLS / GP with periodic potential, $V(x)$ for the case $\sigma = \sigma_{\text{cr}} = 2/d = 2$. Dashed line $P = P_{\text{cr}}$ is the (μ - independent) value of $\mathcal{N}[\psi_\mu]$ for the case $V \equiv 0$. Solid curve is the curve $\mu \mapsto \mathcal{N}[\psi_\mu]$, where $\mu < \mu_\star = \inf(\sigma(-\Delta + V))$ for the case where V is a non-trivial periodic potential. Nontrivial effective mass matrix implies $P_\star = \lim_{\mu \uparrow \mu_\star} \mathcal{N}[\psi_\mu] < P_{\text{cr}}$. See [59].

epochs. Very roughly, these are:

- (i) short and long / intermediate time-scale *classical particle-like dynamics*, which govern motion and deformations of the soliton as it interacts with the defect and exchanges energy with the defect's internal modes (pinned nonlinear defect modes) and
- (ii) the very long time scale, during which the system's energy resolves into outgoing solitons moving away from the defect, small amplitude waves which disperse to infinity and an asymptotically stable nonlinear ground state of supported in the defect. A key process in this *asymptotic resolution* is radiation damping due to coupling of discrete and continuum degrees of freedom. See sections 5 and 6.

Examples of rigorous analyses for *transient / intermediate time-scale* regimes:

- (i) *Soliton scattering from a potential well*: Detailed reflection, transmission and trapping for solitons incident on a potential barrier or potential well [54, 55, 56, 21].
- (ii) *Soliton evolving in a potential well*: In [57] the evolution of an order one soliton in a potential well is considered. The detailed *nonlinear breathing dynamics*, as the soliton relaxes toward its asymptotic state are considered in [58]; see also section 6 and the related work [38, 40, 41].

In [80] the evolution of a weakly nonlinear (small amplitude) NLS / GP solutions in a double-well, for which there is symmetry breaking (see section 3.6) is studied. Figure 4 displays phase portrait of the reduced Hamiltonian dynamical system in [80]. Orbits around the left (respectively, right) equilibrium map to a soliton executing a long-time nearly periodic (back and forth) motion with the left (respectively, right) well of the double well. This analysis has been recently extended to certain orbits outside the separatrix [47].




Figure 6: A soliton interacts with a defect, support around $x = 0$. As time advances, the coherent structure interacts with the defect. Part of the incident energy is scattered to infinity and part of it is trapped in the defect. The trapped energy settles down to a stable nonlinear defect mode.

For $t \gg 1$ the center of mass eventually crystallizes on a stable nonlinear ground state. For $\mathcal{N} > \mathcal{N}_{\text{cr}}$ this asymptotic state will be an asymmetric state (section 3.6) centered in the left or in the right well. See Figure 4 which shows the computed center of mass motion for a solution of NLS / GP; initially there is oscillatory motion among the left and right wells. However, during each cycle some of the soliton's energy radiated away to infinity. The corresponding motion in the reduced phase portrait Figure 4 is damped and the actual center of mass trajectory is transverse to the level energy curves. Eventually the separatrix is crossed and the solution settles down to a stable asymmetric state [19, 96, 97] We emphasize that this picture is only heuristic although there has been considerable progress toward understanding the radiation damping and ground state selection in such systems. We now turn to these phenomena in the next section. We conclude this section noting work on the dynamics of forced / damped NLS and its reduced finite dimensional dynamics, capturing regular and chaotic behavior; see, for example, [10, 90].

5 Resonance, radiation damping and infinite time dynamics

Our next goal is to discuss results concerning the infinite time dynamics of NLS / GP. We begin with an informal discussion of the linear Schrödinger equation (NLS / GP with $g = 0$):

$$i\partial_t\psi = (-\Delta + V(x))\psi \quad (5.1)$$

Assume $V(x)$ is a smooth and sufficiently decaying real-valued potential for which $-\Delta + V$ has




Figure 7: Periodic dynamics of center of mass dynamics in the reduced (approximate) finite-dimensional Hamiltonian system [80]. Equilibria corresponding to stable asymmetric states centered on left and right local minima of double well. Oscillatory orbits decay to stable equilibria for the full, infinite-dimensional NLS/GP.




Figure 8: Numerically computed dynamics of center of mass in the NLS evolution [80]. Corresponding trajectory in the reduced phase portrait of Figure 4 cycles around several times outside the separatrix, crosses the separatrix and then slowly spirals in toward the right stable equilibrium.

bound states :

$$\begin{aligned} (-\Delta + V)\psi_j &= \Omega_j \psi_j, \quad j = 0, 1, \dots, \quad \psi_j \in L^2 \\ \psi(x, t) &= e^{-i\Omega_j t} \psi_j(x) \end{aligned}$$

Then, for sufficiently localized initial conditions, $\psi(0)$, the solution of the initial value problem for (5.1) may be written:

$$\psi(x, t) = e^{-i(-\Delta+V)t} \psi_0 = \sum_j \langle \psi_j, \psi(0) \rangle e^{-i\Omega_j t} \psi_j(x) + R(x, t) \quad (5.2)$$

The sum in (5.2) is quasi-periodic while the second term, $R(x, t)$, radiates to zero in the sense that for some $\alpha > 0$

$$\|R(\cdot, t)\|_{L^p(\mathbb{R}^d)} \lesssim t^{-\alpha}, \text{ as } t \rightarrow \pm\infty \quad (5.3)$$

Thus, as $t \rightarrow \infty$, the solution $\psi(x, t)$ tends to an asymptotic state which is quasi-periodic in time and localized in space.

Question: What is the large time behavior of solutions of the corresponding initial value problem for the nonlinear Schrödinger / Gross-Pitaevskii, $g \neq 0$?

For small amplitude it is natural to attempt expansion of solutions in the basis of localized eigenstates and continuum modes, associated with the unperturbed (solvable) linear Schrödinger equation. In terms of these coordinates, the original Hamiltonian PDE may be written as an equivalent dynamical system comprised of two weakly coupled subsystems:

- a finite or infinite dimensional subsystem with **discrete** degrees of freedom (“oscillators”)
- an infinite dimensional system (wave equation) governing a **continuum** “field” .

These two systems are coupled due to weak nonlinearity. If one “turns off” the nonlinear coupling there are localized in space and time-periodic solutions, corresponding the eigenstates of the linear Schrödinger equation. If nonlinearity is present, new frequencies are generated, and these may lead to resonances among discrete modes or between discrete and continuum radiation modes. The latter type of resonance plays a key role in understanding the radiation damping mechanism central to asymptotic relaxation of solutions to NLS / GP as $t \rightarrow \pm\infty$ and, in particular the phenomena of *ground state selection* and *energy equipartition*.

These results will be discussed in section 6. A variant was studied in the context of the nonlinear Klein Gordon equation in [95, 4]. In the following two subsections we present a very simple example of emergent effective damping in an infinite dimensional Hamiltonian problem.

5.1 Simple model - part 1: Resonant energy exchange between an oscillator and a wave-field

We consider a solvable toy model of an infinite dimensional Hamiltonian system comprised of two subsystems, one governing discrete and the other governing continuum degrees of freedom. We shall see that coupling of these subsystems leads to energy transfer from the discrete to the continuum modes and the emergence of effective damping. Our model is a variation on the work of Lamb (1900) [75] and Weisskopf-Wigner (1930) [111]; see also [93, 14, 92, 95, 94, 67, 68, 69, 18]

Consider a system which couples an “oscillator” with amplitude $a(t)$ to a “field” with amplitude $u(x, t)$, $t \in \mathbb{R}$, $x \in \mathbb{R}$:

$$\frac{da^\varepsilon(t)}{dt} + i\omega a^\varepsilon(t) = -\varepsilon u^\varepsilon(0, t), \quad (5.4)$$

$$\partial_t u^\varepsilon(x, t) + c \partial_x u^\varepsilon(x, t) = \varepsilon \delta(x) a^\varepsilon(t). \quad (5.5)$$

Here, ε and c are taken to be a real parameters, say with $c > 0$, and $\delta(x)$ denotes the Dirac delta function. The amplitudes $a(t)$ and $u(x, t)$ are complex-valued.

Note that the dynamical system (5.4)-(5.5) conserves an energy, $\partial_t \mathcal{E}[a(t), u(\cdot, t)] = 0$.

$$\mathcal{E}[a(t), u(\cdot, t)] \equiv |a(t)|^2 + \int_{\mathbb{R}} |u(x, t)|^2 dx \quad (5.6)$$

We consider the initial value problem for (5.4)-(5.5) with initial data

$$a(0) \text{ arbitrary and } u(x, 0) = 0; \quad (5.7)$$

we perturb the oscillator initially, but not the field.

For $\varepsilon = 0$ the oscillator and wave-field are decoupled and there is an exact time-periodic and finite energy global solution, a *bound state*:

$$\begin{pmatrix} u^{\varepsilon=0}(x, t) \\ a^{\varepsilon=0}(t) \end{pmatrix} = e^{-i\omega t} \begin{pmatrix} 0 \\ a(0) \end{pmatrix}.$$

For $\varepsilon \neq 0$, the oscillator and field are coupled; as time evolves energy can be transfer among the discrete and continuum degrees of freedom. It is simple to solve the initial value problem for any choice of ε and arbitrary initial data, *e.g.* by Laplace transform. In particular, one can proceed by solving the wave equation, (5.5), for $u^\varepsilon = u(x, t; a^\varepsilon)$ as a functional of discrete degree of freedom $a^\varepsilon(t)$ and then substitute the resulting expression for $u^\varepsilon(0, t; a^\varepsilon(t))$ into the oscillator equation (5.4). The result is the closed oscillator

$$\frac{da^\varepsilon(t)}{dt} + i\omega a^\varepsilon(t) = -\frac{\varepsilon^2}{2c} a^\varepsilon(t), \quad t \geq 0. \quad (5.8)$$

with an effective damping term, whose solution is:

$$a^\varepsilon(t) = e^{-i\omega t} e^{-\frac{\varepsilon^2}{2c} t} a_0 = e^{-i(\omega - i\frac{\varepsilon^2}{2c}) t} a_0, \quad t \geq 0. \quad (5.9)$$

That is, in terms of the oscillator, the *closed* infinite dimensional conserved system for $a^\varepsilon(t)$ and $u^\varepsilon(x, t)$ may equivalently be viewed in terms of a reduced *open and damped* finite dimensional system for $a^\varepsilon(t)$.

Solving (5.8) for $a^\varepsilon(t)$ and substituting into the expression for $u^\varepsilon = u(x, t; a^\varepsilon)$, one obtains:

$$u^\varepsilon(x, t) = \begin{cases} 0, & x > ct \geq 0 \\ \frac{\varepsilon}{c} a_0 e^{-i\omega(t-\frac{x}{c})} e^{-\frac{\varepsilon^2}{2c}(t-\frac{x}{c})}, & x < ct \end{cases}$$

To summarize, for $\varepsilon = 0$, the decoupled system has a time-periodic finite energy bound state. For the coupled system, $\varepsilon \neq 0$, the bound state has a finite lifetime; it decays on a timescale of order ε^{-2} . The bound state loses its energy to the continuum degrees of freedom; the lost energy is propagated to spatial infinity.

5.2 Simple model - part 2: Resonance, Effective damping and Perturbations of Eigenvalues in Continuous Spectra

It's physically intuitive that an oscillator coupled to wave propagation in an infinite medium will damp. We now connect this damping to the classical notion of resonance.

We write the oscillator / wave-field model in the form

$$i\partial_t \psi = A(\lambda) \psi, \quad (5.10)$$

where

$$\psi(t) = \begin{pmatrix} u(x, t) \\ a(t) \end{pmatrix}, \quad A(\varepsilon) = \begin{pmatrix} -ic\partial_x & +i\varepsilon\delta(x) \\ -i\varepsilon\langle\delta(\cdot), \cdot\rangle & \omega \end{pmatrix} \quad (5.11)$$

Let

$$\psi(t) = e^{-iEt} \psi_0 = \begin{pmatrix} u(x, t) \\ a(t) \end{pmatrix} = e^{-iEt} \begin{pmatrix} u_0 \\ a_0 \end{pmatrix} \quad (5.12)$$

This yields the spectral problem

$$A(\varepsilon)\psi_0 = E\psi_0, \quad (5.13)$$

or equivalently

$$E \begin{pmatrix} u_0 \\ a_0 \end{pmatrix} = \begin{pmatrix} -ic\partial_x & +i\varepsilon\delta(x) \\ -i\varepsilon\langle\delta(y), \cdot\rangle & \omega \end{pmatrix} \begin{pmatrix} u_0 \\ a_0 \end{pmatrix}. \quad (5.14)$$

This spectral problem may be considered on the Hilbert space:

$$\mathcal{H} = \left\{ \begin{pmatrix} u \\ a \end{pmatrix} \in L^2(\mathbb{R}) \times \mathbb{C} : \|(u, a)\|_{\mathcal{H}} < \infty \right\}, \quad (5.15)$$

where

$$\|(u, a)\|_{\mathcal{H}} = \int_{\mathbb{R}} |U(x)|^2 dx + |a|^2 < \infty \quad (5.16)$$

We note that for $\varepsilon = 0$ (decoupling of oscillator and wave-field), we have

$$E \begin{pmatrix} u_0 \\ a_0 \end{pmatrix} = \begin{pmatrix} -ic\partial_x & 0 \\ 0 & \omega \end{pmatrix} \begin{pmatrix} u_0 \\ a_0 \end{pmatrix} \quad (5.17)$$

$A(\varepsilon = 0)$ is diagonal and has spectrum given by:

- a point eigenvalue at $E = \omega$ with corresponding eigenstate $\psi_\omega = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and
- continuous spectrum $\mathbb{R} = \{E = ck : k \in \mathbb{R}\}$ with eigenstates $\psi_k = \begin{pmatrix} e^{ikx} \\ 0 \end{pmatrix}$, $k \in \mathbb{R}$.

Thus, $A(\varepsilon = 0)$ has an embedded eigenvalue, ω , in the continuous spectrum, \mathbb{R} , and the source the damping is seen to be this resonance. Understanding the coupling of oscillator and wave-field is therefore related to the perturbation theory of an embedded (non-isolated) point in the spectrum; see, for example, [87, 53, 93] and references therein.

In this present simple setting, this perturbation problem can be treated as follows. Note that for $\varepsilon \neq 0$, since $c > 0$ we expect that energy is emitted by the oscillator and it gets carried to positive infinity through $u(x, t)$. Thus we expect that on a fixed compact set, $u(x, t)$, will decay to zero. Alternatively, if we choose $\alpha > 0$ and study $U(x, t) = e^{-\alpha x} u(x, t)$, $U(x, t)$ can be expected to decay to zero as t advances. Indeed, it can be checked that $U(x, t)$ satisfies a *dissipative* PDE, for which the $L^2(\mathbb{R})$ norm of $U(x, t)$ decays as t increases; see the discussion of section 5.1.

This behavior is reflected in the spectral problem. Consider the of change variables $(u, a) \mapsto (e^{\alpha x} U, a)$ in the spectral problem (5.14). This yields

$$E \begin{pmatrix} U_0 \\ a_0 \end{pmatrix} = \begin{pmatrix} -ic\partial_x - ic\alpha & +i\varepsilon e^{-\alpha x} \delta(x) \\ -i\varepsilon \langle \delta(\cdot) e^{\alpha \cdot}, \cdot \rangle & \omega \end{pmatrix} \begin{pmatrix} U_0 \\ a_0 \end{pmatrix} = A_\alpha(\varepsilon) \begin{pmatrix} U_0 \\ a_0 \end{pmatrix}, \quad (5.18)$$

which when considered on $L^2(\mathbb{R})$ is equivalent to the spectral problem (5.14) for $(u(x), a)$ in the weighted function space:

$$\mathcal{H}_\alpha = \{(u(x), a) : \|u\|_{\mathcal{H}_\alpha} < \infty\}, \quad (5.19)$$

where

$$\|u\|_{\mathcal{H}_\alpha} = \int_{\mathbb{R}} |u(x)|^2 e^{-2\alpha x} dx + |a|^2 < \infty, \quad \alpha > 0. \quad (5.20)$$

For $\varepsilon = 0$,

$$E \begin{pmatrix} v_0 \\ a_0 \end{pmatrix} = \begin{pmatrix} -ic\partial_x - ic\alpha & 0 \\ 0 & \omega \end{pmatrix} \begin{pmatrix} v_0 \\ a_0 \end{pmatrix} \quad (5.21)$$

$A_\alpha(0)$ is diagonal and has spectrum:

- point eigenvalue at $E = \omega$ with corresponding eigenstate $\psi_\omega = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$
- continuous spectrum in the lower half plane along the horizontal line $\{E = ck - ic\alpha : k \in \mathbb{R}\}$ with eigenstates $\psi_k = \begin{pmatrix} e^{ikx} \\ 0 \end{pmatrix}$, $k \in \mathbb{R}$

Note that for $\alpha > 0$, ω is an *isolated* eigenvalue of $A_\alpha(\varepsilon = 0)$ and we can therefore implement a standard perturbation theory to calculate the effect of $\varepsilon \neq 0$ on this eigenvalue.

The first equation of (5.18) is

$$(-ic\partial_x - ic\alpha)v_0 + i\varepsilon\delta(x)e^{-\alpha x}a_0 = Ev_0 \quad (5.22)$$

This implies for $x \neq 0$

$$v_0(x) = \begin{cases} e^{i(\frac{E}{c}x-\alpha)x}v_{0+} & x > 0 \\ e^{i(\frac{E}{c}x-\alpha)x}v_{0-} & x < 0 \end{cases}$$

Integration of (5.18) over a small neighborhood of $x = 0$ yields

$$v_{0+} - v_{0-} = +\frac{\varepsilon}{c}a_0 \quad (5.23)$$

The second equation of (5.18) implies

$$(E - \omega)a_0 = -\frac{i\varepsilon}{2}(v_{0+} + v_{0-}) \quad (5.24)$$

Now choose $v_{0+} \neq 0$ and $v_{0-} = 0$. Then,

$$E^\varepsilon = \omega - i\Gamma^\varepsilon, \quad \Gamma^\varepsilon = \frac{\varepsilon^2}{2c} > 0. \quad (5.25)$$

The corresponding eigenstate is

$$v_{0,\alpha}^\varepsilon(x) = \begin{cases} +\frac{\varepsilon}{c}a_0 e^{i\omega\frac{x}{c}}e^{\frac{1}{c}(\frac{\varepsilon^2}{2c}-\alpha c)x}, & x > 0 \\ 0, & x < 0 \end{cases}, \quad (5.26)$$

which is in $L^2(\mathbb{R})$ provided $\alpha > \frac{\varepsilon^2}{2c^2}$.

For $\alpha = 0$, we have

$$v_0^\varepsilon(x) = \begin{cases} +\frac{\varepsilon}{c}a_0 e^{i\omega\frac{x}{c}}e^{\frac{1}{c}\frac{\varepsilon^2}{2c}x}, & x > 0 \\ 0, & x < 0 \end{cases}. \quad (5.27)$$

The solution (5.27) satisfies an *outgoing radiation condition* at $x = +\infty$ and is in $L^2(\mathbb{R}; e^{-2\alpha x}dx)$ provided $\alpha > \frac{\varepsilon^2}{2c^2}$.

To summarize: the $\varepsilon = 0$ eigenvalue problem has discrete eigenvalue, $E^0 = \omega$, embedded in the continuous spectrum, \mathbb{R} with corresponding eigenstate $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$. This eigenvalue perturbs, for $\varepsilon \neq 0$, to a complex energy in the lower half plane, $E^\varepsilon = \omega - i\Gamma^\varepsilon$, $\Gamma^\varepsilon > 0$ with corresponding eigenstate $\begin{pmatrix} v_0^\varepsilon(x) \\ 1 \end{pmatrix}$ which solves the eigenvalue equation with outgoing radiation condition at $x = +\infty$. The complex energy, E^ε , may be viewed as a eigenvalue with normalizable eigenstate, $v_0(x)$ in the weighted function space, \mathcal{H}_α . Since E^ε is in the lower half plane, the corresponding time-dependent state is exponentially decaying as t increases.

Remark 5.1 Finally, we remark that the complex energy E^ε may be viewed as a pole of the Green's function, when analytically continued from the upper half E - plane to the lower half E - plane.

6 Ground state selection and energy equipartition in NLS / GP

In this section we return to the question raised in section 5, here restated. For simplicity we consider NLS / GP for the case with cubic nonlinearity ($\sigma = 1$):

$$i\partial_t\Phi = (-\Delta + V(x)) \Phi + g|\Phi|^2\Phi, \quad x \in \mathbb{R}^3 \quad (6.1)$$

We assume that $-\Delta + V$ has multiple independent localized eigenstates. For simplicity we assume two distinct eigenvalues.

As noted, the linear time-evolution ($g = 0$), for $t \gg 1$, settles down to a quasi-periodic state, a linear superposition of time-periodic and spatially localized solutions, corresponding to the independent eigenstates. Note also, that for $g \neq 0$ there may be multiple co-existing branches of nonlinear defect states; see, for example, the discussion of bound states for the case where V is a double-well potential, discussed in section 3.

Question: What is the long term ($t \uparrow \infty$) behavior of the NLS / GP ($g \neq 0$), (6.1) for initial data of small norm?

Our results show that under reasonable conditions (which have been explored experimentally [78, 97]) we have:

1. *Ground state selection* [96, 97, 42, 43]: The generic large time behavior of the initial value problem is periodic and is, in particular, a nonlinear ground state of the system. See also [8, 7, 104, 105, 9].
2. *Energy equipartition* [43]: For initial data conditions whose nonlinear ground state and excited state components are equal in L^2 , the solution approaches a new nonlinear ground state, whose L^2 norm has gained an amount equal one-half that of the initial excited state energy. The other one-half of the excited state energy is radiated away.

To state our results precisely requires some mathematical setup. For simplicity we assume that the linear operator $-\Delta + V$ has the following properties:

- (V1) V is real-valued and decays sufficiently rapidly, *e.g.* exponentially, as $|x|$ tends to infinity.
- (V2) The linear operator $-\Delta + V$ has two eigenvalues $E_{0\star} < E_{1\star} < 0$. $E_{0\star}$ is the lowest eigenvalue with ground state $\psi_{0\star} > 0$, the eigenvalue $E_{1\star}$ is possibly degenerate with multiplicity $N \geq 1$ and corresponding eigenvectors $\xi_{1\star}, \xi_{2\star}, \dots, \xi_{N\star}$.
- (V3) Resonant coupling assumption

$$\omega_{\star} \equiv 2E_{1\star} - E_{0\star} > 0, \quad . \quad (6.2)$$

We remark on (V3). An important role is played by the mechanism resonant coupling of discrete and continuum modes and energy transfer from localized states to dispersive radiation. In sections 5.1 and 5.2 we introduced and analyzed a simple model of this phenomenon. This model was linear and the resonance was due to a discrete eigenvalue, ω , embedded in the continuous spectrum. For NLS / GP, (6.1) resonant coupling of discrete and continuum modes arises due to higher time-harmonic generation by nonlinearity. The assumption (V3) is made to ensure that this coupling occurs at second order in (small) energy of the solution. However coupling at arbitrary higher order can be studied using the normal form ideas developed in [39, 16, 4, 17].

6.1 Linearization of NLS/GP about the ground state

Recall Theorem 3.2 on the family of nonlinear bound states, $E \mapsto \psi_E$, bifurcating from the ground state. We now consider the linearized NLS / GP, (6.1), time-dynamics about this family of solutions. Let

$$\Phi(x, t) = e^{-iEt} (\psi_E + u + iv),$$

where u and v are real and imaginary parts of the perturbation. Then the linearized perturbation equation about ψ_E , is

$$\frac{\partial}{\partial t} \begin{pmatrix} u \\ v \end{pmatrix} = L(E) \begin{pmatrix} u \\ v \end{pmatrix} = JH(E) \begin{pmatrix} u \\ v \end{pmatrix}, \quad (6.3)$$

where

$$L(E) = \begin{pmatrix} 0 & L_-(E) \\ -L_+(E) & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} L_+(E) & 0 \\ 0 & L_-(E) \end{pmatrix} \equiv JH(E). \quad (6.4)$$

The operators L_+ and L_- are given by:

$$\begin{aligned} L_-(E) &= -\Delta - E + V + g(\psi_E)^2 \\ L_+(E) &= -\Delta - E + V + 3g(\psi_E)^2 \end{aligned} \quad (6.5)$$

The following result on the linearized matrix-operator, $L(E)$, proved by standard perturbation theory [87], is given in [42] (Propositions 4.1 and 5.1):

Lemma 6.1 *Assume (V1), (V2) and (V3). Let $L(E)$ denote the linearized operator about a state on the branch of bifurcating bound states, given by Theorem 3.2:*

$$\psi_E(x) = \rho(E) (\psi_{0\star}(x) + \mathcal{O}(\rho(E)^2)), \quad \text{where } \rho(E) \equiv |E_{0\star} - E|^{\frac{1}{2}} \left(|g| \int \psi_{0\star}^4 \right)^{-\frac{1}{2}} \quad (6.6)$$

For $E = E_{0\star}$, the matrix operator $L(E_{0\star})$ has complex conjugate eigenvalues $\pm i\beta_\star = \pm i(E_{1\star} - E_{0\star})$, each of multiplicity N . For $|E_{0\star} - E|$ and small these perturb to (possibly degenerate) eigenvalues $\pm i\beta_1(E), \dots, \pm i\beta_N(E)$ with corresponding neutral modes (eigenstates)

$$\begin{pmatrix} \xi_1 \\ \pm i\eta_1 \end{pmatrix}, \begin{pmatrix} \xi_2 \\ \pm i\eta_2 \end{pmatrix}, \dots, \begin{pmatrix} \xi_N \\ \pm i\eta_N \end{pmatrix}$$

$$\text{satisfying } \langle \xi_m, \eta_n \rangle = \delta_{m,n}, \quad \langle \xi_m, \psi_E \rangle = \langle \eta_m, \partial_E \psi_E \rangle = 0. \quad (6.7)$$

Moreover,

$$0 \neq \lim_{E \rightarrow E_{0\star}} \xi_n = \lim_{E \rightarrow E_{0\star}} \eta_n \in \text{span}\{\xi_{1\star}, \xi_{2\star}, \dots, \xi_{N\star}\} \quad (6.8)$$

in Sobolev H^k spaces for any $k > 0$.

Furthermore, we note that for $|E_{0\star} - E|$ sufficiently small

$$2\beta_n(E) + E \approx 2(E_{1\star} - E_{0\star}) + E_{0\star} = 2E_{1\star} - E_{0\star} > 0, \quad n = 1, 2, \dots, N, \quad (6.9)$$

Remark 6.1 Equation (6.9) (see also (V3), (6.2)) ensures coupling of discrete to continuous spectrum at second order in $|E - E_{0\star}|$.

6.2 Ground state selection and energy equipartition

In this section we give a detailed description of the long term evolution.

Theorem 6.1 (Ground State Selection) Consider NLS / GP, (6.1), with linear potential satisfying (V1), (V2) and (V3), and cubic nonlinearity ($\sigma = 1$). Assume, most notably, that the non-negative (Fermi golden rule) expression for $\Gamma_0(z, z^*)$ in (6.15) is strictly positive. (See [96, 42, 43] for detailed statement of all technical assumptions. This expression is always non-negative and generically strictly positive due to the assumption (6.2).).

Take initial conditions of the form:

$$\psi_0(x) = e^{i\gamma_0} [\psi_{E_0} + \alpha_0 \cdot \xi + i\beta_0 \cdot \eta + R_0] , \quad (6.10)$$

where γ_0 and $E_0 \in \mathcal{I} = (E_{0\star} - \delta_0, E_{0\star})$ are real constants, α_0 and β_0 are real $1 \times N$ vectors, and $R_0 : \mathbb{R}^3 \rightarrow \mathbb{C}$, are such that

$$|E_0 - E_{0\star}| \ll 1, \quad \frac{|\alpha_0|^2 + |\beta_0|^2}{\|\psi_{E_0}\|_2^2} \ll 1, \quad \|\langle x \rangle^4 R_0\|_{H^2} \lesssim |\alpha_0|^2 + |\beta_0|^2 \quad (6.11)$$

Then there exist smooth functions $E(t) : \mathbb{R}^+ \rightarrow \mathcal{I}$, $\gamma(t) : \mathbb{R}^+ \rightarrow \mathbb{R}$, $z(t) : \mathbb{R}^+ \rightarrow \mathbb{C}^N$ and $R(x, t) : \mathbb{R}^3 \times \mathbb{R}^+ \rightarrow \mathbb{C}$ such that the solution of NLS / GP evolves in the form:

$$\begin{aligned} \psi(x, t) &= e^{-i \int_0^t E(s) ds} e^{i\gamma(t)} \\ &\times [\psi_{E(t)} + a_1(z, \bar{z}) \partial_E \psi_E + ia_2(z, \bar{z}) \psi_E + (Re \tilde{z}[z, \bar{z}]) \cdot \xi + i (Im \tilde{z}[z, \bar{z}]) \cdot \eta + R] , \end{aligned} \quad (6.12)$$

where $\lim_{t \rightarrow \infty} E(t) = E_\infty$, for some $E_\infty \in \mathcal{I}$.

Here, $a_1(z, \bar{z}), a_2(z, \bar{z}) : \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{R}$ and $\tilde{z} - z : \mathbb{C}^N \times \mathbb{C}^N \rightarrow \mathbb{C}^N$ are polynomials of z and \bar{z} , beginning with terms of order $|z|^2$.

(A) The dynamics of mass/energy transfer is captured by the following reduced dynamical system for the key modulating parameters, $E(t)$ and $z(t)$:

$$\frac{d}{dt} \|\psi_{E(t)}\|_2^2 = z^* \Gamma_0(z, \bar{z}) z + \mathcal{S}_E(t), \quad (6.13)$$

$$\frac{d}{dt} |z(t)|^2 = -2z^* \Gamma_0(z, \bar{z}) z + \mathcal{S}_z(t). \quad (6.14)$$

Here, Γ_0 is a Fermi golden rule (damping) matrix given by ¹

$$\Gamma_0(z, z^*) \equiv g^2 \Im \langle [-\Delta + V - \omega_* - i0]^{-1} \psi_E(z \cdot \xi)^2, \psi_E(z \cdot \xi)^2 \rangle \geq c^2 |z|^4 \quad (6.15)$$

and $S_E(t), S_z(t) \lesssim (1+t)^{-\rho}$, with $\int_0^\infty S_E(\tau) d\tau$ and $\int_0^\infty \mathcal{S}_z(\tau) d\tau = o(|z_0|)^2$.

(B) **Estimates on $z(t)$ and the correction $\vec{R}(t)$:** For all $t \geq 0$, we have $\|\vec{R}(t)\|_{H^2} \leq \epsilon_\infty$. Moreover, the following decay estimates hold:

$$\left\| (1+x^2)^{-\nu} \vec{R}(t) \right\|_2 \leq C (\|\langle x \rangle^4 \psi_0\|_{H^2}) (1+t)^{-1}, \quad (6.16)$$

$$|z(t)| \leq C (\|\langle x \rangle^4 \psi_0\|_{H^2}) (1+t)^{-\frac{1}{2}} \quad (6.17)$$

Theorem 6.2 (Mass / Energy equipartition) Consider NLS / GP, (6.1), under the technical hypotheses of Theorem 6.1 and assumptions on the initial data (6.11), where $|\alpha_0|^2 + |\beta_0|^2$ is a measure of the neutral modes' perturbation of the ground state. Recall that E_0 is the energy of the initial nonlinear ground state and E_∞ is the asymptotic energy, guaranteed by Theorem 6.1. Then, in the limit as $t \rightarrow \infty$, one half of the neutral modes' mass contributes to forming a more massive asymptotic ground state and one half is radiated away as dispersive waves:

$$\|\psi_{E_\infty}\|_2^2 = \|\psi_{E_0}\|_2^2 + \frac{1}{2} [|\alpha_0|^2 + |\beta_0|^2] + o(|\alpha_0|^2 + |\beta_0|^2). \quad (6.18)$$

Figure 9 illustrates the phenomena of ground state selection and mass / energy equipartition in NLS / GP. These simulations are for the case of double-well potentials (see section 3.6). Asymptotic energy equipartition has been verified for the case when the stable ground state is symmetric ($\mathcal{N} < \mathcal{N}_{cr}$) and the case when the stable ground states is asymmetric ($\mathcal{N} > \mathcal{N}_{cr}$).

A result on equipartition of energy holds as well, for NLS-GP with a general power nonlinearity, (2.1), under more restrictive hypotheses [43]. Finally, we remark that it would be of interest to establish detailed results on energy-transfer in systems with multiple bound states for subcritical

¹ $\Gamma_0(z, z^*)$ assumed to be strictly positive, is non-negative by the following:

$$\begin{aligned} \Im [-\Delta + V - \omega_* - i0]^{-1} &= \frac{1}{2i} \lim_{\delta \downarrow 0} ([-\Delta + V - \omega_* - i\delta]^{-1} - [-\Delta + V - \omega_* + i\delta]^{-1}), \\ &= \pi \delta(-\Delta + V - \omega_*), \text{ where } \omega_* \in \sigma_{cont}(-\Delta + V). \end{aligned}$$

Note: $\Im [-\Delta + V - \omega_* - i0]^{-1}$ projects onto the generalized mode at energy $\omega_* \in \sigma_{cont}(-\Delta + V)$.




Figure 9: Numerical verification ground state selection and asymptotic energy equipartition. Computations performed by E. Shlizerman .

nonlinearities. In this setting the current perturbative treatment of small amplitude dispersive waves does not apply. A step in this direction is in the work [70, 71, 66], where asymptotic stability for systems with a single family of nonlinear bound states is treated by a novel time-dependent linearization procedure.

7 A nonlinear toy model of nonlinearity-induced energy transfer

In this section we explain the idea behind the results on ground state selection and energy equipartition. In the style of sections 5.1 and 5.2 we introduce a toy minimal model, here nonlinear, which captures the essential mechanisms and is, by comparison, simpler to analyze.

Our model is for the interaction of three amplitudes:

1. a “ground state” complex amplitude, $\alpha_0(t)$ with associated frequency $E_{0\star}$
2. an “excited state” complex amplitude, $\alpha_1(t)$ with associated frequency $E_{1\star}$, and
3. a “continuum wave-field” complex amplitude, $R(x, t)$, with spectrum of frequencies given by the positive real-line $[0, \infty)$.

As with NLS / GP, our model has cubic nonlinearity and the assumption

$$\omega_\star \equiv 2E_{1\star} - E_{0\star} > 0 \tag{7.1}$$

is assumed to assure coupling of discrete and continuum modes at second order in the solution amplitude. We also introduce a function, $\chi(x)$, sufficiently rapidly decaying at spatial infinity. (Recall that in section 5.1 we took $\chi(x) = \delta(x)$.)

Our nonlinear model is the following:

$$i\partial_t \alpha_0(t) - \Omega_0 \alpha_0 = g \langle \chi, \overline{R(t)} \rangle \alpha_1^2(t) \quad (7.2)$$

$$i\partial_t \alpha_1(t) - \Omega_1 \alpha_1 = 2g \langle \chi, R(t) \rangle \alpha_0(t) \overline{\alpha_1(t)} \quad (7.3)$$

$$i\partial_t R(x, t) = -\Delta R(x, t) + g \chi(x) \alpha_1^2(t) \overline{\alpha_0(t)} \quad (7.4)$$

A first observation is that the system (7.2)-(7.4) is Hamiltonian and has the time-invariant quantity

$$\frac{d}{dt} \left(|\alpha_0(t)|^2 + |\alpha_1(t)|^2 + \int_{\mathbb{R}^d} |R(x, t)|^2 dx \right) = 0$$

To solve (7.2)-(7.4), we first separate fast and slow scales by introducing slowly varying amplitudes, A_0 and A_1 :

$$\alpha_0(t) = e^{-i\Omega_0 t} A_0(t), \quad \alpha_1(t) = e^{-i\Omega_1 t} A_1(t)$$

Then, the system for A_0, A_1 and R becomes:

$$i\partial_t A_0(t) = g \langle \chi, \overline{R(t)} \rangle e^{-i\omega_\star t} A_1^2(t) \quad (7.5)$$

$$i\partial_t A_1(t) = 2g \langle \chi, R(t) \rangle e^{i\omega_\star t} A_0(t) \overline{A_1(t)} \quad (7.6)$$

$$i\partial_t R(x, t) = -\Delta R(x, t) + g \chi(x) e^{-i\omega_\star t} A_1^2(t) \overline{A_0(t)} \quad (7.7)$$

Note that since the spectrum of $-\Delta = [0, \infty)$ and since $\omega_\star > 0$ (by assumption) the Schrödinger wave equation, (7.7) is resonantly forced. We proceed now to solve for $R(x, t)$, keeping its dominant contributions. Duhamel's principle and a regularization, motivated by the local decay estimate (7.9) below, gives:

$$\begin{aligned} R(t) &= e^{i\Delta t} R(0) - ig e^{i\Delta t} \int_0^t e^{-i(-\Delta + \omega_\star)s} \chi \overline{A_0(s)} A_1^2(s) ds \\ &\stackrel{\delta \downarrow 0}{=} e^{i\Delta t} R(0) - ig e^{i\Delta t} \int_0^t \frac{1}{i(-\Delta + \omega_\star - i\delta)} \frac{d}{ds} \left[e^{i(-\Delta - \omega_\star)s} \right] \chi \overline{A_0(s)} A_1^2(s) ds \end{aligned}$$

Next we have, by integration by parts,

$$\begin{aligned} R(x, t) &\sim e^{i\Delta t} R(0) - g e^{-i\omega_\star t} \frac{1}{-\Delta - \omega_\star - i0} \chi \overline{A_0(t)} A_1^2(t) \\ &\quad + \overline{A_0(0)} A_1^2(0) \frac{e^{i\Delta t}}{-\Delta - \omega_\star - i0} \chi \\ &\quad + \int_0^t \frac{e^{i\Delta(t-s)}}{-\Delta - \omega_\star - i0} \chi \frac{d}{ds} \left[\overline{A_0(s)} A_1^2(s) \right] ds, \end{aligned}$$

which may be written as

$$R(t) \sim -ge^{-i\omega_*t} \frac{1}{-\Delta - \omega_* - i0} \chi \overline{A_0(t)} A_1^2(t) + \mathcal{O}\left(t^{-\frac{3}{2}}\right) \quad (7.8)$$

where the error term is obtained using the *local energy decay estimate*:

$$\left\| \langle x \rangle^{-\nu} \frac{e^{i\Delta t}}{-\Delta - \omega_* - i0} \langle x \rangle^{-\nu} \right\|_{L^2(\mathbb{R}^3) \rightarrow L^2(\mathbb{R}^3)} = \mathcal{O}(t^{-\frac{3}{2}}), \quad t \rightarrow +\infty, \quad (7.9)$$

where ν is sufficiently large and positive. Substitution of the leading order terms of $R(t)$ in the expansion (7.8) into (7.5)-(7.6) and use of the distributional identity

$$(-\Delta - \omega_* - i0)^{-1} = \text{P.V.} \frac{1}{-\Delta - \omega_*} + i\pi \delta(-\Delta - \omega_*)$$

yields the dissipative system for the amplitudes A_0 and A_1 :

$$\partial_t A_0 \approx +\Gamma |A_1|^4 A_0, \quad \partial_t A_1 \approx -2\Gamma |A_0|^2 |A_1|^2 A_1 \quad (7.10)$$

where $\Gamma \sim g^2 |\widehat{\chi}(\omega_*)|^2 \geq 0$ and is generically strictly positive. This is the analogue of the strict positivity discussed in the statement of Theorem 6.2. Here, $\widehat{\chi}$ denotes the Fourier transform of χ .

The omitted correction terms in (7.10) can be controlled in terms of the quantities $\|R(t)\|_{L^p}$ ($p > 2$ and sufficiently large), $\|\langle x \rangle^{-\nu} R(t)\|_{L^2}$ and $\|\langle x \rangle^{-\nu} e^{i\Delta t} (-\Delta - \omega_* - i0)^{-1} \langle x \rangle^{-\nu}\|_{L^2 \rightarrow L^2}$.

Finally, ground state selection with energy equipartition can be seen through the reduction (7.10). Indeed, setting $P_0 = |\alpha_0|^2$, $P_1 = |\alpha_1|^2$ we obtain

$$\frac{dP_0}{dt} \sim \Gamma P_1^2 P_0, \quad \frac{dP_1}{dt} \sim -2\Gamma P_1^2 P_0$$

It follows that $2P_0(t) + P_1(t) \sim 2P_0(0) + P_1(0)$. Sending $t \rightarrow \infty$ and using that $P_1(t) \rightarrow 0$, (indeed, $\Gamma > 0$), we obtain $P_0(\infty) = P_0(0) + \frac{1}{2}P_1(0)$.

8 Concluding remarks

A fundamental question in the theory of nonlinear waves for non-integrable PDEs is whether and in what sense arbitrary finite-energy initial conditions evolve toward the family of available nonlinear bound states and radiation. This question is often referred to as the *Soliton Resolution Conjecture*; see, for example, [103]. The asymptotic stability / nonlinear scattering results of the type discussed in section 6, being based on normal forms ideas, spectral theory of linearized operators, dispersive estimates and low-energy (perturbative) scattering methods, give a local picture of the phase space near families of nonlinear bound states. In contrast, integrable nonlinear systems which can be mapped to exactly solvable linear motions, allow for a global description of the dynamics [23, 24, 22, 20]. We also note the important line of research by Merle *et. al.* which is based on the monotone

evolution properties of appropriately designed local energies (see, for example, in [79, 81]) and does not rely on dispersive time-decay estimates of the linearized flow. It would be of great interest to adapt these methods for use in tandem with dynamical systems ideas to situations where the background linear medium, *i.e.* a potential $V(x)$, gives rise to structures (defect modes) which interact with “free solitons”.

With a view toward understanding asymptotic resolution in non-integrable Hamiltonian PDEs, one of the simplest global questions to ask is the following. Let V denote a smooth, radially-symmetric and uni-modal potential well which decays very rapidly at spatial infinity. Assume further that $-\Delta + V$ has a finite number of bound states. Consider NLS / GP with a repulsive nonlinear potential ($g = +1$) with radially symmetric initial conditions:

$$i\partial_t \Psi = -\Delta \Psi + V(x)\Psi + |\Psi|^{p-1}\Psi, \quad \Psi(x, 0) = \Psi_0(|x|) \in H_{radial}^1(\mathbb{R}^d)$$

It is natural to expect that energy which remains spatially localized must reside the family of nonlinear defect modes, *i.e.* bound states, localized within the potential well, which lie on solution branches bifurcating from the linear bound states of $-\Delta + V$ [88]; see section 3. Indeed, any concentration of energy outside the well should disperse to zero, since for $|x|$ large the equation is a translation invariant NLS equation with repulsive (defocusing) nonlinearity having no bound states. We conjecture that for arbitrary initial data solutions either disperse to zero or approach a nonlinear defect state, and furthermore that generic initial conditions approach a stable nonlinear defect state.

A step in this direction is the result of Tao [102] for $1 + \frac{4}{d} < p < 2^*$, in very high spatial dimensions. *For spatial dimensions $d \geq 11$ there is a compact attractor. That is, there exists K , a compact subset of $H_{radial}^1(\mathbb{R}^d)$, such that K is invariant under the the NLS / GP flow. Moreover, if $u \in C_t^0 H_x^1(\mathbb{R} \times \mathbb{R}^d)$ is a global-in-time solution of NLS / GP, then there exists $u_+ \in H^1(\mathbb{R}_{radial}^d)$ such that $\text{dist}_{H^1}(\Psi(t) - e^{i\Delta t} u_+, K) \rightarrow 0$, as $t \rightarrow +\infty$.*

It is natural to conjecture that K is the set of nonlinear bound (defect) states.

Finally we mention that the detailed dynamical picture of energy transfer from discrete to radiation modes has connections with important questions in applied physics. Note that the Fermi golden rule damping matrix, $\Gamma_0(z, z)$, appearing in Theorems 6.1 and 6.2, which controls the rate of decay of excited states (neutral modes) is controlled by the density of states of the linearized operator near the resonant frequency ω_* ; see (6.2).

Control problem: Can one design the potential, $V(x)$, in NLS / GP in order to inhibit / enhance energy transfer and relaxation (decay) to the system’s asymptotic state?

This relates to the problem of controlling the spontaneous emission rate of atoms by modifying the background environment of the atom (for us, the background linear potential), thereby

controlling the density of states [13]. A problem of this type was investigated analytically and computationally for a closely related parametrically forced linear Schrödinger equation [83]. Such a study for NLS / GP could inform the design of experiments, such as those reported on in [78].

References

- [1] G. P. Agrawal. *Nonlinear Fiber Optics*. Optics and Photonics. Academic Press, 2nd edition, 1995.
- [2] W.H. Aschbacher, J. Fröhlich, G.M. Graf, K. Schnee, and M. Troyer. Symmetry breaking regime in the nonlinear Hartree equation. *J. Math. Phys.*, 43:3879—3891, 2002.
- [3] N. W. Ashcroft and N. D. Mermin. *Solid State Physics*. Saunders College Publishing, Harcourt Brace Publishers, 1976.
- [4] D. Bambusi and C. Cuccagna. On dispersion of small energy solutions of the nonlinear Klein Gordon equation with a potential. *Am. J. Math.*, 133(5):1421–1468, 2011.
- [5] H. Berestycki and P.-L. Lions. Nonlinear scalar field equations, . Existence of a ground state. *Arch. Rat. Mech. Anal.*, 1983.
- [6] J. Bourgain. *Global Solutions of Nonlinear Schrödinger Equations*, volume 46 of *Colloquium Publications*. AMS, 1999.
- [7] V. S. Buslaev and G. Perel'man. *On the stability of solitary waves for nonlinear Schrödinger equations*, volume 164 of *Amer. Math. Soc. Transl. Ser. 2*. Amer. Math. Soc., Providence, RI, 1995.
- [8] V. S. Buslaev and G. S. Perelman. Scattering for the nonlinear Schrödinger equation: states that are close to a soliton. *St. Petersburg Math. J.*, 4(6):1111–1142, 1993.
- [9] V. S. Buslaev and C. Sulem. On asymptotic stability of solitary waves for nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(3):419—475, 2003.
- [10] D. Cai, D.W. McLaughlin, and K.T.R. McLaughlin. The nonlinear Schrödinger equation as both a PDE and a dynamical system. In *Handbook of Dynamical Systems*, volume 2, pages 599–675. North-Holland, 2002.
- [11] X. D. Cao and B. A. Malomed. Soliton-defect collisions in the nonlinear schrodinger equation. *Physics Letters A*, 206:177–182, 1995.
- [12] T. Cazenave and P.L. Lions. Orbital stability of standing waves for some nonlinear Schrödinger equations. *Comm. Math. Phys.*, 85:549–561, 1982.

- [13] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg. *Atom-Photon Interactions - Basic Processes and Applications*. Wiley-Interscience, 1992.
- [14] O. Costin and A. Soffer. Resonance theory for Schrödinger operators. *Comm. Math. Phys.*, 224:133–152, 2001.
- [15] R. Courant and D. Hilbert. *Methods of Mathematical Physics*. Interscience Publishers, Inc., New York, N.Y., 1953.
- [16] S. Cuccagna. The Hamiltonian structure of the nonlinear Schrödinger equation and the asymptotic stability of its ground states. *Comm. Math. Phys.*, 305(2):279–331, 2011.
- [17] S. Cuccagna. On the Darboux and Birkhoff steps in the asymptotic stability of solitons. *Rend. Istit. Mat. Univ. Trieste*, 44:197–257, 2012.
- [18] S. Cuccagna, E. Kirr, and D. Pelinovsky. Parametric resonance of ground states in the nonlinear Schrödinger equation. *J. Differential Equations*, 220(1):85–120, 2006.
- [19] S. Cuccagna and J. Marzuola. On instability for the quintic nonlinear schrödinger equation of some approximate periodic solutions. *Indiana J. Math.*, 61(6):2053–2083, 2013.
- [20] S. Cuccagna and D. Pelinovsky. The asymptotic stability of solitons in the cubic NLS equation on the line. *Applicable Analysis*, 93(791-822), 2014.
- [21] K. Datchen and J. Holmer. Fast soliton scattering by attractive delta impurities. *Commun. Partial Differential Equations*, 34:1074–1113, 2009.
- [22] P. Deift and J. Park. Long-time asymptotics for solutions of the NLS equation with a delta potential and even initial data. *Int. Math. Res. Not. IMRN*, 24:5505–5624, 2011.
- [23] P.A. Deift, A.R. Its, and X. Zhou. Long-time asymptotics for integrable nonlinear wave equations,. In *Important developments in soliton theory*, Springer Ser. Nonlinear Dynam., pages 181–204. Springer, Berlin, 1993.
- [24] P.A. Deift and X. Zhou. Long-time asymptotics for solutions of the nls equation with initial-data in weighted sobolev spaces. *Comm. Pure Appl. Math.*, 56:1029–1077, 2003.
- [25] T. Dohnal, D. Pelinovsky, and G. Schneider. Coupled-mode equations and gap solitons in a two-dimensional nonlinear elliptic problem with a separable potential. *J. Nonlin. Sci.*, 19:95–131, 2009.
- [26] T. Dohnal and H. Uecker. Coupled-mode equations and gap solitons for he 2d Gross-Pitaevskii equation with a non-separable periodic potential. *Physica D*, 238:860–879, 2009.

- [27] V. Duchêne, J.L. Marzuola, and M.I. Weinstein. Wave operator bounds for one-dimensional schrödinger operators with singular potentials and applications. *J. Math. Phys.*, 52:013505, 2011.
- [28] V. Duchene, I. Vukicevic, and M.I. Weinstein. Oscillatory and localized perturbations of periodic structures and the bifurcation of defect modes. <http://arxiv.org/abs/1407.8403>, 2014.
- [29] V. Duchene, I. Vukicevic, and M.I. Weinstein. Homogenized description of defect modes in periodic structures with localized defects. *Communications in Mathematical Sciences*, 2015.
- [30] L. Erdős, B. Schlein, and H.T. Yau. Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems. *Invent. Math.*, 59(12):1659–1741, 2007.
- [31] C. L. Fefferman and M. I. Weinstein. Wave packets in honeycomb lattice structures and two-dimensional Dirac equations. *Commun. Math. Phys.*, 326:251–286, 2014.
- [32] C.L. Fefferman, J.P. Lee-Thorp, and M.I. Weinstein. Topologically protected states in one-dimensional continuous systems. *Proc. Nat. Acad. Sci.*, 111(24):8759–8763, 2014.
- [33] C.L. Fefferman, J.P. Lee-Thorp, and M.I. Weinstein. Topologically protected states in one-dimensional systems. *to appear in Memoirs of the American Mathematical Society*, 2015.
- [34] C.L. Fefferman and M.I. Weinstein. Honeycomb lattice potentials and Dirac points. *J. Amer. Math. Soc.*, 25:1169–1220, 2012.
- [35] C.L. Fefferman and M.I. Weinstein. Waves in honeycomb structures. In D. Lannes, editor, *Proceedings of Journees Equations aux derivees partielles, Biarritz, 3-7 juin 2012*, volume GDR 243 4 (CNRS), – <http://arxiv.org/abs/1212.6684>, 2012.
- [36] G. Fibich. Some modern aspects of self-focusing theory. In Y.R. Shen R.W. Boyd, S.G. Lukishova, editor, *Self-Focusing: Past and Present*. Springer, 2009.
- [37] G. Fibich. *The Nonlinear Schrödinger Equation: Singularity Solutions and Optical Collapse*, volume 192 of *Applied Mathematical Sciences*. Springer-Verlag, 2014.
- [38] J. Fröhlich, S. Gustafson, B. L. G. Jonsson, and I. M. Sigal. Solitary wave dynamics in an external potential. *Comm. Math. Phys.*, 250(3):613–642, 2004.
- [39] Z. Gang. Perturbation expansion and Nth order Fermi golden rule of the nonlinear Schrödinger equations. *J. Math. Phys.*, 48(5):053509, 23, 2007.

- [40] Z. Gang and I. M. Sigal. On soliton dynamics in nonlinear Schrödinger equations. *Geom. Funct. Anal.*, 16(6):1377–1390, 2006.
- [41] Z. Gang and I.M. Sigal. Relaxation of solitons in nonlinear Schrödinger equations with potential. *Adv. Math.*, 216(2):443–490, 2007.
- [42] Z. Gang and M.I. Weinstein. Dynamics of nonlinear Schrödinger – Gross-Pitaevskii equations; mass transfer in systems with solitons and degenerate neutral modes. *Analysis and PDE*, 1(3), 2008.
- [43] Z. Gang and M.I. Weinstein. Equipartition of energy in Nonlinear Schrödinger / Gross-Pitaevskii Equations. *Applied Math. Research Express (AMRX)*, 2011.
- [44] B. Gidas, W.-M. Ni, and L. Nirenberg. Symmetry and related properties via the maximum principle. *Comm. Math. Phys.*, 68(3):209–243, 1979.
- [45] M. Golubitsky, I. Stewart, and D. Schaefer. *Singularities and Groups in Bifurcation Theory - Volume 2*. Springer-Verlag, 1988.
- [46] R.H. Goodman, P.J. Holmes, and M.I. Weinstein. Strong NLS soliton-defect interactions. *Physica D*, 161:21–44, 2004.
- [47] R.H. Goodman, J. Marzuola, and M.I. Weinstein. Self-trapping and josephson tunneling solutions to the nonlinear Schrödinger – Gross-Pitaevskii equation. *Discrete and Continuous Dynamical Systems - A*, 35(1):225–246, 2015.
- [48] R.H. Goodman, R.E. Slusher, and M.I. Weinstein. Stopping light on a defect. *J. Opt. Soc. B*, 19(7):1635–1652, 2002.
- [49] M. Grillakis, M. Machedon, and D. Margetis. Second order corrections to weakly interacting bosons, i. *Communications in Mathematical Physics*, 294(1):273–301, 2010.
- [50] M.G. Grillakis. Analysis of the linearization around a critical point of an infinite-dimensional hamiltonian system. *Comm. Pure Appl. Math.*, 43(3):299–333, 1990.
- [51] M.G. Grillakis, J. Shatah, and W.A. Strauss. Stability theory of solitary waves in the presence of symmetry. I. *J. Func. Anal.*, 74(1):160–197, 1987.
- [52] E. M. Harrell. Double wells. *Comm. Math. Phys.*, 75:239–261, 1980.
- [53] P. D. Hislop and I. M. Sigal. *Introduction to Spectral Theory: With applications to Schrödinger Operators*, volume 113 of *Applied Mathematical Sciences*. Springer, 1996.

- [54] J. Holmer, J. Marzuola, and M. Zworski. Fast soliton scattering by delta impurities. *Comm. Math. Phys.*, 274:349–367, 2007.
- [55] J. Holmer, J. Marzuola, and M. Zworski. Soliton splitting by delta impurities. *J. Nonlinear Sci.*, 7:349–367, 2007.
- [56] J. Holmer and M. Zworski. Slow soliton interaction with delta impurities. *J. Mod. Dyn.*, 1(4):689–718, 2007.
- [57] J. Holmer and M. Zworski. Soliton interaction with slowly varying potentials. *IMRN Internat. Math. Res. Notices*, 2008.
- [58] J. Holmer and M. Zworski. Breathing patterns in nonlinear relaxation. *Nonlinearity*, 22:1259–1301, 2009.
- [59] B. Ilan and M. I. Weinstein. Band edge solitons, nonlinear Schroedinger / Gross-Pitaevskii equations and effective media. *Multiscale Model. and Simul.*, 8(4):1055–1101, 2010.
- [60] R.K. Jackson and M.I. Weinstein. Geometric analysis of bifurcation and symmetry breaking in a Gross-Pitaevskii equation. *J. Stat. Phys.*, 116:881–905, 2004.
- [61] M. Jenkinson and M.I. Weinstein. On-site and off-site solitary waves of the discrete nonlinear Schroedinger equation in multiple dimensions. <http://arxiv.org/abs/1405.3892>, 2014.
- [62] C.K.R.T. Jones. An instability mechanism for radially symmetric standing waves of a nonlinear Schrödinger equation. *J. Diff. Eqns*, 71(1):34–62, 1988.
- [63] T. Kapitula, P. G. Kevrekidis, and Z. Chen. Three is a crowd: Solitary waves in photorefractive media with three potential wells. *SIAM Journal of Applied Dynamical Systems*, 5:598–633, 2006.
- [64] T. Kato. On nonlinear Schrödinger equations. *Ann. Inst. H. Poincaré Phys. Theor.*, 46:113–129, 1987.
- [65] E. Kirr, P. G. Kevrekidis, and D. E. Pelinovsky. Symmetry breaking in the nonlinear Schrödinger equation with a symmetric potential. *Comm. Math. Phys.*, 308(3):795–844, 2011.
- [66] E. Kirr and O. Mizrak. Asymptotic stability of ground states in 3D nonlinear Schrödinger equation including subcritical cases. *J. Func. Anal.*, 257(12):3691–3747, 2009.
- [67] E. Kirr and M.I. Weinstein. Parametrically excited Hamiltonian partial differential equations. *SIAM. J. Math. Anal.*, 33(1):16–52, 2001.

- [68] E. Kirr and M.I. Weinstein. Metastable states in parametrically excited multimode Hamiltonian systems. *Commun. Math. Phys.*, 236(2):335–372, 2003.
- [69] E. Kirr and M.I. Weinstein. Diffusion of power in randomly perturbed hamiltonian partial differential equations. *Comm. Math. Phys.*, 255(2):293–328, 2005.
- [70] E. Kirr and A. Zarnescu. On the asymptotic stability of bound states in 2D cubic Schrödinger equation. *Comm. Math. Phys.*, 272(2):443–468, 2007.
- [71] E. Kirr and A. Zarnescu. Asymptotic stability of ground states in 2D nonlinear Schrödinger equation including subcritical cases. *J. Differential Equations*, 247(3):710–735, 2009.
- [72] E.W. Kirr, P. G. Kevrekidis, E. Shlizerman, and M.I. Weinstein. Symmetry breaking in the nonlinear Schrödinger / Gross-Pitaevskii equation. *SIAM J. Math. Anal.*, 40(2):566–604, 2008.
- [73] W. Kirsch and B. Simon. Comparison theorems for the gap of Schrödinger operators. *J. Func. Anal.*, 75:396–410, 1987.
- [74] M. K. Kwong. Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in R^n . *Arch. Rat. Mech. Anal.*, 105:243–266, 1989.
- [75] H. Lamb. On a peculiarity of the wave-system due to the free vibrations of a nucleus in an extended medium. *Proc. London Math. Soc.*, 32:208–211, 1900.
- [76] E.H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. AMS, Providence, Rhode Island, 2nd edition, 1997.
- [77] P.-L. Lions. The concentration compactness principle in the calculus of variations i,ii. *Anal. I.H.P. Anal. Nonlin.*, 1:109–145, 223–283, 1984.
- [78] D. Mandelik, Y. Lahini, and Y. Silberberg. Nonlinearly induced relaxation to the ground state in a two-level system. *Phys. Rev. Lett.*, 95:073902, 2005.
- [79] Y. Martel and F. Merle. Asymptotic stability of solitons for subcritical generalized KdV equations. *Arch. Rat. Mech. Anal.*, 157:219–254, 2001.
- [80] J. Marzuola and M.I. Weinstein. Long time dynamics near the symmetry breaking bifurcation for nonlinear Schrödinger / Gross-Pitaevskii equation. *Discrete Contin. Dyn. Syst. A*, 28(4):1505–1554, 2010.
- [81] F. Merle and P. Raphael. On the universality of blow-up profile for the l^2 critical nonlinear Schrödinger equation. *Invent. Math.*, 156(565-672), 2004.

- [82] L. Nirenberg. *Topics in Nonlinear Functional Analysis*, volume 6 of *Courant Institute Lecture Notes*. American Mathematical Society, Providence, Rhode Island, 1974.
- [83] B. Osting and M.I. Weinstein. Emergence of periodic structure from maximizing the lifetime of a bound state coupled to radiation. *SIAM J. Multiscale Model. and Simul.*, 9(2):654–685, 2011.
- [84] D. E. Pelinovsky, A. A. Sukhorukov, and Y. S. Kivshar. Bifurcations and stability of gap solitons in periodic potentials. *Phys. Rev. E*, 70:036618, 2004.
- [85] C.A. Pillet and C.E. Wayne. Invariant manifolds for a class of dispersive Hamiltonian partial differential equations. *J. Diff. Eqns*, 1997.
- [86] L.P. Pitaevskii and S. Stringari. *Bose-Einstein Condensation*, volume 116 of *International Series of Monographs on Physics*. Clarendon Press, 2003.
- [87] M. Reed and B. Simon. *Modern Methods of Mathematical Physics, IV*. Academic Press, 1978.
- [88] H. A. Rose and M. I. Weinstein. On the bound states of the nonlinear Schrödinger equation with a linear potential. *Physica D*, 30:207–218, 1988.
- [89] Z. Shi and J. Yang. Solitary waves bifurcated from Bloch-band edges in two-dimensional periodic media. *Phys. Rev. E*, 75:056602, 2007.
- [90] E. Shlizerman and V. Rom-Kedar. Classification of solutions of the forced periodic nonlinear Schrödinger equation. *Nonlinearity*, 23:2183, 2010.
- [91] Y. Sivan, G. Fibich, B. Ilan, and M.I. Weinstein. Qualitative and quantitative analysis of stability and instability dynamics of positive lattice solitons. *Phys. Rev. E*, 78:046602, 2008.
- [92] A. Soffer and M. I. Weinstein. Nonautonomous Hamiltonians. *J. Stat. Phys.*, 93, 1998.
- [93] A. Soffer and M. I. Weinstein. Time dependent resonance theory. *Geom. Func. Anal.*, 8, 1998.
- [94] A. Soffer and M.I. Weinstein. Ionization and scattering for short lived potentials. *Lett. Math. Phys.*, 48, 1999.
- [95] A. Soffer and M.I. Weinstein. Resonances, radiation damping, and instability of Hamiltonian nonlinear waves. *Inventiones Mathematicae*, 136:9–74, 1999.
- [96] A. Soffer and M.I. Weinstein. Selection of the ground state for nonlinear Schrödinger equations. *Reviews in Mathematical Physics*, 16(8):977–1071, 2004.

- [97] A. Soffer and M.I. Weinstein. Theory of nonlinear dispersive waves and selection of the ground state. *Phys. Rev. Lett.*, 95:213905, 2005.
- [98] C. Sparber. Effective mass theorems for NLS equations. *SIAM J. Appl. Math.*, 66:820–842, 2006.
- [99] W.A. Strauss. Existence of solitary waves in higher dimensions. *Comm. Math. Phys.*, 55:149–162, 1979.
- [100] C. Sulem and P.L. Sulem. *The Nonlinear Schrödinger Equation: Self-Focusing and Wave Collapse*, volume 139 of *Series in Mathematical Science*. Springer-Verlag, 1999.
- [101] T. Tao. *Nonlinear Dispersive Equations: Local and Global Analysis*. Number 106 in CBMS Regional Conference Series. AMS, 2006.
- [102] T. Tao. A global compact attractor for high-dimensional defocusing non-linear Schrödinger equations with potential. *Dyn. PDE*, 5:101–116, 2008.
- [103] T. Tao. Why are solitons stable? *Bulletin and the AMS*, 46(1):1–33, 2009.
- [104] T.-P. Tsai and H.-T. Yau. Asymptotic dynamics of nonlinear Schrödinger equations: resonance-dominated and dispersion-dominated solutions. *Comm. Pure Appl. Math.*, 55(2):153–216, 2002.
- [105] T.-P. Tsai and H.-T. Yau. Classification of asymptotic profiles for nonlinear Schrödinger equations with small initial data. *Adv. Theor. Math. Phys.*, 6(1):107–139, 2002.
- [106] M. Vakhitov and A. Kolokolov. Stationary solutions of the wave equation in a medium with nonlinearity saturation. *Radiophys. Quant. Elec.*, 16:783, 1973.
- [107] M. I. Weinstein. Resonance problems in photonics. In D-Y Hsieh, M Zhang, and W Sun, editors, *Frontiers of Applied Mathematics- Proceedings of the 2nd International Symposium*. World Scientific, 2007.
- [108] M.I. Weinstein. Nonlinear Schrödinger equations and sharp interpolation estimates. *Comm. Math. Phys.*, 87:567–576, 1983.
- [109] M.I. Weinstein. Modulational stability of ground states of nonlinear Schrödinger equations. *SIAM J. Math. Anal.*, 16:472–490, 1985.
- [110] M.I. Weinstein. Lyapunov stability of ground states of nonlinear dispersive evolution equations. *Comm. Pure Appl. Math.*, 39:51–68, 1986.

- [111] V. Weisskopf and E. Wigner. Berechnung der natürlichen Linienbreite auf Grund der Dirac-schen Lichttheorie. *Z. Phys.*, 63:54–73, 1930.
- [112] G.B. Whitham. *Linear and Nonlinear Waves*. Wiley-Interscience, 1974.