

Decays of a neutral particle with zero spin and arbitrary CP parity into two off-mass-shell Z bosons

T.V. Zagoskin^{1,*} and A.Yu. Korchin^{1,2,†}

¹*NSC “Kharkov Institute of Physics and Technology”, 61108 Kharkov, Ukraine*

²*V.N. Karazin Kharkov National University, 61022 Kharkov, Ukraine*

Effects are investigated of CP symmetry violation in the decay of a scalar particle X (the Higgs boson) into two off-mass-shell Z bosons both decaying into a fermion-antifermion pair, $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$. The most general form of the amplitude of the transition $X \rightarrow Z_1^* Z_2^*$, wherein the boson X may not have definite CP parity, is considered. Limits of applicability of the narrow- Z -width approximation used when obtaining differential widths of the decay under consideration are determined. Various observables connected with the structure of the amplitude of the decay $X \rightarrow Z_1^* Z_2^*$ are studied. These observables are analyzed in the Standard Model, as well as in models conceding indefinite CP parity of the Higgs boson. An experimental measurement at the LHC of angular and invariant mass distributions of the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ can give information about the CP properties of the Higgs boson and its interaction with the Z boson.

I. INTRODUCTION

In 2012 the ATLAS and CMS collaborations detected [1] a neutral boson h with a mass of about 126 GeV. At the present time, detailed study of properties of this particle, called the Higgs boson, is an important task. The Standard Model (SM) Higgs boson is a state with $J^{CP} = 0^{++}$, and all the available experimental data about properties of the particle h are close to the corresponding theoretical predictions about the SM Higgs boson (see, for example, [2–4]). In particular, the spin of the boson h is equal to zero or two, and many hypotheses in which the spin of h is two are excluded with probability 95% or higher [3]. At the same time, the situation may be more complicated. For example, some supersymmetric models predict existence of neutral bosons with negative or even indefinite CP parity [5–7].

The issue of the CP parity of the Higgs boson is also related to the search for CP symmetry breaking sources which are additional to the mechanism built into the Cabibbo-Kobayashi-Maskawa quark-mixing matrix. Such sources of CP violation, for example, in the Higgs sector, could help in explaining the known problem of the matter-antimatter asymmetry in the Universe [8].

*Electronic address: taras.zagoskin@gmail.com

†Electronic address: korchin@kipt.kharkov.ua

It has been suggested [9, 10] that the CP properties of the Higgs boson be studied by investigation of decays into two photons, $h \rightarrow \gamma\gamma$, via measurement of the polarization characteristics of the photons. In Refs. [11] the decay to the photon and the Z boson, $h \rightarrow Z^*\gamma \rightarrow f\bar{f}\gamma$, has been examined while [12, 13] study the decay to the photon and a lepton pair, $h \rightarrow \gamma l^+ l^-$. In these papers it has been shown that the “forward-backward” escape asymmetry for the final fermions carries information about the CP properties of the h boson and physics beyond the SM.

Investigation into the decay of the Higgs boson into two Z bosons with their consequent decay to fermions is another opportunity to ascertain the CP properties of h . Such a cascade decay wherein the final fermions are leptons, along with the two-photon decay channel, has allowed the determination [1] of the mass of the particle h with the highest accuracy. In Refs. [14–17] theoretical distributions of the decay $h \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ have been studied at various values of the spin of h and in case of various CP properties of this boson. In [14] it has been reported what properties of experimental distributions testify about a particular spin and a particular CP parity of h . In [15–17] asymmetries measurement of which allows clarification of the mentioned properties of the Higgs boson are suggested and investigated. Finally, papers [18] put forward various methodologies on getting constraints on the Higgs boson couplings to ZZ , $W^- W^+$, $\gamma\gamma$ and $Z\gamma$ from experimental data.

Besides, various theories with spontaneous breaking of the conformal invariance (for example, theories of technicolor) assume the existence of one more neutral zero-spin particle which interacts with the gauge bosons – the dilaton. At present, the mass of the dilaton is not determined, but according to estimates performed in Ref. [19], in some models the mass can exceed 10^4 GeV. Along with that, in [20–22] it has been shown that the variant in which the boson h is the dilaton is not excluded.

In order to clarify the CP properties of the particle h and the hypothetical dilaton we consider a neutral particle X with zero spin and arbitrary CP parity. We examine the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ in case of the non-identical fermions, $f_1 \neq f_2$, and study in detail the differential width of this decay with respect to the three angles of the fermions in the helicity frame and with respect to the invariant masses of the fermion pairs $f_1 \bar{f}_1$ and $f_2 \bar{f}_2$. The most general $X \rightarrow Z_1^* Z_2^*$ vertex, which generalizes the corresponding SM vertex and contains a term corresponding to the negative CP parity of the particle X , is used.

We also find limits of applicability of the narrow-width approximation for the Z boson for the presented calculation of differential widths of the given decay. By means of this approximation we derive a formula for the total width of the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ (the formula is valid also in case $f_1 = f_2$) and a formula for the total width of the decay $h \rightarrow Z_1^* Z_2^*$. These formulas are more general and more precise than those obtained in Ref. [23].

Next we find observables connected with the structure of the amplitude of the decay $X \rightarrow Z_1^* Z_2^*$. The formula for the fully differential decay width contains nine coefficients related to the amplitude $X \rightarrow Z_1^* Z_2^*$. For each of them one or two observables linear in this coefficient are defined. Note that some of these observables, as well as different ones, have been studied in [15–17, 24], however we also obtain new experimentally measurable quantities

and analyze the dependences of the observables on the mass of one of the Z bosons (Z_2^*) in much more detail than it has been done in the mentioned papers. This analysis is carried out within the framework of the SM as well as in certain SM extensions wherein the boson h is a mixture of a CP -even state and a CP -odd one. Measurement of the suggested observables at the LHC can yield important information about the CP properties of the Higgs boson and its interaction with the Z boson.

II. FORMALISM FOR THE DECAYS $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$

A. The amplitude of the decay $X \rightarrow Z_1^* Z_2^*$ and the fully differential decay width for $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$

Let us consider the decay of a neutral spin-zero particle X with arbitrary CP parity into two off-mass-shell Z bosons (Z_1^* and Z_2^*) each of which decays to a fermion-antifermion pair, $f_1 \bar{f}_1$ and $f_2 \bar{f}_2$,

$$X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2, \quad (1)$$

where $m_X > 2(m_{f_1} + m_{f_2})$ (to satisfy the law of conservation of energy in a rest frame of X), m_X is the mass of the particle X , m_{f_j} is the mass of the fermion f_j . We will consider this decay at tree level. If $m_X \in (4m_b, 2m_t]$ (m_b is the mass of the b quark, m_t is the mass of the t quark), which holds true if $X = h$, then $f_j = e^-, \mu^-, \tau^-, \nu_e, \nu_\mu, \nu_\tau, u, c, d, s, b$. If $m_X > 4m_t$, which is possible [19] if X is the dilaton, then f_j can be the top quark as well.

From the energy-momentum conservation we find that a_1 and a_2 (a_j is the mass squared of the boson Z_j^* , i.e. the invariant mass squared of the pair $f_j \bar{f}_j$) lie within limits

$$4m_{f_1}^2 < a_1 < (m_X - 2m_{f_2})^2, \quad 4m_{f_2}^2 < a_2 < (m_X - \sqrt{a_1})^2. \quad (2)$$

The amplitude $A_{X \rightarrow Z_1^* Z_2^*}(\lambda_1, \lambda_2)$ of the decay of X into Z_1^* and Z_2^* is equal to [15–17, 24]

$$\begin{aligned} A_{X \rightarrow Z_1^* Z_2^*}(\lambda_1, \lambda_2) = & 2\sqrt{\sqrt{2}G_F m_Z^2} \left(a_Z(e_1^* \cdot e_2^*) + \frac{b_Z}{m_X^2}(e_1^* \cdot (p_1 + p_2))(e_2^* \cdot (p_1 + p_2)) + \right. \\ & \left. + i \frac{c_Z}{m_X^2} \varepsilon_{\mu\nu\rho\sigma} (p_1^\mu + p_2^\mu)(p_1^\nu - p_2^\nu)(e_1^\rho)^*(e_2^\sigma)^* \right), \end{aligned} \quad (3)$$

where λ_j , e_j , p_j are respectively the helicity, the polarization 4-vector and the 4-momentum of the boson Z_j^* , G_F is the Fermi constant, m_Z is the mass of the Z boson, a_Z , b_Z , c_Z are complex-valued dimensionless functions of a_1 and a_2 , $\varepsilon_{\mu\nu\rho\sigma}$ is the Levi-Civita symbol ($\varepsilon_{0123} = 1$). Note that at tree level

- if X is the SM Higgs boson, then $a_Z = 1$, $b_Z = c_Z = 0$;
- if the CP parity of X is -1, then $a_Z = b_Z = 0$ and $c_Z \neq 0$;
- if the CP parity of X is indefinite, then $a_Z \neq 0$, $c_Z \neq 0$ and/or $b_Z \neq 0$, $c_Z \neq 0$.

Calculating the Lorentz-invariant amplitude $A_{X \rightarrow Z_1^* Z_2^*}(\lambda_1, \lambda_2)$ in a reference frame in which $\mathbf{p}_1 + \mathbf{p}_2 = 0$, we derive that

$$\begin{aligned}
A_{X \rightarrow Z_1^* Z_2^*}(-1, -1) &= 2\sqrt{\sqrt{2}G_F m_Z^2} \left(a_Z - c_Z \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2)}{m_X^2} \right), \\
A_{X \rightarrow Z_1^* Z_2^*}(0, 0) &= -2\sqrt{\sqrt{2}G_F m_Z^2} \left(a_Z \frac{m_X^2 - a_1 - a_2}{2\sqrt{a_1 a_2}} + b_Z \frac{\lambda(m_X^2, a_1, a_2)}{4m_X^2 \sqrt{a_1 a_2}} \right), \\
A_{X \rightarrow Z_1^* Z_2^*}(1, 1) &= 2\sqrt{\sqrt{2}G_F m_Z^2} \left(a_Z + c_Z \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2)}{m_X^2} \right), \\
A_{X \rightarrow Z_1^* Z_2^*}(\lambda_1, \lambda_2) &= 0, \quad \lambda_1 \neq \lambda_2,
\end{aligned} \tag{4}$$

where the function $\lambda(x, y, z)$ is defined in the standard way: $\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$.




Fig. 1: The kinematics of the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$. The momenta of Z_1^* and Z_2^* are shown in a rest frame of X , the momenta of f_1 and \bar{f}_1 (f_2 and \bar{f}_2) are shown in a rest frame of Z_1^* (Z_2^*).

To describe the decay (1), let us introduce the following angles (see Fig. 1): θ_1 (θ_2) is the angle between the momentum of Z_1^* (Z_2^*) in a rest frame of X and the momentum of f_1 (f_2) in a rest frame of Z_1^* (Z_2^*) and φ is the azimuthal angle between the planes of the decays $Z_1^* \rightarrow f_1 \bar{f}_1$ and $Z_2^* \rightarrow f_2 \bar{f}_2$. Further we go into the case of the non-identical fermions, $f_1 \neq f_2$. Using the helicity formalism (see, for example, [25]), we obtain that in the approximation of the massless fermions, $m_{f_1} = m_{f_2} = 0$, the differential decay width of (1) with respect to a_1 , a_2 , θ_1 , θ_2 , φ appears as follows:

$$\begin{aligned}
\frac{d^5 \Gamma}{da_1 da_2 d\theta_1 d\theta_2 d\varphi} &= \frac{\sqrt{2}G_F^3 m_Z^8}{(4\pi)^6 m_X^3} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2) a_1 a_2}{D(a_1)D(a_2)} \\
&\quad \times \sin \theta_1 \sin \theta_2 [(|A_{\parallel}|^2 + |A_{\perp}|^2) ((1 + \cos^2 \theta_1)(1 + \cos^2 \theta_2) + 4A_{f_1} A_{f_2} \cos \theta_1 \cos \theta_2) \\
&\quad + 4|A_0|^2 \sin^2 \theta_1 \sin^2 \theta_2 - 4 \operatorname{Re}(A_{\parallel}^* A_{\perp})(A_{f_1} \cos \theta_1 (1 + \cos^2 \theta_2) + A_{f_2} \cos \theta_2 (1 + \cos^2 \theta_1)) \\
&\quad + 4\sqrt{2} \sin \theta_1 \sin \theta_2 ((\operatorname{Re}(A_0^* A_{\parallel}) \cos \varphi - \operatorname{Im}(A_0^* A_{\perp}) \sin \varphi)(A_{f_1} A_{f_2} + \cos \theta_1 \cos \theta_2) \\
&\quad - (\operatorname{Re}(A_0^* A_{\perp}) \cos \varphi - \operatorname{Im}(A_0^* A_{\parallel}) \sin \varphi)(A_{f_1} \cos \theta_2 + A_{f_2} \cos \theta_1)) \\
&\quad + \sin^2 \theta_1 \sin^2 \theta_2 ((|A_{\parallel}|^2 - |A_{\perp}|^2) \cos 2\varphi - 2 \operatorname{Im}(A_{\parallel}^* A_{\perp}) \sin 2\varphi)],
\end{aligned} \tag{5}$$

where a_f is the projection of the weak isospin of a fermion f , $v_f \equiv a_f - \frac{2q_f}{e} \sin^2 \theta_W$, q_f is the electric charge of the fermion f , e is the electric charge of the positron, θ_W is the weak mixing angle, $D(a_{1,2}) \equiv (a_{1,2} - m_Z^2)^2 + (m_Z \Gamma_Z)^2$,

Γ_Z is the total width of the Z boson, $A_f \equiv \frac{2a_f v_f}{a_f^2 + v_f^2}$,

$$\begin{aligned} A_{\parallel}(a_1, a_2) &\equiv \frac{A_{X \rightarrow Z_1^* Z_2^*}(1, 1) + A_{X \rightarrow Z_1^* Z_2^*}(-1, -1)}{2^{\frac{7}{4}} \sqrt{G_F} m_Z^2} = \sqrt{2} a_Z, \\ A_{\perp}(a_1, a_2) &\equiv \frac{A_{X \rightarrow Z_1^* Z_2^*}(1, 1) - A_{X \rightarrow Z_1^* Z_2^*}(-1, -1)}{2^{\frac{7}{4}} \sqrt{G_F} m_Z^2} = \sqrt{2} c_Z \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2)}{m_X^2}, \\ A_0(a_1, a_2) &\equiv \frac{A_{X \rightarrow Z_1^* Z_2^*}(0, 0)}{2^{\frac{5}{4}} \sqrt{G_F} m_Z^2} = - \left(a_Z \frac{m_X^2 - a_1 - a_2}{2\sqrt{a_1 a_2}} + b_Z \frac{\lambda(m_X^2, a_1, a_2)}{4m_X^2 \sqrt{a_1 a_2}} \right). \end{aligned} \quad (6)$$

Futher the approximation $m_{f_1} = m_{f_2} = 0$ is used. Using Eq. (5), one can connect the ratios of quantities $|A_0|^2$, $|A_{\parallel}|^2 + |A_{\perp}|^2$, $|A_{\parallel}|^2 - |A_{\perp}|^2$, $\text{Re}(A_0^* A_{\parallel})$, $\text{Im}(A_0^* A_{\parallel})$, $\text{Re}(A_0^* A_{\perp})$, $\text{Im}(A_0^* A_{\perp})$, $\text{Re}(A_{\parallel}^* A_{\perp})$, $\text{Im}(A_{\parallel}^* A_{\perp})$ to $|A_0|^2 + |A_{\parallel}|^2 + |A_{\perp}|^2$ with functions of a_1, a_2 which can be measured in experiment. We will call these ratios the helicity coefficients of the decay $X \rightarrow Z_1^* Z_2^*$.

B. A differential width $\frac{d^2 \Gamma}{da_1 da_2}$

The number of the decays

$$h \rightarrow Z_1^* Z_2^* \rightarrow l_1^- l_1^+ l_2^- l_2^+ \quad (l_j = e, \mu), \quad (7)$$

detected in the ATLAS experiment [2] wherein the invariant mass of the four leptons was in the interval [120 GeV, 130 GeV], is equal to 32. The number of the decays (7) detected in the CMS experiment [3] in which the four-lepton invariant mass was within [121.5 GeV, 130.5 GeV], is equal to 25. In view of the insignificant amount of data, at the present time an experimental dependence of the distribution $\frac{1}{\Gamma} \frac{d^5 \Gamma}{da_1 da_2 d\theta_1 d\theta_2 d\varphi}$ (Γ is the total width of the decay (1)) for any of the decays (7) is not available. Let us consider differential decay widths of (1) with respect to four and fewer variables. Integrating Eq. (5) with respect to $\theta_1, \theta_2, \varphi$, we obtain

$$\frac{d^2 \Gamma}{da_1 da_2} = \frac{\sqrt{2} G_F^3 m_Z^8}{9(2\pi)^5 m_X^3} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2) a_1 a_2}{D(a_1) D(a_2)} \sum_{p=0, \parallel, \perp} |A_p|^2. \quad (8)$$

It follows from Eqs. (8), (6) that the dependence of the differential width $\frac{d^2 \Gamma}{da_1 da_2}$ on a_Z, b_Z, c_Z boils down only to the dependence on $|a_Z|, |b_Z|, |c_Z|$ and on $\cos(\arg b_Z - \arg a_Z)$.

The available experimental data on properties of the particle h are close to the corresponding theoretical predictions about the SM Higgs boson (see, for example, [2–4]). That is why $a_{hZ} \approx 1$, $b_{hZ} \approx 0$, $c_{hZ} \approx 0$, where

$$a_{hZ} \equiv a_Z|_{X=h}, \quad b_{hZ} \equiv b_Z|_{X=h}, \quad c_{hZ} \equiv c_Z|_{X=h}.$$

In Fig. 2 we show the differential decay width (8) for $X \rightarrow Z_1^* Z_2^* \rightarrow l_1^- l_1^+ l_2^- l_2^+$ ($l_j = e, \mu, \tau, l_1 \neq l_2$) as a function of $\sqrt{a_1}, \sqrt{a_2}$ in the SM for $|a_Z| = 1$, $b_Z = c_Z = 0$ and $m_X = m_h$, where m_h is the mass of the Higgs boson h . The range of $\sqrt{a_1}, \sqrt{a_2}$ in this plot is determined by the inequalities (2) in the approximation of the massless fermions. In calculations and when plotting graphs the experimental data listed in Table I are used, and $\sin^2 \theta_W = 1 - m_W^2/m_Z^2$, where m_W is the mass of the W boson.




Fig. 2: The dependence of the differential decay width $\frac{d^2\Gamma}{da_1 da_2}$ of $X \rightarrow Z_1^* Z_2^* \rightarrow l_1^- l_1^+ l_2^- l_2^+$ ($l_j = e, \mu, \tau, l_1 \neq l_2$) on $\sqrt{a_1}$ and $\sqrt{a_2}$ in the SM for $m_X = m_h$.

Table I: Values of the Fermi constant, of the masses of h , Z , W and of the total width of Z [26].

$$G_F = 1.1663787(6) \times 10^{-5} \text{ GeV}^{-2}$$

$$m_h = 125.7(4) \text{ GeV}$$

$$m_Z = 91.1876(21) \text{ GeV}$$

$$m_W = 80.385(15) \text{ GeV}$$

$$\Gamma_Z = 2.4952(23) \text{ GeV}$$

As one can see from Fig. 2, in the SM the function $\frac{d^2\Gamma}{da_1 da_2}$ has peaks at $\sqrt{a_1} = m_Z$ and $\sqrt{a_2} = m_Z$, resulting from the quantities $D(a_1)$ and $D(a_2)$ in (8).

Let us calculate the ratio of a typical value of $\frac{d^2\Gamma}{da_1 da_2}$ in the SM on the peaks to its typical value in an area in which $\sqrt{a_1}$ and $\sqrt{a_2}$ significantly differ from m_Z (we will call this area “plateau”). As indicative values of $\sqrt{a_1}$ and $\sqrt{a_2}$ on the peaks we take $\sqrt{a_1} = m_Z$, $\sqrt{a_2} = \frac{1}{2}(m_h - m_Z)$ and $\sqrt{a_1} = \frac{1}{2}(m_h - m_Z)$, $\sqrt{a_2} = m_Z$ (see (2)), and values on the “plateau” are chosen $\sqrt{a_1} = \sqrt{a_2} = \frac{1}{2}m_Z$. It follows from (8) that in the SM for any f_1, f_2 values of $\frac{d^2\Gamma}{da_1 da_2}$ at $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$ are approximately 100 times as great as values of this function on the “plateau”.

If $m_X \neq m_h$ but just greater than m_Z , then $\sqrt{a_1}$ and/or $\sqrt{a_2}$ can be equal to m_Z (according to (2)), and, consequently, in this case the behavior of the function $\frac{d^2\Gamma}{da_1 da_2}$ in the SM is similar to that in case $m_X = m_h$. That is why for any $m_X > m_Z$ and for any final fermions the differential width $\frac{d^2\Gamma}{da_1 da_2}$ in the SM has a sharp maximum at $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$. Therefore, if $|a_Z| \approx 1$, $b_Z \approx 0$, $c_Z \approx 0$ (which is the case of a small distinction between the couplings and their SM values), $\frac{d^2\Gamma}{da_1 da_2}$ also has a sharp maximum at $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$, provided that $m_X > m_Z$.

C. Limits of applicability of the narrow- Z -width approximation

In Refs. [27–29] the accuracy of the narrow-width approximation has been studied for calculation of the total widths of various decays along with the total and differential cross sections of various processes. It is shown that in many cases (especially for processes beyond the SM) this approximation is not applicable. In this connection the question arises whether the narrow- Z -width approximation is applicable for obtaining the differential width $\frac{d\Gamma}{da_2}$ by means of integrating $\frac{d^2\Gamma}{da_1 da_2}$. In this subsection we find the interval of all the a_2 -values for which the approximate integration is valid.

We consider the m_X -values such that $m_X > m_Z$ and the dependences of $a_Z(a_1, a_2)$, $b_Z(a_1, a_2)$, $c_Z(a_1, a_2)$ such that for any f_1 and f_2 $\frac{d^2\Gamma}{da_1 da_2}$ has a sharp maximum when $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$ (an example of such dependences is $|a_Z| \approx 1$, $b_Z \approx 0$, $c_Z \approx 0$). Then while calculating the differential width $\frac{d\Gamma}{da_2}$ one may use the narrow- Z -width approximation:

$$\begin{aligned} \frac{d\Gamma}{da_2} &= \int_0^{(m_X - \sqrt{a_2})^2} da_1 \frac{d^2\Gamma}{da_1 da_2} \approx \int_0^{(m_X - \sqrt{a_2})^2} da_1 \frac{\pi}{m_Z \Gamma_Z} \delta(a_1 - m_Z^2) f(a_1, a_2) \\ &= \frac{\pi}{m_Z \Gamma_Z} f(m_Z^2, a_2) \quad \forall \sqrt{a_2} \in (0, m_X - m_Z - \Delta], \end{aligned} \quad (9)$$

where Δ is some positive quantity and

$$f(a_1, a_2) \equiv \frac{\sqrt{2} G_F^3 m_Z^8}{9(2\pi)^5 m_X^3} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \frac{\lambda^{\frac{1}{2}}(m_X^2, a_1, a_2) a_1 a_2}{D(a_2)} \sum_{p=0, \parallel, \perp} |A_p|^2. \quad (10)$$

$\Delta > 0$ since in Eq. (9) one may use the approximation $\frac{d^2\Gamma}{da_1 da_2} = \frac{\pi}{m_Z \Gamma_Z} \delta(a_1 - m_Z^2) f(a_1, a_2)$ only when $\sqrt{a_2} < m_X - m_Z$, because if $\sqrt{a_2}$ approaches $m_X - m_Z$, the peak of $\frac{d^2\Gamma}{da_1 da_2}$ at $\sqrt{a_1} = m_Z$ gets less sharp and at $\sqrt{a_2} = m_X - m_Z$ the peak disappears (see Fig. 2 and Eq. (8)). However, the derivation (9) does not allow one to estimate the accuracy of the formula $\frac{d\Gamma}{da_2} \approx \frac{\pi}{m_Z \Gamma_Z} f(m_Z^2, a_2)$ at a given value of $\sqrt{a_2}$, and for this reason it is not clear what value of Δ should be chosen.

To clarify this point, let us derive the formula for $\frac{d\Gamma}{da_2}$ in the following way:

$$\begin{aligned} \frac{d\Gamma}{da_2} &= \int_0^{(m_X - \sqrt{a_2})^2} da_1 \frac{d^2\Gamma}{da_1 da_2} \approx \int_{m_Z^2 - \varepsilon_1}^{m_Z^2 + \varepsilon_2} da_1 \frac{d^2\Gamma}{da_1 da_2} = \int_{m_Z^2 - \varepsilon_1}^{m_Z^2 + \varepsilon_2} da_1 \frac{f(a_1, a_2)}{(a_1 - m_Z^2)^2 + (m_Z \Gamma_Z)^2} \\ &\approx \int_{m_Z^2 - \varepsilon_1}^{m_Z^2 + \varepsilon_2} da_1 \frac{f(m_Z^2, a_2)}{(a_1 - m_Z^2)^2 + (m_Z \Gamma_Z)^2} = \frac{\arctan \frac{\varepsilon_2}{m_Z \Gamma_Z} + \arctan \frac{\varepsilon_1}{m_Z \Gamma_Z}}{m_Z \Gamma_Z} f(m_Z^2, a_2) \\ &\approx \frac{\pi}{m_Z \Gamma_Z} f(m_Z^2, a_2), \end{aligned} \quad (11)$$

where ε_1 and ε_2 are some positive quantities such that $m_Z \Gamma_Z \ll \varepsilon_j \ll m_Z^2$, the variable a_2 takes values in the interval $\left(0, \left(m_X - \sqrt{m_Z^2 + \varepsilon_2}\right)^2\right]$.

One of the approximations used in Eq. (11) is the switch from the integration over an interval $(0, (m_X - \sqrt{a_2})^2)$ to the integration over an interval $(m_Z^2 - \varepsilon_1, m_Z^2 + \varepsilon_2)$. Thus, $m_Z^2 - \varepsilon_1$ has to be greater than or equal to $4m_{f_1}^2$ (which holds true since $\varepsilon_1 \ll m_Z^2$) and $m_Z^2 + \varepsilon_2$ has to be less than or equal to $(m_X - \sqrt{a_2})^2$, i.e. $a_2 \leq (m_X - \sqrt{m_Z^2 + \varepsilon_2})^2$. The latter inequality restricts the interval of all the a_2 -values for which these approximations are applicable. Consequently, in order to apply them for as long an interval of a_2 -values as possible, one should use the minimal ε_2 -value at which the approximations are valid.

While obtaining (11), we also used an approximation $A \approx \pi$ ($A \equiv \arctan \frac{\varepsilon_2}{m_Z \Gamma_Z} + \arctan \frac{\varepsilon_1}{m_Z \Gamma_Z}$). Let us define ε_1 as $\varepsilon_1 \equiv m_Z \sqrt{m_Z \Gamma_Z}$ (so as $\frac{\varepsilon_1}{m_Z \Gamma_Z} = \frac{m_Z^2}{\varepsilon_1}$). The values of quantities A and $m_h - \sqrt{m_Z^2 + \varepsilon_2}$ which are listed in Table II specify for the considered ε_2 -values the accuracy of the approximation $A \approx \pi$ and the maximal value of $\sqrt{a_2}$ at which the narrow- Z -width approximation is applicable in case $X = h$.

Table II: Values of A ($\varepsilon_1 \equiv m_Z \sqrt{m_Z \Gamma_Z}$) and of $m_h - \sqrt{m_Z^2 + \varepsilon_2}$ at various values of ε_2 .

ε_2	A	$m_h - \sqrt{m_Z^2 + \varepsilon_2}$ (GeV)
0	0.45π	34.51
$m_Z \Gamma_Z$	0.70π	33.27
$2m_Z \Gamma_Z$	0.80π	32.05
$3m_Z \Gamma_Z$	0.85π	30.84
$4m_Z \Gamma_Z$	0.87π	29.65

According to Table II, if $\varepsilon_2 < 3m_Z \Gamma_Z$, then $A < 0.85\pi$ and, in view of the big difference between A and π , we will not apply the approximations (11) for such values of ε_2 . Hence we will use $\varepsilon_2 = 3m_Z \Gamma_Z$. It follows from (11) that

$$\frac{d\Gamma}{da_2} \approx \frac{\sqrt{2}G_F^3 m_Z^9}{9 \cdot 2^5 \pi^4 m_X^3 \Gamma_Z} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \frac{\lambda^{\frac{1}{2}}(m_X^2, m_Z^2, a_2)a_2}{D(a_2)} \sum_{p=0, \parallel, \perp} |A'_p|^2 \quad (12)$$

$$\forall \sqrt{a_2} \in \left(0, m_X - \sqrt{m_Z^2 + \varepsilon_2}\right],$$

where $A'_p \equiv A_p(m_Z^2, a_2)$ ($p = 0, \parallel, \perp$).

Note that in Refs. [14, 16, 17] when plotting dependences of $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$ on $\sqrt{a_2}$, formulas for $\frac{d\Gamma}{da_2}$ which correspond to (12) have been used, but these graphs have been plotted for $\sqrt{a_2} \leq m_X - m_Z$, despite the fact that Eq. (11) is not valid at $\varepsilon_2 = 0$ (see Table II), and, therefore, the plotted dependences significantly differ from the true ones in the interval $\sqrt{a_2} \in (m_X - \sqrt{m_Z^2 + 3m_Z \Gamma_Z}, m_X - m_Z]$.

D. An inequality constraining a'_{hZ} , b'_{hZ} , c'_{hZ} from CMS data

According to [3],

$$\frac{\sigma(pp \rightarrow h) \frac{\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)}{\Gamma_h}}{\sigma_{SM}(pp \rightarrow h) \frac{\Gamma_{SM}(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)}{\Gamma_{hSM}}} = 0.93^{+0.26}_{-0.23}(\text{stat})^{+0.13}_{-0.09}(\text{syst}), \quad (13)$$

where $\sigma(pp \rightarrow h)$ is the cross section for production of h in pp collisions,

$$\begin{aligned} \Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l) &\equiv \Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4e) + \Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4\mu) + \Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 2e2\mu) \\ &= 2\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4e) + \Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 2e2\mu), \end{aligned} \quad (14)$$

Γ_h is the total width of the boson h , $\sigma_{SM}(pp \rightarrow h)$, $\Gamma_{SM}(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)$, Γ_{hSM} are the predictions of the SM for respectively $\sigma(pp \rightarrow h)$, $\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)$, Γ_h at $m_h = 125.6$ GeV. Obtaining (13), the CMS collaboration has combined data from pp collisions corresponding to an integrated luminosity of 5.1 fb^{-1} at a center-of-mass energy $\sqrt{s} = 7$ TeV and 19.7 fb^{-1} at $\sqrt{s} = 8$ TeV.

We consider the case in which the functions $|a'_{hZ}|$, $|b'_{hZ}|$, $|c'_{hZ}|$, $\cos(\arg b'_{hZ} - \arg a'_{hZ})$ do not depend on a_2 . Here we define

$$a'_{hZ} \equiv a_{hZ}(m_Z^2, a_2), \quad b'_{hZ} \equiv b_{hZ}(m_Z^2, a_2), \quad c'_{hZ} \equiv c_{hZ}(m_Z^2, a_2).$$

Then using the approximation

$$\frac{\sigma(pp \rightarrow h)}{\Gamma_h} \approx \frac{\sigma_{SM}(pp \rightarrow h)}{\Gamma_{hSM}} \quad (15)$$

and Eqs. (13) (within one standard deviation), (A10), (A13) (see Appendix A), we derive the relation

$$|a'_{hZ}|^2 + 0.015 |b'_{hZ}|^2 + 0.177 \text{Re}(a'^*_{hZ} b'_{hZ}) + 0.037 |c'_{hZ}|^2 \in [0.68, 1.22]. \quad (16)$$

While obtaining (16) we plugged the central values of m_h , m_Z , Γ_Z listed in Table I into Eq. (A10). Note that the latter equation is derived at tree level and without allowance for the interference term connected with the permutation of the identical fermions in case $f_1 = f_2$. The interference contribution to $\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)$ at tree level is expected to be negligible since in the SM at $m_h = 140$ GeV it amounts to 2.99% (see Table 1 in Ref. [30]).

Using the data of Table III and considering two sigma errors where available, we obtain that at $\sqrt{s} = 8$ TeV

$$\frac{\sigma(pp \rightarrow h)/\Gamma_h}{\sigma_{SM}(pp \rightarrow h)/\Gamma_{hSM}} \in (0.17, \infty), \quad (17)$$

which means that the approximation (15) does not contradict the experimental limits.

Moreover, assuming that all the couplings of the Higgs boson except for a_{hZ} , b_{hZ} and c_{hZ} are equal to their SM values, we can verify (15). In this case the only anomalous contribution to Γ_h comes from $\Gamma(h \rightarrow Z_1^* Z_2^*)$, which makes up, in the SM, only about 2.81% [34] of the total Higgs boson width, and therefore Γ_h is unlikely to

Table III: Experimental and theoretical results for the total production cross-section of the Higgs boson in pp collisions and for its total width.

$\sigma(pp \rightarrow h) = 33.0 \pm 5.3(\text{stat}) \pm 1.6(\text{syst}) \text{ pb at } \sqrt{s} = 8 \text{ TeV [31]}$
$\sigma_{SM}(pp \rightarrow h) = 22.09 \text{ pb (uncertainties not available) at } \sqrt{s} = 8 \text{ TeV [32]}$
$\Gamma_h < 22 \text{ MeV at 95\% confidence level (CL) [33]}$
$\Gamma_{hSM} = 4.15 \pm 0.16 \text{ MeV [34]}$

substantially differ from its SM prediction. Besides, the inequality (16) means that $|\Gamma(h \rightarrow Z_1^* Z_2^*) - \Gamma_{SM}(h \rightarrow Z_1^* Z_2^*)|/\Gamma_{SM}(h \rightarrow Z_1^* Z_2^*) \in [0, 0.32]$ because its left-hand side is

$$\frac{\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)}{\Gamma_{SM}(h \rightarrow Z_1^* Z_2^* \rightarrow 4l)} = \frac{\Gamma(h \rightarrow Z_1^* Z_2^*)}{\Gamma_{SM}(h \rightarrow Z_1^* Z_2^*)}$$

(see (13), (A10)). For this reason (16) implies that the relative change of Γ_h is less than $2.81\% \cdot 0.32 \approx 0.90\%$, and, consequently, (16) is consistent with the approximation $\Gamma_h \approx \Gamma_{hSM}$.

The dominant contribution to the Higgs boson production cross section $\sigma_{SM}(pp \rightarrow h)$ comes from the gluon fusion process $gg \rightarrow h$, which is independent of the hZZ vertex. The processes involving the hZZ interaction, i.e. the Higgs-strahlung Zh and the Z boson fusion, constitute much less parts of $\sigma_{SM}(pp \rightarrow h)$. Specifically, at $\sqrt{s} = 8 \text{ TeV}$ they can be estimated as 0.41 pb and 0.70 pb respectively [32]. The total production cross section at this energy is 22.09 pb (see Table III), so the processes of interest contribute about 5% of the total cross section. That is why it seems improbable that the couplings a_{hZ} , b_{hZ} and c_{hZ} provide a significant difference between $\sigma(pp \rightarrow h)$ and $\sigma_{SM}(pp \rightarrow h)$. However, a derivation of the dependence of the total production cross section on the hZZ couplings would require a separate study.

Summarizing the discussion of the approximation (15), we can infer, firstly, that it is consistent with the available data [31, 33] and, secondly, under the assumption that the only anomalous Higgs boson couplings are related to the hZZ vertex, Eq. (15) is most likely to be valid due to the small contributions of the hZZ vertex to Γ_h and $\sigma(pp \rightarrow h)$.

E. Constraints on a'_{hZ} , b'_{hZ} , c'_{hZ}

The inequality (16) constrains the whole six-dimensional space formed by the real and imaginary parts of the couplings a'_{hZ} , b'_{hZ} and c'_{hZ} to the set of ellipsoids allowed by (13). Note that a similar interpretation has been suggested in Ref. [35].

From (16) it follows that the variant $a'_{hZ} = 0, b'_{hZ} = 0, c'_{hZ} = 1$ (negative CP parity of the boson h) is excluded. Now let us find constraints on the values of b'_{hZ} and c'_{hZ} , assuming that a'_{hZ} is taken from the SM, i.e. $|a'_{hZ}| = 1$

or $a'_{hZ} = 1$. Then

$$|a'_{hZ}| = 1 \quad \text{and} \quad b'_{hZ} = 0 \Rightarrow |c'_{hZ}| \in [0, 2.44]; \quad (18a)$$

$$a'_{hZ} = 1 \quad \text{and} \quad c'_{hZ} = 0 \quad \text{and} \quad \text{Im } b'_{hZ} = 0 \Rightarrow b'_{hZ} \in ([-12.66, -9.31] \cup [-2.22, 1.14]), \quad (18b)$$

$$a'_{hZ} = 1 \quad \text{and} \quad c'_{hZ} = 0 \quad \text{and} \quad \text{Re } b'_{hZ} = 0 \Rightarrow \text{Im } b'_{hZ} \in [-3.84, 3.84]. \quad (18c)$$

Let us compare (18) with the hZZ coupling constraints obtained by the CMS [36] and ATLAS [37] collaborations. For this purpose we first express our XZZ couplings in terms of the CMS ones $\tilde{a}_1, \tilde{a}_2, \tilde{a}_3$ (we denote a_1, a_2, a_3 from [36] as $\tilde{a}_1, \tilde{a}_2, \tilde{a}_3$ to avoid confusion):

$$a_Z = \tilde{\alpha} \left(\tilde{a}_1 - \exp(i\phi_{\Lambda_1}) \frac{a_1 + a_2}{\Lambda_1^2} + \frac{m_X^2 - a_1 - a_2}{m_Z^2} \tilde{a}_2 \right), \quad (19a)$$

$$b_Z = -2\tilde{\alpha} \frac{m_X^2}{m_Z^2} \tilde{a}_2, \quad (19b)$$

$$c_Z = -i\tilde{\alpha} \frac{m_X^2}{m_Z^2} \tilde{a}_3, \quad (19c)$$

where $\tilde{\alpha} \equiv \alpha_0 v/2$, α_0 is the proportionality factor of the amplitude $A(HZZ)$ of the transition $X \rightarrow Z_1^* Z_2^*$ (see Eq. (1) in [36]), $v \equiv 1/\sqrt{\sqrt{2}G_F}$ is the vacuum expectation value of the Higgs field, Λ_1 is a scale of physics beyond the SM, ϕ_{Λ_1} is the phase in the term with Λ_1 . In general $\tilde{a}_1, \tilde{a}_2, \tilde{a}_3$ may depend on a_1 and a_2 , however in [36] they are set to be constant. The ATLAS XZZ couplings $\alpha, \kappa_{SM}, \kappa_{HZZ}, \kappa_{AZZ}$ are related to the CMS ones in the following way:

$$\tilde{\alpha} \left(\tilde{a}_1 - \exp(i\phi_{\Lambda_1}) \frac{a_1 + a_2}{\Lambda_1^2} \right) = \kappa_{SM} \cos \alpha, \quad \tilde{\alpha} \tilde{a}_2 = \frac{v}{4\Lambda} \kappa_{HZZ} \cos \alpha, \quad \tilde{\alpha} \tilde{a}_3 = \frac{v}{4\Lambda} \kappa_{AZZ} \sin \alpha, \quad (20)$$

where Λ is the EFT energy scale. Note that comparing the Lagrangian (1) in [37] with the one describing the interaction of the SM Higgs field with ZZ and W^-W^+ , one can deduce that the coupling g_{HZZ} from (1) in [37] is equal to $2m_Z^2/v$. In [37] the couplings $\alpha, \kappa_{SM}, \kappa_{HZZ}, \kappa_{AZZ}$ are considered constant and real.

In Refs. [36, 37] 95% CL allowed regions for hZZ couplings are reported (see Table IV). Note that

$$\tilde{\kappa}_{HZZ} \equiv \frac{v}{4\Lambda} \kappa_{HZZ}, \quad \tilde{\kappa}_{AZZ} \equiv \frac{v}{4\Lambda} \kappa_{AZZ}, \quad (21)$$

$$\frac{\tilde{\kappa}_{HZZ}}{\kappa_{SM}} = -\frac{m_Z^2 b_Z}{2m_X^2 a_Z + (m_X^2 - a_1 - a_2) b_Z}, \quad \frac{\tilde{\kappa}_{AZZ}}{\kappa_{SM}} \tan \alpha = \frac{2im_Z^2 c_Z}{2m_X^2 a_Z + (m_X^2 - a_1 - a_2) b_Z} \quad (22)$$

and in the limit $\Lambda_1 \rightarrow \infty$ the CMS and ATLAS ratios coincide:

$$\frac{\tilde{a}_2}{\tilde{a}_1} = \lim_{\Lambda_1 \rightarrow \infty} \frac{\tilde{\kappa}_{HZZ}}{\kappa_{SM}}, \quad \frac{\tilde{a}_3}{\tilde{a}_1} = \lim_{\Lambda_1 \rightarrow \infty} \left(\frac{\tilde{\kappa}_{AZZ}}{\kappa_{SM}} \tan \alpha \right). \quad (23)$$

Following [37], we assume the ATLAS hZZ couplings to be constant. Then considering the case $\kappa_{HZZ} = 0$, we find that our couplings a_{hZ}, b_{hZ}, c_{hZ} are constant as well (see (19), (20)), and using (18a) we obtain an allowed interval for $\tilde{\kappa}_{AZZ} \tan \alpha / \kappa_{SM}$ (see Table V). However, in case $\kappa_{HZZ} = 0$ the results (18b) and (18c) only show that h may be the SM Higgs boson, and thus they do not constrain any hZZ couplings.

Table IV: The CMS [36] and ATLAS [37] 95% CL allowed regions for hZZ couplings. The last row shows the conditions under which these regions have been derived.

CMS		ATLAS	
$\frac{\tilde{a}_2}{\tilde{a}_1}$	$\frac{\tilde{a}_3}{\tilde{a}_1}$	$\frac{\tilde{\kappa}_{HZZ}}{\kappa_{SM}}$	$\frac{\tilde{\kappa}_{AZZ}}{\kappa_{SM}} \tan \alpha$
$[-2.28, -1.88] \cup [-0.69, \infty)$	$[-2.05, 2.19]$	$(-0.75, 2.45)$	$(-2.85, 0.95)$
$\text{Im} \frac{\tilde{a}_2}{\tilde{a}_1} = 0, \phi_{\Lambda_1} = 0 \text{ or } \pi$		$\kappa_{AZZ} = 0$	$\kappa_{HZZ} = 0$

If $\kappa_{HZZ} \neq 0$, then a_{hZ} acquires a dependence on the invariant masses squared a_1 and a_2 , and therefore the constraints (16) and (18) get invalid since they have been derived under the assumption that $|a'_{hZ}|$ is independent of a_2 . Therefore to constrain the ATLAS couplings in case $\kappa_{HZZ} \neq 0$, we start with Eqs. (13) and (15), which demonstrate that within one standard deviation

$$\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4l) / \Gamma_{SM}(h \rightarrow Z_1^* Z_2^* \rightarrow 4l) \in [0.68, 1.22].$$

To obtain Γ we have to calculate the integral (A1) for a'_Z depending on a_2 . Taking into account the limits of the integration, we substitute a_2 with $(m_X - m_Z)^2/2$ in the expression for a'_Z (see (19a), (20)) and therefore derive Eq. (A10) where a_Z has the expression (19a) with $a_2 = (m_X - m_Z)^2/2$. It means that if κ_{HZZ} is not zero, we may use (16) and (18b), (18c) with a'_{hZ} determined by Eq. (19a) where a_2 is replaced by $(m_h - m_Z)^2/2$. This conclusion allows us to constrain $\tilde{\kappa}_{HZZ}/\kappa_{SM}$ and $\text{Im} \tilde{\kappa}_{HZZ}/\text{Re} \kappa_{SM}$, as one can see in Table V.

Table V: Our allowed regions for the ATLAS hZZ couplings. The last two rows show the conditions under which these regions have been derived.

$\frac{\tilde{\kappa}_{HZZ}}{\kappa_{SM}}$	$\frac{\tilde{\kappa}_{AZZ}}{\kappa_{SM}} \tan \alpha$	$\frac{\text{Im} \tilde{\kappa}_{HZZ}}{\text{Re} \kappa_{SM}}$
$[-2.38, -1.89] \cup [-0.24, 1.13]$	$[-1.28, 1.28]$	$[-1.01, 1.01]$
$a_2 = (m_h - m_Z)^2/2$ in (19a),	$ a'_{hZ} = 1, \kappa_{HZZ} = 0, \text{Im} \frac{\tilde{\kappa}_{AZZ}}{\kappa_{SM}} = 0$	$a_2 = (m_h - m_Z)^2/2$ in (19a),
$a'_{hZ} = 1, \kappa_{AZZ} \sin \alpha = 0, \text{Im} \kappa_{HZZ} = 0$		$a'_{hZ} = 1, \kappa_{AZZ} \sin \alpha = 0, \text{Re} \kappa_{HZZ} = 0$

Note that the results (16), (18) along with the regions shown in Table V are estimated with consideration of the one sigma interval in (13), with the approximation (15), the central values of m_h, m_Z, Γ_Z from Table I and Eq. (A10). Comparing Tables IV and V, one notices significant overlaps between the constraints reported in papers [36, 37] and our ones. In addition, we present an allowed interval for the ratio $\text{Im} \tilde{\kappa}_{HZZ}/\text{Re} \kappa_{SM}$ unconstrained in Refs. [36, 37].




Fig. 3: The distribution $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$ as a function of $\sqrt{a_2}$ for the decay $h \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ in case $|a'_{hZ}| = 1, b'_{hZ} = 0, c'_{hZ} = 0$ (solid line); $|a'_{hZ}| = 1, b'_{hZ} = 0, |c'_{hZ}| = 0.5$ (dashed line); $a'_{hZ} = 1, b'_{hZ} = -0.5, c'_{hZ} = 0$ (dash-dotted line).

We choose the following sets of values of a'_{hZ} , b'_{hZ} and c'_{hZ} :

$$\begin{aligned}
 |a'_{hZ}| &= 1, \quad b'_{hZ} = 0, \quad c'_{hZ} = 0, \\
 a'_{hZ} &= 1, \quad b'_{hZ} = 0, \quad c'_{hZ} = 0.5, \\
 a'_{hZ} &= 1, \quad b'_{hZ} = 0, \quad c'_{hZ} = 0.5i, \\
 a'_{hZ} &= 1, \quad b'_{hZ} = -0.5, \quad c'_{hZ} = 0
 \end{aligned} \tag{24}$$

and

$$a'_{hZ} = 1, \quad b'_{hZ} = -0.5i, \quad c'_{hZ} = 0, \tag{25}$$

which are consistent with the constraints (18). The sets (24) and (25) will be used for examination of further results.

Regarding the selected values in (24) and (25) one should mention that even in the SM the couplings b_{hZ} and c_{hZ} acquire small values due to electroweak radiative corrections where $\text{Im } b_{hZ}$ and $\text{Im } c_{hZ}$ come from the absorptive parts of the corresponding loop diagrams. In Eqs. (24), (25) we assume that the hZZ vertex may be significantly modified by physics beyond the SM.

It is of interest to study the distribution $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$ as a function of $\sqrt{a_2}$ for various sets of a'_Z , b'_Z , c'_Z . Here $a'_Z \equiv a_Z(m_Z^2, a_2)$, $b'_Z \equiv b_Z(m_Z^2, a_2)$, $c'_Z \equiv c_Z(m_Z^2, a_2)$. In accordance with (8), the function $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$ is independent of the final fermion state. Figure 3 shows this observable in case $X = h$.

As one can see from Fig. 3, the function $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$ is sensitive to b'_{hZ} and almost insensitive to c'_{hZ} . For this reason, having measured this distribution with sufficient accuracy, one can get significant constraints on the values of b'_{hZ} . However, one should keep in mind that this conclusion is obtained for the case in which $|a'_{hZ}|$, $|b'_{hZ}|$, $|c'_{hZ}|$ and

$\cos(\arg b'_{hZ} - \arg a'_{hZ})$ are independent of a_2 , and their a_2 -dependence can considerably modify the dependence of $\frac{1}{\Gamma} \frac{d\Gamma}{da_2}$. In Sec. II F we develop methods of getting constraints on the dependences of a'_Z, b'_Z, c'_Z on a_2 .

F. Connection between the helicity coefficients of the decay $X \rightarrow Z_1^* Z_2^*$ and observables

Let us consider now arbitrary dependences of $a_Z(a_1, a_2), b_Z(a_1, a_2), c_Z(a_1, a_2)$ such that the differential width $\frac{d^5\Gamma}{da_1 da_2 d\theta_1 d\theta_2 d\varphi}$ has a sharp maximum as a function of a_1 and a_2 at $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$ for any f_1, f_2 . From Eq. (5), using approximations analogous to those used when deriving the formulas (11), (A1), we carry out integration over a_1 and some of the angular variables. Then we obtain the following relations between observables $O_i(a_2)$ and the helicity coefficients:

$$\begin{aligned} O_1^{(1)}(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\theta_1 \frac{d^2\Gamma}{da_2 d\theta_1} - \int_{\frac{\pi}{2}}^{\pi} d\theta_1 \frac{d^2\Gamma}{da_2 d\theta_1} \right) = -\frac{3}{2} A_{f_1} \frac{\text{Re}(A'_{\parallel}{}^* A'_{\perp})}{\sum_p |A'_p|^2}, \\ O_1^{(2)}(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\theta_2 \frac{d^2\Gamma}{da_2 d\theta_2} - \int_{\frac{\pi}{2}}^{\pi} d\theta_2 \frac{d^2\Gamma}{da_2 d\theta_2} \right) = -\frac{3}{2} A_{f_2} \frac{\text{Re}(A'_{\parallel}{}^* A'_{\perp})}{\sum_p |A'_p|^2}, \end{aligned} \quad (26)$$

under the condition $\sqrt{a_2} \in (0, m_X - \sqrt{m_Z^2 + \varepsilon_2}]$.

One can write these two formulas in the following way:

$$O_1^{(1,2)}(a_2) \equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\theta_{1,2} \frac{d^2\Gamma}{da_2 d\theta_{1,2}} - \int_{\frac{\pi}{2}}^{\pi} d\theta_{1,2} \frac{d^2\Gamma}{da_2 d\theta_{1,2}} \right) = -\frac{3}{2} A_{f_{1,2}} \frac{\text{Re}(A'_{\parallel}{}^* A'_{\perp})}{\sum_p |A'_p|^2}. \quad (27)$$

Then we deduce that

$$\begin{aligned} O_2(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_{\frac{\pi}{2}-\beta}^{\frac{\pi}{2}-\alpha} d\theta_2 \frac{d^2\Gamma}{da_2 d\theta_2} + \int_{\frac{\pi}{2}+\alpha}^{\frac{\pi}{2}+\beta} d\theta_2 \dots \right) = \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_{\frac{\pi}{2}-\beta}^{\frac{\pi}{2}-\alpha} d\theta_1 \frac{d^2\Gamma}{da_2 d\theta_1} + \int_{\frac{\pi}{2}+\alpha}^{\frac{\pi}{2}+\beta} d\theta_1 \dots \right) \\ &= \frac{1}{4} \left((\sin \beta - \sin \alpha)(3 + \sin^2 \alpha + \sin^2 \beta + \sin \alpha \sin \beta) \right. \\ &\quad \left. + 3 \frac{|A'_0|^2}{\sum_p |A'_p|^2} (\sin \beta \cos^2 \beta - \sin \alpha \cos^2 \alpha) \right), \\ &\quad 0 \leq \alpha < \beta \leq \frac{\pi}{2}, \end{aligned} \quad (28)$$

$$\begin{aligned} O_3(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\theta_2 \left(\int_0^{\frac{\pi}{2}} d\theta_1 \frac{d^3\Gamma}{da_2 d\theta_1 d\theta_2} - \int_{\frac{\pi}{2}}^{\pi} d\theta_1 \frac{d^3\Gamma}{da_2 d\theta_1 d\theta_2} \right) - \int_{\frac{\pi}{2}}^{\pi} d\theta_2 \dots \right) \\ &= \frac{9}{16} A_{f_1} A_{f_2} \frac{|A'_{\parallel}|^2 + |A'_{\perp}|^2}{\sum_p |A'_p|^2}, \end{aligned} \quad (29)$$

$$\begin{aligned} O_4(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{4}} d\varphi \frac{d^2\Gamma}{da_2 d\varphi} - \int_{\frac{\pi}{4}}^{\frac{3}{4}\pi} d\varphi \dots + \int_{\frac{3}{4}\pi}^{\frac{5}{4}\pi} d\varphi \dots - \int_{\frac{5}{4}\pi}^{\frac{7}{4}\pi} d\varphi \dots + \int_{\frac{7}{4}\pi}^{2\pi} d\varphi \dots \right) \\ &= \frac{1}{2\pi} \frac{|A'_{\parallel}|^2 - |A'_{\perp}|^2}{\sum_p |A'_p|^2}, \end{aligned} \quad (30)$$

$$O_5(a_2) \equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\varphi \frac{d^2\Gamma}{da_2 d\varphi} - \int_{\frac{\pi}{2}}^{\pi} d\varphi \dots + \int_{\pi}^{\frac{3}{2}\pi} d\varphi \dots - \int_{\frac{3}{2}\pi}^{2\pi} d\varphi \dots \right) = -\frac{1}{\pi} \frac{\text{Im}(A_0'^* A_\perp')}{\sum_p |A_p'|^2}, \quad (31)$$

$$O_6(a_2) \equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\varphi \frac{d^2\Gamma}{da_2 d\varphi} - \int_{\frac{\pi}{2}}^{\frac{3}{2}\pi} d\varphi \dots + \int_{\frac{3}{2}\pi}^{2\pi} d\varphi \dots \right) = \frac{9}{32} \sqrt{2} \pi A_{f_1} A_{f_2} \frac{\text{Re}(A_0'^* A_\parallel')}{\sum_p |A_p'|^2}, \quad (32)$$

$$\begin{aligned} O_7^{(1,2)}(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\pi} d\varphi \left(\int_0^{\frac{\pi}{2}} d\theta_{1,2} \frac{d^3\Gamma}{da_2 d\theta_{1,2} d\varphi} - \int_{\frac{\pi}{2}}^{\pi} d\theta_{1,2} \frac{d^3\Gamma}{da_2 d\theta_{1,2} d\varphi} \right) - \int_{\pi}^{2\pi} d\varphi \dots \right) \\ &= \frac{3}{8} \sqrt{2} A_{f_{2,1}} \frac{\text{Im}(A_0'^* A_\parallel')}{\sum_p |A_p'|^2}, \end{aligned} \quad (33)$$

$$\begin{aligned} O_8^{(1,2)}(a_2) &\equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\frac{\pi}{2}} d\varphi \left(\int_0^{\frac{\pi}{2}} d\theta_{1,2} \frac{d^3\Gamma}{da_2 d\theta_{1,2} d\varphi} - \int_{\frac{\pi}{2}}^{\pi} d\theta_{1,2} \frac{d^3\Gamma}{da_2 d\theta_{1,2} d\varphi} \right) - \int_{\frac{\pi}{2}}^{\frac{3}{2}\pi} d\varphi \dots + \int_{\frac{3}{2}\pi}^{2\pi} d\varphi \dots \right) \\ &= -\frac{3}{8} \sqrt{2} A_{f_{2,1}} \frac{\text{Re}(A_0'^* A_\perp')}{\sum_p |A_p'|^2}, \end{aligned} \quad (34)$$

$$O_9(a_2) \equiv \left(\frac{d\Gamma}{da_2}\right)^{-1} \left(\int_0^{\pi} d\varphi \frac{d^2\Gamma}{da_2 d\varphi} - \int_{\pi}^{2\pi} d\varphi \dots \right) = -\frac{9}{32} \sqrt{2} \pi A_{f_1} A_{f_2} \frac{\text{Im}(A_0'^* A_\perp')}{\sum_p |A_p'|^2}. \quad (35)$$

From the measured observables $O_i(a_2)$ one can get constraints on the dependences of the couplings $a_Z'(a_2)$, $b_Z'(a_2)$ and $c_Z'(a_2)$. As for $O_2(a_2)$, it can be measured at a fixed value of $\sqrt{a_2}$ and at various values of the parameters β and α . Then, after obtaining central values and uncertainties of a quantity $|A_0'|^2/\sum_p |A_p'|^2$ from Eq. (28) at several sets of values of β , α , one can combine these central values and uncertainties and thereby get a value of $|A_0'|^2/\sum_p |A_p'|^2$ with greater precision than in case of any particular values of β , α .

As an illustration of the behavior of these observables, in Fig. 4 we show their $\sqrt{a_2}$ -dependence with the constant a_{hZ}' , b_{hZ}' , c_{hZ}' from the sets (24). The observable $O_2(a_2)$ is presented for $\beta = 90^\circ$ and $\alpha = 70^\circ$.

As one can see, for each of the observables $O_2(a_2)$, $O_3(a_2)$, $O_4(a_2)$, $O_6(a_2)$ their dependences on $\sqrt{a_2}$ for all the four sets (24) are very close. The observables $O_2(a_2)$ (at $\beta = 90^\circ$, $\alpha = 70^\circ$) and $O_4(a_2)$ are relatively large with the maximum values greater than 0.05, and thus to measure these observables a relatively small amount of data is needed, while $O_3(a_2)$ and $O_6(a_2)$ are smaller, which complicates their experimental observation.

Further, $O_1^{(1,2)}(a_2)$, $O_5(a_2)$, $O_8^{(1,2)}(a_2)$, $O_9(a_2)$ vanish for $c_{hZ}'(a_2) = 0$, according to Eqs. (27), (31), (34), (35) and Fig. 4. Therefore, these observables can give significant constraints on the CP -odd coupling $c_{hZ}'(a_2)$, although their moduli are relatively small.

The functions $O_7^{(1,2)}(a_2)$ are proportional to $\text{Im}(a_{hZ}' b_{hZ}')$ (see (33), (6)), and, consequently, they are equal to zero for any set from (24). Among all the observables under consideration, $O_7^{(1,2)}(a_2)$ are the only ones vanishing in case $b_{hZ}'(a_2) = 0$ for any $a_{hZ}'(a_2)$ and $c_{hZ}'(a_2)$. Therefore, knowing the dependences $O_7^{(1,2)}(a_2)$ allows one to get notable constraints on the function $b_{hZ}'(a_2)$. Although in case (25) these observables turn out to be relatively small in absolute value (see Fig. 5).




Fig. 4: The observables $O_1^{(1,2)}$, O_2 (at $\beta = 90^\circ$, $\alpha = 70^\circ$), O_3 , O_4 , O_5 , O_6 , $O_8^{(1,2)}$, O_9 for the decay $h \rightarrow Z_1^* Z_2^* \rightarrow l_1^- l_1^+ l_2^- l_2^+$ ($l_j = e, \mu, \tau, l_1 \neq l_2$) as functions of $\sqrt{a_2}$ in case $|a'_{hZ}| = 1, b'_{hZ} = 0, c'_{hZ} = 0$ (solid lines); $a'_{hZ} = 1, b'_{hZ} = 0, c'_{hZ} = 0.5$ (dashed lines); $a'_{hZ} = 1, b'_{hZ} = 0, c'_{hZ} = 0.5i$ (dash-dotted lines); $a'_{hZ} = 1, b'_{hZ} = -0.5, c'_{hZ} = 0$ (dotted lines).




Fig. 5: The observables $O_7^{(1,2)}$ for the decay $h \rightarrow Z_1^* Z_2^* \rightarrow l_1^- l_1^+ l_2^- l_2^+$ ($l_j = e, \mu, \tau, l_1 \neq l_2$) versus $\sqrt{a_2}$ in case $a'_{hZ} = 1, b'_{hZ} = -0.5i, c'_{hZ} = 0$.

Note that from (27)-(35), regardless of the values of the couplings $a'_Z(a_2)$, $b'_Z(a_2)$ and $c'_Z(a_2)$, it follows that for any a_2

$$\begin{aligned} O_1^{(1,2)} \in [-\frac{3}{4}A_{f_{1,2}}, \frac{3}{4}A_{f_{1,2}}], \quad O_2 \in [0, 1], \quad O_3 \in [0, \frac{9}{16}A_{f_1}A_{f_2}], \quad O_4, O_5 \in [-\frac{1}{2\pi}, \frac{1}{2\pi}], \\ O_6, O_9 \in [-\frac{9}{64}\sqrt{2\pi}A_{f_1}A_{f_2}, \frac{9}{64}\sqrt{2\pi}A_{f_1}A_{f_2}], \quad O_7^{(1,2)}, O_8^{(1,2)} \in [-\frac{3}{16}\sqrt{2}A_{f_{2,1}}, \frac{3}{16}\sqrt{2}A_{f_{2,1}}]. \end{aligned} \quad (36)$$

Since $A_{e^-} = A_{\mu^-} = A_{\tau^-} \approx 0.214$, $A_{\nu_e} = A_{\nu_\mu} = A_{\nu_\tau} = 1$, $A_u = A_c = A_t \approx 0.697$, $A_d = A_s = A_b \approx 0.941$, the moduli of $O_1^{(1,2)}(a_2)$, $O_7^{(1,2)}(a_2)$, $O_8^{(1,2)}(a_2)$, $O_3(a_2)$, $O_6(a_2)$, $O_9(a_2)$ for the decays (1) with quarks and/or neutrinos in the final states are greater than those for the decays (1) to leptons, therefore, the former processes seem more feasible for experimental study. On the other hand, detection of leptons is much simpler. That is why the study of each decay channel of the type (1) has advantages and disadvantages which strongly depend on experimental methods and parameters of detectors. Consequently, measurement of the observables $O_1^{(1,2)}(a_2)$, ..., $O_9(a_2)$ for various decay channels and for various invariant masses of the fermion pair ($\sqrt{a_2}$) may help to put constraints on the XZZ couplings $a'_Z(a_2)$, $b'_Z(a_2)$ and $c'_Z(a_2)$.

III. CONCLUSIONS

In the present paper the decay of a neutral particle X with zero spin and arbitrary CP parity into two off-mass-shell Z bosons (Z_1^* and Z_2^*) each of which decays to a fermion-antifermion pair, i.e. the decay $X \rightarrow Z_1^*Z_2^* \rightarrow f_1\bar{f}_1f_2\bar{f}_2$, has been considered. The given decay has been examined at tree level for the non-identical fermions, $f_1 \neq f_2$. In the approximation of the massless fermions a formula for the fully differential width has been obtained. It has been established that the narrow- Z -width approximation is applicable for finding differential decay widths of $X \rightarrow Z_1^*Z_2^* \rightarrow f_1\bar{f}_1f_2\bar{f}_2$ only if the invariant mass $\sqrt{a_2}$ of the pair $f_2\bar{f}_2$ lies in an interval $(0, m_X - \sqrt{m_Z^2 + \varepsilon_2}]$. If the parameter ε_2 gets larger, the accuracy of the used approximation increases, but the interval in which the approximation is valid reduces. As an optimal value of ε_2 we have chosen $\varepsilon_2 = 3m_Z\Gamma_Z$.

In the narrow- Z -width approximation, but without the neglect of Γ_Z in the propagator of Z_2^* , a formula for the total width of the decay (1) and the total width of $h \rightarrow Z_1^*Z_2^*$ have been derived. The former formula is valid in case $f_1 = f_2$ as well. Note that in Ref. [23] within the framework of the SM the total width of the decay $X \rightarrow ZZ^* \rightarrow Zf\bar{f}$ has been found in the approximation $\Gamma_Z \approx 0$ in the propagator of Z^* . In an analogous way one can obtain the total width of the decay (1) in the SM after the neglect of Γ_Z in the propagator of Z_2^* , however the formula (A2), derived in the present paper, is more general and more precise.

Using the CMS data [3], we have found constraints on the couplings a'_{hZ} , b'_{hZ} , c'_{hZ} , which determine the hZZ interaction and the CP properties of the boson h detected in the experiments [1]. Comparing our constraints with those reported in Refs. [36, 37], one can notice appreciable overlaps between the three results. Besides, we have derived an allowed interval for a ratio not studied in [36, 37]. Taking our allowed regions into account, we have

selected several sets of values of the couplings q'_{hZ} ($q = a, b, c$) and analyzed results for these sets.

The observables $O_1^{(1,2)}(a_2), \dots, O_9(a_2)$, measurement of which will allow one to get constraints on the dependences of q'_Z on $\sqrt{a_2}$, are defined. It is shown that the observables $O_1^{(1,2)}(a_2), O_5(a_2), O_8^{(1,2)}(a_2), O_9(a_2)$ become zero in case $c'_Z(a_2) = 0$, and therefore their experimental dependences on $\sqrt{a_2}$ can put significant constraints on the CP -odd coupling $c'_Z(a_2)$. The observables $O_7^{(1,2)}(a_2)$ vanish if $b'_Z(a_2) = 0$, and, therefore, their measurement is important for finding the CP -even coupling $b'_Z(a_2)$.

Note that the absolute values of $O_1^{(1,2)}(a_2), O_7^{(1,2)}(a_2), O_8^{(1,2)}(a_2), O_3(a_2), O_6(a_2)$ and $O_9(a_2)$ for the decays (1) where f_1 and/or f_2 is a quark or a neutrino are greater than those for the processes in which the fermions are leptons. At the same time, the processes with the leptons are much more convenient from the experimental point of view.

Thus, measurement of the observables $O_1^{(1,2)}(a_2), \dots, O_9(a_2)$ for the decays (1) can help to clarify the CP properties of the particle X and the structure of the amplitude of the decay $X \rightarrow Z_1^* Z_2^*$.

The authors thank Sergiy Ivashyn for useful discussions. The work is partially supported by the National Academy of Sciences of Ukraine (project IIQ-15-1/2015) and the Ministry of Education and Science of Ukraine (project 0115U00473).

Appendix A: Calculation of the total widths of the decays $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ and $h \rightarrow Z_1^* Z_2^*$

In this Appendix we calculate the total width of the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$ for the m_X -values such that $m_X > m_Z$ and for the dependences of $a_Z(a_1, a_2), b_Z(a_1, a_2), c_Z(a_1, a_2)$ such that the differential width $\frac{d^2\Gamma}{da_1 da_2}$ has a sharp maximum when $\sqrt{a_1} = m_Z$ or $\sqrt{a_2} = m_Z$ and the functions $|a'_Z|, |b'_Z|, |c'_Z|, \cos(\arg b'_Z - \arg a'_Z)$ ($q'_Z \equiv q_Z(m_Z^2, a_2)$; $q = a, b, c$) are independent of a_2 . For example, $|a_Z| \approx 1, b_Z \approx 0, c_Z \approx 0$ are such dependences (see Sec. II B). Then we calculate the total decay width of $h \rightarrow Z_1^* Z_2^*$ and examine the applicability of an approximation $\Gamma_Z \approx 0$ for derivation of the total widths.

1. The total width of the decay $X \rightarrow Z_1^* Z_2^* \rightarrow f_1 \bar{f}_1 f_2 \bar{f}_2$

Analogously to the derivation of Eq. (11), we find that

$$\Gamma \approx \frac{2\pi}{m_Z \Gamma_Z} \int_0^{(m_X - m_Z)^2} da_2 f(m_Z^2, a_2). \quad (\text{A1})$$

Let us consider the case wherein $|a'_Z|, |b'_Z|, |c'_Z|, \cos(\arg b'_Z - \arg a'_Z)$ are independent of a_2 . Having exactly calculated the integral in Eq. (A1) with allowance for Eqs. (10) and (6), we obtain:

$$\Gamma \approx (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, s), \quad (\text{A2})$$

where

$$\begin{aligned}
f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, s) &\equiv \frac{\sqrt{2}G_F^3 m_Z^7 m_X}{2^{10} 3^3 \pi^4 \Gamma_Z} \\
&\times \left[(1-\alpha) \left(-24(23\alpha-5)|a'_Z|^2 + (3\alpha^3 - 37\alpha^2 - \alpha(235+6\beta^2) + 77 - 54\beta^2)|b'_Z|^2 \right. \right. \\
&- 16(2\alpha^2 + 26\alpha - 13 + 3\beta^2)\text{Re}(a'_Z{}^* b'_Z) + 64\alpha(\alpha^2 + 40\alpha - 11 + 6\beta^2)|c'_Z|^2 \Big) + 6 \ln \left(\frac{1}{\alpha} \right) \\
&\times \left(4(12\alpha^2 - 18\alpha + 3 - \beta^2)|a'_Z|^2 + (30\alpha^2 - 10\alpha(3 - \beta^2) + 5 - 10\beta^2 + \beta^4)|b'_Z|^2 \right. \\
&+ 8(6\alpha^2 - \alpha(9 - \beta^2) + 2 - 2\beta^2)\text{Re}(a'_Z{}^* b'_Z) - 32\alpha(6\alpha^2 - \alpha(9 - \beta^2) + 1 - 3\beta^2)|c'_Z|^2 \Big) \\
&+ s \frac{3\sqrt{2}}{\beta} \left(P(\alpha, \beta, a'_Z, b'_Z, c'_Z, r_+, -4\beta r_-) \right. \\
&\times \ln \frac{(1-\alpha)^2 \sqrt{(4\alpha-1+\beta^2)^2 + 4\beta^2} + (3\alpha-1)^2 + \beta^2(\alpha+1)^2 + s\sqrt{2}(1-\alpha)((3\alpha-1)r_- - \beta(\alpha+1)r_+)}{4\alpha(\alpha^2 + \beta^2)} \\
&+ 2P(\alpha, \beta, a'_Z, b'_Z, c'_Z, r_-, 4\beta r_+) \left(\pi - \arg(-\alpha(3\alpha-1+\beta^2) - \beta^2 + s \frac{1-\alpha}{\sqrt{2}}(\beta r_+ - \alpha r_-)) \right. \\
&\left. \left. + i(1-\alpha)(s \frac{\alpha r_+ + \beta r_-}{\sqrt{2}} - \beta(1-\alpha)) \right) \right) \Big], \tag{A3}
\end{aligned}$$

$$\alpha \equiv \left(\frac{m_Z}{m_X} \right)^2, \quad \beta \equiv \frac{m_Z \Gamma_Z}{m_X^2},$$

$$\begin{aligned}
P(\alpha, \beta, a'_Z, b'_Z, c'_Z, x, y) &\equiv 2(2x(12\alpha^2 - 4\alpha + 1 - \beta^2) + y(6\alpha - 1))|a'_Z|^2 + (x(16\alpha^2 - 8\alpha(1 - \beta^2) + 1 \\
&- 6\beta^2 + \beta^4) + y(4\alpha - 1 + \beta^2))|b'_Z|^2 + (4x(8\alpha^2 - 2\alpha(3 - \beta^2) + 1 - 3\beta^2) \\
&+ y(8\alpha - 3 + \beta^2))\text{Re}(a'_Z{}^* b'_Z) - 8\alpha(4x(4\alpha^2 - \alpha(1 - \beta^2) - 2\beta^2) \\
&+ y(6\alpha - 1 + \beta^2))|c'_Z|^2, \tag{A4}
\end{aligned}$$

$$r_{\pm} \equiv \sqrt{\sqrt{(4\alpha-1+\beta^2)^2 + 4\beta^2} \pm (4\alpha-1+\beta^2)}. \tag{A5}$$

In place of s one may take 1 or -1 ($f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, -1) = f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, 1)$). In this article the argument $\arg z$ of a complex number z is defined as follows:

$$\begin{aligned}
\arg z &= \arctan \frac{\text{Im } z}{\text{Re } z} + \pi n(\text{Re } z, \text{Im } z) \quad \forall z \in C | \text{Re } z \neq 0, \\
\arg z &= \pi \left(\frac{1}{2} + \Theta(-\text{Im } z) \right) \quad \forall z \in C | (\text{Re } z = 0 \text{ and } \text{Im } z \neq 0), \tag{A6}
\end{aligned}$$

where $n(x, y) \equiv \Theta(-x) + 2\Theta(x)\Theta(-y) \quad \forall x \neq 0$,

$$\Theta(x) \equiv 0 \quad \forall x \in (-\infty, 0], \quad \Theta(x) \equiv 1 \quad \forall x \in (0, +\infty). \tag{A7}$$

From the definition (A6) it follows that $\arg z$ is the angle counted clockwise on the complex plane from the vector $(\text{Re } z, \text{Im } z)$ towards the vector $(1, 0)$ and $\arg z \in [0, 2\pi)$. Sometimes in literature a different function

$$\arg' z \equiv \arg z - 2\pi\Theta(-\text{Im } z) \tag{A8}$$

is used as the argument of z . From (A8) it follows that $\arg' z \in (-\pi, \pi]$. Note that we have already used $\arg z$ above in the expression $\cos(\arg b_Z - \arg a_Z)$, but since $\cos(\arg b_Z - \arg a_Z) = \cos(\arg' b_Z - \arg' a_Z)$, at that point the distinction between $\arg z$ and $\arg' z$ was irrelevant.

Calculating the integral over a_2 in Eq. (A1), one finds an antiderivative of $f(m_Z^2, a_2)$ on the interval $[0, (m_X - m_Z)^2]$. In this antiderivative the function $\arg u_1(a_2)$ naturally appears, where $u_1(a_2)$ is a complex-valued dimensionless function such that

$$\begin{aligned} \forall a_2 \in [0, (m_X - m_Z)^2) \quad \text{Im } u_1(a_2) &\neq 0, \\ \text{Im } u_1((m_X - m_Z)^2) &= 0, \quad \text{Re } u_1((m_X - m_Z)^2) < 0. \end{aligned} \quad (\text{A9})$$

$\arg' u_1(a_2)$ does not emerge in place of $\arg u_1(a_2)$ since, according to (A8), the function $\arg' z$ has a discontinuity on the half-line $\text{Im } z = 0, \text{Re } z < 0$ and thus $\arg' u_1(a_2)$ has a discontinuity at the point $a_2 = (m_X - m_Z)^2$. To avoid this drawback it is convenient to use $\arg z$ in Eq. (A3).

Note that in case of the Higgs boson, i.e. $X = h$, Eq. (A3) can also be written in terms of the function \arg' : for this one has to substitute in Eq. (A3) $\pi - \arg \dots$ by $s\pi - \arg' \dots$, since according to (A8) and to data of Table I, $\pi - \arg \dots = s\pi - \arg' \dots$.

In case of the identical fermions, $f_1 = f_2$, one may neglect the interference term and then in order to obtain a formula for Γ one has to multiply the right-hand side of the relation (A2) by $\frac{1}{2!2!}$ (in view of the identity of the final fermions) and by 2 (since the contribution of the diagram with the permutation of the particles to Γ is equal to that of the diagram without the permutation), i.e. to multiply the right-hand side by $\frac{1}{2}$. Consequently, for any f_1 and f_2

$$\Gamma \approx (1 - \frac{1}{2}\delta_{f_1 f_2})(a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2)f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, s) \equiv \Gamma_{\Gamma_Z}, \quad (\text{A10})$$

where $\delta_{f_1 f_2} \equiv 0(1)$ at $f_1 \neq f_2$ ($f_1 = f_2$). The neglected interference term seems small based on qualitative arguments of Ref. [38]. For a quantitative estimate we can use Ref. [30] (see Table 1 there), according to which the interference contribution to $\Gamma(h \rightarrow Z_1^* Z_2^* \rightarrow 4e)$ in the SM at tree level is 5.80% for $m_h = 140$ GeV.

In Ref. [23] the width of the decay $h \rightarrow ZZ^* \rightarrow Zf\bar{f}$ has been derived at tree level in the SM after the neglect of Γ_Z in the propagator of Z^* . Following [23], when calculating the integral in Eq. (A1), in the expression for $f(m_Z^2, a_2)$ we may also neglect Γ_Z , and then we obtain the following approximate formula for Γ in the SM:

$$\begin{aligned} \Gamma|_{SM} &\approx (1 - \frac{1}{2}\delta_{f_1 f_2}) \frac{\sqrt{2}G_F^3 m_Z^7 m_X}{2^7 3^2 \pi^4 \Gamma_Z} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \\ &\times \left(6 \frac{1 - 8\alpha + 20\alpha^2}{\sqrt{4\alpha - 1}} \arccos\left(\frac{3\alpha - 1}{2\alpha^{\frac{3}{2}}}\right) - \frac{1 - \alpha}{\alpha} (2 - 13\alpha + 47\alpha^2) + 3(1 - 6\alpha + 4\alpha^2) \ln \frac{1}{\alpha} \right) \\ &\equiv \Gamma_0|_{SM}. \end{aligned} \quad (\text{A11})$$

From (A11) and (A10) we obtain that at $m_X = m_h$

$$\Gamma_0|_{SM} \approx 1.001 \times \Gamma_{\Gamma_Z}|_{SM}. \quad (\text{A12})$$

Besides, $\Gamma_0 > \Gamma_{\Gamma_Z}$ (for any $a'_Z, b'_Z, c'_Z, f_1, f_2$) since when deriving the formula for Γ_0 one neglects the width Γ_Z in $f(m_Z^2, a_2)$ and the value of the integral increases. Still according to (A12), the difference between $\Gamma_0|_{SM}$ and $\Gamma_{\Gamma_Z}|_{SM}$ is about one per mille.

Finally, note that at $m_X = m_h$ we can represent the dependence of the function f_0 on the XZZ couplings a'_Z, b'_Z, c'_Z in the convenient form:

$$f_0(a'_Z, b'_Z, c'_Z, m_Z, \Gamma_Z, s) \approx (3.359|a'_Z|^2 + 0.052|b'_Z|^2 + 0.594 \text{Re}(a'_Z b'_Z) + 0.125|c'_Z|^2) \text{ keV}. \quad (\text{A13})$$

2. The total width of the decay $h \rightarrow Z_1^* Z_2^*$

The total decay width $\Gamma(h \rightarrow Z_1^* Z_2^*)$ is

$$\Gamma(h \rightarrow Z_1^* Z_2^*) = \sum_{f_1} \sum_{f_2 \geq f_1} \Gamma|_{m_X=m_h}, \quad (\text{A14})$$

where the sums run over the fermions $e^-, \mu^-, \tau^-, \nu_e, \nu_\mu, \nu_\tau, u_i, c_i, d_i, s_i, b_i$ (since $m_h \in (4m_b, 2m_t]$), $i = r, g, b$ is an index of quark color. It follows from Eqs. (A14), (A12) that in the SM

$$\Gamma_0(h \rightarrow Z_1^* Z_2^*) \approx 1.001 \times \Gamma_{\Gamma_Z}(h \rightarrow Z_1^* Z_2^*). \quad (\text{A15})$$

Further we use Eq. (A10) since it is more precise than Eq. (A11) and consider the case wherein $|a'_{hZ}|, |b'_{hZ}|, |c'_{hZ}|, \cos(\arg b'_{hZ} - \arg a'_{hZ})$ do not depend on a_2 . From Eqs. (A14), (A10), (A3) we derive that

$$\begin{aligned} \Gamma(h \rightarrow Z_1^* Z_2^*) &\approx f_0(a'_{hZ}, b'_{hZ}, c'_{hZ}, m_Z, \Gamma_Z, s)|_{m_X=m_h} \left(\frac{1}{2} \sum_{f_1} (a_{f_1}^2 + v_{f_1}^2)^2 + \right. \\ &\quad \left. + \frac{1}{2} \sum_{f_1} \sum_{f_2 \neq f_1} (a_{f_1}^2 + v_{f_1}^2)(a_{f_2}^2 + v_{f_2}^2) \right) = \frac{f_0(\dots)}{2} \left(\sum_f (a_f^2 + v_f^2) \right)^2 = \\ &= \frac{f_0(\dots)}{18} \left(\frac{103}{2} - 100 \left(\frac{m_W}{m_Z} \right)^2 + 80 \left(\frac{m_W}{m_Z} \right)^4 \right)^2. \end{aligned} \quad (\text{A16})$$

Carrying out calculations, we find the total decay width for the sets (24), (25):

$$\begin{aligned} |a'_{hZ}| = 1, b'_{hZ} = 0, c'_{hZ} = 0 &\Rightarrow \Gamma(h \rightarrow Z_1^* Z_2^*) \approx 91.16^{+16.66}_{-14.50} \text{ keV}, \\ |a'_{hZ}| = 1, b'_{hZ} = 0, |c'_{hZ}| = 0.5 &\Rightarrow \Gamma(h \rightarrow Z_1^* Z_2^*) \approx 92.01^{+16.85}_{-14.67} \text{ keV}, \\ a'_{hZ} = 1, b'_{hZ} = -0.5, c'_{hZ} = 0 &\Rightarrow \Gamma(h \rightarrow Z_1^* Z_2^*) \approx 83.45^{+15.06}_{-13.14} \text{ keV}, \\ a'_{hZ} = 1, b'_{hZ} = \pm 0.5i, c'_{hZ} = 0 &\Rightarrow \Gamma(h \rightarrow Z_1^* Z_2^*) \approx 91.51^{+16.74}_{-14.57} \text{ keV}. \end{aligned} \quad (\text{A17})$$

The uncertainties shown in Eqs. (A17) are calculated by finding the maximum and minimum values of the function $\Gamma(h \rightarrow Z_1^* Z_2^*)$ in the region $v \in [v_0 - 3\sigma_v, v_0 + 3\sigma_v]$ ($v = G_F, m_h, m_Z, m_W, \Gamma_Z$). Here v_0 is the central value of a quantity v , σ_v is the 1-standard-deviation uncertainty of v ; according to the data of Table I, $G_{F0} =$

$1.1663787 \times 10^{-5} \text{ GeV}^{-2}$, $\sigma_{GF} = 6 \times 10^{-12} \text{ GeV}^{-2}$, $m_{h0} = 125.7 \text{ GeV}$, $\sigma_{m_h} = 0.4 \text{ GeV}$ etc.

- [1] G. Aad et al. (ATLAS Collaboration), Phys. Lett. B **716**, 1 (2012);
S. Chatrchyan et al. (CMS Collaboration), Phys. Lett. B **716**, 30 (2012).
- [2] G. Aad et al. (ATLAS Collaboration), Phys. Lett. B **726**, 88 (2013).
- [3] S. Chatrchyan et al. (CMS Collaboration), Phys. Rev. D **89**, 092007 (2014).
- [4] S. Chatrchyan et al. (CMS Collaboration), Phys. Rev. Lett. **110**, 081803 (2013).
- [5] A. Pilaftsis, C.E.M. Wagner, Nucl. Phys. B **553**, 3 (1999).
- [6] V. Barger, P. Langacker, M. McCaskey et al., Phys. Rev. D **79**, 015018 (2009).
- [7] G.C. Branco, P.M. Ferreira, L. Lavoura et al., Phys. Rep. **516**, 1 (2012).
- [8] D. Bailin, A. Love, *Cosmology in Gauge Field Theory and String Theory* (Institute of Physics Publishing, Bristol-Philadelphia, 2004).
- [9] M.B. Voloshin, Phys. Rev. D **86**, 093016 (2012).
- [10] F. Bishara, Y. Grossman, R. Harnik et al., JHEP **1404**, 084 (2014).
- [11] A.Yu. Korchin, V.A. Kovalchuk, Phys. Rev. D **88**, 036009 (2013);
A.Yu. Korchin, V.A. Kovalchuk, Acta Phys. Polon. B **44**, 2121 (2013).
- [12] J. S. Gainer, W. Y. Keung, I. Low and P. Schwaller, Phys. Rev. D **86**, 033010 (2012).
- [13] A.Y. Korchin and V.A. Kovalchuk, Eur. Phys. J. C **74**, 3141 (2014).
- [14] S.Y. Choi, D.J. Miller, M.M. Mühlleitner et al., Phys. Lett. B **553**, 61 (2003).
- [15] V.A. Kovalchuk, J. Exp. Theor. Phys. **107**, 774 (2008).
- [16] A. Menon, T. Modak, D. Sahoo et al., Phys. Rev. D **89**, 095021 (2014).
- [17] Y. Sun, X.-F. Wang, and D.-N. Gao, Int. J. Mod. Phys. A **29**, 1450086 (2014).
- [18] A. De Rujula, J. Lykken, M. Pierini et al., Phys. Rev. D **82**, 013003 (2010);
Y. Gao, A. V. Gritsan, Z. Guo et al., Phys. Rev. D **81**, 075022 (2010);
S. Bolognesi, Y. Gao, A. V. Gritsan et al., Phys. Rev. D **86**, 095031 (2012);
D. Stolarski and R. Vega-Morales, Phys. Rev. D **86**, 117504 (2012);
P. Avery, D. Bourilkov, M. Chen et al., Phys. Rev. D **87**, no. 5, 055006 (2013);
M. Chen, T. Cheng, J. S. Gainer et al., Phys. Rev. D **89**, no. 3, 034002 (2014);
B. Bhattacharjee, T. Modak, S. K. Patra et al., arXiv:1503.08924 [hep-ph].
- [19] M. Gasperini, Phys. Lett. B **327**, 214 (1994).
- [20] Z. Chacko, R. Franceschini, and R.K. Mishra, JHEP **1304**, 015 (2013).
- [21] B. Bellazzini, C. Csáki, J. Hubisz et al., Eur. Phys. J. C **73**, 2333 (2013).
- [22] J. Serra, EPJ Web Conf. **60**, 17005 (2013).
- [23] W.-Y. Keung and W.J. Marciano, Phys. Rev. D **30**, 248 (1984).
- [24] R. M. Godbole, D. J. Miller and M. M. Mühlleitner, JHEP **0712**, 031 (2007).

- [25] W.-Y. Keung, I. Low, and J. Shu, Phys. Rev. Lett. **101**, 091802 (2008);
T.L. Trueman, Phys. Rev. D **18**, 3423 (1978).
J. R. Dell'Aquila and C. A. Nelson, Phys. Rev. D **33**, 80 (1986).
- [26] K.A. Olive et al. (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
- [27] N.N. Achasov and V.V. Gubin, JETP Lett. **62**, 191 (1995) [Pisma Zh. Eksp. Teor. Fiz. **62**, 182 (1995)].
- [28] D. Berdine, N. Kauer, and D. Rainwater, Phys. Rev. Lett. **99**, 111601 (2007).
- [29] C.F. Uhlemann and N. Kauer, Nucl. Phys. B **814**, 195 (2009).
- [30] A. Bredenstein, A. Denner, S. Dittmaier and M. M. Weber, Phys. Rev. D **74**, 013004 (2006).
- [31] G. Aad et al. (ATLAS Collaboration), arXiv:1504.05833 [hep-ex].
- [32] LHC Higgs Cross Section Working Group,
<https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageAt8TeV>.
- [33] V. Khachatryan et al. (CMS Collaboration), Phys. Lett. B **736**, 64 (2014).
- [34] S. Heinemeyer et al. (LHC Higgs Cross Section Working Group), arXiv:1307.1347v2 [hep-ph].
- [35] J. S. Gainer, J. Lykken, K. T. Matchev et al., Phys. Rev. Lett. **111**, 041801 (2013).
- [36] V. Khachatryan et al. (CMS Collaboration), Phys. Rev. D **92**, no. 1, 012004 (2015).
- [37] G. Aad et al. (ATLAS Collaboration), arXiv:1506.05669v1 [hep-ex].
- [38] J. C. Romão and S. Andringa, Eur. Phys. J. C **7**, 631 (1999).