

Effective field theory approach to LHC Higgs data

Adam Falkowski

Laboratoire de Physique Théorique, Bat. 210, Université Paris-Sud, 91405 Orsay, France.

Abstract

I review the effective field theory approach to LHC Higgs data.

Invited review prepared for Pramana - Journal of Physics

1 Introduction

The standard model (SM) of particle physics was proposed back in the 60s as a theory of quarks and leptons interacting via strong, weak, and electromagnetic forces [1]. It is build on the following principles

- #1 The basic framework is that of a relativistic quantum field theory, with interactions between particles described by a local Lagrangian.
- #2 The Lagrangian is invariant under the linearly realized local $SU(3) \times SU(2) \times U(1)$ symmetry.
- #3 The vacuum state of the theory preserves only $SU(3) \times U(1)$ local symmetry, as a result of the Brout-Englert-Higgs mechanism [2–4]. The spontaneous breaking of the $SU(2) \times U(1)$ symmetry down to $U(1)$ arises due to a vacuum expectation value (VEV) of a scalar field transforming as $(1, 2)_{1/2}$ under the local symmetry.
- #4 Interactions are *renormalizable*, which means that only interactions up to the canonical mass dimension 4 are allowed in the Lagrangian.

Given the experimentally observed matter content (3 families of quarks and leptons), these rules completely specify the theory up to 19 free parameters. The local symmetry implies the presence of spin-1 vector bosons which mediate the strong and electroweak forces. The breaking pattern of the local symmetry ensures that the carriers of the strong and electromagnetic force are massless, whereas the carriers of the weak force are massive. Finally, the particular realization of the Brout-Englert-Higgs mechanism in the SM leads to the emergence of exactly one spin-0 scalar boson - the famous Higgs boson [5–7].

The SM passed an incredible number of experimental tests. It correctly describes the rates and differential distributions of particles produced in high-energy collisions; a robust deviation from the SM predictions has never been observed. It predicts very accurately many properties of elementary particles, such as the magnetic and electric dipole moments, as well as certain properties of simple enough composite particles, such as atomic energy levels. The discovery of a 125 GeV boson at the Large Hadron Collider (LHC) [8,9] nails down the last propagating degree of freedom predicted by the SM. Measurements of its production and decay rates vindicates the simplest realization of the Brout-Englert-Higgs mechanism, in which a VEV of a *single* $SU(2)$ doublet field spontaneously breaks the electroweak symmetry. Last not least, the SM is a consistent quantum theory, whose validity range extends to energies all the way up to the Planck scale (at which point the gravitational interactions become strong and can no longer be neglected).

Yet we know that the SM is not the ultimate theory. It cannot account for dark matter, neutrino masses, matter/anti-matter asymmetry, and cosmic inflation, which are all experimental facts. In addition, some theoretical or esthetic arguments (the strong CP problem, flavor hierarchies, unification, the naturalness problem) suggest that the SM should be extended. This justifies the ongoing searches for *new physics*, that is particles or interactions not predicted by the SM.

In spite of good arguments for the existence of new physics, a growing body of evidence suggests that, at least up to energies of a few hundred GeV, the fundamental degrees of freedom are those of the SM. Given the absence of any direct or indirect collider signal of new physics, it is reasonable to assume that new particles from beyond the SM are much heavier than the SM particles. If that

is correct, physics at the weak scale can be adequately described using *effective field theory* (EFT) methods.

In the EFT framework adopted here the assumptions #1...#3 above continue to be valid.¹ Thus, much as in the SM, the Lagrangian is constructed from gauge invariant operators involving the SM fermion, gauge, and Higgs fields. The difference is that the assumption #4 is dropped and interactions with arbitrary large mass dimension D are allowed. These interactions can be organized in a systematic expansion in D . The leading order term in this expansion is the SM Lagrangian with operators up to $D = 4$. All possible effects of heavy new physics are encoded in operators with $D > 4$, which are suppressed in the Lagrangian by appropriate powers of the mass scale Λ . Since all $D = 5$ operators violate lepton number and are thus stringently constrained by experiment, the leading corrections to the Higgs observables are expected from $D = 6$ operators suppressed by Λ^2 [14]. I will assume that the operators with $D > 6$ can be ignored, which is always true for $v \ll \Lambda$.

This review discusses the interpretation of the LHC data on the Higgs boson production and decay in the framework of an EFT beyond the SM. For practical reasons, three more assumptions about higher-dimensional operators are adopted:

- The baryon and lepton numbers are conserved.
- The coefficients of operators involving fermions are flavor conserving and universal, except for Yukawa-type operators, which are aligned with the corresponding SM Yukawa matrices.²
- The corrections from $D = 6$ operators to the Higgs signal strength are subleading compared to the SM contribution.

Other than that, the discussion will be model-independent.

In the following section I review the SM Lagrangian, in order to prepare the ground and fix the notation. The part of the $D = 6$ effective Lagrangian relevant for Higgs studies is discussed in Section 3. The dependence of the Higgs signal strength measured at the LHC on the effective Lagrangian parameters is summarized in Section 4. The experimental results and the current model-independent constraints on the $D = 6$ parameters are discussed in Section 5. The bibliography contains a number of references where an EFT-inspired approach to physics of the 125 GeV Higgs at the LHC is exercised; citation complaints are welcome for omitted papers belonging to that category.

2 Standard Model Lagrangian

I start by summarizing the SM Lagrangian and defining my notation.

The SM Lagrangian is invariant under the global Poincaré symmetry (Lorentz symmetry + translations), and a local symmetry with the gauge group $G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$. The

¹One could consider a more general EFT where the assumptions #2 and #3 are also relaxed and the electroweak symmetry is realized *non-linearly* [10–13]. In that case, the Higgs boson is introduced as a singlet of the local symmetry, rather than as a part of an $SU(2)$ doublet.

²This assumption is largely practical, because there is little experimental information about Higgs couplings to the 1st and 2nd generation fermions. Currently, these couplings are probed indirectly [15, 16], while in the future some may be probed directly via exclusive Higgs decays to a photon and a meson [17, 18].

	$SU(3)_C$	$SU(2)_L$	$U(1)_Y$
$q = \begin{pmatrix} u \\ d \end{pmatrix}$	3	2	1/6
u^c	$\bar{3}$	1	-2/3
d^c	3	1	1/3
$\ell = \begin{pmatrix} \nu \\ e \end{pmatrix}$	1	2	-1/2
e^c	1	1	1
H	1	2	1/2

Table 1: Representation of the SM scalar and fermion fields under the SM gauge group.

fields building the SM Lagrangian fill representations of these symmetries. The field content of the SM is the following:

- Vector fields G_μ^a , W_μ^i , B_μ , where $i = 1 \dots 3$ and $a = 1 \dots 8$. They transform as four-vectors under the Lorentz symmetry and are the gauge fields of the G_{SM} group.
- 3 generations of fermionic fields $q = (u, V_{\text{CKM}}d)$, u^c , d^c , $\ell = (\nu, e)$, e^c . They transform as 2-component spinors under the Lorentz symmetry.³ The transformation properties under G_{SM} are listed in Table 1.
- Scalar field $H = (H^+, H^0)$ transforming as $(1, 2)_{1/2}$ under G_{SM} . I also define $\tilde{H}_i = \epsilon_{ij} H_j^*$ that transforms as $(1, 2)_{-1/2}$.

The SM Lagrangian can be split as

$$\mathcal{L}^{\text{SM}} = \mathcal{L}_V^{\text{SM}} + \mathcal{L}_F^{\text{SM}} + \mathcal{L}_H^{\text{SM}} + \mathcal{L}_Y^{\text{SM}}. \quad (2.1)$$

The first term above contains gauge invariant kinetic terms for the vector fields:

$$\mathcal{L}_V^{\text{SM}} = -\frac{1}{4g_s^2} G_{\mu\nu}^a G_{\mu\nu}^a - \frac{1}{4g_L^2} W_{\mu\nu}^i W_{\mu\nu}^i - \frac{1}{4g_Y^2} B_{\mu\nu} B_{\mu\nu}, \quad (2.2)$$

where g_s , g_L , g_Y are gauge couplings of $SU(3)_C \times SU(2)_L \times U(1)_Y$, here defined as the normalization of the appropriate gauge kinetic term. I also define the electromagnetic coupling $e = g_L g_Y / \sqrt{g_L^2 + g_Y^2}$, and the Weinberg angle $s_\theta = g_Y / \sqrt{g_L^2 + g_Y^2}$. The field strength tensors are given by

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu, \quad W_{\mu\nu}^i = \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + \epsilon^{ijk} W_\mu^j W_\nu^k, \quad G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + f^{abc} G_\mu^b G_\nu^c. \quad (2.3)$$

where ϵ^{ijk} and f^{abc} are the totally anti-symmetric structure tensors of $SU(2)$ and $SU(3)$.

The second term in Eq. (2.1) contains covariant kinetic terms of the fermion fields:

$$\mathcal{L}_F^{\text{SM}} = i\bar{q}\sigma_\mu D_\mu q + iu^c\sigma_\mu D_\mu \bar{u}^c + id^c\sigma_\mu D_\mu \bar{d}^c + i\bar{\ell}\sigma_\mu D_\mu \ell + ie^c\sigma_\mu D_\mu \bar{e}^c. \quad (2.4)$$

³Throughout this review I use the 2-component spinor notation for fermions; in all instances I follow the conventions of Ref. [19].

Each fermion field is a 3-component vector in the generation space. I assume all the rotations needed to put fermions in the mass eigenstate basis have already been made; in the SM the only residue of these rotations is the CKM matrix appearing in the definition of the quark doublet components. The covariant derivatives are defined as

$$D_\mu f = \left(\partial_\mu - iG_\mu^a T_f^a - iW_\mu^i T_f^i - iY_f B_\mu \right) f. \quad (2.5)$$

Here $T_f^a = (\lambda^a, -\lambda^a, 0)$ for f in the triplet/anti-triplet/singlet representation of SU(3), where λ^a are Gell-Mann matrices; $T_f^i = (\sigma^i/2, 0)$ for f in the doublet/singlet representation of SU(2); Y_f is the U(1) hypercharge. The electric charge is given by $Q_f = T_f^3 + Y_f$.

The third term in Eq. (2.1) contains Yukawa interactions between the Higgs field and the fermions:

$$\mathcal{L}_Y^{\text{SM}} = -\tilde{H}^\dagger u^c y_u q - H^\dagger d^c y_d q - H^\dagger e^c y_e \ell + \text{h.c.}, \quad (2.6)$$

where y_f are 3×3 diagonal matrices.

The last term in Eq. (2.1) are the Higgs kinetic and potential terms:

$$\mathcal{L}_H^{\text{SM}} = D_\mu H^\dagger D_\mu H + \mu_H^2 H^\dagger H - \lambda (H^\dagger H)^2, \quad (2.7)$$

where the covariant derivative acting on the Higgs field is

$$D_\mu H = \left(\partial_\mu - \frac{i}{2} W_\mu^i \sigma^i - \frac{i}{2} B_\mu \right) H. \quad (2.8)$$

Because of the negative mass squared term μ_H^2 in the Higgs potential the Higgs field gets a VEV,

$$\langle H \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad \mu_H^2 = \lambda v^2. \quad (2.9)$$

This generates mass terms for W_μ^i and B_μ and a field rotation is needed to diagonalize the mass matrix. The mass eigenstates are defined to the electroweak vector fields by

$$\begin{aligned} W_\mu^1 &= \frac{g_L}{\sqrt{2}} (W_\mu^+ + W_\mu^-), & W_\mu^3 &= \frac{g_L}{\sqrt{g_L^2 + g_Y^2}} (g_L Z_\mu + g_Y A_\mu), \\ W_\mu^2 &= \frac{i g_L}{\sqrt{2}} (W_\mu^+ - W_\mu^-), & B_\mu &= \frac{g_Y}{\sqrt{g_L^2 + g_Y^2}} (-g_Y Z_\mu + g_L A_\mu). \end{aligned} \quad (2.10)$$

With this definition, the mass eigenstates such that their quadratic terms are canonically normalized and their mass terms are diagonal:

$$\mathcal{L}_{V,\text{kin}}^{\text{SM}} = -\frac{1}{2} W_{\mu\nu}^+ W_{\mu\nu}^- - \frac{1}{4} Z_{\mu\nu} Z_{\mu\nu} - \frac{1}{4} A_{\mu\nu} A_{\mu\nu} + m_W^2 W_\mu^+ W_\mu^- + \frac{m_Z^2}{2} Z_\mu Z_\mu, \quad (2.11)$$

where the W and Z boson masses are

$$m_W = \frac{g_L v}{2}, \quad m_Z = \frac{\sqrt{g_L^2 + g_Y^2} v}{2}. \quad (2.12)$$

The SM fermions (except for the neutrinos) also acquire masses after electroweak symmetry breaking via the Yukawa interactions in Eq. (2.6). I choose a basis in the fermion flavor space where the Yukawa interactions are diagonal, in which case the fermion masses are given by

$$m_{f_i} = \frac{v}{\sqrt{2}}[y_f]_{ii}. \quad (2.13)$$

Interactions of the gauge boson mass eigenstates with fermions are given by

$$\begin{aligned} \mathcal{L}_{vff}^{\text{SM}} &= eA_\mu \sum_{f \in u,d,e} Q_f (\bar{f} \bar{\sigma}_\mu f + f^c \sigma_\mu f^c) + g_s G_\mu^a \sum_{f \in u,d} (\bar{f} \bar{\sigma}_\mu T^a f + f^c \sigma_\mu T^a f^c) \\ &+ \frac{g_L}{\sqrt{2}} (W_\mu^+ \bar{u} \bar{\sigma}_\mu V_{\text{CKM}} d + W_\mu^+ \bar{\nu} \bar{\sigma}_\mu e + \text{h.c.}) \\ &+ \sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f \in u,d,e,\nu} \bar{f} \bar{\sigma}_\mu (T_f^3 - s_\theta^2 Q_f) f + \sum_{f \in u,d,e} f^c \sigma_\mu (-s_\theta^2 Q_f) f^c \right]. \end{aligned} \quad (2.14)$$

Finally, I move to the Higgs sector. After electroweak symmetry breaking, the Higgs doublet field can be conveniently written as

$$H = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} G^+ \\ v + h + iG^0 \end{pmatrix}. \quad (2.15)$$

The fields G^0 and G^+ do not correspond to new physical degrees of freedom (they kinetically mix with the massive gauge bosons and can be gauged away). From now on, I will work in the unitary gauge and set $G^\pm = 0 = G^0$. The star of this review - the scalar field h - is called *the Higgs boson*. Its mass can be expressed by the parameters of the Higgs potential as

$$m_h^2 = 2\mu_H^2 = 2\lambda v^2. \quad (2.16)$$

The interactions in the SM Lagrangian involving a single Higgs boson are the following

$$\mathcal{L}_h^{\text{SM}} = \frac{h}{v} \left[\frac{g_L^2 v^2}{2} W_\mu^+ W_\mu^- + \frac{(g_L^2 + g_Y^2) v^2}{4} Z_\mu Z_\mu \right] - \frac{h}{v} \sum_f m_f (f f^c + \text{h.c.}). \quad (2.17)$$

Roughly speaking, the Higgs boson couples to mass, in the sense that it couples to pairs of SM particles with the strength proportional to their masses (for fermions) or masses squared (for bosons). Since all the masses have been measured by experiment, the strength of Higgs boson interactions in the SM is precisely predicted and contains no free parameters.

I conclude this section with a summary of the SM parameters used in this review. For the Higgs boson mass I take $m_h = 125.09$ GeV, which is the central value of the recent ATLAS and CMS combination of mass measurements [20]. The gauge boson masses are $m_W = 80.385$ GeV [21], and $m_Z = 91.1875$ GeV [22]. The Higgs VEV is calculated at from the muon lifetime (equivalently, from the Fermi constant $G_F = 1/\sqrt{2}v^2 = 1.16637 \times 10^{-5}$ GeV⁻² [23]), corresponding to $v = 246.221$ GeV. The electroweak couplings at the Z boson mass scale are extracted from m_Z and the electromagnetic structure constant $\alpha(m_Z) = 7.755 \times 10^{-3}$ [24], and the strong coupling from $\alpha_s(m_Z) = 1.172 \times 10^{-3}$ [23]. To evaluate corrections to the Higgs observables I will use the couplings run up to the scale m_h : $g_s = 1.187$, $g_L = 0.643$, $g_Y = 0.358$. The light fermion masses are also evaluated at the scale m_h : the relevant ones are $m_b = 2.76$ GeV, $m_\tau = 1.78$ GeV, and $m_c = 0.62$ GeV. For the top mass I take $m_t = 173.2$ GeV.

3 Dimension Six Lagrangian

We consider the effective Lagrangian of the form,

$$\mathcal{L}_{\text{eff}} = \mathcal{L}^{\text{SM}} + \mathcal{L}^{D=6}, \quad \mathcal{L}^{D=6} = \frac{1}{v^2} \sum_{\alpha} c_{\alpha} O_{\alpha}, \quad (3.1)$$

where \mathcal{L}^{SM} is the SM Lagrangian discussed in Section 2, and O_{α} is a complete basis of $SU(3) \times SU(2) \times U(1)$ invariant $D = 6$ operators constructed out of the SM fields. In general, such a basis contains 2499 independent operators after imposing baryon and lepton number conservation [25]. One of the assumptions in this review is that coefficients of $D = 6$ operators are flavor universal, which brings the number of independent parameters down to 76. Furthermore, only 9 combinations of these operators will be relevant for a completely general description of the Higgs signal strength measurements considered later in this review.

One can choose a complete, non-redundant basis of operators in many distinct (though equivalent) ways. Here we work with the so-called *Higgs basis* introduced in Ref. [26] and inspired by Refs. [27, 28].⁴ The basis is spanned by particular combinations of $D = 6$ operators. Each of these combinations maps to an interaction term of the SM mass-eigenstates in the tree-level effective Lagrangian. The coefficients multiplying these combinations in the Lagrangian are called the *independent couplings*. The single Higgs couplings to pairs of gauge bosons and fermions are chosen among the independent couplings. The advantage of this basis is that the independent couplings are related in a simple way to observables in Higgs physics.

Most often, an $SU(3) \times SU(2) \times U(1)$ invariant operator gives rise to more than one interaction term of mass eigenstates. This leads to relations between various couplings in the effective Lagrangian. Therefore, several of these couplings are not free but can be expressed in terms of the independent couplings; they are called the *dependent couplings*. For example, at the level of the $D = 6$ Lagrangian, the W boson couplings to fermions are dependent couplings, as they can be expressed in terms of the Z boson couplings to fermions. Of course, the choice which couplings are chosen as independent and which are dependent is subjective and dictated by convenience.

Below I review the part of $D = 6$ Lagrangian in the Higgs basis that is relevant for LHC Higgs observables; see Ref. [26] for the full set of independent couplings and the algorithm to construct the complete $D = 6$ Lagrangian. In this formalism, by construction, all kinetic terms are canonically normalized, there is no kinetic mixing between the Z boson and the photon, and there is no correction to the Z boson mass term. While, in general, $D = 6$ operators do generate mixing and mass corrections, the canonical form can always be recovered by using equations of motion, integration by parts, and redefinition of fields and couplings. Thus, the kinetic and mass terms for the electroweak gauge bosons are those in Eq. (2.11), except for the correction to the W boson mass term: $\Delta\mathcal{L}_{\text{kinetic}}^{D=6} = 2\delta m \frac{g^2 v^2}{4} W_{\mu}^{+} W_{\mu}^{-}$. The independent coupling δm is a free parameter from the EFT point of view, however it is very well constrained by experiment: $\delta m = (2.6 \pm 1.9) \cdot 10^{-4}$ [32]. Given the precision of LHC data, effects proportional to δm are currently not relevant for Higgs searches and will be ignored.

We move to interactions of a single Higgs boson with pairs of SM gauge bosons and fermions. The SM interactions of this type were given in Eq. (2.17) and they contain no free parameters. Dimension six operators lead to shifts of the couplings in Eq. (2.17), as well as to the appearance

⁴Other popular choices in the Higgs-related literature are the Warsaw basis [25, 29], and the SILH basis [30, 31].

of 2-derivative Higgs couplings to gauge bosons. In the Higgs basis, these effects are parametrized by the following independent couplings:

$$\begin{aligned} & \delta c_z, c_{zz}, c_{z\Box}, c_{\gamma\gamma}, c_{z\gamma}, c_{gg}, \tilde{c}_{gg}, \tilde{c}_{zz}, \tilde{c}_{\gamma\gamma}, \tilde{c}_{z\gamma}, \\ & \delta y^u, \delta y^d, \delta y^e, \sin \phi^u, \sin \phi^d, \sin \phi^\ell. \end{aligned} \quad (3.2)$$

The couplings in the first line are defined via the Higgs boson couplings to gauge bosons:

$$\begin{aligned} \Delta\mathcal{L}_{\text{hvv}}^{D=6} = & \frac{h}{v} \left[2\delta c_w m_W^2 W_\mu^+ W_\mu^- + \delta c_z m_Z^2 Z_\mu Z_\mu \right. \\ & + c_{ww} \frac{g_L^2}{2} W_{\mu\nu}^+ W_{\mu\nu}^- + \tilde{c}_{ww} \frac{g_L^2}{2} W_{\mu\nu}^+ \tilde{W}_{\mu\nu}^- + c_{w\Box} g_L^2 (W_\mu^- \partial_\nu W_\mu^+ + \text{h.c.}) \\ & + c_{gg} \frac{g_s^2}{4} G_{\mu\nu}^a G_{\mu\nu}^a + c_{\gamma\gamma} \frac{e^2}{4} A_{\mu\nu} A_{\mu\nu} + c_{z\gamma} \frac{eg_L}{2c_\theta} Z_{\mu\nu} A_{\mu\nu} + c_{zz} \frac{g_L^2}{4c_\theta^2} Z_{\mu\nu} Z_{\mu\nu} \\ & + c_{z\Box} g_L^2 Z_\mu \partial_\nu Z_{\mu\nu} + c_{\gamma\Box} g_L g_Y Z_\mu \partial_\nu A_{\mu\nu} \\ & \left. + \tilde{c}_{gg} \frac{g_s^2}{4} G_{\mu\nu}^a \tilde{G}_{\mu\nu}^a + \tilde{c}_{\gamma\gamma} \frac{e^2}{4} A_{\mu\nu} \tilde{A}_{\mu\nu} + \tilde{c}_{z\gamma} \frac{eg_L}{2c_\theta} Z_{\mu\nu} \tilde{A}_{\mu\nu} + \tilde{c}_{zz} \frac{g_L^2}{4c_\theta^2} Z_{\mu\nu} \tilde{Z}_{\mu\nu} \right], \end{aligned} \quad (3.3)$$

where the dependent couplings δc_w , c_{ww} , \tilde{c}_{ww} , $c_{w\Box}$, and $c_{\gamma\Box}$ can be expressed by the independent couplings as

$$\begin{aligned} \delta c_w &= \delta c_z + 4\delta m, \\ c_{ww} &= c_{zz} + 2s_\theta^2 c_{z\gamma} + s_\theta^4 c_{\gamma\gamma}, \\ \tilde{c}_{ww} &= \tilde{c}_{zz} + 2s_\theta^2 \tilde{c}_{z\gamma} + s_\theta^4 \tilde{c}_{\gamma\gamma}, \\ c_{w\Box} &= \frac{1}{g_L^2 - g_Y^2} [g_L^2 c_{z\Box} + g_Y^2 c_{zz} - e^2 s_\theta^2 c_{\gamma\gamma} - (g_L^2 - g_Y^2) s_\theta^2 c_{z\gamma}], \\ c_{\gamma\Box} &= \frac{1}{g_L^2 - g_Y^2} [2g_L^2 c_{z\Box} + (g_L^2 + g_Y^2) c_{zz} - e^2 c_{\gamma\gamma} - (g_L^2 - g_Y^2) c_{z\gamma}]. \end{aligned} \quad (3.4)$$

The coupling in the second line of Eq. (3.2) are defined via the Higgs boson couplings to fermions:

$$\Delta\mathcal{L}_{\text{hff}}^{D=6} = -\frac{h}{v} \sum_{f \in u,d,e} \delta y_f e^{i\phi_f} m_f f^c f + \text{h.c.} \quad (3.5)$$

Following my assumption of flavor universal coefficients of dimension-6 operators, each δy_f and ϕ_f is a real number. Moreover, the couplings in Eq. (3.5) are diagonal in the generation space, therefore flavor violating Higgs decays are absent (see Refs. [33,34] for a discussion of such decays in the EFT language).

The complete Higgs interaction Lagrangian relevant for this review is given by $\mathcal{L}_h^{\text{SM}} + \mathcal{L}_{vff}^{\text{SM}} + \Delta\mathcal{L}_{\text{hvv}}^{D=6} + \Delta\mathcal{L}_{\text{hff}}^{D=6}$ and is parametrized by the independent couplings in Eq. (3.2). The effect of these couplings on the LHC Higgs observables will be discussed in the following sections. But before that, a comment is in order on other effects of $D = 6$ operators that could, a priori, be relevant. First, in the Higgs basis there are corrections to the Z and W boson interactions in Eq. (2.14), parametrized by vertex corrections δg . These would feed indirectly into Higgs observables, such as,

for example, the vector boson fusion (VBF) production cross section or the $h \rightarrow VV^* \rightarrow 4$ fermions decays. However, there are model-independent constraints on these vertex corrections [32] which ensure that their effects on Higgs observables are too small to be currently observable. For this reason I will ignore the vertex corrections in this review. Next, $D = 6$ operators may induce two classes of Higgs boson interactions that could affect $h \rightarrow VV^* \rightarrow 4$ fermions decays. One class is the $hVff$ vertex-like contact interactions:

$$\begin{aligned} \Delta\mathcal{L}_{hvf}^{D=6} &= \sqrt{2}g_L \frac{h}{v} W_\mu^+ \left(\delta g_L^{W\ell} \bar{\nu} \bar{\sigma}_\mu e + \delta g_L^{Wq} \bar{u} \bar{\sigma}_\mu d + \delta g_R^{Wq} u^c \sigma_\mu \bar{d}^c + \text{h.c.} \right) \\ &+ 2 \frac{h}{v} \sqrt{g_L^2 + g_Y^2} Z_\mu \left[\sum_{f=u,d,e,\nu} \delta g_L^{Zf} \bar{f} \bar{\sigma}_\mu f + \sum_{f=u,d,e} \delta g_R^{Zf} f^c \sigma_\mu \bar{f}^c \right]. \end{aligned} \quad (3.6)$$

In the Higgs basis, the parameters δg above are equal to the corresponding vertex corrections to the SM couplings in Eq. (2.14). Given the constraints on the δg 's in Ref. [32], the LHC Higgs studies cannot be currently sensitive to the vertex-like Higgs interactions, therefore they are neglected in this analysis. The other class is the dipole-like contact interactions:

$$\begin{aligned} \Delta\mathcal{L}_{hdvff}^{D=6} &= -\frac{h}{4v^2} \left[g_s \sum_{f \in u,d} d_{Gf} f^c \bar{\sigma}_{\mu\nu} T^a f G_{\mu\nu}^a + e \sum_{f \in u,d,e} d_{Af} f^c \bar{\sigma}_{\mu\nu} f A_{\mu\nu} \right. \\ &+ \left. \sqrt{g_L^2 + g_Y^2} \sum_{f \in u,d,e} d_{Zf} f^c \bar{\sigma}_{\mu\nu} f Z_{\mu\nu} + \sqrt{2} g_L d_{Wq} d^c \bar{\sigma}_{\mu\nu} u W_{\mu\nu}^- + \text{h.c.} \right] \\ &- \frac{h}{4v^2} \left[\sum_{f \in u,d} \tilde{d}_{Gf} f^c \bar{\sigma}_{\mu\nu} T^a f \tilde{G}_{\mu\nu}^a + e \sum_{f \in u,d,e} \tilde{d}_{Af} f^c \bar{\sigma}_{\mu\nu} f \tilde{A}_{\mu\nu} \right. \\ &+ \left. \sqrt{g_L^2 + g_Y^2} \sum_{f \in u,d,e} \tilde{d}_{Zf} f^c \bar{\sigma}_{\mu\nu} f \tilde{Z}_{\mu\nu} + \sqrt{2} g_L \tilde{d}_{Wq} d^c \bar{\sigma}_{\mu\nu} u \tilde{W}_{\mu\nu}^- + \text{h.c.} \right]. \end{aligned} \quad (3.7)$$

For Higgs decays into four light fermions, the dipole-like contributions do not interfere with the SM amplitudes due to the different helicity structure. Therefore, corrections to the decay width enter quadratically in d_{Vf} , and should be neglected. Furthermore, as a consequence of the linearly realized electroweak symmetry in the $D = 6$ Lagrangian, the parameters d_{Vf} are proportional to the respective dipole moments which are stringently constrained by experiment, especially for light fermions. For these two reasons, it is safe to neglect the dipole-like Higgs interactions for the sake of LHC analyses.

Finally, $D = 6$ operators produce several more interactions involving the single Higgs boson field, for example Higgs couplings to 3 gauge bosons. Observable effects of these couplings are extremely suppressed, therefore they are not listed here. Moreover, new interactions involving two (or three) Higgs boson fields appear in the Lagrangian, and they are relevant for an EFT description of double Higgs production [35–41]. This review is focused on single Higgs production, therefore multi-Higgs couplings are not listed; see Ref. [26] for the relevant expressions in the Higgs basis.

I close this section with a brief discussion of the validity range of this approach. Formally, EFT is an expansion in powers of the scale Λ suppressing higher-dimensional operators. Since the

independent couplings in Eq. (3.2) arise from $D = 6$ operators, they are formally of order v^2/Λ^2 . The rule of thumb is that the EFT approach to Higgs physics is valid if $\Lambda \gtrsim v$, which translates to $|c_i| \lesssim 1$ and $\delta y_f \lesssim v/m_f$ for the independent couplings. However, a detailed analysis of this issue is much more tricky and depends on the kinematic region probed by a given observable. For example, for observables probing the high \sqrt{s} or high p_T tail of differential distributions the validity range will be different than for inclusive observables. See Ref. [42] for a more in-depth discussion of these issues. In this review I restrict to the Higgs signal strength observables in various production modes, which are typically dominated by $\sqrt{s} \sim m_h$. Moreover, I am dodging the question of the validity range because it is assumed from the onset that higher-dimensional operators provide small corrections on top of SM contributions. Consequently, I will only take into account corrections to the observables that are linear in the parameters in Eq. (3.2), which corresponds to retaining only $\mathcal{O}(\Lambda^{-2})$ effects in the EFT expansion.⁵ Incidentally, the LHC so far confirms that the SM is a decent first approximation of the Higgs sector, and deviations due to new physics are small.

4 Observables

Consider the Higgs boson produced at the LHC via the process X , and subsequently decaying to the final state Y . It is possible, to an extent, to isolate experimentally different Higgs boson production modes and decays channels. The LHC collaborations typically quote the Higgs signal strength relative to the SM one in a given channel, here denoted as $\mu_{X;Y}$. Thanks to the narrow width of the Higgs boson, the production and decay can be separated:⁶

$$\mu_{X;Y} = \frac{\sigma(pp \rightarrow X)}{\sigma(pp \rightarrow X)_{\text{SM}}} \frac{\Gamma(h \rightarrow Y)}{\Gamma(h \rightarrow Y)_{\text{SM}}} \frac{\Gamma(h \rightarrow \text{all})_{\text{SM}}}{\Gamma(h \rightarrow \text{all})}. \quad (4.1)$$

Below I summarize how the Higgs production and decays depend on the parameters in the effective Lagrangian. These formulas allow one to derive experimental constraints on the EFT parameters. This kind of approach to LHC Higgs data was pioneered in Refs. [48, 49] and perfected in Refs. [50–87]. As discussed at the end of Section 3, only linear corrections in the independent couplings are kept, while quadratic corrections are ignored. For this reason only CP-even couplings appear in these formulas (the CP-odd ones enter inclusive observables only at the quadratic level). Moreover, I only include $D = 6$ corrections at the *tree level* and I ignore new physics effects suppressed by a loop factor. The exception is the gluon fusion production process which is computed at the next-to-leading order in the $D = 6$ parameters. Unless noted otherwise, I give the inclusive production and decay rates. Note that the signal strength quoted by experiments may depend on analysis-specific cuts, which may slightly change the dependence on the effective theory parameters.

⁵Typically, $\mathcal{O}(\Lambda^{-4})$ effects should be neglected in the context of $D = 6$ effective Lagrangian, as they may receive contributions from $D = 8$ operators. The exception is the observables where the SM contribution is suppressed or vanishes, in which case $D = 6$ operators contribute at $\mathcal{O}(\Lambda^{-4})$, while contributions of higher-order operators are suppressed by more powers of Λ . One example is the lepton-flavor violating Higgs decays into 2 fermions where the SM contribution is exactly zero. In this review I focus on the observables where the SM contribution is dominant.

⁶Except in off-shell Higgs processes [43]. However, given the current precision, these processes do not impose any meaningful constraints within the EFT framework [44–47].

Production

For the relevant partonic processes of Higgs production at the LHC, the cross section relative to the SM one depends on the effective theory parameters as follows:

- Gluon fusion (ggh), $gg \rightarrow h$:

$$\frac{\sigma_{ggh}}{\sigma_{ggh}^{\text{SM}}} \simeq \left| 1 + \frac{\hat{c}_{gg}}{c_{gg}^{\text{SM}}} \right|^2, \quad (4.2)$$

where

$$\begin{aligned} \hat{c}_{gg} &\simeq c_{gg} + \frac{1}{12\pi^2} \left[\delta y_u A_f \left(\frac{m_h^2}{4m_t^2} \right) + \delta y_d A_f \left(\frac{m_h^2}{4m_b^2} \right) \right], \\ c_{gg}^{\text{SM}} &\simeq \frac{1}{12\pi^2} \left[A_f \left(\frac{m_h^2}{4m_t^2} \right) + A_f \left(\frac{m_h^2}{4m_b^2} \right) \right], \\ A_f(\tau) &\equiv \frac{3}{2\tau^2} [(\tau - 1)f(\tau) + \tau], \\ f(\tau) &\equiv \begin{cases} \arcsin^2 \sqrt{\tau} & \tau \leq 1 \\ -\frac{1}{4} \left[\log \frac{1+\sqrt{1-\tau^{-1}}}{1-\sqrt{1-\tau^{-1}}} - i\pi \right]^2 & \tau > 1 \end{cases}. \end{aligned} \quad (4.3)$$

As discussed in Ref. [88], in this case it is appropriate to calculate c_{gg}^{SM} at the leading order in QCD because then the large k-factors, approximately common for c_{gg} and δy_u , cancel in the ratio.⁷ Numerically,

$$\hat{c}_{gg} \simeq c_{gg} + (8.7\delta y_u - (0.3 - 0.3i)\delta y_d) \times 10^{-3}, \quad c_{gg}^{\text{SM}} \simeq (8.4 + 0.3i) \times 10^{-3}, \quad (4.4)$$

$$\frac{\sigma_{ggh}}{\sigma_{ggh}^{\text{SM}}} \simeq 1 + 237c_{gg} + 2.06\delta y_u - 0.06\delta y_d. \quad (4.5)$$

- Vector boson fusion (VBF), $qq \rightarrow hqq$:

$$\begin{aligned} \frac{\sigma_{VBF}}{\sigma_{VBF}^{\text{SM}}} &\simeq 1 + 1.49\delta c_w + 0.51\delta c_z - \begin{pmatrix} 1.08 \\ 1.11 \\ 1.23 \end{pmatrix} c_{w\Box} - 0.10c_{ww} - \begin{pmatrix} 0.35 \\ 0.35 \\ 0.40 \end{pmatrix} c_{z\Box} \\ &\quad - 0.04c_{zz} - 0.10c_{z\Box} - 0.02c_{z\gamma} \\ &\rightarrow 1 + 2\delta c_z - 2.25c_{z\Box} - 0.83c_{zz} + 0.30c_{z\gamma} + 0.12c_{\gamma\gamma}. \end{aligned} \quad (4.6)$$

The numbers in the columns multiplying $c_{w\Box}$ and $c_{z\Box}$ refer to the LHC collision energy of $\sqrt{s} = 7, 8, \text{ and } 13$ TeV; for other parameters the dependence is weaker. The expression after the arrow arises due to replacing the dependent couplings by the independent ones in Eq. (3.2). Each LHC Higgs analysis uses somewhat different cuts to isolate the VBF signal, and the relative cross section slightly depends on these cuts. The result in Eq. (4) has been computed numerically by simulating the parton-level process in MadGraph5 [90] at the tree level with the cuts $p_{T,q} > 20$ GeV, $|\eta_q| < 5$ and $m_{qq} > 250$ GeV. Replacing the last cut by $m_{qq} > 500$ GeV affects the numbers at the level of 5%.

⁷Accidentally, with the SM parameters used in this review, the dependence on δy_d is also captured with a decent accuracy by this procedure. One can compare Eq. (4.5) to NLO QCD results in Ref. [89], where the coefficient in front of δy_d is found to be -0.06 for $\sqrt{s} = 8$ TeV, and -0.05 for $\sqrt{s} = 14$ TeV.

- Vector boson associated production (Vh), $q\bar{q} \rightarrow Vh$, where $V = W, Z$,

$$\begin{aligned}
\frac{\sigma_{Wh}}{\sigma_{Wh}^{\text{SM}}} &\simeq 1 + 2\delta c_w + \begin{pmatrix} 6.39 \\ 6.51 \\ 6.96 \end{pmatrix} c_{w\Box} + \begin{pmatrix} 1.49 \\ 1.49 \\ 1.50 \end{pmatrix} c_{ww} \\
&\rightarrow 1 + 2\delta c_z + \begin{pmatrix} 9.26 \\ 9.43 \\ 10.08 \end{pmatrix} c_{z\Box} + \begin{pmatrix} 4.35 \\ 4.41 \\ 4.63 \end{pmatrix} c_{zz} - \begin{pmatrix} 0.81 \\ 0.84 \\ 0.93 \end{pmatrix} c_{z\gamma} - \begin{pmatrix} 0.43 \\ 0.44 \\ 0.48 \end{pmatrix} c_{\gamma\gamma} \\
\frac{\sigma_{Zh}}{\sigma_{Zh}^{\text{SM}}} &\simeq 1 + 2\delta c_z + \begin{pmatrix} 5.30 \\ 5.40 \\ 5.72 \end{pmatrix} c_{z\Box} + \begin{pmatrix} 1.79 \\ 1.80 \\ 1.82 \end{pmatrix} c_{zz} + \begin{pmatrix} 0.80 \\ 0.82 \\ 0.87 \end{pmatrix} c_{\gamma\Box} + \begin{pmatrix} 0.22 \\ 0.22 \\ 0.22 \end{pmatrix} c_{z\gamma}, \\
&\rightarrow 1 + 2\delta c_z + \begin{pmatrix} 7.61 \\ 7.77 \\ 8.24 \end{pmatrix} c_{z\Box} + \begin{pmatrix} 3.31 \\ 3.35 \\ 3.47 \end{pmatrix} c_{zz} - \begin{pmatrix} 0.58 \\ 0.60 \\ 0.65 \end{pmatrix} c_{z\gamma} + \begin{pmatrix} 0.27 \\ 0.28 \\ 0.30 \end{pmatrix} c_{\gamma\gamma}.
\end{aligned} \tag{4.7}$$

The numbers in the columns refer to the LHC collision energy of $\sqrt{s} = 7, 8, \text{ and } 13$ TeV.

- Top pair associated production, $gg \rightarrow h\bar{t}t$:

$$\frac{\sigma_{t\bar{t}h}}{\sigma_{t\bar{t}h}^{\text{SM}}} \simeq 1 + 2\delta y_u. \tag{4.8}$$

Decay

- $h \rightarrow f\bar{f}$. Higgs boson decays into 2 fermions occur at the tree level in the SM via the Yukawa couplings in Eq. (2.17). In the presence of $D = 6$ operators they are affected via the corrections to the Yukawa couplings in Eq. (3.5):

$$\frac{\Gamma_{cc}}{\Gamma_{cc}^{\text{SM}}} \simeq 1 + 2\delta y_u, \quad \frac{\Gamma_{bb}}{\Gamma_{bb}^{\text{SM}}} \simeq 1 + 2\delta y_d, \quad \frac{\Gamma_{\tau\tau}}{\Gamma_{\tau\tau}^{\text{SM}}} \simeq 1 + 2\delta y_e, \tag{4.9}$$

where I abbreviate $\Gamma(h \rightarrow Y) \equiv \Gamma_Y$.

- $h \rightarrow \mathbf{V}\mathbf{V}$. In the SM, Higgs decays into on-shell gauge bosons: gluon pairs gg , photon pairs $\gamma\gamma$, and $Z\gamma$ occur only at the one-loop level. In the presence of $D = 6$ operators these decays are corrected already at the tree level by the 2-derivative contact interactions of the Higgs boson with two vector bosons in Eq. (3.3). The relative decay widths are given by

$$\frac{\Gamma_{VV}}{\Gamma_{VV}^{\text{SM}}} \simeq \left| 1 + \frac{\hat{c}_{vv}}{c_{vv}^{\text{SM}}} \right|^2, \quad vv \in \{gg, \gamma\gamma, z\gamma\}, \tag{4.10}$$

where

$$\begin{aligned}
\hat{c}_{\gamma\gamma} &= c_{\gamma\gamma}, & c_{\gamma\gamma}^{\text{SM}} &\simeq -8.3 \times 10^{-2}, \\
\hat{c}_{z\gamma} &= c_{z\gamma}, & c_{z\gamma}^{\text{SM}} &\simeq -5.9 \times 10^{-2},
\end{aligned} \tag{4.11}$$

while \hat{c}_{gg} and c_{gg}^{SM} are defined in Eq. (4.3). Note that contributions to $\Gamma_{\gamma\gamma}$ and $\Gamma_{z\gamma}$ arising due to corrections to the SM Higgs couplings to the W bosons and fermions are not included in Eq. (4.11), unlike in Eq. (4.3). The reason is that, for these processes, corrections from $D = 6$ operators are included at the tree level only. If these particular one-loop corrections were included, one should also consistently include *all* one-loop corrections to this process arising at the $D = 6$ level, some of which are divergent and require renormalization. The net result would be to redefine $\hat{c}_{\gamma\gamma} = c_{\gamma\gamma}^{\text{ren.}} - 0.11\delta c_w + 0.02\delta y_u + \dots$, and $\hat{c}_{z\gamma} = c_{z\gamma}^{\text{ren.}} - 0.06\delta c_w + 0.003\delta y_t + \dots$. Here "ren." stands for "renormalized" and the dots stand for a dependence on other Lagrangian parameters (c_{ww} , $c_{w\Box}$, and corrections to triple gauge couplings). A full next-to-leading order computation of these processes have not been yet attempted in the literature.

- **$h \rightarrow 4\ell$.** The decay process $h \rightarrow 2\ell 2\nu$ (where ℓ here stands for charged leptons) proceeds via intermediate W bosons. The relative width is given by

$$\begin{aligned} \frac{\Gamma_{2\ell 2\nu}}{\Gamma_{2\ell 2\nu}^{\text{SM}}} &\simeq 1 + 2\delta c_w + 0.46c_{w\Box} - 0.15c_{ww} \\ &\rightarrow 1 + 2\delta c_z + 0.67c_{z\Box} + 0.05c_{zz} - 0.17c_{z\gamma} - 0.05c_{\gamma\gamma}. \end{aligned} \quad (4.12)$$

In the SM, the decay process $h \rightarrow 4\ell$ proceeds at the tree-level via intermediate Z bosons. In the presence $D = 6$ operators, intermediate photon contributions may also arise at the tree level. If that is the case, the decay width diverges due to the photon pole. Below I quote the relative width $\bar{\Gamma}(h \rightarrow 4\ell)$ regulated by imposing the cut $m_{\ell\ell} > 12$ GeV on the invariant mass of same-flavor lepton pairs:

$$\begin{aligned} \frac{\bar{\Gamma}_{4\ell}}{\bar{\Gamma}_{4\ell}^{\text{SM}}} &\simeq 1 + 2\delta c_z + \begin{pmatrix} 0.41 \\ 0.39 \end{pmatrix} c_{z\Box} - \begin{pmatrix} 0.15 \\ 0.14 \end{pmatrix} c_{zz} + \begin{pmatrix} 0.07 \\ 0.05 \end{pmatrix} c_{z\gamma} - \begin{pmatrix} 0.02 \\ 0.02 \end{pmatrix} c_{\gamma\Box} + \begin{pmatrix} < 0.01 \\ 0.03 \end{pmatrix} c_{\gamma\gamma} \\ &\rightarrow 1 + 2\delta c_z + \begin{pmatrix} 0.35 \\ 0.32 \end{pmatrix} c_{z\Box} - \begin{pmatrix} 0.19 \\ 0.19 \end{pmatrix} c_{zz} + \begin{pmatrix} 0.09 \\ 0.08 \end{pmatrix} c_{z\gamma} + \begin{pmatrix} 0.01 \\ 0.02 \end{pmatrix} c_{\gamma\gamma}. \end{aligned} \quad (4.13)$$

The numbers in the columns correspond to the $2e2\mu$ and $4e/\mu$ final states, respectively. The difference between these two is numerically irrelevant in the total width, but may be important for differential distributions, especially regarding the $c_{\gamma\gamma}$ dependence [91]. The dependence on the $m_{\ell\ell}$ cut is weak; very similar numbers are obtained if $m_{\ell\ell} > 4$ GeV is imposed instead.

Given the partial widths, the branching fractions can be computed as $\text{Br}_Y = \Gamma_Y/\Gamma(h \rightarrow \text{all})$, where the total decay width is given by

$$\frac{\Gamma(h \rightarrow \text{all})}{\Gamma(h \rightarrow \text{all})} \simeq \frac{\Gamma_{bb}}{\Gamma_{bb}^{\text{SM}}} \text{Br}_{bb}^{\text{SM}} + \frac{\Gamma_{cc}}{\Gamma_{cc}^{\text{SM}}} \text{Br}_{cc}^{\text{SM}} + \frac{\Gamma_{\tau\tau}}{\Gamma_{\tau\tau}^{\text{SM}}} \text{Br}_{\tau\tau}^{\text{SM}} + \frac{\Gamma_{WW^*}}{\Gamma_{WW^*}^{\text{SM}}} \text{Br}_{WW^*}^{\text{SM}} + \frac{\Gamma_{ZZ^*}}{\Gamma_{ZZ^*}^{\text{SM}}} \text{Br}_{ZZ^*}^{\text{SM}} + \frac{\Gamma_{gg}}{\Gamma_{gg}^{\text{SM}}} \text{Br}_{gg}^{\text{SM}}. \quad (4.14)$$

Note that, in line with the basic assumption of no new light particles, there is no additional contributions to the Higgs width other than from the SM decay channels. In particular, the invisible Higgs width is absent in this EFT framework (except for the small SM contribution arising via $h \rightarrow ZZ^* \rightarrow 4\nu$).

5 Current Constraints

In this section I present the constraints on the independent couplings characterizing the Higgs boson couplings in the dimension six EFT Lagrangian. A disclaimer is in order. The objective is to illustrate what is the constraining power of the present data. As we will see, the existing data is not yet good enough to even constrain all the couplings inside the EFT validity range. In the future, as the measurements become more precise and more information is available, this kind of analysis will become fully consistent.

5.1 Data

I first review the experimental data used to constrain the effective theory parameters. In the best of all worlds, the LHC collaborations would quote a multi-dimensional likelihood function for the signal strength $\mu_{X,Y}$ for all production modes and decay channels, separately for each LHC collision energy. This would allow one to consistently use available experimental information, including non-trivial correlations between the different μ 's. Although the manner in which the LHC data is presented has been constantly improving, we are not yet in the ideal world. For these reasons, constraining Higgs couplings from existing data involves inevitably somewhat arbitrary assumptions and approximations. Nevertheless, thanks to the fact that the experimental uncertainties are still statistics-dominated in most cases, one should expect that these approximations do not affect the results in a dramatic way.

The measurements of the Higgs signal strength μ included in this analysis are summarized in Table 2. They are separated according to the final state (channel) and the production mode. For the all-inclusive production mode (total) I use the value of μ quoted in the table.⁸ The same is true for μ in a specific production mode (Wh, Zh, tth), in which case I ignore correlations with other production channels. In the remaining cases μ is quoted for illustration only, and more information is included in the analysis. 2D stands for two-dimensional likelihood functions in the plane $\mu_{ggh+tth}-\mu_{VBF+Vh}$. Since, the contribution of the Vh production mode is subleading with respect to the VBF one, I combine the separate measurements of the Vh signal strength (whenever it is given) with the 2D likelihood, ignoring the correlation between the two. For the diphoton final state I construct five-dimensional likelihood function in the space spanned by $(\mu_{ggh}, \mu_{tth}, \mu_{VBF}, \mu_{Wh}, \mu_{Zh})$ using the signal strength in all diphoton event categories (cats.), using the known contribution of each production mode to each category. In many channels there is a certain degree of arbitrariness as to which set of results (inclusive, 1D, 2D, or cats.) to include in the fit; here the strategy is to choose the set that maximizes the available information about various EFT couplings.

⁸CMS does not quote the best-fit μ in the $Z\gamma$ channel. The value in Table 2 was obtained by digitizing the plot showing the expected and observed 95% CL limits on μ in function of m_h , extracting the values at $m_h = 125$ GeV, and using these to calculate the best-fit μ assuming the uncertainties are Gaussian. This is a dire reminder of how Higgs fits had to be done back in the early 2010s.

ATLAS				CMS			
Channel	μ	Production	Ref.	Channel	μ	Production	Ref.
$\gamma\gamma$	$1.17^{+0.28}_{-0.26}$	cats.	[92]	$\gamma\gamma$	$1.12^{+0.25}_{-0.22}$	cats.	[101]
$Z\gamma$	$2.7^{+4.6}_{-4.5}$	total	[93]	$Z\gamma$	$-0.2^{+4.9}_{-4.9}$	total	[102]
ZZ^*	$1.46^{+0.40}_{-0.34}$	2D	[94]	ZZ^*	$1.00^{+0.29}_{-0.29}$	2D	[103]
WW^*	$1.18^{+0.24}_{-0.21}$	2D	[95]	WW^*	$0.83^{+0.21}_{-0.21}$	2D	[103]
	$2.1^{+1.9}_{-1.6}$	Wh	[96]		$0.80^{+1.09}_{-0.93}$	Vh	[103]
	$5.1^{+4.3}_{-3.1}$	Zh	[96]	$\tau\tau$	$0.91^{+0.28}_{-0.28}$	2D	[103]
$\tau\tau$	$1.44^{+0.42}_{-0.37}$	2D	[97]		$0.87^{+1.00}_{-0.88}$	Vh	[103]
bb	$1.11^{+0.65}_{-0.61}$	Wh	[98]		$-1.3^{+6.3}_{-5.5}$	tth	[104]
	$0.05^{+0.52}_{-0.49}$	Zh	[98]	bb	$0.89^{+0.47}_{-0.44}$	Vh	[103]
	$1.5^{+1.1}_{-1.1}$	tth	[99]		$1.2^{+1.6}_{-1.5}$	tth	[105]
$\mu\mu$	$-0.7^{+3.7}_{-3.7}$	total	[93]	$\mu\mu$	$0.8^{+3.5}_{-3.4}$	total	[106]
multi- ℓ	$2.1^{+1.4}_{-1.2}$	tth	[100]	multi- ℓ	$3.8^{+1.4}_{-1.4}$	tth	[104]

Table 2: The LHC Higgs results used in the fit. See Section 5.1 for explanations.

5.2 Fit

Using the dependence of the signal strength on EFT parameters worked out in Section 4 and the LHC data in Table 2 one can constrain all *CP-even* independent Higgs couplings in Eq. (3.2).⁹ In the Gaussian approximation near the best fit point I find the following constraints:

$$\begin{pmatrix} \delta c_z \\ c_{zz} \\ c_{z\Box} \\ c_{\gamma\gamma} \\ c_{z\gamma} \\ c_{gg} \\ \delta y_u \\ \delta y_d \\ \delta y_e \end{pmatrix} = \begin{pmatrix} -0.12 \pm 0.20 \\ 0.5 \pm 1.8 \\ -0.21 \pm 0.82 \\ 0.014 \pm 0.029 \\ 0.01 \pm 0.10 \\ -0.0056 \pm 0.0028 \\ 0.55 \pm 0.30 \\ -0.42 \pm 0.45 \\ -0.17 \pm 0.35 \end{pmatrix}, \quad (5.1)$$

⁹To constrain the CP-odd couplings $\sin\phi_f$ and $\tilde{c}_{\nu\nu}$ within the EFT framework one should study the differential distributions in multi-body Higgs decays where these couplings enter at the linear level [107–114].

where the uncertainties correspond to 1σ . The correlation matrix is

$$\rho = \begin{pmatrix} 1. & -0.23 & 0.17 & -0.62 & -0.18 & 0.16 & 0.09 & 0.88 & 0.63 \\ \cdot & 1. & -0.997 & 0.85 & 0.23 & 0.13 & 0.17 & -0.47 & -0.81 \\ \cdot & \cdot & 1. & -0.82 & -0.21 & -0.15 & -0.17 & 0.41 & 0.78 \\ \cdot & \cdot & \cdot & 1. & 0.27 & 0.02 & 0.09 & -0.79 & -0.92 \\ \cdot & \cdot & \cdot & \cdot & 1. & 0.01 & 0.02 & -0.22 & -0.26 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1. & -0.81 & 0.21 & 0.03 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1. & 0.05 & -0.06 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1. & 0.82 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1. \end{pmatrix}. \quad (5.2)$$

Using the above central values c_0 , uncertainties δc , and the correlation matrix ρ , one can reconstruct the 9-dimensional likelihood function near the best fit point:

$$\chi^2 \simeq \sum_{ij} [c - c_0]_i \sigma_{ij}^{-2} [c - c_0]_j, \quad \sigma_{ij}^{-2} \equiv [[\delta c]_i \rho_{ij} [\delta c]_j]^{-1}. \quad (5.3)$$

5.3 Discussion

As one can see from Eq. (5.1), certain EFT parameters are very weakly constrained by experiment, with order one deviations from the SM being allowed. In other words, the current data cannot even constrain all the parameters to be within the EFT validity range. This violates the initial assumption that the $D = 6$ operators give a small correction on top of the SM. For this reason, the results in Eq. (5.1) should not be taken at face value. In particular, one should conclude that there is currently no model-independent constraints at all on c_{zz} and $c_{z\Box}$. Indeed, including corrections to observables that are quadratic in these parameters would completely change the central values and the uncertainties. This signals a sensitivity of the fit to operators with $D > 6$. Furthermore, the experimental constraints in the $Z\gamma$ channel are still too weak to justify the linear approximation. Again, including quadratic EFT corrections would significantly affect the constraints on $c_{z\gamma}$. To a lesser extent, the sensitivity to higher-order EFT corrections is also true for deformations along the δy_f directions.

Nevertheless, the results in Eq. (5.1) are of some value. First of all, they demonstrate that *certain* EFT parameters are strongly constrained. This is true especially for c_{gg} and $c_{\gamma\gamma}$ who are constrained at the 10^{-3} level. Next, the fit in Eq. (5.1) identifies “blind” directions in the space of the EFT parameters that are weakly constrained by current data. The most dramatic example is the approximate degeneracy along the line $c_{zz} \approx -2.3c_{z\Box}$, as witnessed by the ≈ 1 entry in the correlation matrix in Eq. (5.2). More data is needed to lift this degeneracy. To this end, extremely helpful pieces of information can be extracted from differential distributions in $h \rightarrow 4\ell$ decays [108, 115–119], as well as in the Vh [42, 120–125] and VBF [126–128] production. A consistent, model-independent EFT approach to Higgs differential distributions has not yet been implemented in LHC analyses, but the CMS collaboration made first steps in this direction [129]. Note also the large correlations between δy_d and other parameters. This happens because δy_d strongly affects the total Higgs width (via the $h \rightarrow b\bar{b}$ partial width) and this way it affects the signal strength in all Higgs decay channels. More precise measurements of the signal strength in the $h \rightarrow b\bar{b}$ channel should soon alleviate this degeneracy. Finally, there is the strong correlation

between c_{gg} and δy_t which has been extensively discussed in the Higgs fits literature. In the future, that degeneracy will be lifted by better measurements of the $t\bar{t}$ signal strength, and by measurements of the Higgs p_T distribution in the gluon fusion production mode [130–134] (see also [135] for an earlier work in this direction).

Finally, the importance of the fit is in the fact that the likelihood in Eq. (5.3) can be combined with other datasets that constrain the same EFT parameters. In this case, one may obtain stronger bounds that will push the parameters into the EFT validity range. For example, one can use constraints on cubic self-couplings of electroweak gauge bosons [83, 86, 119, 136–139]. These are customarily parametrized by 3 parameters $\delta g_{1,z}$, $\delta\kappa_\gamma$, λ_z [140] which characterize deviations of these self-couplings from the SM predictions. Now, in the EFT Lagrangian with $D = 6$ operators the first two parameters are related to the Higgs couplings. In the Higgs basis one finds [26]:

$$\begin{aligned}\delta g_{1,z} &= \frac{1}{2(g_L^2 - g_Y^2)} [c_{\gamma\gamma} e^2 g_Y^2 + c_{z\gamma} (g_L^2 - g_Y^2) g_Y^2 - c_{zz} (g_L^2 + g_Y^2) g_Y^2 - c_{z\Box} (g_L^2 + g_Y^2) g_L^2], \\ \delta\kappa_\gamma &= -\frac{g_L^2}{2} \left(c_{\gamma\gamma} \frac{e^2}{g_L^2 + g_Y^2} + c_{z\gamma} \frac{g_L^2 - g_Y^2}{g_L^2 + g_Y^2} - c_{zz} \right).\end{aligned}\tag{5.4}$$

Therefore, model-independent constraints on triple gauge couplings imply additional constraint on the EFT parameters characterizing the Higgs couplings. In particular, Ref. [138] argues that, after marginalizing over λ_z , the single and pair W boson production in LEP-2 implies the bounds $\delta g_{1,z} = -0.83 \pm 0.34$, $\delta\kappa_\gamma = 0.14 \pm 0.05$ with the correlation coefficient $[\rho]_{\delta g_{1,z}}^{\delta\kappa_\gamma} = -0.71$. Combining this bound with the likelihood in Eq. (5.3) the degeneracy between c_{zz} and $c_{z\Box}$ is lifted, and one obtains much stronger bounds: $c_{zz} = 0.22 \pm 0.18$, $c_{z\Box} = -0.08 \pm 0.09$, $[\rho]_{c_{zz}}^{c_{z\Box}} = -0.76$. More constraints of this type, for example model-independent constraints on triple gauge couplings from the LHC, could further improve the limits on Higgs couplings within the EFT approach. As soon as more precise Higgs and di-boson data from the 13 TeV LHC run start arriving, it should be possible to constrain all the 9 parameters in Eq. (5.1) safely within the EFT validity range.

6 Closing Words

The Higgs boson has been discovered, and for the remainder of this century we will study its properties. Precision measurements of Higgs couplings and determination of their tensor structure is an important part of the physics program at the LHC and future colliders. Given that no slightest hint for a particular scenario beyond the SM has emerged so far, it is important to (also) perform these studies in a model-independent framework. The EFT approach described here, with the SM extended by dimension six operators, provides a perfect tool to this end.

One should be aware that Higgs precision measurements cannot probe new physics at very high scales. For example, LHC Higgs measurements are sensitive to new physics at $\Lambda \sim 1$ TeV at the most. This is not too impressive, especially compared to the new physics reach of flavor observables or even electroweak precision tests. However, Higgs physics probes a subset of operators that are often not accessible by other searches. For example, for most of the 9 parameters in Eq. (5.1) the *only* experimental constraints come from Higgs physics. It is certainly conceivable that new physics talks to the SM via the Higgs portal, and it will first manifest itself within this particular class of $D = 6$ operators. If this is the case, we must not miss it.

Acknowledgements

I am supported by the ERC Advanced Grant Higgs@LHC.

References

- [1] S. Weinberg, *A Model of Leptons*, *Phys.Rev.Lett.* **19** (1967) 1264–1266.
- [2] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys.Rev.Lett.* **13** (1964) 321–323.
- [3] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Phys.Lett.* **12** (1964) 132–133.
- [4] G. Guralnik, C. Hagen, and T. Kibble, *Global Conservation Laws and Massless Particles*, *Phys.Rev.Lett.* **13** (1964) 585–587.
- [5] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, *Phys.Rev.Lett.* **13** (1964) 508–509.
- [6] P. W. Higgs, *Spontaneous Symmetry Breakdown without Massless Bosons*, *Phys.Rev.* **145** (1966) 1156–1163.
- [7] T. Kibble, *Symmetry breaking in nonAbelian gauge theories*, *Phys.Rev.* **155** (1967) 1554–1561.
- [8] **ATLAS** Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys.Lett.* **B716** (2012) 1–29, [[arXiv:1207.7214](#)].
- [9] **CMS** Collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys.Lett.* **B716** (2012) 30–61, [[arXiv:1207.7235](#)].
- [10] B. Grinstein and M. Trott, *A Higgs-Higgs bound state due to new physics at a TeV*, *Phys.Rev.* **D76** (2007) 073002, [[arXiv:0704.1505](#)].
- [11] R. Alonso, M. Gavela, L. Merlo, S. Rigolin, and J. Yepes, *The Effective Chiral Lagrangian for a Light Dynamical "Higgs Particle"*, *Phys.Lett.* **B722** (2013) 330–335, [[arXiv:1212.3305](#)].
- [12] G. Isidori, A. V. Manohar, and M. Trott, *Probing the nature of the Higgs-like Boson via $h \rightarrow V\mathcal{F}$ decays*, *Phys.Lett.* **B728** (2014) 131–135, [[arXiv:1305.0663](#)].
- [13] G. Buchalla, O. Catà, and C. Krause, *Complete Electroweak Chiral Lagrangian with a Light Higgs at NLO*, *Nucl.Phys.* **B880** (2014) 552–573, [[arXiv:1307.5017](#)].
- [14] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl.Phys.* **B268** (1986) 621–653.

- [15] F. Goertz, *Indirect Handle on the Down-Quark Yukawa Coupling*, *Phys.Rev.Lett.* **113** (2014), no. 26 261803, [arXiv:1406.0102].
- [16] W. Altmannshofer, J. Brod, and M. Schmaltz, *Experimental constraints on the coupling of the Higgs boson to electrons*, arXiv:1503.04830.
- [17] G. T. Bodwin, F. Petriello, S. Stoynev, and M. Velasco, *Higgs boson decays to quarkonia and the $H\bar{c}c$ coupling*, *Phys.Rev.* **D88** (2013), no. 5 053003, [arXiv:1306.5770].
- [18] A. L. Kagan, G. Perez, F. Petriello, Y. Soreq, S. Stoynev, et al., *Exclusive Window onto Higgs Yukawa Couplings*, *Phys.Rev.Lett.* **114** (2015), no. 10 101802, [arXiv:1406.1722].
- [19] H. K. Dreiner, H. E. Haber, and S. P. Martin, *Two-component spinor techniques and Feynman rules for quantum field theory and supersymmetry*, *Phys.Rept.* **494** (2010) 1–196, [arXiv:0812.1594].
- [20] **CMS s** Collaboration, G. Aad et al., *Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments*, arXiv:1503.07589.
- [21] **CDF Collaboration, D0 Collaboration** Collaboration, T. E. W. Group, *2012 Update of the Combination of CDF and D0 Results for the Mass of the W Boson*, arXiv:1204.0042.
- [22] **ALEPH, DELPHI, L3, OPAL, SLD, LEP Electroweak Working Group, SLD Electroweak Group, SLD Heavy Flavour Group** Collaboration, S. Schael et al., *Precision electroweak measurements on the Z resonance*, *Phys.Rept.* **427** (2006) 257–454, [hep-ex/0509008].
- [23] **Particle Data Group** Collaboration, J. Beringer et al., *Review of Particle Physics (RPP)*, *Phys.Rev.* **D86** (2012) 010001.
- [24] H. Burkhardt and B. Pietrzyk, *Recent BES measurements and the hadronic contribution to the QED vacuum polarization*, *Phys.Rev.* **D84** (2011) 037502, [arXiv:1106.2991].
- [25] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **1404** (2014) 159, [arXiv:1312.2014].
- [26] **LHC Higgs Cross Section Working Group 2** Collaboration, *Higgs Basis: Proposal for an EFT basis choice for LHC HXSWG*, LHCHSWG-INT-2015-001 cds.cern.ch/record/2001958.
- [27] R. S. Gupta, A. Pomarol, and F. Riva, *BSM Primary Effects*, *Phys.Rev.* **D91** (2015), no. 3 035001, [arXiv:1405.0181].
- [28] A. Pomarol, *Higgs Physics*, arXiv:1412.4410.
- [29] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **1010** (2010) 085, [arXiv:1008.4884].

- [30] G. Giudice, C. Grojean, A. Pomarol, and R. Rattazzi, *The Strongly-Interacting Light Higgs*, *JHEP* **0706** (2007) 045, [[hep-ph/0703164](#)].
- [31] R. Contino, M. Ghezzi, C. Grojean, M. Muhlleitner, and M. Spira, *Effective Lagrangian for a light Higgs-like scalar*, *JHEP* **1307** (2013) 035, [[arXiv:1303.3876](#)].
- [32] A. Efrati, A. Falkowski, and Y. Soreq, *Electroweak constraints on flavorful effective theories*, [arXiv:1503.07872](#).
- [33] G. Blankenburg, J. Ellis, and G. Isidori, *Flavour-Changing Decays of a 125 GeV Higgs-like Particle*, *Phys.Lett.* **B712** (2012) 386–390, [[arXiv:1202.5704](#)].
- [34] R. Harnik, J. Kopp, and J. Zupan, *Flavor Violating Higgs Decays*, *JHEP* **1303** (2013) 026, [[arXiv:1209.1397](#)].
- [35] R. Contino, M. Ghezzi, M. Moretti, G. Panico, F. Piccinini, et al., *Anomalous Couplings in Double Higgs Production*, *JHEP* **1208** (2012) 154, [[arXiv:1205.5444](#)].
- [36] M. J. Dolan, C. Englert, and M. Spannowsky, *Higgs self-coupling measurements at the LHC*, *JHEP* **1210** (2012) 112, [[arXiv:1206.5001](#)].
- [37] M. McCullough, *An Indirect Model-Dependent Probe of the Higgs Self-Coupling*, *Phys.Rev.* **D90** (2014), no. 1 015001, [[arXiv:1312.3322](#)].
- [38] C. Englert, F. Krauss, M. Spannowsky, and J. Thompson, *Di-Higgs phenomenology in $t\bar{t}hh$: The forgotten channel*, *Phys.Lett.* **B743** (2015) 93–97, [[arXiv:1409.8074](#)].
- [39] F. Goertz, A. Papaefstathiou, L. L. Yang, and J. Zurita, *Higgs boson pair production in the $D=6$ extension of the SM*, [arXiv:1410.3471](#).
- [40] A. Azatov, R. Contino, G. Panico, and M. Son, *Effective field theory analysis of double Higgs production via gluon fusion*, [arXiv:1502.00539](#).
- [41] R. Grober, M. Muhlleitner, M. Spira, and J. Streicher, *NLO QCD Corrections to Higgs Pair Production including Dimension-6 Operators*, [arXiv:1504.06577](#).
- [42] A. Biekötter, A. Knochel, M. Kraemer, D. Liu, and F. Riva, *Vices and Virtues of Higgs EFTs at Large Energy*, [arXiv:1406.7320](#).
- [43] F. Caola and K. Melnikov, *Constraining the Higgs boson width with ZZ production at the LHC*, *Phys.Rev.* **D88** (2013) 054024, [[arXiv:1307.4935](#)].
- [44] C. Englert and M. Spannowsky, *Limitations and Opportunities of Off-Shell Coupling Measurements*, *Phys.Rev.* **D90** (2014), no. 5 053003, [[arXiv:1405.0285](#)].
- [45] G. Cacciapaglia, A. Deandrea, G. Drieu La Rochelle, and J.-B. Flament, *Higgs couplings: disentangling New Physics with off-shell measurements*, *Phys.Rev.Lett.* **113** (2014), no. 20 201802, [[arXiv:1406.1757](#)].

- [46] A. Azatov, C. Grojean, A. Paul, and E. Salvioni, *Taming the off-shell Higgs boson*, *Zh.Eksp.Teor.Fiz.* **147** (2015) 410–425, [arXiv:1406.6338].
- [47] C. Englert, Y. Soreq, and M. Spannowsky, *Off-Shell Higgs Coupling Measurements in BSM scenarios*, arXiv:1410.5440.
- [48] D. Carmi, A. Falkowski, E. Kuflik, and T. Volansky, *Interpreting LHC Higgs Results from Natural New Physics Perspective*, *JHEP* **1207** (2012) 136, [arXiv:1202.3144].
- [49] A. Azatov, R. Contino, and J. Galloway, *Model-Independent Bounds on a Light Higgs*, *JHEP* **1204** (2012) 127, [arXiv:1202.3415].
- [50] J. Espinosa, C. Grojean, M. Muhlleitner, and M. Trott, *Fingerprinting Higgs Suspects at the LHC*, *JHEP* **1205** (2012) 097, [arXiv:1202.3697].
- [51] M. Rauch, *Determination of Higgs-boson couplings (SFitter)*, arXiv:1203.6826.
- [52] P. P. Giardino, K. Kannike, M. Raidal, and A. Strumia, *Reconstructing Higgs boson properties from the LHC and Tevatron data*, *JHEP* **1206** (2012) 117, [arXiv:1203.4254].
- [53] J. Ellis and T. You, *Global Analysis of Experimental Constraints on a Possible Higgs-Like Particle with Mass ~ 125 GeV*, *JHEP* **1206** (2012) 140, [arXiv:1204.0464].
- [54] A. Azatov, R. Contino, D. Del Re, J. Galloway, M. Grassi, et al., *Determining Higgs couplings with a model-independent analysis of $h \rightarrow \gamma\gamma$* , *JHEP* **1206** (2012) 134, [arXiv:1204.4817].
- [55] M. Farina, C. Grojean, and E. Salvioni, *(Dys)Zphilia or a custodial breaking Higgs at the LHC*, *JHEP* **1207** (2012) 012, [arXiv:1205.0011].
- [56] M. Klute, R. Lafaye, T. Plehn, M. Rauch, and D. Zerwas, *Measuring Higgs Couplings from LHC Data*, *Phys.Rev.Lett.* **109** (2012) 101801, [arXiv:1205.2699].
- [57] T. Corbett, O. Eboli, J. Gonzalez-Fraile, and M. Gonzalez-Garcia, *Constraining anomalous Higgs interactions*, *Phys.Rev.* **D86** (2012) 075013, [arXiv:1207.1344].
- [58] P. P. Giardino, K. Kannike, M. Raidal, and A. Strumia, *Is the resonance at 125 GeV the Higgs boson?*, *Phys.Lett.* **B718** (2012) 469–474, [arXiv:1207.1347].
- [59] J. Ellis and T. You, *Global Analysis of the Higgs Candidate with Mass ~ 125 GeV*, *JHEP* **1209** (2012) 123, [arXiv:1207.1693].
- [60] M. Montull and F. Riva, *Higgs discovery: the beginning or the end of natural EWSB?*, *JHEP* **1211** (2012) 018, [arXiv:1207.1716].
- [61] J. Espinosa, C. Grojean, M. Muhlleitner, and M. Trott, *First Glimpses at Higgs' face*, *JHEP* **1212** (2012) 045, [arXiv:1207.1717].
- [62] D. Carmi, A. Falkowski, E. Kuflik, T. Volansky, and J. Zupan, *Higgs After the Discovery: A Status Report*, *JHEP* **1210** (2012) 196, [arXiv:1207.1718].

- [63] S. Banerjee, S. Mukhopadhyay, and B. Mukhopadhyaya, *New Higgs interactions and recent data from the LHC and the Tevatron*, *JHEP* **1210** (2012) 062, [[arXiv:1207.3588](#)].
- [64] F. Bonnet, T. Ota, M. Rauch, and W. Winter, *Interpretation of precision tests in the Higgs sector in terms of physics beyond the Standard Model*, *Phys.Rev.* **D86** (2012) 093014, [[arXiv:1207.4599](#)].
- [65] T. Plehn and M. Rauch, *Higgs Couplings after the Discovery*, *Europhys.Lett.* **100** (2012) 11002, [[arXiv:1207.6108](#)].
- [66] A. Djouadi, *Precision Higgs coupling measurements at the LHC through ratios of production cross sections*, *Eur.Phys.J.* **C73** (2013) 2498, [[arXiv:1208.3436](#)].
- [67] B. Batell, S. Gori, and L.-T. Wang, *Higgs Couplings and Precision Electroweak Data*, *JHEP* **1301** (2013) 139, [[arXiv:1209.6382](#)].
- [68] T. Corbett, O. Eboli, J. Gonzalez-Fraile, and M. Gonzalez-Garcia, *Robust Determination of the Higgs Couplings: Power to the Data*, *Phys.Rev.* **D87** (2013) 015022, [[arXiv:1211.4580](#)].
- [69] D. Choudhury, R. Islam, and A. Kundu, *Anomalous Higgs Couplings as a Window to New Physics*, *Phys.Rev.* **D88** (2013), no. 1 013014, [[arXiv:1212.4652](#)].
- [70] G. Belanger, B. Dumont, U. Ellwanger, J. Gunion, and S. Kraml, *Higgs Couplings at the End of 2012*, *JHEP* **1302** (2013) 053, [[arXiv:1212.5244](#)].
- [71] K. Cheung, J. S. Lee, and P.-Y. Tseng, *Higgs Precision (Higgcision) Era begins*, *JHEP* **1305** (2013) 134, [[arXiv:1302.3794](#)].
- [72] A. Falkowski, F. Riva, and A. Urbano, *Higgs at last*, *JHEP* **1311** (2013) 111, [[arXiv:1303.1812](#)].
- [73] P. P. Giardino, K. Kannike, I. Masina, M. Raidal, and A. Strumia, *The universal Higgs fit*, *JHEP* **1405** (2014) 046, [[arXiv:1303.3570](#)].
- [74] J. Ellis and T. You, *Updated Global Analysis of Higgs Couplings*, *JHEP* **1306** (2013) 103, [[arXiv:1303.3879](#)].
- [75] A. Djouadi and G. Moreau, *The couplings of the Higgs boson and its CP properties from fits of the signal strengths and their ratios at the 7+8 TeV LHC*, *Eur.Phys.J.* **C73** (2013), no. 9 2512, [[arXiv:1303.6591](#)].
- [76] B. Dumont, S. Fichet, and G. von Gersdorff, *A Bayesian view of the Higgs sector with higher dimensional operators*, *JHEP* **1307** (2013) 065, [[arXiv:1304.3369](#)].
- [77] P. Bechtle, S. Heinemeyer, O. Stl, T. Stefaniak, and G. Weiglein, *HiggsSignals: Confronting arbitrary Higgs sectors with measurements at the Tevatron and the LHC*, *Eur.Phys.J.* **C74** (2014), no. 2 2711, [[arXiv:1305.1933](#)].

- [78] G. Belanger, B. Dumont, U. Ellwanger, J. Gunion, and S. Kraml, *Global fit to Higgs signal strengths and couplings and implications for extended Higgs sectors*, *Phys.Rev.* **D88** (2013) 075008, [[arXiv:1306.2941](#)].
- [79] M. Ciuchini, E. Franco, S. Mishima, and L. Silvestrini, *Electroweak Precision Observables, New Physics and the Nature of a 126 GeV Higgs Boson*, *JHEP* **1308** (2013) 106, [[arXiv:1306.4644](#)].
- [80] P. Artoisenet, P. de Aquino, F. Demartin, R. Frederix, S. Frixione, et al., *A framework for Higgs characterisation*, *JHEP* **1311** (2013) 043, [[arXiv:1306.6464](#)].
- [81] **LHC Higgs Cross Section Working Group** Collaboration, S. Heinemeyer et al., *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, [arXiv:1307.1347](#).
- [82] S. Choi, S. Jung, and P. Ko, *Implications of LHC data on 125 GeV Higgs-like boson for the Standard Model and its various extensions*, *JHEP* **1310** (2013) 225, [[arXiv:1307.3948](#)].
- [83] A. Pomarol and F. Riva, *Towards the Ultimate SM Fit to Close in on Higgs Physics*, *JHEP* **1401** (2014) 151, [[arXiv:1308.2803](#)].
- [84] H. Belusca-Maito, *Effective Higgs Lagrangian and Constraints on Higgs Couplings*, [arXiv:1404.5343](#).
- [85] M. Baak, J. Cuth, J. Haller, A. Hoecker, R. Kogler, et al., *The global electroweak fit at NNLO and prospects for the LHC and ILC*, [arXiv:1407.3792](#).
- [86] J. Ellis, V. Sanz, and T. You, *The Effective Standard Model after LHC Run I*, *JHEP* **1503** (2015) 157, [[arXiv:1410.7703](#)].
- [87] J.-B. Flament, *Higgs couplings and BSM physics: Run I Legacy constraints*, [arXiv:1504.07919](#).
- [88] S. Gori and I. Low, *Precision Higgs Measurements: Constraints from New Oblique Corrections*, *JHEP* **1309** (2013) 151, [[arXiv:1307.0496](#)].
- [89] R. Harlander, M. Mhleitner, J. Rathsman, M. Spira, and O. Stl, *Interim recommendations for the evaluation of Higgs production cross sections and branching ratios at the LHC in the Two-Higgs-Doublet Model*, [arXiv:1312.5571](#).
- [90] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **1407** (2014) 079, [[arXiv:1405.0301](#)].
- [91] Y. Chen, R. Harnik, and R. Vega-Morales, *New Opportunities in $h \rightarrow 4\ell$* , [arXiv:1503.05855](#).
- [92] **ATLAS** Collaboration, G. Aad et al., *Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector*, *Phys.Rev.* **D90** (2014), no. 11 112015, [[arXiv:1408.7084](#)].

- [93] **ATLAS** Collaboration, *Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $s = 7$ and 8 TeV in the ATLAS experiment*, *ATLAS-CONF-2015-007* (2015).
- [94] **ATLAS Collaboration** Collaboration, G. Aad et al., *Measurements of Higgs boson production and couplings in the four-lepton channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector*, *Phys.Rev.* **D91** (2015), no. 1 012006, [[arXiv:1408.5191](#)].
- [95] **ATLAS Collaboration** Collaboration, G. Aad et al., *Observation and measurement of Higgs boson decays to WW^* with the ATLAS detector*, [arXiv:1412.2641](#).
- [96] **ATLAS** Collaboration, *Study of the Higgs boson decaying to WW^* produced in association with a weak boson with the ATLAS detector at the LHC*, *ATLAS-CONF-2015-005* (2015).
- [97] **ATLAS** Collaboration, G. Aad et al., *Evidence for the Higgs-boson Yukawa coupling to tau leptons with the ATLAS detector*, [arXiv:1501.04943](#).
- [98] **ATLAS** Collaboration, G. Aad et al., *Search for the $b\bar{b}$ decay of the Standard Model Higgs boson in associated $(W/Z)H$ production with the ATLAS detector*, *JHEP* **1501** (2015) 069, [[arXiv:1409.6212](#)].
- [99] **ATLAS** Collaboration, G. Aad et al., *Search for the Standard Model Higgs boson produced in association with top quarks and decaying into $b\bar{b}$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, [arXiv:1503.05066](#).
- [100] **ATLAS** Collaboration, *Search for the associated production of the Higgs boson with a top quark pair in multi-lepton final states with the ATLAS detector*, *ATLAS-CONF-2015-006* (2015).
- [101] **CMS** Collaboration, V. Khachatryan et al., *Observation of the diphoton decay of the Higgs boson and measurement of its properties*, *Eur.Phys.J.* **C74** (2014), no. 10 3076, [[arXiv:1407.0558](#)].
- [102] **CMS** Collaboration, S. Chatrchyan et al., *Search for a Higgs boson decaying into a Z and a photon in pp collisions at $\sqrt{s} = 7$ and 8 TeV*, *Phys.Lett.* **B726** (2013) 587–609, [[arXiv:1307.5515](#)].
- [103] **CMS** Collaboration, V. Khachatryan et al., *Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV*, [arXiv:1412.8662](#).
- [104] **CMS** Collaboration, V. Khachatryan et al., *Search for the associated production of the Higgs boson with a top-quark pair*, *JHEP* **1409** (2014) 087, [[arXiv:1408.1682](#)].
- [105] **CMS** Collaboration, V. Khachatryan et al., *Search for a standard model Higgs boson produced in association with a top-quark pair and decaying to bottom quarks using a matrix element method*, [arXiv:1502.02485](#).

- [106] CMS Collaboration, V. Khachatryan et al., *Search for a standard model-like Higgs boson in the $\mu^+\mu^-$ and e^+e^- decay channels at the LHC*, arXiv:1410.6679.
- [107] F. Bishara, Y. Grossman, R. Harnik, D. J. Robinson, J. Shu, et al., *Probing CP Violation in $h \rightarrow \gamma\gamma$ with Converted Photons*, *JHEP* **1404** (2014) 084, [arXiv:1312.2955].
- [108] Y. Chen, E. Di Marco, J. Lykken, M. Spiropulu, R. Vega-Morales, et al., *8D likelihood effective Higgs couplings extraction framework in $h \rightarrow 4\ell$* , *JHEP* **1501** (2015) 125, [arXiv:1401.2077].
- [109] M. J. Dolan, P. Harris, M. Jankowiak, and M. Spannowsky, *Constraining CP-violating Higgs Sectors at the LHC using gluon fusion*, *Phys.Rev.* **D90** (2014), no. 7 073008, [arXiv:1406.3322].
- [110] Y. Chen, A. Falkowski, I. Low, and R. Vega-Morales, *New Observables for CP Violation in Higgs Decays*, *Phys.Rev.* **D90** (2014), no. 11 113006, [arXiv:1405.6723].
- [111] Y. Chen, R. Harnik, and R. Vega-Morales, *Probing the Higgs Couplings to Photons in $h \rightarrow 4\ell$ at the LHC*, *Phys.Rev.Lett.* **113** (2014), no. 19 191801, [arXiv:1404.1336].
- [112] M. Beneke, D. Boito, and Y.-M. Wang, *Anomalous Higgs couplings in angular asymmetries of $H \rightarrow Z\ell^+\ell^-$ and $e^+e^- \rightarrow HZ$* , *JHEP* **1411** (2014) 028, [arXiv:1406.1361].
- [113] F. Demartin, F. Maltoni, K. Mawatari, B. Page, and M. Zaro, *Higgs characterisation at NLO in QCD: CP properties of the top-quark Yukawa interaction*, *Eur.Phys.J.* **C74** (2014), no. 9 3065, [arXiv:1407.5089].
- [114] S. Berge, W. Bernreuther, and S. Kirchner, *Determination of the Higgs CP mixing angle in the tau decay channels at the LHC including the Drell-Yan background*, *Eur.Phys.J.* **C74** (2014), no. 11 3164, [arXiv:1408.0798].
- [115] D. Stolarski and R. Vega-Morales, *Directly Measuring the Tensor Structure of the Scalar Coupling to Gauge Bosons*, *Phys.Rev.* **D86** (2012) 117504, [arXiv:1208.4840].
- [116] Y. Chen, N. Tran, and R. Vega-Morales, *Scrutinizing the Higgs Signal and Background in the $2e2\mu$ Golden Channel*, *JHEP* **1301** (2013) 182, [arXiv:1211.1959].
- [117] Y. Chen and R. Vega-Morales, *Extracting Effective Higgs Couplings in the Golden Channel*, *JHEP* **1404** (2014) 057, [arXiv:1310.2893].
- [118] M. Gonzalez-Alonso, A. Greljo, G. Isidori, and D. Marzocca, *Pseudo-observables in Higgs decays*, *Eur.Phys.J.* **C75** (2015), no. 3 128, [arXiv:1412.6038].
- [119] M. Gonzalez-Alonso, A. Greljo, G. Isidori, and D. Marzocca, *Electroweak bounds on Higgs pseudo-observables and $h \rightarrow 4\ell$ decays*, arXiv:1504.04018.
- [120] J. Ellis, V. Sanz, and T. You, *Associated Production Evidence against Higgs Impostors and Anomalous Couplings*, *Eur.Phys.J.* **C73** (2013) 2507, [arXiv:1303.0208].

- [121] R. Godbole, D. J. Miller, K. Mohan, and C. D. White, *Boosting Higgs CP properties via VH Production at the Large Hadron Collider*, *Phys.Lett.* **B730** (2014) 275–279, [arXiv:1306.2573].
- [122] C. Englert, M. McCullough, and M. Spannowsky, *Gluon-initiated associated production boosts Higgs physics*, *Phys.Rev.* **D89** (2014), no. 1 013013, [arXiv:1310.4828].
- [123] G. Isidori and M. Trott, *Higgs form factors in Associated Production*, *JHEP* **1402** (2014) 082, [arXiv:1307.4051].
- [124] R. M. Godbole, D. J. Miller, K. A. Mohan, and C. D. White, *Jet substructure and probes of CP violation in Vh production*, arXiv:1409.5449.
- [125] J. Ellis, V. Sanz, and T. You, *Complete Higgs Sector Constraints on Dimension-6 Operators*, *JHEP* **1407** (2014) 036, [arXiv:1404.3667].
- [126] A. Djouadi, R. Godbole, B. Mellado, and K. Mohan, *Probing the spin-parity of the Higgs boson via jet kinematics in vector boson fusion*, *Phys.Lett.* **B723** (2013) 307–313, [arXiv:1301.4965].
- [127] F. Maltoni, K. Mawatari, and M. Zaro, *Higgs characterisation via vector-boson fusion and associated production: NLO and parton-shower effects*, *Eur.Phys.J.* **C74** (2014), no. 1 2710, [arXiv:1311.1829].
- [128] R. Edezhath, *Dimension-6 Operator Constraints from Boosted VBF Higgs*, arXiv:1501.00992.
- [129] **CMS** Collaboration, V. Khachatryan et al., *Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV*, arXiv:1411.3441.
- [130] A. Azatov and A. Paul, *Probing Higgs couplings with high p_T Higgs production*, *JHEP* **1401** (2014) 014, [arXiv:1309.5273].
- [131] C. Grojean, E. Salvioni, M. Schlaffer, and A. Weiler, *Very boosted Higgs in gluon fusion*, *JHEP* **1405** (2014) 022, [arXiv:1312.3317].
- [132] M. Buschmann, C. Englert, D. Goncalves, T. Plehn, and M. Spannowsky, *Resolving the Higgs-Gluon Coupling with Jets*, *Phys.Rev.* **D90** (2014), no. 1 013010, [arXiv:1405.7651].
- [133] S. Dawson, I. Lewis, and M. Zeng, *Effective field theory for Higgs boson plus jet production*, *Phys.Rev.* **D90** (2014), no. 9 093007, [arXiv:1409.6299].
- [134] S. Dawson, I. Lewis, and M. Zeng, *The Usefulness of EFT for Boosted Higgs Production*, arXiv:1501.04103.
- [135] C. Arnesen, I. Z. Rothstein, and J. Zupan, *Smoking Guns for On-Shell New Physics at the LHC*, *Phys.Rev.Lett.* **103** (2009) 151801, [arXiv:0809.1429].

- [136] T. Corbett, O. Eboli, J. Gonzalez-Fraile, and M. Gonzalez-Garcia, *Determining Triple Gauge Boson Couplings from Higgs Data*, *Phys.Rev.Lett.* **111** (2013) 011801, [arXiv:1304.1151].
- [137] E. Masso, *An Effective Guide to Beyond the Standard Model Physics*, *JHEP* **1410** (2014) 128, [arXiv:1406.6376].
- [138] A. Falkowski and F. Riva, *Model-independent precision constraints on dimension-6 operators*, *JHEP* **1502** (2015) 039, [arXiv:1411.0669].
- [139] C. Bobeth and U. Haisch, *Anomalous triple gauge couplings from B-meson and kaon observables*, arXiv:1503.04829.
- [140] K. Hagiwara, R. Peccei, D. Zeppenfeld, and K. Hikasa, *Probing the Weak Boson Sector in $e^+ e^- \rightarrow W^+ W^-$* , *Nucl.Phys.* **B282** (1987) 253.