

Towards stability and optimality in stochastic gradient descent

Panos Toulis
Harvard University

Dustin Tran
Harvard University

Edoardo M. Airolidi
Harvard University

Abstract

Iterative procedures for parameter estimation based on stochastic gradient descent (SGD) allow the estimation to scale to massive data sets. However, in both theory and practice, they suffer from numerical instability. Moreover, they are statistically inefficient as estimators of the true parameter value. To address these two issues, we propose a new iterative procedure termed *averaged implicit* SGD (AI-SGD). For statistical efficiency, AI-SGD employs averaging of the iterates, which achieves the optimal Cramér-Rao bound under strong convexity, i.e., it is an optimal unbiased estimator of the true parameter value. For numerical stability, AI-SGD employs an implicit update at each iteration, which is related to proximal operators in optimization. In practice, AI-SGD achieves competitive performance with other state-of-the-art procedures. Furthermore, it is more stable than averaging procedures that do not employ proximal updates, and is simple to implement as it requires fewer tunable hyperparameters than procedures that do employ proximal updates.

1 Introduction

The majority of problems in statistical estimation can be cast as finding the parameter value $\theta_\star \in \Theta$ such that

$$\theta_\star = \arg \min_{\theta \in \Theta} \mathbb{E} (L(\theta, \xi)), \quad (1)$$

where the expectation is with respect to the random variable $\xi \in \Xi \subseteq \mathbb{R}^d$ that represents the data, $\Theta \subseteq \mathbb{R}^p$ is the parameter space, and $L : \Theta \times \Xi \rightarrow \mathbb{R}$ is a loss function. A popular procedure for solving Eq.(12) is stochastic gradient descent (SGD) (Zhang, 2004; Bottou, 2004), where a sequence θ_n approximates θ_\star , and is updated iteratively, one data point at a time, through the iteration

$$\theta_n = \theta_{n-1} - \gamma_n \nabla L(\theta_{n-1}, \xi_n), \quad (2)$$

where $\{\xi_1, \xi_2, \dots\}$ is a stream of i.i.d. realizations of ξ , and $\{\gamma_n\}$ is a non-increasing sequence of positive real numbers, known as the learning rate. The n th iterate θ_n in SGD (2) can be viewed as an *estimator* of θ_\star . To evaluate such iterative estimators it is typical to consider three properties: convergence rate and numerical stability, by studying the mean-squared errors $\mathbb{E} (\|\theta_n - \theta_\star\|^2)$; and statistical efficiency, by studying the limit $n \text{Var} (\theta_n)$, as $n \rightarrow \infty$.

While computationally efficient, the SGD procedure (2) suffers from numerical instability and statistical inefficiency. Regarding stability, SGD is sensitive to specification of the learning rate γ_n , since the mean-squared errors can diverge arbitrarily when γ_n is misspecified with the respect to problem parameters, e.g., the convexity and Lipschitz parameters of the loss function (Benveniste et al., 1990; Moulines and Bach, 2011). Regarding statistical efficiency, SGD loses statistical information. In fact, the amount of information loss depends on the misspecification of γ_n with respect to the spectral gap of the matrix $\mathbb{E} (\nabla^2 L(\theta_\star, \xi))$ (Toulis et al., 2014), also known as the Fisher information matrix. Several solutions have been proposed to resolve these two issues, e.g., using projections and gradient clipping. However, they are usually heuristic and hard to generalize.

In this paper, we aim for the ideal combination of computational efficiency, numerical stability, and statistical efficiency using the following procedure:

$$\theta_n = \theta_{n-1} - \gamma_n \nabla L(\theta_n, \xi_n), \quad (3)$$

AI-SGD

$$\bar{\theta}_n = (1/n) \sum_{i=1}^n \theta_i. \quad (4)$$

Our proposed procedure, termed *averaged implicit* SGD (AI-SGD), is comprised of two inner procedures. The first procedure employs updates given in Eq.(3), which are *implicit* because the iterate θ_n appears on both sides of the equation. Procedure (3), also known as *implicit SGD* (Toulis et al., 2014), aims to stabilize the updates of the classic SGD procedure (2). In fact, implicit SGD can be motivated as the limit of a sequence of improved classic SGD procedures. To see this, first fix the sample history $\mathcal{F}_{n-1} = \{\theta_0^s, \xi_1, \xi_2, \dots, \xi_{n-1}\}$, where we use the superscript “s” in the classic SGD procedure in order to distinguish from implicit SGD. Then, $\theta_n^s = \theta_{n-1}^s - \gamma_n \nabla L(\theta_{n-1}^s, \xi_n) \triangleq \theta_n^{(1)}$.

If we “trust” $\theta_n^{(1)}$ to be a better estimate of θ_* than θ_{n-1}^s , then we can use $\theta_n^{(1)}$ instead of θ_{n-1}^s in computing the loss function at data point ξ_n . This leads to a revised update $\theta_n^s = \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^{(1)}, \xi_n) \triangleq \theta_n^{(2)}$. Likewise, we can use $\theta_n^{(2)}$ instead of $\theta_n^{(1)}$, and so on. If we repeat this argument ad infinitum, then we get the following sequence of improved SGD procedures,

$$\begin{aligned}\theta_n^s &= \theta_{n-1}^s - \gamma_n \nabla L(\theta_{n-1}, \xi_n), \\ \theta_n^s &= \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^{(1)}, \xi_n), \\ \theta_n^s &= \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^{(2)}, \xi_n), \\ &\dots \\ \theta_n^s &= \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^{(\infty)}, \xi_n),\end{aligned}\quad (5)$$

where $\theta_n^{(i)} = \theta_{n-1}^s - \nabla L(\theta_n^{(i-1)}, \xi_n)$, with initial condition $\theta_n^{(0)} = \theta_{n-1}^s$. In the limit, assuming a unique fixed point is reached almost surely, the final procedure of sequence (5) satisfies $\theta_n^s = \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^{(\infty)}, \xi_n) = \theta_n^{(\infty)}$. This can be rewritten as $\theta_n^s = \theta_{n-1}^s - \gamma_n \nabla L(\theta_n^s, \xi_n)$, which is equivalent to implicit SGD. Thus, implicit SGD can be viewed as a repeated application of classic SGD, where we keep updating the same iterate θ_{n-1}^s using the same data point ξ_n , until a fixed-point is reached. Nesterov’s accelerated gradient, a popular improvement of classic SGD, is only one application of this procedure.

The stability improvement achieved by implicit updates can be motivated by the following argument. Assume for simplicity that L is strongly convex, almost surely, with parameter $\mu > 0$. Then for the implicit SGD procedure (3),

$$\begin{aligned}\theta_n + \gamma_n \nabla L(\theta_n, \xi_n) &= \theta_{n-1}, \\ \|\theta_n - \theta_*\|^2 + 2\gamma_n(\theta_n - \theta_*)^\top \nabla L(\theta_n, \xi_n) &\leq \|\theta_{n-1} - \theta_*\|^2, \\ (1 + \gamma_n \mu) \|\theta_n - \theta_*\|^2 &\leq \|\theta_{n-1} - \theta_*\|^2, \\ \|\theta_n - \theta_*\|^2 &\leq \frac{1}{1 + \gamma_n \mu} \|\theta_{n-1} - \theta_*\|^2,\end{aligned}$$

which implies that $\|\theta_n - \theta_*\|^2$ is contracting almost surely. In contrast, the classic SGD procedure does not share this contracting property.

While the implicit updates of Eq.(3) aim to achieve stability, the averaging of the iterates in Eq.(4) aims to achieve statistical optimality. Ruppert (1988) gave a nice intuition on why iterate averaging can lead to statistical optimality. When the learning rate is $\gamma_n \propto n^{-1}$, then $\bar{\theta}_n - \theta_*$ is a weighted average of n error variables $\nabla L(\theta_{i-1}, \xi_i)$, which therefore are significantly autocorrelated. However, when $\gamma_n \propto n^{-\gamma}$ with $\gamma \in (0, 1)$, then $\bar{\theta}_n - \theta_*$ is the average of $n^\gamma \log n$ error variables, which become uncorrelated in the limit. Thus, averaging improves the estimation accuracy.

1.1 Related work

The implicit update (3) is equivalent to

$$\theta_n = \arg \min_{\theta \in \Theta} \left\{ \frac{1}{2\gamma_n} \|\theta - \theta_{n-1}\|^2 + L(\theta, \xi_n) \right\}. \quad (6)$$

Arguably, the first method that used an update similar to (6) for estimation was the normalized least-mean squares filter of Nagumo and Noda (1967), used in signal processing. This update is also used by the *incremental proximal method* in optimization (Bertsekas, 2011), and has shown superior performance to classic SGD both in theory and applications (Bertsekas, 2011; Toulis et al., 2014; Défossez and Bach, 2015; Toulis and Airoldi, 2015). In particular, implicit updates lead to similar convergence rates as classic SGD updates, but are significantly more stable. This stability can also be motivated from a Bayesian interpretation of Eq.(6), where θ_n is the posterior mode of a model with the standard multivariate normal $\mathcal{N}(\theta_{n-1}, \gamma_n I)$ as the prior, $L(\theta, \cdot)$ as the log-likelihood, and ξ_n as the observation.

A statistical analysis of procedure (3) without averaging was done by Toulis et al. (2014) who derived the asymptotic variance $\text{Var}(\theta_n)$ of θ_n , and provided an algorithm to efficiently solve the fixed-point equation (3) for θ_n in the family of generalized linear models, which we generalize in this current work. In the online learning literature, Kivinen et al. (2006) and Kulis and Bartlett (2010) have also analyzed implicit updates; Schuurmans and Caelli (2007) have further applied implicit procedures on learning with kernels. Notably the implicit update (6) is related to the importance weight updates proposed by Karampatziakis and Langford (2010), but the two update forms are not equivalent, and are usually combined in practice (Karampatziakis and Langford, 2010, Section 5).

Assuming that the expected loss ℓ is known, instead of update (6) we could use the update

$$\theta_n^+ = \arg \min_{\theta \in \Theta} \left\{ \frac{1}{2\gamma_n} \|\theta - \theta_{n-1}\|^2 + \ell(\theta) \right\}. \quad (7)$$

In optimization, this mapping from θ_{n-1} to θ_n^+ in Eq. (7) is known as a *proximal operator*, and is a special instance of the proximal point algorithm (Rockafellar, 1976). Thus implicit SGD involves mappings that are stochastic versions of mappings from proximal operators. The stochastic proximal gradient algorithm (Singer and Duchi, 2009; Parikh and Boyd, 2013; Rosasco et al., 2014) is related but different to implicit SGD. In contrast to implicit SGD, the stochastic proximal gradient algorithm first makes a classic SGD update (forward step), and then an implicit update (backward step). Only the forward step is stochastic whereas the backward proximal step is not. This may increase convergence speed but may also introduce instability due to the forward step.

Interest on proximal operators has surged in recent years because they are non-expansive and converge with minimal assumptions. Furthermore, they can be applied on non-smooth objectives, and can easily be combined in modular algorithms for optimization in large-scale and distributed settings (Parikh and Boyd, 2013). The idea has also been generalized through splitting algorithms (Lions and Mercier, 1979; Beck and Teboulle, 2009; Singer and Duchi, 2009; Duchi et al., 2011). Krakowski et al. (2007) and Nemirovski et al. (2009) have shown that proximal methods can fit better in the geometry of the parameter space Θ , and Toulis and Airolidi (2014) have made a connection to shrinkage methods in statistics.

Two recent procedures based on stochastic proximal updates are PROX-SVRG (Xiao and Zhang, 2014) and PROX-SAG (Schmidt et al., 2013, Section 6). The main idea in both methods is to periodically compute an estimate of the full gradient averaged over all data points in order to reduce the variance of stochastic gradients. This requires a finite data setting, whereas AI-SGD also applies to streaming data. Moreover, the periodic calculations in PROX-SVRG are controlled by additional hyperparameters, and the periodic calculations in PROX-SAG require storage of the full gradient at every iteration. AI-SGD differs because it employs averaging to achieve statistical efficiency, has no additional hyperparameters or major storage requirements, and thus it has a simpler implementation.

Averaging of the iterates in Eq.(4) is the other key component of AI-SGD. Averaging was proposed and analyzed in the stochastic approximation literature by Ruppert (1988) and Bather (1989). Polyak and Juditsky (1992) substantially expanded the scope of the averaging method by proving asymptotic optimality of the classic SGD procedure with averaging, under suitable assumptions. Their results showed clearly that slowly-convergent stochastic approximations (achieved when the learning rates are large) need to be averaged. Recent work has analyzed classic SGD with averaging (Zhang, 2004; Xu, 2011; Shamir and Zhang, 2012; Bach and Moulines, 2013) and has shown their superiority in numerous learning tasks.

1.2 Overview of results

In this paper, we study the iterates θ_n and use the results to study $\bar{\theta}_n$ as an estimator of θ_* . Under strong convexity of the expected loss, we derive upper bounds for the squared errors $\mathbb{E}(\|\theta_n - \theta_*\|^2)$ and $\mathbb{E}(\|\bar{\theta}_n - \theta_*\|^2)$ in Theorem 3 and Theorem 2, respectively. In the supplementary material, we also give bounds for $\mathbb{E}(\|\theta_n - \theta_*\|^4)$.

Two main results are derived from our theoretical analysis. First, $\bar{\theta}_n$ achieves the Cramér-Rao bound, i.e., no other unbiased estimator of θ_* can do better in the limit, which is equivalent to the optimal $\mathcal{O}(1/n)$ rate of convergence for first-order procedures. Second, AI-SGD is significantly

more stable to misspecification of the learning rate relative to classic averaged SGD procedures, with respect to the learning problem parameters, e.g., convexity and Lipschitz constants. Finally, we perform experiments on several standard machine learning tasks, which show that AI-SGD comes closer to combining stability, optimality, and simplicity than other competing methods.

2 Preliminaries

Notation. Let $\mathcal{F}_n = \{\theta_0, \xi_1, \xi_2, \dots, \xi_n\}$ denote the filtration that process θ_n (3) is adapted to. The norm $\|\cdot\|$ will denote the L_2 norm. The symbol \triangleq indicates a definition, and the symbol $\stackrel{\text{def}}{=}$ denotes “equal by definition”. For example, $x \triangleq y$ defines x as equal to known variable y , whereas $x \stackrel{\text{def}}{=} y$ denotes that x is equal to y by definition. We will not use this formalism when defining constants. For two positive sequences a_n, b_n , we write $b_n = \mathcal{O}(a_n)$ if there exists a fixed $c > 0$ such that $b_n \leq ca_n$, for all n ; also, $b_n = o(a_n)$ if $b_n/a_n \rightarrow 0$. When a positive scalar sequence a_n is monotonically decreasing to zero, we write $a_n \downarrow 0$. Similarly, for a sequence X_n of vectors or matrices, $X_n = \mathcal{O}(a_n)$ denotes that $\|X_n\| = \mathcal{O}(a_n)$, and $X_n = o(a_n)$ denotes that $\|X_n\| = o(a_n)$. For two matrices A, B , $A \preceq B$ denotes that $B - A$ is nonnegative-definite; $\text{tr}(A)$ denotes the trace of A .

We now introduce the main assumptions pertaining to the theory of this paper.

Assumption 1. *The loss function $L(\theta, \xi)$ is almost-surely differentiable. The random vector ξ can be decomposed as $\xi = (x, y)$, $x \in \mathbb{R}^p$, $y \in \mathbb{R}^d$, such that*

$$L(\theta, \xi) = L(x^\top \theta, y). \quad (8)$$

Assumption 2. *The learning rate sequence $\{\gamma_n\}$ is defined as $\gamma_n = \gamma_1 n^{-\gamma}$, where $\gamma_1 > 0$ and $\gamma \in (1/2, 1]$.*

Assumption 3 (Lipschitz conditions). *For all $\theta_1, \theta_2 \in \Theta$, a combination of the following conditions is satisfied almost-surely:*

(a) *The loss function L is Lipschitz-continuous with parameter λ_0 , i.e.,*

$$|L(\theta_1, \xi) - L(\theta_2, \xi)| \leq \lambda_0 \|\theta_1 - \theta_2\|,$$

(b) *The map ∇L is Lipschitz-continuous with parameter λ_1 , i.e.,*

$$\|\nabla L(\theta_1, \xi) - \nabla L(\theta_2, \xi)\| \leq \lambda_1 \|\theta_1 - \theta_2\|,$$

(c) *The map $\nabla^2 L$ is Lipschitz-continuous with parameter λ_2 , i.e.,*

$$\|\nabla^2 L(\theta_1, \xi) - \nabla^2 L(\theta_2, \xi)\| \leq \lambda_2 \|\theta_1 - \theta_2\|.$$

Assumption 4. *The observed Fisher information matrix, $\hat{\mathcal{I}}(\theta) \triangleq \nabla^2 L(\theta, \xi)$, has non-vanishing trace, i.e., there exists $\phi > 0$ such that $\text{tr}(\hat{\mathcal{I}}(\theta)) \geq \phi$, almost-surely, for all $\theta \in \Theta$. The expected Fisher information matrix, $\mathcal{I}(\theta) \triangleq \mathbb{E}(\hat{\mathcal{I}}(\theta))$, has minimum eigenvalue $0 < \underline{\lambda}_f \leq \phi$, for all $\theta \in \Theta$.*

Assumption 5. *The zero-mean random variable $W_\theta \triangleq \nabla L(\theta, \xi) - \nabla \ell(\theta)$ is square-integrable, such that, for a fixed positive-definite Σ ,*

$$\mathbb{E}(W_{\theta_*} W_{\theta_*}^\top) \preceq \Sigma.$$

Remarks. Assumption 6 puts a constraint on the loss function, but it is not very restrictive because the majority of machine learning models indeed depend on the parameter θ through a linear combination with features. A notable exception includes loss functions with a regularization term. Although it is easy to add regularization to AI-SGD we will not do so in this paper because AI-SGD works well without it, since the proximal operator (6) already regularizes the estimate θ_n towards θ_{n-1} . In experiments, regularization neither improved nor worsened AI-SGD (see supplementary material for more details). Assumption 7 on learning rates and Assumption 10 are standard in the literature of stochastic approximations, dating back to the original paper of Robbins and Monro (1951) in the one-dimensional parameter case.

Assumptions on Lipschitz gradients (Assumption 8(b), Assumption 8(c)) can be relaxed; for example, Benveniste et al. (1990) relax this assumption using $\|\theta_1 - \theta_2\|^q$. However, these two Lipschitz conditions are commonly used in order to simplify the non-asymptotic analysis (Moulines and Bach, 2011). Assumption 8(a) is less standard in classic SGD literature but has so far been standard in the limited literature on implicit SGD (Bertsekas, 2011). We can forgo this assumption and still maintain identical rates for the errors, although at the expense of a more complicated analysis. It is also an open problem whether a nice stability result similar to Theorem 3 can be derived under Assumption 8(b) instead of Assumption 8(a). We discuss this issue after the proof of Theorem 3 in the supplementary material.

Assumption 9 makes two claims. The first claim on the observed Fisher information matrix is a relaxed form of strong convexity for the loss $L(\theta, \xi)$. However, in contrast to strong convexity, this claim allows several eigenvalues of $\nabla^2 L$ to be zero. The second claim of Assumption 9 is equivalent to strong convexity of the expected loss $\ell(\theta)$. From a statistical perspective, strong convexity posits that there is information in the data for all elements of θ_* . This assumption is necessary to derive bounds on the errors $\mathbb{E}(\|\theta_n - \theta_*\|^2)$, and has been used to show optimality of classic SGD with averaging (Polyak and Juditsky, 1992; Ljung et al., 1992; Xu, 2011; Moulines and Bach, 2011).

Overall, our assumptions are weaker than the assumptions in the limited literature on implicit SGD. For example,

Bertsekas (2011, Assumptions 3.1, 3.2) assumes almost-sure bounded gradients $\nabla L(\theta, \xi)$ in addition to Assumption 8(a); Ryu and Boyd (2014) assume strong convexity of $L(\theta, \xi)$, in expectation, which can simplify the analysis significantly. We discuss more details in the supplementary material after the proof of Theorem 3.

3 Theory

In this section we present our theoretical analysis of AI-SGD. All proofs are given in the supplementary material. The main technical challenge in analyzing implicit SGD (3) is that unlike typical analysis with classic SGD (2), the error ξ_n is not conditionally independent of θ_n . This implies that $\mathbb{E}(\nabla L(\theta_n, \xi_n) | \theta_n) \neq \nabla \ell(\theta_n)$, which makes it no longer possible to use the convexity properties of ℓ to analyze the errors $\mathbb{E}(\|\theta_n - \theta_*\|^2)$, as it is common in the literature.

As mentioned earlier, to circumvent this issue other authors have made strict almost-sure assumptions on the implicit procedure (3) (Bertsekas, 2011; Ryu and Boyd, 2014). In this paper, we rely on weaker conditions, namely the Lipschitz assumptions 8(a)-8(c), which are also used in non-implicit procedures. Our proof strategy relies on a master lemma (Lemma 3 in supplementary material) for the analysis of recursions that appear to be typical in implicit procedures. This result is novel to our best knowledge, and it can be useful in future research on implicit procedures.

3.1 Computational efficiency

Our first result enables efficient computation of the implicit update (3). In general, this can be expensive due to solving a fixed-point equation in many dimensions, at every iteration. We reduce this multi-dimensional equation to an equation of only one dimension. Furthermore, under almost-sure convexity of the loss function, efficient search bounds for the one-dimensional fixed-point equation are available. This result generalizes an earlier result in efficient computation of implicit updates on generalized linear models (Toullis et al., 2014, Algorithm 1).

Definition 1. *Suppose that Assumption 6 holds. For observation $\xi = (x, y)$, the first derivative with respect to the natural parameter $x^\top \theta$ is denoted by $L'(\theta, \xi)$, and is defined as*

$$L'(\theta, \xi) \triangleq \frac{\partial L(\theta, \xi)}{\partial (x^\top \theta)} \stackrel{\text{def}}{=} \frac{\partial L(x^\top \theta, y)}{\partial (x^\top \theta)}. \quad (9)$$

Similarly, $L''(\xi, \theta) \triangleq \frac{\partial L'(\theta, \xi)}{\partial (x^\top \theta)}$.

Lemma 1. *Suppose that Assumption 6 holds, and consider functions L', L'' from Definition 2. Then, almost-surely,*

$$\nabla L(\theta_n, \xi_n) = s_n \nabla L(\theta_{n-1}, \xi_n); \quad (10)$$

the scalar s_n satisfies the fixed-point equation,

$$s_n \kappa_{n-1} = L'(\theta_{n-1} - s_n \gamma_n \kappa_{n-1} x_n, \xi_n), \quad (11)$$

where $\kappa_{n-1} \triangleq L'(\theta_{n-1}, \xi_n)$. Moreover, if $L''(\theta, \xi) \geq 0$ almost-surely for all $\theta \in \Theta$, then

$$s_n \in \begin{cases} [\kappa_{n-1}, 0] & \text{if } \kappa_{n-1} < 0, \\ [0, \kappa_{n-1}] & \text{otherwise.} \end{cases}$$

Remarks. Lemma 2 has two parts. First, it shows that the implicit update can be performed by obtaining s_n from the fixed-point Eq.(18), and then using $\nabla L(\theta_n, \xi_n) = s_n \nabla L(\theta_{n-1}, \xi_n)$ in the implicit update (3). The fixed-point equation can be solved through a numerical root-finding procedure (Kivinen et al., 2006; Kulis and Bartlett, 2010; Toulis et al., 2014). Second, when the loss function is convex, then narrow search bounds for s_n are available. This property holds, for example, when the loss function is the negative log-likelihood in an exponential family.

3.2 Non-asymptotic analysis

Our next result is on the mean-squared errors $\mathbb{E}(\|\theta_n - \theta_\star\|^2)$. These errors show the stability and convergence rates of implicit SGD and are used in combination with bounds on errors $\mathbb{E}(\|\theta_n - \theta_\star\|^4)$ to derive bounds on the errors $\mathbb{E}(\|\bar{\theta}_n - \theta_\star\|^2)$ of the averaged procedure.¹

Theorem 1. Suppose that Assumptions 6, 7, 8(a), and 9 hold. Define $\delta_n \triangleq \mathbb{E}(\|\theta_n - \theta_\star\|^2)$, and constants $\Gamma^2 = 4\lambda_0^2 \sum \gamma_i^2 < \infty$, $\epsilon = (1 + \gamma_1(\phi - \lambda_f))^{-1}$, and $\lambda = 1 + \gamma_1 \lambda_f \epsilon$. Then, there exists constant $n_0 > 0$ such that, for all $n > 0$,

$$\delta_n \leq (8\lambda_0^2 \gamma_1 \lambda / \lambda_f \epsilon) n^{-\gamma} + e^{-\log \lambda \cdot n^{1-\gamma}} [\delta_0 + \lambda^{n_0} \Gamma^2].$$

Remarks. According to Theorem 3, the convergence rate of the implicit iterates θ_n is $\mathcal{O}(n^{-\gamma})$. This matches earlier results on rates of classic SGD (Benveniste et al., 1990; Moulines and Bach, 2011). The most important difference, however, is that the implicit procedure discounts the initial conditions δ_0 at an exponential rate, regardless of the specification of the learning rate. As shown by Moulines and Bach (2011, Theorem 1), in classic SGD there exists a term $\exp(\lambda_1^2 \gamma_1^2 n^{1-2\gamma})$ in front of the initial conditions, which can be catastrophic if the learning rate parameter γ_1 is misspecified. In contrast, the implicit iterates are unconditionally stable, i.e., any specification of the learning rate will lead to a stable discounting of the initial conditions.

¹ The bounds for the fourth moments $\mathbb{E}(\|\theta_n - \theta_\star\|^4)$ are given in the supplementary material because they rely on the same intermediate results as $\mathbb{E}(\|\theta_n - \theta_\star\|^2)$.

Theorem 2. Consider the AI-SGD procedure (4), and suppose that Assumptions 7, 8(a), 8(c), 9, and 10 hold. Then,

$$\begin{aligned} (\mathbb{E}(\|\bar{\theta}_n - \theta_\star\|^2))^{1/2} &\leq \frac{1}{\sqrt{n}} (\text{tr}(\nabla^2 \ell(\theta_\star)^{-1} \Sigma \nabla^2 \ell(\theta_\star)^{-1}))^{1/2} \\ &\quad + \mathcal{O}(n^{-1+\gamma/2}) + \mathcal{O}(n^{-\gamma}) \\ &\quad + \mathcal{O}(\exp(-\log \lambda \cdot n^{1-\gamma}/2)). \end{aligned}$$

Remarks. The full version of Theorem 2, which includes all constants, is given in the supplementary material. Even in its shortened form, Theorem 2 delivers three main results. First, the iterates $\bar{\theta}_n$ attain the Cramér-Rao lower bound, i.e., any other unbiased estimator of θ_\star cannot have lower MSE than $\bar{\theta}_n$. From an optimization perspective, $\bar{\theta}_n$ attains the rate $\mathcal{O}(1/n)$, which is optimal for first-order methods (Nesterov, 2004). This result matches the asymptotic optimality of averaged iterates from classic SGD procedures, which has been proven by Polyak and Juditsky (1992).

Second, the remaining rates are $\mathcal{O}(n^{-2+\gamma})$ and $\mathcal{O}(n^{-2\gamma})$. This implies the optimal choice $\gamma = 2/3$ for the exponent of the learning rate. It extends the results of Ruppert (1988), and more recently by Xu (2011), and Moulines and Bach (2011), on optimal exponents for classic SGD procedures.

Third, as with non-averaged implicit iterates in Theorem 3, the averaged iterates $\bar{\theta}_n$ have a decay of the initial conditions regardless of the specification of the learning rate parameter. This stability property is inherited from the underlying implicit SGD procedure (3) that is being averaged. In contrast, averaged iterates of classic SGD procedures can diverge numerically because arbitrarily large terms can appear in front of initial conditions (Moulines and Bach, 2011, Theorem 3).

4 Experiments

In this section, we show that AI-SGD achieves comparable, and sometimes superior, results to other methods while combining statistical efficiency, stability, and simplicity. In our experiments, we compare our procedure to the following procedures:

- **SGD:** Classic stochastic gradient descent in its standard formulation (Sakrison, 1965; Zhang, 2004), which employs the update $\theta_n = \theta_{n-1} - \gamma_n \nabla L(\theta_{n-1}, \xi_n)$.
- **IMPLICIT SGD:** Stochastic gradient descent procedure introduced in Toulis et al. (2014) which employs implicit update (3) without averaging. It is robust to misspecification of the learning rate but also exhibits slower convergence in practice relative to classic SGD.
- **ASGD:** Averaged stochastic gradient descent procedure with classic updates of the iterates (Xu, 2011; Shamir

and Zhang, 2012; Bach and Moulines, 2013). This is equivalent to AI-SGD where the update (3) is replaced by the classic step $\theta_n = \theta_{n-1} - \gamma_n \nabla L(\theta_{n-1}, \xi_n)$.

- PROX-SVRG: A proximal version of the stochastic gradient descent procedure with progressive variance reduction (SVRG) (Xiao and Zhang, 2014).
- PROX-SAG: A proximal version of the stochastic average gradient (SAG) procedure (Schmidt et al., 2013). While its theory has not been formally established, PROX-SAG has shown similar convergence properties to PROX-SVRG in practice.
- ADAGRAD: A stochastic gradient descent procedure with a form of diagonal scaling to adapt the learning rate (Duchi et al., 2011).

Note that PROX-SVRG and PROX-SAG are applicable only to fixed data sets and not to the streaming setting. Therefore the theoretical linear convergence rate of these methods refers to convergence to an empirical minimizer (e.g., maximum likelihood, or maximum a-posteriori if there is regularization), and not to the ground truth θ_* . On the other hand, AI-SGD can be applied to both data settings.

We also note that ADAGRAD, and similar adaptive schedules, (Tieleman and Hinton, 2012; Kingma and Ba, 2015) effectively approximate the natural gradient $\mathcal{I}(\theta)^{-1} \nabla L(\theta, \xi)$ by using a multi-dimensional learning rate. These learning rates have the added advantage of being less sensitive than one-dimensional rates to tuning of hyperparameters; they can be combined in practice with AI-SGD.

4.1 Statistical efficiency and stability

We first demonstrate the theoretical results on the stability and statistical optimality of AI-SGD. To do so, we follow a simple normal linear regression example from Bach and Moulines (2013). Let $N = 10^6$ be the number of observations, and $p = 20$ be the number of features. Let $\theta_* = (0, 0, \dots, 0)^\top$ be the ground truth. The random variable ξ is decomposed as $\xi_n = (x_n, y_n)$, where the feature vectors $x_1, \dots, x_N \sim \mathcal{N}_p(0, H)$ are i.i.d. normal random variables, and H is a randomly generated symmetric matrix with eigenvalues $1/k$, for $k = 1, \dots, p$. The outcome y_n is sampled from a normal distribution as $y_n | x_n \sim \mathcal{N}(x_n^\top \theta_*, 1)$, for $n = 1, \dots, N$. Our loss function is defined as the squared residual, i.e., $L(\theta, \xi_n) = (y_n - x_n^\top \theta)^2$, and thus $\ell(\theta) = \mathbb{E}(L(\theta, \xi)) = (\theta - \theta_*)^\top H (\theta - \theta_*)$.

We choose a constant learning rate $\gamma_n \equiv \gamma$ according to the average radius of the data $R^2 = \text{trace}(H)$, and for both ASGD and AI-SGD we collect iterates θ_n , $n = 1, \dots, N$, and keep the average $\bar{\theta}_n$. In Figure 1, we plot $\ell(\bar{\theta}_n)$ for each iteration for a maximum of N iterations in log-log space.




Figure 1: Loss of AI-SGD, ASGD, and IMPLICIT SGD, on simulated multivariate normal data with $N = 10^6$ observations, $d = 20$ features. The plot shows that AI-SGD achieves stability regardless of the specification of the learning rate $\gamma_n \equiv \gamma$. In contrast, ASGD diverges when the learning rate is only slightly misspecified (e.g., solid, blue line).

Figure 1 shows that AI-SGD performs on par with ASGD for the rates at which ASGD is known to be optimal. However, the benefit of the implicit procedure (3) in AI-SGD becomes clear as the learning rate increases. Notably, AI-SGD remains stable for learning rates that are above the theoretical threshold, i.e., when $\gamma > 1/R^2$, whereas ASGD diverges above that threshold, e.g., when $\gamma = 2/R^2$. This stable behavior is also exhibited in IMPLICIT SGD, but IMPLICIT SGD converges at a slower rate than AI-SGD, and thus does not combine stability with statistical efficiency. This behavior is also reflected for AI-SGD when using decaying learning rates, e.g., $\gamma_n \propto 1/n$.

4.2 Classification error

We now conduct a study of AI-SGD’s empirical performance on standard benchmarks of large-scale linear classification. For brevity, we display results on four data sets, although we have seen similar results on eight additional ones (see the supplementary material for more details).

Table 2 displays a summary of the data sets. The COVTYPE data set (Blackard, 1998) consists of forest cover types in which the task is to classify class 2 among 7 forest cover types. DELTA is synthetic data offered in the PASCAL Large Scale Challenge (Sonnenburg et al., 2008) and we apply the default processing offered by the challenge organizers. The task in RCV1 is to classify documents belonging




Figure 2: Large scale linear classification with log loss on four data sets. Each plot indicates the test error of various stochastic gradient methods over a single pass of the data.

to class CCAT in the text dataset (Lewis et al., 2004), where we apply the standard preprocessing provided by Bottou (2012). In the MNIST data set (Le Cun et al., 1998) of images of handwritten digits, the task is to classify digit 9 against all others.

For AI-SGD and ASGD, we use the learning rate $\gamma_n = \eta_0(1 + \eta_0 n)^{-3/4}$ prescribed in Xu (2011), where the constant η_0 is determined through preprocessing on a small subset of the data. Hyperparameters for other methods are set based on a computationally intensive grid search over the entire hyperparameter space: this includes step sizes for PROX-SAG, PROX-SVRG, and ADAGRAD, and the inner iteration count for PROX-SVRG. For all methods we use L_2 regularization with parameter λ which varies for each data set, and which is also used in Xu (2011).

The results are shown in Figure 2. We see that AI-SGD achieves comparable performance with the tuned proximal methods PROX-SVRG and PROX-SAG, as well as ADAGRAD. All methods have a comparable convergence rate and take roughly a single pass in order to converge. Interestingly, ADAGRAD exhibits a larger variance in its estimate than the proximal methods. This comes from the less known fact that the learning rate in ADAGRAD is a suboptimal approximation of the Fisher information, and hence it is statistically inefficient.

4.3 Sensitivity analysis

We examine the inherent stability of the aforementioned procedures by perturbing their hyperparameters. That is, we perform sensitivity analysis by varying any hyperparameters that the user must tweak in order to fine tune the convergence of each procedure. We do so for hyperparameters in ASGD (the learning rate), PROX-SVRG (proximal step size η and inner iteration m), and AI-SGD (the learning rate).




Figure 3: Top: Logistic regression on the RCV1 dataset, performing sensitivity analysis of AI-SGD and ASGD for the choice of regularization parameter λ . Bottom: linear SVM on the covtype dataset, performing sensitivity analysis of AI-SGD and PROX-SVRG, in which PROX-SVRG has additional hyperparameters η according to the step size of the proximal update and m according to the inner iteration count.

The results are shown in Figure 3. When we decrease the regularization parameter, ASGD performs increasingly worse. While it may converge, the test error can be arbitrarily large. On the other hand, AI-SGD always achieves convergence and is not affected by the choice of the hyperparameter. When the regularization parameter is about $1/N$, e.g., when $\lambda < 1E-6$, ASGD remains stable and achieves the same performance as AI-SGD. Similar results hold when perturbing the hyperparameters η and m in PROX-SVRG, as

	description	type	features	training set	test set	λ
covtype	forest cover type	sparse	54	464,809	116,203	10^{-6}
delta	synthetic data	dense	500	450,000	50,000	10^{-2}
rcv1	text data	sparse	47,152	781,265	23,149	10^{-5}
mnist	digit image features	dense	784	60,000	10,000	10^{-3}

Table 1: Summary of data sets and the L_2 regularization parameter, following the settings in Xu (2011).

AI-SGD does not require specification of such hyperparameters.

5 Conclusion

We propose a statistical learning procedure, termed AI-SGD, and investigate its theoretical and empirical properties. AI-SGD combines simple stochastic proximal steps, also known as implicit updates, with iterate averaging and larger step-sizes. The proximal steps allow AI-SGD to be significantly more stable compared to classic SGD procedures, with or without averaging of the iterates; this stability comes at virtually no computational cost for a large family of machine learning models. Furthermore, the averaging of the iterates lead AI-SGD to be statistically optimal, i.e., the variance of the iterate $\bar{\theta}_n$ of AI-SGD achieves the minimum Cramér-Rao lower bound, under strong convexity. Last but not least, AI-SGD is as simple to implement as classic SGD. In comparison, other stochastic proximal procedures, such as PROX-SVRG or PROX-SAG, require tuning of hyperparameters that control periodic calculations over the entire dataset, and possibly storage of the full gradient.

References

- Francis Bach and Eric Moulines. Non-strongly-convex smooth stochastic approximation with convergence rate $O(1/n)$. In *Advances in Neural Information Processing Systems*, pages 773–781, 2013.
- JA Bather. *Stochastic approximation: A generalisation of the Robbins-Monro procedure*, volume 89. Mathematical Sciences Institute, Cornell University, 1989.
- Amir Beck and Marc Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM Journal on Imaging Sciences*, 2(1):183–202, 2009.
- Albert Benveniste, Pierre Priouret, and Michel Métivier. Adaptive algorithms and stochastic approximations. 1990.
- Dimitri P Bertsekas. Incremental proximal methods for large scale convex optimization. *Mathematical programming*, 129(2):163–195, 2011.
- Jock Blackard. *Comparison of neural networks and discriminant analysis in predicting forest cover types*. PhD thesis, Department of Forest Sciences, Colorado State University, 1998.
- Léon Bottou. Stochastic learning. In *Advanced lectures on machine learning*, pages 146–168. Springer, 2004.
- Leon Bottou. Stochastic Gradient Descent Tricks. In *Neural Networks: Tricks of the Trade*, volume 1, pages 421–436. 2012.
- Alexandre Défossez and Francis Bach. Averaged least-mean-squares: Bias-variance trade-offs and optimal sampling distributions. In *Proceedings of the Eighteenth International Conference on Artificial Intelligence and Statistics*, pages 205–213, 2015.
- John Duchi, Elad Hazan, and Yoram Singer. Adaptive sub-gradient methods for online learning and stochastic optimization. *The Journal of Machine Learning Research*, 12:2121–2159, 2011.
- Nikos Karampatziakis and John Langford. Online importance weight aware updates. *arXiv preprint arXiv:1011.1576*, 2010.
- Diederik Kingma and Jimmy Ba. Adam: A Method for Stochastic Optimization. In *International Conference on Learning Representations*, 2015.
- Jyrki Kivinen, Manfred K Warmuth, and Babak Hassibi. The p-norm generalization of the lms algorithm for adaptive filtering. *Signal Processing, IEEE Transactions on*, 54(5):1782–1793, 2006.
- Krzysztof A Krakowski, Robert E Mahony, Robert C Williamson, and Manfred K Warmuth. A geometric view of non-linear on-line stochastic gradient descent. *Author website*, 2007.
- Brian Kulis and Peter L Bartlett. Implicit online learning. In *Proceedings of the 27th International Conference on Machine Learning (ICML-10)*, pages 575–582, 2010.
- Yann Le Cun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of IEEE*, 86(11):2278–2324, 1998.
- David Lewis, Yiming Yang, Tony Rose, and Fan Li. Rcv1: A new benchmark collection for text categorization research. *The Journal of Machine Learning Research*, 5: 361–397, 2004.

- Pierre-Louis Lions and Bertrand Mercier. Splitting algorithms for the sum of two nonlinear operators. *SIAM Journal on Numerical Analysis*, 16(6):964–979, 1979.
- Lennart Ljung, Georg Ch Pflug, and Harro Walk. *Stochastic approximation and optimization of random systems*, volume 17. Springer, 1992.
- Eric Moulines and Francis R Bach. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In *Advances in Neural Information Processing Systems*, pages 451–459, 2011.
- Jin-Ichi Nagumo and Atsuhiko Noda. A learning method for system identification. *Automatic Control, IEEE Transactions on*, 12(3):282–287, 1967.
- Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.
- Yurii Nesterov. *Introductory lectures on convex optimization*, volume 87. Springer Science & Business Media, 2004.
- Neal Parikh and Stephen Boyd. Proximal algorithms. *Foundations and Trends in optimization*, 1(3):123–231, 2013.
- Boris T Polyak and Anatoli B Juditsky. Acceleration of stochastic approximation by averaging. *SIAM Journal on Control and Optimization*, 30(4):838–855, 1992.
- Herbert Robbins and Sutton Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.
- R Tyrrell Rockafellar. Monotone operators and the proximal point algorithm. *SIAM journal on control and optimization*, 14(5):877–898, 1976.
- Lorenzo Rosasco, Silvia Villa, and Bang Công Vũ. Convergence of stochastic proximal gradient algorithm. *arXiv preprint arXiv:1403.5074*, 2014.
- David Ruppert. Efficient estimators from a slowly convergent robbins-monro process. Technical report, School of Operations Research and Industrial Engineering, Cornell University, 1988.
- Ernest K Ryu and Stephen Boyd. Stochastic proximal iteration: A non-asymptotic improvement upon stochastic gradient descent. *Author website, early draft*, 2014.
- David J Sakrison. Efficient recursive estimation; application to estimating the parameters of a covariance function. *International Journal of Engineering Science*, 3(4):461–483, 1965.
- Mark Schmidt, Nicolas Le Roux, and Francis Bach. Minimizing finite sums with the stochastic average gradient. Technical report, HAL 00860051, 2013.
- Li Cheng SVN Schuurmans and SW Caelli. Implicit online learning with kernels. *Advances in neural information processing systems*, 19:249, 2007.
- Ohad Shamir and Tong Zhang. Stochastic gradient descent for non-smooth optimization: Convergence results and optimal averaging schemes. *arXiv preprint arXiv:1212.1824*, 2012.
- Yoram Singer and John C Duchi. Efficient learning using forward-backward splitting. In *Advances in Neural Information Processing Systems*, pages 495–503, 2009.
- Soeren Sonnenburg, Vojtech Franc, Elad Yom-Tov, and Michele Sebag. Pascal large scale learning challenge, 2008. URL <http://largescale.first.fraunhofer.de>.
- T. Tieleman and G. Hinton. Lecture 6.5—RmsProp: Divide the Gradient by a Running Average of its Recent Magnitude. COURSERA: Neural Networks for Machine Learning, 2012.
- Panos Toulis and Edoardo M Airolidi. Implicit stochastic gradient descent. *arXiv preprint arXiv:1408.2923*, 2014.
- Panos Toulis and Edoardo M Airolidi. Implicit stochastic approximation. *arXiv preprint arXiv:1510.00967*, 2015.
- Panos Toulis, Jason Rennie, and Edoardo Airolidi. Statistical analysis of stochastic gradient methods for generalized linear models. In *31st International Conference on Machine Learning*, 2014.
- Lin Xiao and Tony Zhang. A proximal stochastic gradient method with progressive variance reduction. *SIAM Journal on Optimization*, 24:2057–2075, 2014.
- Wei Xu. Towards optimal one pass large scale learning with averaged stochastic gradient descent. *arXiv preprint arXiv:1107.2490*, 2011.
- Tong Zhang. Solving large scale linear prediction problems using stochastic gradient descent algorithms. In *Proceedings of the twenty-first international conference on Machine learning*, page 116. ACM, 2004.

A Note

Lemmas 1, 2, 3 and 4, and Corollary 1, were originally derived by [Toulis and Airoldi \(2014\)](#). These intermediate results (and Theorem 1) provide the necessary foundation to derive Lemma 5 (only in this supplement) and Theorem 2 on the asymptotic optimality of $\bar{\theta}_n$, which is the key result of the main paper. We fully state these intermediate results here for convenience but we point the reader to the aforementioned reference for the proofs and for more details on the theory of (non-averaged) implicit stochastic gradient descent (implicit SGD).

B Introduction

Consider a random variable $\xi \in \Xi$, a parameter space Θ that is convex and compact, and a loss function $L : \Theta \times \Xi \rightarrow \mathbb{R}$. We wish to solve the following stochastic optimization problem:

$$\theta_\star = \arg \min_{\theta \in \Theta} \mathbb{E} (L(\theta, \xi)), \quad (12)$$

where the expectation is with respect to ξ . Define the expected loss,

$$\ell(\theta) = \mathbb{E} (L(\theta, \xi)), \quad (13)$$

where L is differentiable almost-surely. In this work we study a stochastic approximation procedure to solve (12) defined through the iterations

$$\theta_n = \theta_{n-1} - \gamma_n \nabla L(\theta_n, \xi_n), \quad \theta_0 \in \Theta, \quad (14)$$

$$\bar{\theta}_n = \frac{1}{n} \sum_{i=1}^n \theta_i, \quad (15)$$

where $\{\xi_1, \xi_2, \dots\}$ are i.i.d. realizations of ξ , and $\nabla L(\theta, \xi_n)$ is the gradient of the loss function with respect to θ given realized value ξ_n . The sequence $\{\gamma_n\}$ is a non-increasing sequence of positive real numbers. We will refer to procedure defined by (14) and (15) as *averaged implicit stochastic gradient descent*, or AI-SGD for short. Procedure AI-SGD combines two ideas, namely an implicit update in Eq. (14) as θ_n appears on both sides of the update, and averaging of the iterates θ_n in Eq. (15).

C Notation and assumptions

Let $\mathcal{F}_n = \{\theta_0, \xi_1, \xi_2, \dots, \xi_n\}$ denote the filtration that process θ_n (14) is adapted to. The norm $\|\cdot\|$ will denote the L_2 norm. The symbol \triangleq indicates a definition, and the symbol $\stackrel{\text{def}}{=}$ denotes “equal by definition”. For example, $x \triangleq y$ defines x as equal to known variable y , whereas $x \stackrel{\text{def}}{=} y$ denotes that x is equal to y by definition. We will not use this formalism when defining constants. For two positive sequences a_n, b_n , we write $b_n = \mathcal{O}(a_n)$ if there exists a fixed $c > 0$ such that $b_n \leq ca_n$, for all n ; also, $b_n = o(a_n)$ if $b_n/a_n \rightarrow 0$. When a positive scalar sequence a_n is monotonically decreasing to zero, we write $a_n \downarrow 0$. Similarly, for a sequence X_n of vectors or matrices, $X_n = \mathcal{O}(a_n)$ denotes that $\|X_n\| = \mathcal{O}(a_n)$, and $X_n = o(a_n)$ denotes that $\|X_n\| = o(a_n)$. For two matrices A, B , $A \preceq B$ denotes that $B - A$ is nonnegative-definite; $\text{tr}(A)$ denotes the trace of A .

We now introduce the main assumptions pertaining to the theory of this paper.

Assumption 6. *The loss function $L(\theta, \xi)$ is almost-surely differentiable. The random vector ξ can be decomposed as $\xi = (x, y)$, $x \in \mathbb{R}^p, y \in \mathbb{R}^d$, such that*

$$L(\theta, \xi) = L(x^\top \theta, y). \quad (16)$$

Assumption 7. *The learning rate sequence $\{\gamma_n\}$ is defined as $\gamma_n = \gamma_1 n^{-\gamma}$, where $\gamma_1 > 0$ and $\gamma \in (1/2, 1]$.*

Assumption 8 (Lipschitz conditions). *For all $\theta_1, \theta_2 \in \Theta$, a combination of the following conditions is satisfied almost-surely:*

(a) *The loss function L is Lipschitz-continuous with parameter λ_0 , i.e.,*

$$|L(\theta_1, \xi) - L(\theta_2, \xi)| \leq \lambda_0 \|\theta_1 - \theta_2\|,$$

(b) The map ∇L is Lipschitz-continuous with parameter λ_1 , i.e.,

$$\|\nabla L(\theta_1, \xi) - \nabla L(\theta_2, \xi)\| \leq \lambda_1 \|\theta_1 - \theta_2\|,$$

(c) The map $\nabla^2 L$ is Lipschitz-continuous with parameter λ_2 , i.e.,

$$\|\nabla^2 L(\theta_1, \xi) - \nabla^2 L(\theta_2, \xi)\| \leq \lambda_2 \|\theta_1 - \theta_2\|.$$

Assumption 9. The observed Fisher information matrix, $\hat{\mathcal{I}}(\theta) \triangleq \nabla^2 L(\theta, \xi)$, has non-vanishing trace, i.e., there exists $\phi > 0$ such that $\text{tr}(\hat{\mathcal{I}}(\theta)) \geq \phi$, almost-surely, for all $\theta \in \Theta$. The expected Fisher information matrix, $\mathcal{I}(\theta) \triangleq \mathbb{E}(\hat{\mathcal{I}}(\theta))$, has minimum eigenvalue $0 < \lambda_f \leq \phi$, for all $\theta \in \Theta$.

Assumption 10. The zero-mean random variable $W_\theta \triangleq \nabla L(\theta, \xi) - \nabla \ell(\theta)$ is square-integrable, such that, for a fixed positive-definite Σ ,

$$\mathbb{E}(W_{\theta_*} W_{\theta_*}^\top) \preceq \Sigma.$$

D Proof of Lemma 2

Definition 2. Suppose that Assumption 6 holds. For observation $\xi = (x, y)$, the first derivative with respect to the natural parameter $x^\top \theta$ is denoted by $L'(\theta, \xi)$, and is defined as

$$L'(\theta, \xi) \triangleq \frac{\partial L(\theta, \xi)}{\partial (x^\top \theta)} \stackrel{\text{def}}{=} \frac{\partial L(x^\top \theta, y)}{\partial (x^\top \theta)}. \quad (17)$$

Similarly, $L''(\xi, \theta) \triangleq \frac{\partial L'(\theta, \xi)}{\partial (x^\top \theta)}$.

Lemma 2. Suppose that Assumption 6 holds, and consider functions L', L'' from Definition 2. Then, almost-surely,

$$\nabla L(\theta_n, \xi_n) = s_n \nabla L(\theta_{n-1}, \xi_n); \quad (18)$$

the scalar s_n satisfies the fixed-point equation,

$$s_n \kappa_{n-1} = L'(\theta_{n-1} - s_n \gamma_n \kappa_{n-1} x_n, \xi_n), \quad (19)$$

where $\kappa_{n-1} \triangleq L'(\theta_{n-1}, \xi_n)$. Moreover, if $L''(\theta, \xi) \geq 0$ almost-surely for all $\theta \in \Theta$, then

$$s_n \in \begin{cases} [\kappa_{n-1}, 0) & \text{if } \kappa_{n-1} < 0, \\ [0, \kappa_{n-1}] & \text{otherwise.} \end{cases}$$

Proof. See Toulis and Airoidi (2014, Theorem 4.1). □

E Proof of Theorem 3

E.1 Useful lemmas

In this section, we will present the intermediate lemmas on recursions that will be useful for the non-asymptotic analysis of the implicit procedures.

Lemma 3. Consider a sequence b_n such that $b_n \downarrow 0$ and $\sum_{i=1}^{\infty} b_i = \infty$. Then, there exists a positive constant $K > 0$, such that

$$\prod_{i=1}^n \frac{1}{1 + b_i} \leq \exp(-K \sum_{i=1}^n b_i). \quad (20)$$

Proof. See [Toulis and Airoidi \(2014, Lemma B.1\)](#). \square

Lemma 4. Consider scalar sequences $a_n \downarrow 0$, $b_n \downarrow 0$, and $c_n \downarrow 0$ such that, $a_n = o(b_n)$, and $A \triangleq \sum_{i=1}^{\infty} a_i < \infty$. Suppose there exists n' such that $c_n/b_n < 1$ for all $n > n'$. Define,

$$\delta_n \triangleq \frac{1}{a_n}(a_{n-1}/b_{n-1} - a_n/b_n) \text{ and } \zeta_n \triangleq \frac{c_n}{b_{n-1}} \frac{a_{n-1}}{a_n}, \quad (21)$$

and suppose that $\delta_n \downarrow 0$ and $\zeta_n \downarrow 0$. Fix $n_0 > 0$ such that $\delta_n + \zeta_n < 1$ and $(1 + c_n)/(1 + b_n) < 1$, for all $n \geq n_0$.

Consider a positive sequence $y_n > 0$ that satisfies the recursive inequality,

$$y_n \leq \frac{1 + c_n}{1 + b_n} y_{n-1} + a_n. \quad (22)$$

Then, for every $n > 0$,

$$y_n \leq K_0 \frac{a_n}{b_n} + Q_1^n y_0 + Q_{n_0+1}^n (1 + c_1)^{n_0} A, \quad (23)$$

where $K_0 = (1 + b_1)(1 - \delta_{n_0} - \zeta_{n_0})^{-1}$, and $Q_i^n = \prod_{j=i}^n (1 + c_j)/(1 + b_j)$, such that $Q_i^n = 1$ if $n < i$, by definition.

Proof. See [Toulis and Airoidi \(2014, Lemma B.2\)](#). \square

Corollary 1. In Lemma 4 assume $a_n = a_1 n^{-\alpha}$ and $b_n = b_1 n^{-\beta}$, and $c_n = 0$, where $a_1, b_1, \beta > 0$ and $\max\{\beta, 1\} < \alpha < 1 + \beta$, and $\beta \neq 1$. Then,

$$y_n \leq 2 \frac{a_1(1 + b_1)}{b_1} n^{-\alpha+\beta} + \exp(-\log(1 + b_1)n^{1-\beta})[y_0 + (1 + b_1)^{n_0} A], \quad (24)$$

where $n_0 > 0$ and $A = \sum_i a_i < \infty$. If $\beta = 1$ then the above inequality holds by replacing the term $n^{1-\beta}$ with $\log n$.

Proof. See [Toulis and Airoidi \(2014, Corollary B.1\)](#). \square

Lemma 5. Suppose Assumptions 6, 8(a), and 9 hold. Then, almost surely,

$$s_n \geq \frac{1}{1 + \gamma_n \phi}, \quad (25)$$

$$\|\theta_n - \theta_{n-1}\|^2 \leq 4\lambda_0^2 \gamma_n^2, \quad (26)$$

where s_n is defined in Lemma 2, and θ_n is the n th iterate of implicit SGD (14).

Proof. See [Toulis and Airoidi \(2014, Lemma B.3\)](#). \square

Theorem 3. Suppose that Assumptions 6, 7, 8(a), and 9 hold. Define $\delta_n \triangleq \mathbb{E}(\|\theta_n - \theta_\star\|^2)$, and constants $\Gamma^2 = 4\lambda_0^2 \sum \gamma_i^2 < \infty$, $\epsilon = (1 + \gamma_1(\phi - \underline{\lambda}_f))^{-1}$, and $\lambda = 1 + \gamma_1 \underline{\lambda}_f \epsilon$. Then, there exists constant $n_0 > 0$ such that, for all $n > 0$,

$$\delta_n \leq (8\lambda_0^2 \gamma_1 \lambda / \underline{\lambda}_f \epsilon) n^{-\gamma} + e^{-\log \lambda \cdot n^{1-\gamma}} [\delta_0 + \lambda^{n_0} \Gamma^2].$$

Proof. See [Toulis and Airoidi \(2014, Theorem 3.1\)](#). \square

Remarks. #1. Assuming Lipschitz continuity of the gradient ∇L instead of function L , i.e., Assumption 8(b) over Assumption 8(a) would not alter the main result of Theorem 3 about the $\mathcal{O}(n^{-\gamma})$ rate of the mean-squared error. Assuming Lipschitz continuity with constant λ_1 of ∇L and boundedness of $\mathbb{E}(\|\nabla L(\theta_\star, \xi_n)\|^2) \leq \sigma^2$, as it is typical in the literature,

would simply add a term $\gamma_n^2 \lambda_1^2 \mathbb{E}(\|\theta_n^{\text{im}} - \theta_\star\|^2) + \gamma_n^2 \sigma^2$ in the corresponding recursive inequality. Specifically, by Lemma 2, $s_n \leq 1$, and thus

$$\begin{aligned} \mathbb{E}(\|\nabla L(\theta_n, \xi_n)\|^2) &= \mathbb{E}(s_n^2 \|\nabla L(\theta_{n-1}, \xi_n)\|^2) \leq \mathbb{E}(\|\nabla L(\theta_{n-1}, \xi_n)\|^2) \\ &= \mathbb{E}(\|\nabla L(\theta_{n-1}, \xi_n) - \nabla L(\theta_\star, \xi_n) + \nabla L(\theta_\star, \xi_n)\|^2) \\ &\leq \lambda_1^2 \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^2) + \gamma_n^2 \mathbb{E}(\|\nabla L(\theta_\star, \xi_n)\|^2) \\ &\leq \lambda_1^2 \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^2) + \gamma_n^2 \sigma^2. \end{aligned} \quad (27)$$

The recursion for the implicit errors would then be

$$\mathbb{E}(\|\theta_n^{\text{im}} - \theta_\star\|^2) \leq \left(\frac{1}{1 + \gamma_n \lambda_f \epsilon} + \lambda_1^2 \gamma_n^2 \right) \mathbb{E}(\|\theta_{n-1}^{\text{im}} - \theta_\star\|^2) + \gamma_n^2 \sigma^2,$$

which also implies the $\mathcal{O}(n^{-\gamma})$ convergence rate. However, it is an open problem whether it is possible to derive a nice stability property for implicit SGD under Assumption 8(b) similar to the result of Theorem 3 under Assumption 8(a).

Remarks. #2. An assumption of almost-sure convexity can simplify the analysis significantly. For example, similar to the assumption of Ryu and Boyd (2014), assume that $L(\theta, \xi)$ is convex almost surely such that

$$(\theta_n - \theta_\star)^\top \nabla L(\theta_n, \xi_n) \geq \frac{\mu_n}{2} \|\theta_n^{\text{im}} - \theta_\star\|^2, \quad (28)$$

where $\mu_n \geq 0$ and $\mathbb{E}(\mu_n) = \mu > 0$. Then,

$$\begin{aligned} \theta_n + 2\gamma_n \nabla L(\theta_n, \xi_n) &= \theta_{n-1} \quad [\text{by definition of implicit SGD (14)}] \\ \|\theta_n - \theta_\star\|^2 + 2\gamma_n (\theta_n - \theta_\star)^\top \nabla L(\theta_n, \xi_n) &\leq \|\theta_{n-1} - \theta_\star\|^2. \\ (1 + \gamma_n \mu_n) \|\theta_n - \theta_\star\|^2 &\leq \|\theta_{n-1} - \theta_\star\|^2. \\ \mathbb{E}(\|\theta_n - \theta_\star\|^2) &\leq \frac{1}{1 + \gamma_n \mu} \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^2) + \text{SD}(1 + \gamma_n \mu_n) \text{SD}(\|\theta_n - \theta_\star\|^2), \end{aligned} \quad (29)$$

where the last inequality follows from the identity $\mathbb{E}(XY) \geq \mathbb{E}(X) \mathbb{E}(Y) - \text{SD}(X) \text{SD}(Y)$. However, $\text{SD}(1 + \gamma_n \mu_n) = \mathcal{O}(\gamma_n)$, and assuming bounded θ_n we get

$$\mathbb{E}(\|\theta_n - \theta_\star\|^2) \leq \frac{1}{1 + \gamma_n \mu} \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^2) + \mathcal{O}(\gamma_n), \quad (30)$$

which indicates a fast convergence towards θ_\star . It is also possible to work with the recursion

$$\|\theta_n - \theta_\star\|^2 \leq \frac{1}{1 + \gamma_n \mu_n} \|\theta_{n-1} - \theta_\star\|^2, \quad (31)$$

and then use a stochastic version of Lemma 4 although the analysis would be more complex in this case.

F Proof of Theorem 5

In this section, we prove Theorem 5. To do so, we need bounds for $\mathbb{E}(\|\theta_n - \theta_\star\|^2)$, which are available through Theorem 3, but also bounds for $\mathbb{E}(\|\theta_n - \theta_\star\|^4)$, which are established in the following lemma.

Theorem 4. Suppose that Assumptions 6, 7, 8(a), and 9 hold. For a constant $K_3 > 0$, define $\zeta_n \triangleq \mathbb{E}(\|\theta_n - \theta_\star\|^2)$, and constants $\Delta^3 \triangleq K_3 \sum \gamma_i^3 < \infty$, $\epsilon \triangleq (1 + \gamma_1(\phi - \lambda_f))^{-1}$, and $\lambda \triangleq 1 + \gamma_1 \lambda_f \epsilon$. Then, there exists constant n_0 such that, for all $n > 0$,

$$\zeta_n \leq (2K_3 \gamma_1^2 \lambda / \lambda_f \epsilon) n^{-2\gamma} + e^{-\log \lambda \cdot n^{1-\gamma}} [\zeta_0 + \lambda^{n_0} \Delta^3].$$

Proof. Define $W_n \triangleq s_n(\theta_{n-1}^{\text{im}} - \theta_\star)^\top \nabla L(\theta_{n-1}, \xi_n)$ for compactness, and proceed as follows,

$$\begin{aligned}
 \|\theta_n^{\text{im}} - \theta_\star\|^2 &= \|\theta_{n-1}^{\text{im}} - \theta_\star\|^2 - 2\gamma_n s_n(\theta_{n-1}^{\text{im}} - \theta_\star)^\top \nabla L(\theta_{n-1}, \xi_n) + \gamma_n^2 \|\nabla L(\theta_{n-1}, \xi_n)\|^2 \\
 \|\theta_n^{\text{im}} - \theta_\star\|^2 &= \|\theta_{n-1}^{\text{im}} - \theta_\star\|^2 - 2\gamma_n W_n + \gamma_n^2 \|\nabla L(\theta_{n-1}, \xi_n)\|^2 \quad [\text{by definition}] \\
 \|\theta_n^{\text{im}} - \theta_\star\|^2 &\leq \|\theta_{n-1}^{\text{im}} - \theta_\star\|^2 - 2\gamma_n W_n + 4\lambda_0^2 \gamma_n^2, \\
 \|\theta_n^{\text{im}} - \theta_\star\|^4 &\leq \|\theta_{n-1}^{\text{im}} - \theta_\star\|^4 + 4\gamma_n^2 W_n^2 + 16\lambda_0^4 \gamma_n^4 \\
 &\quad - 2\gamma_n \|\theta_{n-1} - \theta_\star\|^2 W_n + 4\lambda_0^2 \gamma_n^2 \|\theta_{n-1} - \theta_\star\|^2 - 8\lambda_0^2 \gamma_n^3 W_n.
 \end{aligned} \tag{32}$$

By Lemma 4 we have

$$\mathbb{E}(W_n | \mathcal{F}_{n-1}) \geq \frac{\lambda_f}{2(1 + \gamma_n \phi)} \|\theta_{n-1} - \theta_\star\|^2. \tag{33}$$

Furthermore,

$$\begin{aligned}
 \mathbb{E}(W_n^2 | \mathcal{F}_{n-1}) &\stackrel{\text{def}}{=} \mathbb{E}([s_n(\theta_{n-1} - \theta_\star)^\top \nabla L(\theta_{n-1}, \xi_n)]^2 | \mathcal{F}_{n-1}) \\
 &\stackrel{\text{def}}{=} \mathbb{E}([\theta_{n-1} - \theta_\star]^\top \nabla L(\theta_{n-1}, \xi_n) [\theta_{n-1} - \theta_\star] | \mathcal{F}_{n-1}) \quad [\text{by Lemma 1}] \\
 &\leq \|\theta_{n-1} - \theta_\star\|^2 \mathbb{E}(\|\nabla L(\theta_{n-1}, \xi_n)\|^2 | \mathcal{F}_{n-1}) \quad [\text{by Cauchy-Schwartz inequality}] \\
 &\leq 4\lambda_0^2 \|\theta_{n-1} - \theta_\star\|^2 \quad [\text{by Lemma 4}]
 \end{aligned} \tag{34}$$

Define $B_n \triangleq \mathbb{E}(\|\theta_n - \theta_\star\|^2)$ for notational brevity. We use results (33) and (34) to get

$$\begin{aligned}
 \mathbb{E}(\|\theta_n - \theta_\star\|^4) &\leq \left(1 - \frac{\gamma_n \lambda_f}{1 + \gamma_n \phi}\right) \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^4) + 4\lambda_0^2 \gamma_n^2 \left(5 - \frac{\gamma_n \lambda_f}{1 + \gamma_n \phi}\right) B_{n-1} + 16\lambda_0^4 \gamma_n^4 \\
 \mathbb{E}(\|\theta_n - \theta_\star\|^4) &\leq \left(1 - \frac{\gamma_n \lambda_f}{1 + \gamma_n \phi}\right) \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^4) + 20\lambda_0^2 \gamma_n^2 B_{n-1} + 16\lambda_0^4 \gamma_n^4 \\
 \mathbb{E}(\|\theta_n - \theta_\star\|^4) &\leq \frac{1}{1 + \gamma_n \lambda_f \epsilon} \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^4) + 20\lambda_0^2 \gamma_n^2 B_{n-1} + 16\lambda_0^4 \gamma_n^4 \quad [\text{by Assumption 9}] \\
 \mathbb{E}(\|\theta_n - \theta_\star\|^4) &\leq \frac{1}{1 + \gamma_n \lambda_f \epsilon} \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^4) + K_0 \gamma_n^3 + e^{-\log \lambda \cdot n^{1-\gamma}} K_1 + K_2 \gamma_n^4, \quad [\text{by Theorem 3}]
 \end{aligned} \tag{35}$$

where $\lambda = (1 + \gamma_1(\phi - \lambda_f))^{-1}$ and $\Gamma^2 = 4\lambda_0^2 \sum \gamma_i^2$, (as in Theorem 3), $K_0 \triangleq 160\lambda_0^4 \lambda / \lambda_f$, $K_1 \triangleq 20\lambda_0^2 (\mathbb{E}(\|\theta_0 - \theta_\star\|^2) + \lambda^{n_0} \Gamma^2)$, and $K_2 \triangleq 16\lambda_0^4$, and n_0 is a constant defined in the proof of Theorem 3.

Now, define

$$K_3 \triangleq K_0 + K_2 \gamma_1 + \max\left\{\frac{e^{-\log \lambda \cdot \rho_\gamma(n) K_1}}{\gamma_n^3}\right\}, \tag{36}$$

which exists and is finite. Through simple algebra it is easy to verify that

$$K_0 \gamma_n^3 + e^{-\log \lambda \cdot \rho_\gamma(n) K_1} + K_2 \gamma_n^4 \leq K_3 \gamma_n^3, \tag{37}$$

for all n . Therefore, we can simplify Ineq. (35) as

$$\mathbb{E}(\|\theta_n - \theta_\star\|^4) \leq \frac{1}{1 + \gamma_n \lambda_f \epsilon} \mathbb{E}(\|\theta_{n-1} - \theta_\star\|^4) + K_3 \gamma_n^3. \tag{38}$$

We can now apply Corollary 1 with $a_n \equiv K_3 \gamma_n^3$ and $b_n \equiv \gamma_n \lambda_f \epsilon$ to derive the final bounds for $\mathbb{E}(\|\theta_n - \theta_\star\|^4)$. \square

We now evaluate the mean squared error of the averaged iterates, $\bar{\theta}_n$.

Theorem 5. Consider the AI-SGD procedure 15 and suppose that Assumptions 6, 7, 8(a), 8(c), 9, and 10 hold. Then,

$$\begin{aligned}
 (\mathbb{E}(\|\bar{\theta}_n - \theta_\star\|^2))^{1/2} &\leq \frac{1}{\sqrt{n}} (\text{trace}(\nabla^2 \ell(\theta_\star)^{-1} \Sigma \nabla^2 \ell(\theta_\star)^{-1}))^{1/2} \\
 &\quad + \frac{2\gamma + 1}{\lambda_f^{1/2} \gamma_1} (8\lambda_0^2 \gamma_1 \lambda / \lambda_f \epsilon)^{1/2} n^{-1+\gamma/2} \\
 &\quad + \frac{2\gamma + 1}{\lambda_f^{1/2} n \gamma_n} [\delta_0 + \lambda^{n_{0,1}} \Gamma^2]^{1/2} e^{-\log \lambda \cdot n^{1-\gamma}/2} \\
 &\quad + \frac{\lambda_2}{2\lambda_f^{1/2}} (2K_3 \gamma_1^2 \lambda / \lambda_f \epsilon)^{1/2} n^{-\gamma} \\
 &\quad + \frac{\lambda_2}{2n\lambda_f^{1/2}} [\zeta_0 + \lambda^{n_{0,2}} \Delta^3]^{1/2} K_2(n).
 \end{aligned} \tag{39}$$

where $K_2(n) = \sum_{i=1}^n \exp(-\log \lambda \cdot i^{1-\gamma}/2)$, and constants $\lambda, \epsilon, n_{0,1}, \delta_0, \Gamma^2$ are defined in Theorem 3 (substituting n_0 for $n_{0,1}$), and $\zeta_0, n_{0,2}, \Delta^3$ are defined in Theorem 4, substituting $(n_0$ for $n_{0,2})$.

Proof. We leverage a result shown for averaged explicit stochastic gradient descent. In particular, it has been shown that the squared error for the averaged iterate satisfies:

$$\begin{aligned}
 (\mathbb{E}(\|\bar{\theta}_n - \theta_\star\|^2))^{1/2} &\leq \frac{1}{\sqrt{n}} (\text{trace}(\nabla^2 \ell(\theta_\star)^{-1} \Sigma \nabla^2 \ell(\theta_\star)^{-1}))^{1/2} \\
 &\quad + \frac{2\gamma + 1}{\lambda_f^{1/2} n \gamma_n} (\mathbb{E}(\|\theta_n - \theta_\star\|^2))^{1/2} \\
 &\quad + \frac{\lambda_2}{2n\lambda_f^{1/2}} \sum_{i=1}^n (\mathbb{E}(\|\theta_i - \theta_\star\|^4))^{1/2}.
 \end{aligned} \tag{40}$$

The proof technique for (40) was first devised by Polyak and Juditsky (1992), but was later refined by Xu (2011), and Moulines and Bach (2011). In this paper, we follow the formulation of Moulines and Bach (2011, Theorem 3, page 20); the derivation of Ineq.(40) for the implicit procedure is identical to the derivation for the explicit one, however the two procedures differ in the terms that appear in the bound (40).

All such terms in (40) have been bounded in the previous sections. In particular, we can use Theorem 3 for $\mathbb{E}(\|\theta_n - \theta_\star\|^2)$; we can also use Theorem 5 and the concavity of the square-root to derive

$$\begin{aligned}
 \sum_{i=1}^n (\mathbb{E}(\|\theta_i - \theta_\star\|^4))^{1/2} &\leq \sum_{i=1}^n \left((2K_3 \gamma_1^2 \lambda / \lambda_f \epsilon)^{1/2} i^{-\gamma} + e^{-\log \lambda \cdot i^{1-\gamma}/2} [\zeta_0 + \lambda^{n_{0,2}} \Delta^3]^{1/2} \right) \\
 &\leq (2K_3 \gamma_1^2 \lambda / \lambda_f \epsilon)^{1/2} n^{1-\gamma} + K_2(n) [\zeta_0 + \lambda^{n_{0,2}} \Delta^3]^{1/2},
 \end{aligned} \tag{41}$$

where $K_2(n) = \sum_{i=1}^n \exp(-\frac{\log \lambda}{2} i^{1-\gamma})$, $\zeta_0 = \mathbb{E}(\|\theta_0 - \theta_\star\|^4)$, and $\Delta^3, n_{0,2}$ are defined in Lemma ??, substituting n_0 for $n_{0,2}$. Similarly, using Theorem 3,

$$(\mathbb{E}(\|\theta_n - \theta_\star\|^2))^{1/2} \leq (8\lambda_0^2 \gamma_1 \lambda / \lambda_f \epsilon)^{1/2} n^{-\gamma/2} + e^{-\log \lambda \cdot n^{1-\gamma}/2} [\delta_0 + \lambda^{n_{0,1}} \Gamma^2]^{1/2},$$

where $\delta_0 = \mathbb{E}(\|\theta_n - \theta_\star\|^2)$, and $n_{0,1}, \Gamma^2$ are defined in Theorem 3, substituting $n_{0,1}$ for n_0 . These two bounds can be used in Ineq.(40) and thus yield the result of Theorem 5. \square

G Data sets used in experiments

Table 2 includes a full summary of all data sets considered in our experiments. The majority of regularization parameters are set according to Xu (2011).

	description	type	features	training set	test set	λ
covtype	forest cover type	sparse	54	464,809	116,203	10^{-6}
delta	synthetic data	dense	500	450,000	50,000	10^{-2}
rcv1	text data	sparse	47,152	781,265	23,149	10^{-5}
mnist	digit image features	dense	784	60,000	10,000	10^{-3}
sido	molecular activity	dense	4,932	10,142	2,536	10^{-3}
alpha	synthetic data	dense	500	400k	50k	10^{-5}
beta	synthetic data	dense	500	400k	50k	10^{-4}
gamma	synthetic data	dense	500	400k	50k	10^{-3}
epsilon	synthetic data	dense	2000	400k	50k	10^{-5}
zeta	synthetic data	dense	2000	400k	50k	10^{-5}
fd	character image	dense	900	1000k	470k	10^{-5}
ocr	character image	dense	1156	1000k	500k	10^{-5}
dna	DNA sequence	sparse	800	1000k	1000k	10^{-3}

Table 2: Summary of data sets and the L_2 regularization parameter λ used