

Theoretical analysis of direct CP violation and differential decay width in $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ in phase space around the resonances $\rho^0(770)$ and $f_0(500)$

Zhen-Hua Zhang^{1,2,a}, Ren Song¹, Yu-Mo Su¹, Gang Lü^{3,b}, and Bo Zheng^{1,2,c}

¹School of Nuclear Science and Technology, University of South China, Hengyang, Hunan 421001, China.

²Cooperative Innovation Centre for Nuclear Fuel Cycle Technology & Equipment, University of South China, Hengyang, Hunan 421001, China.

³ College of Science, Henan University of Technology, Zhengzhou 450001, China.
email: ^azhangzh@usc.edu.cn, ^bganglv@haut.edu.cn, ^czhengb@ihep.ac.cn

January 28, 2022

Abstract

We perform a theoretical study on direct CP violation in $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ in phase space around the intermediate states $\rho^0(770)$ and $f_0(500)$. The possible interference between the amplitudes corresponding to the two resonances is taken into account, and the relative strong phase of the two amplitudes is treated as a free parameter. Our analysis shows that by properly chosen the strong phase, both the CP violation strength and differential decay width accommodate to the experimental results.

1 Introduction

Charge-Parity (CP) violation has gain extensive attentions ever since its first discovery in $K^0 - \bar{K}^0$ systems in 1964 [1]. Within the Standard Model (SM), CP violation is originated from a complex phase in Cabibbo-Kobayashi-Maskawa (CKM) matrix, which describes the mixing of weak and mass eigenstates of quarks [2].

Although it is a small effect in general, CP violation can be relatively large in some decay channels of B and B_s mesons [3, 4, 5]. In fact, large CP violation has been confirmed in some two-body decay channel of B and B_s meson, such as $B^\pm \rightarrow D_{CP(+1)} \pi^\pm$ [6], $B^\pm \rightarrow \rho^0 K^\pm$ [7], $B^\pm \rightarrow \rho^0 K^*(892)^\pm$ [8], $B^\pm \rightarrow f^0(1370) \pi^\pm$ [9], $B^0 \rightarrow \rho^- K^+$ [10], and $B_s \rightarrow \pi^+ K^-$ [11, 12]. In recent years, even larger CP violation which localized in three-body decay phase space of B meson was observed by LHCb collaboration in channels such as $B^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ and $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ [13, 14]. In view of its anisotropy property for small invariant mass of $\pi^+ \pi^-$ pair (no larger than the mass of $\rho^0(770)$) in phase space, the large localized CP violation was first interpreted as a consequence of the interference of the decay amplitudes

corresponding to nearby resonances with different spins [15, 16]. Some other explanations such as the re-scattering effects of final states [17, 18] were also proposed thereafter.

Since it is believed to be very small within SM, CP violation in the charm sector provides a good place for searching for New Physics. However, due mainly to the pollution of non-perturbative effects of strong interactions, an order of 1% CP asymmetries in D meson decay are still understandable in SM [19, 20, 21]. To date, no CP violation in charm sector is established. Though there were some hints of CP violation for the channels $D \rightarrow \pi\pi$ and $D \rightarrow KK$ [22, 23], the latest result from LHCb collaboration showed however, no evidence of CP violation in these channels [24].

A measurement of CP violation in the three-body decay $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ has also been performed by LHCb [25]. With very high statistics, no localized or overall CP asymmetries are found. As has been shown in some aforementioned three-body decay channels of B meson, localized CP asymmetries in phase space can be enhanced by the interference of the decay amplitudes corresponding to two intermediate resonances with different spins, whereas the same mechanism should also apply to D meson decays. In this paper, we will perform a theoretical analysis of CP violation and differential decay width in the channel $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$, and will pay special attention to effects caused by the interference of intermediate state $\rho^0(770)$ and $f_0(500)$.

2 Formalism of decay amplitudes and CP asymmetries

In the region of the phase space around the resonances $\rho^0(770)$ and $f_0(500)$, the process $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ is dominated by two cascade decays, $D^\pm \rightarrow \pi^\pm f_0(500) \rightarrow \pi^\pm \pi^+ \pi^-$ and $D^\pm \rightarrow \pi^\pm \rho^0(770) \rightarrow \pi^\pm \pi^+ \pi^-$. For the two weak decays $D^\pm \rightarrow \pi^\pm f_0(500)$ and $D^\pm \rightarrow \pi^\pm \rho^0(770)$, the corresponding effective Hamiltonian can be expressed as [26, 27]

$$H_{\Delta C=1} = \frac{G_F}{\sqrt{2}} \left\{ \left[\sum_{q=d,s} V_{uq} V_{cq}^* (c_1 O_1^q + c_2 O_2^q) \right] - V_{ub} V_{cb}^* \sum_{i=3}^6 c_i O_i \right\} + \text{h.c.}, \quad (1)$$

where G_F is the Fermi constant, $V_{q_1 q_2}$ (q_1 and q_2 represent quarks) is the CKM matrix element, c_i ($i = 1, \dots, 6$) is the Wilson coefficient, and O_i is the four quark operator, which can be written as

$$\begin{aligned} O_1^q &= \bar{u}_\alpha \gamma_\mu (1 - \gamma_5) q_\beta \bar{q}_\beta \gamma^\mu (1 - \gamma_5) c_\alpha, \\ O_2^q &= \bar{u} \gamma_\mu (1 - \gamma_5) q \bar{q} \gamma^\mu (1 - \gamma_5) c, \\ O_3 &= \bar{u} \gamma_\mu (1 - \gamma_5) c \sum_{q'} \bar{q}' \gamma^\mu (1 - \gamma_5) q', \\ O_4 &= \bar{u}_\alpha \gamma_\mu (1 - \gamma_5) c_\beta \sum_{q'} \bar{q}'_\beta \gamma^\mu (1 - \gamma_5) q'_\alpha, \\ O_5 &= \bar{u} \gamma_\mu (1 - \gamma_5) c \sum_{q'} \bar{q}' \gamma^\mu (1 + \gamma_5) q', \\ O_6 &= \bar{u}_\alpha \gamma_\mu (1 - \gamma_5) c_\beta \sum_{q'} \bar{q}'_\beta \gamma^\mu (1 + \gamma_5) q'_\alpha, \end{aligned} \quad (2)$$

with O_1^q and O_2^q being tree operators, $O_3 - O_6$ being QCD penguin operators, α and β being color indices, and q' running through all the light flavour quarks.

The effective Hamiltonian for the strong decays $\rho^0 \rightarrow \pi^+\pi^-$ and $f_0(500) \rightarrow \pi^+\pi^-$ can be formally expressed as

$$\mathcal{H}_{\rho^0\pi\pi} = ig_{\rho\pi\pi}\rho_\mu^0(\pi^-\partial^\mu\pi^+ - \pi^+\partial^\mu\pi^-), \quad (3)$$

$$\mathcal{H}_{f_0\pi\pi} = g_{f_0\pi\pi}f_0(2\pi^+\pi^- + \pi^0\pi^0), \quad (4)$$

where ρ_μ^0 , f_0 and π^\pm are the field operators for ρ^0 , $f_0(500)$ and π mesons, respectively, $g_{\rho\pi\pi}$ and $g_{f_0\pi\pi}$ are the effective coupling constants, which can be expressed in terms of the decay widths as

$$g_{\rho\pi\pi}^2 = \frac{48\pi}{\left(1 - \frac{4m_\pi^2}{m_\rho^2}\right)^{3/2}} \cdot \frac{\Gamma_{\rho^0 \rightarrow \pi^+\pi^-}}{m_\rho}, \quad (5)$$

$$g_{f_0\pi\pi}^2 = \frac{4\pi m_{f_0} \Gamma_{f_0 \rightarrow \pi^+\pi^-}}{\left(1 - \frac{4m_\pi^2}{m_{f_0}^2}\right)^{1/2}}. \quad (6)$$

Both $f_0(500)$ and $\rho^0(770)$ decay into one pion pair dominantly through strong interaction, and the isospin symmetry of the strong interaction tells us that $\Gamma_{\rho^0} \simeq \Gamma_{\rho^0 \rightarrow \pi^+\pi^-}$, and $\Gamma_{f_0} \simeq \frac{3}{2}\Gamma_{f_0 \rightarrow \pi^+\pi^-}$.

The decay amplitudes for the cascade decays $D^+ \rightarrow \pi^+ f_0(500) \rightarrow \pi^+\pi^+\pi^-$ and $D^+ \rightarrow \pi^+ \rho^0(770) \rightarrow \pi^+\pi^+\pi^-$ take the form

$$\begin{aligned} \mathcal{M}_{\rho^0}(s_{\text{low}}, s_{\text{high}}) &= \frac{\sqrt{2}G_F g_{\rho\pi\pi} m_\rho (s_{\text{high}} - \Sigma_{s_{\text{low}}})}{s_{\text{low}} - m_{\rho^2} + im_\rho \Gamma_\rho} \cdot \left\{ V_{ud} V_{cd}^* \left(-\frac{1}{\sqrt{2}} a_1 f_\rho F_1 + a_2 f_\pi A_0 \right) \right. \\ &\quad \left. - V_{ub} V_{cb}^* \left[\left(a_4 - \frac{2m_\pi^2 a_6}{(m_c + m_d)(m_u + m_d)} \right) f_\pi A_0 \right] \right\}, \end{aligned} \quad (7)$$

$$\begin{aligned} \mathcal{M}_{f_0}(s_{\text{low}}, s_{\text{high}}) &= \frac{\sqrt{2}G_F (m_D^2 - m_\pi^2) g_{f_0\pi\pi} f_\pi F_0}{s_{\text{low}} - m_{f_0}^2 + im_{f_0} \Gamma_{f_0}} \cdot \left\{ V_{ud} V_{cd}^* a_2 \right. \\ &\quad \left. - V_{ub} V_{cb}^* \left[a_4 - \frac{2m_\pi^2 a_6}{(m_c + m_d)(m_u + m_d)} \right] \right\}, \end{aligned} \quad (8)$$

respectively, where s_{low} and s_{high} are the invariant mass squared of $\pi^+\pi^-$ pairs with lower and higher invariant masses, respectively, $\Sigma_{s_{\text{low}}} = (s_{\text{high,max}} + s_{\text{high,min}})/2$, with $s_{\text{high,max(min)}}$ being the maximum (minimum) value of s_{high} allowed by phase space for each s_{low} , F_0 , F_1 and A_0 are short for the form factors $F_0^{D \rightarrow f_0}(m_\pi^2)$, $F_1^{(D \rightarrow \pi)}(s_{\text{low}})$ and $A_0^{(D \rightarrow \rho)}(m_\pi^2)$, respectively. All the a_i 's are built up from the Wilson coefficients c_i 's, and take the form $a_i = c_i + c_{i+1}/N_c$ for odd i and $a_i = c_i + c_{i-1}/N_c$ for even i .

In the phase space that we are considering, the total decay amplitude for $D^+ \rightarrow \pi^+\pi^+\pi^-$ is dominated by \mathcal{M}_{f_0} and \mathcal{M}_{ρ^0} . As a result, it can be expressed as

$$\mathcal{M} = \left[\mathcal{M}_{f_0}(s_{\text{low}}, s_{\text{high}}) + \mathcal{M}_{\rho^0}(s_{\text{low}}, s_{\text{high}}) e^{i\delta} \right] + [s_{\text{low}} \leftrightarrow s_{\text{high}}], \quad (9)$$

where δ is the relative strong phase between the two amplitudes \mathcal{M}_{f_0} and \mathcal{M}_{ρ^0} , which in principle, arises from long distance effect, $[s_{\text{low}} \leftrightarrow s_{\text{high}}]$ represents a term which take the same form as that in the first square bracket except an interchange between s_{low} and s_{high} . For the calculation of the decay amplitude of the CP conjugate process $D^- \rightarrow \pi^+\pi^-\pi^-$, which will be denoted as $\overline{\mathcal{M}}$, all one need to do is to replace the CKM matrix elements in \mathcal{M} with their complex conjugates.

The differential CP asymmetry for $D^\pm \rightarrow \pi^\pm\pi^+\pi^-$ is defined as

$$A_{CP} = \frac{|\mathcal{M}|^2 - |\overline{\mathcal{M}}|^2}{|\mathcal{M}|^2 + |\overline{\mathcal{M}}|^2}, \quad (10)$$

while the localized CP asymmetry can be expressed as

$$A_{CP}^R = \frac{\int_R ds_{\text{high}} ds_{\text{low}} (|\mathcal{M}|^2 - |\overline{\mathcal{M}}|^2)}{\int_R ds_{\text{high}} ds_{\text{low}} (|\mathcal{M}|^2 + |\overline{\mathcal{M}}|^2)}, \quad (11)$$

where, R represents certain region of the phase space that we are considering.

3 Numerical analysis

Table 1 list the input parameters and the corresponding references we used in this paper. In the following, we give some comments on these input parameters. We use the Wolfenstein parameterization for the CKM matrix elements, which up to the order of λ^8 , can be expressed as [28, 29]

$$\begin{aligned} V_{ud} &= 1 - \frac{\lambda^2}{2} - \frac{\lambda^4}{8} - \frac{\lambda^6}{16}[1 + 8A^2(\rho^2 + \eta^2)] - \frac{\lambda^8}{128}[5 - 32A^2(\rho^2 + \eta^2)], \\ V_{cd} &= -\lambda + \frac{\lambda^5}{2}A^2[1 - 2(\rho + i\eta)] + \frac{\lambda^7}{2}A^2(\rho + i\eta), \\ V_{ub} &= \lambda^3A(\rho - i\eta), \\ V_{cb} &= A\lambda^2 - \frac{\lambda^8}{2}A^3(\rho^2 + \eta^2), \end{aligned} \quad (12)$$

with A, ρ, η , and λ being the Wolfenstein parameters. To all orders in λ , the relation between ρ, η and $\overline{\rho}, \overline{\eta}$ can be expressed as [29]

$$\rho + i\eta = \frac{\sqrt{1 - A^2\lambda^4}(\overline{\rho} + i\overline{\eta})}{\sqrt{1 - \lambda^2}[1 - A^2\lambda^4(\overline{\rho} + i\overline{\eta})]}. \quad (13)$$

For the invariant mass dependence of the form factors $F_1^{D \rightarrow \pi}$ and $A_0^{D \rightarrow \rho}$, we use a model from Ref. [30], which take the form

$$F(s) = \frac{F(0)}{1 - a^X \cdot \frac{s}{m_D^2} + b^X \cdot (\frac{s}{m_D^2})^2}, \quad (14)$$

Table 1: Input parameters used in this paper.

Parameters	Input data	References
Fermi constant (in GeV^{-2})	$G_F = 1.16638 \times 10^{-5}$	[32]
Wilson coefficients	$c_1 = -0.6941$, $c_2 = 1.3777$, $c_3 = 0.0652$, $c_4 = -0.0627$, $c_5 = 0.0206$, $c_6 = -0.1355$,	[27]
Masses and decay widths (in GeV)	$m_{D^\pm} = 1.86961$, $\tau_{D^\pm} = 1.040 \times 10^{-12} s$ $BR(D^+ \rightarrow \pi^+ \pi^- \pi^+) = 3.18 \times 10^{-3}$ $m_{\rho^0(770)} = 0.775$, $\Gamma_{\rho^0(770)} = 0.150$, $m_{f_0(500)} = 0.5$, $\Gamma_{f_0(500)} = 0.5$, $m_\pi = 0.13957$, $m_u = 0.0023$, $m_d = 0.0048$, $m_s = 0.095$, $m_c = 1.275$,	[32]
Form factors	$F_1^{D \rightarrow \pi}(0) = 0.67$, $A_0^{D \rightarrow \rho}(0) = 0.64$, $a^\pi = 1.19$, $b^\pi = 0.36$ $a^\rho = 1.07$, $b^\rho = 0.54$	[30]
	$F_0^{F \rightarrow f_0}(m_\pi^2) = 0.33$,	—
Decay constants (in GeV)	$f_\pi = 0.13041$, $f_K = 0.1562$,	[32]
	$f_\rho = 0.216$,	[33]
Wolfenstein parameters of CKM matrix	$\lambda = 0.22548^{+0.00068}_{-0.00034}$, $A = 0.810^{+0.018}_{-0.024}$, $\bar{\rho} = 0.145^{+0.013}_{-0.007}$, $\bar{\eta} = 0.343^{+0.011}_{-0.012}$,	[34]

where F and X can be $F_1^{D \rightarrow \pi}$ and π , or $A_0^{D \rightarrow \rho}$ and ρ , respectively. The form factor $F^{D \rightarrow f_0}(m_\pi^2)$ which we use here is a rough estimation, and is consistent with branching ratio of $D^+ \rightarrow f_0(500)\pi^+$ extracted from Dalitz analysis of Data [31].

As is observed by LHCb, the CP asymmetries around the vicinities of $f_0(500)$ and $\rho^0(770)$ have opposite signs for small and large values of s_{high} in the case of B meson decay channel $B^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ [14]. In view of the above, for the case of $D^\pm \rightarrow \pi^\pm \pi^+ \pi^-$, we will focus on CP asymmetries of two regions, denoted as Ω^+ and Ω^- , where Ω^+ (Ω^-) represents phase space satisfying $s_{\text{high}} > (<) \Sigma_{s_{\text{low}}}$, and $s_{\text{high}} > m_\rho^2$. The CP asymmetry difference of the aforementioned two regions is

$$\Delta A_{CP} = A_{CP}^{\Omega^+} - A_{CP}^{\Omega^-}. \quad (15)$$

In Fig. 1, the CP asymmetries $A_{CP}^{\Omega^+}$, $A_{CP}^{\Omega^-}$ and their difference ΔA_{CP} are shown as a function of the strong phase δ , where the strong phase δ is assumed as a constant with respect to s_{low} and s_{high} . It can be seen from Fig. 1 that ΔA_{CP} is negative(positive) when δ is around 0 (π). The magnitude of ΔA_{CP} can reach as large as 0.5×10^{-4} for some values of δ . Especially, Fig. 1 shows that our mechanism indicates possibilities for ΔA_{CP} being zero. This is interesting because that the experimental result from LHCb collaboration shows no CP violation in this channel. One can read off two zero points for ΔA_{CP} from Fig. 1, which are $\delta_1 = 4.50$ and $\delta_2 = 1.06$.

Figures 2 and 3 present in the phase space for $\delta = 4.50$ and $\delta = 1.06$, respectively, the relative differential decay width γ of $D^+ \rightarrow \pi^+ \pi^- \pi^+$, which is defined as

$$\gamma(s_{\text{low}}, s_{\text{high}}) \equiv \frac{1}{\Gamma_{D^+ \rightarrow \pi^+ \pi^- \pi^+}} \cdot \frac{d^2 \Gamma_{D^+ \rightarrow \pi^+ \pi^- \pi^+}}{ds_{\text{low}} ds_{\text{high}}} = \frac{1}{256\pi^3 m_D^3 \Gamma_{D^+ \rightarrow \pi^+ \pi^- \pi^+}} |\mathcal{M}|^2. \quad (16)$$

For comparison, we also present the relative differential decay width in Fig. 4 for $\delta = 0$, and that in Fig. 5 for the situation only resonance $\rho^0(770)$ is taken into account. Experimental data from LHCb shows that symmetries of event distribution around the $\rho^0(770)$ resonance are badly destroyed. The number of events around the $\rho^0(770)$ resonance for $s_{\text{high}} < \Sigma_{s_{\text{low}}}$ are much larger than that for $s_{\text{high}} > \Sigma_{s_{\text{low}}}$, as is shown in Ref. [25]. Besides, LHCb results also shows an enhancement of event distributions in the region of phase space where $\sqrt{s_{\text{low}}}$ and $\sqrt{s_{\text{high}}}$ are around the masses of $f_0(500)$ and $\rho^0(770)$, respectively. These behaviours are roughly the same as those shown in Fig. 2, which indicate that our mechanism is consistent with experimental data when $\delta = 4.50$.

4 Discussion

We used a naive factorization approach for the weak decay processes in the calculation of the decay amplitudes. The reason is simply because that large part of the region in phase space that we focused on is off the mass shells of $\rho^0(770)$ and $f_0(500)$, the advantages of factorization approaches such as perturbative QCD [35], QCD factorization [36], Soft Collinear Effective Theory [37] are smeared out by the off shell effect.

In determining the strong phase δ , we used the CP asymmetry difference of two regions of phase space instead of the differential CP asymmetry. The reason is because that the use of the differential CP asymmetry as a tool to determine the strong phase δ is not an

appropriate approach at all. On one hand, if use the differential CP asymmetry, one would find that no strong phase can accommodate that with the data. On the other hand, if one check the nonzero differential CP asymmetries distributed in phase space for $\delta = 4.50$, one would find that the large differential CP asymmetries goes always with very small differential decay amplitude \mathcal{M} , indicating a cancellation between $\mathcal{M}_\rho e^{i\delta}$ and \mathcal{M}_{f_0} . In this situation, the dominance of these two amplitudes is no longer valid. Consequently, in order to deduce the differential CP asymmetries in this kind of regions, we should in principle consider other contributions to the decay amplitude \mathcal{M} in these phase space regions, which is out of the scope of this paper.

We choose the right boundary of the two regions Ω^+ and Ω^- in phase space to be m_ρ^2 . Although it is not an unique choice, the boundary should not be far away from the vicinity of $\rho^0(770)$, in which case, the allowed strong phase δ is not sensitive to the choice. On the other hand, either it is too large or too small than m_ρ^2 , the dominance of two resonance ρ^0 and $f_0(500)$ of the total decay amplitude is no longer valid.

Other resonances can also contribute to the decay amplitude. For resonance such as $f_0(980)$, only a small part of the total resonance lies in the region. As a result, this contribution is small compared with $\rho^0(770)$. Resonance ω can enter in the amplitude through an isospin breaking effect, which is called $\rho^0 - \omega$ mixing mechanism. This mechanism can generate large differential CP asymmetries in the vicinity of Ω . However, the width of ω is small, its contribution to regional CP violation is small. More importantly, the contribution of ω to CP asymmetry is independent of s_{high} , and hence has no contribution to ΔA_{CP} ¹.

From Fig. 1, one can see that the CP asymmetries of the two regions Ω^+ and Ω^- are nonzero for $\delta = 4.50$, which seems to be a disadvantage of our work. However, these two CP asymmetries are small, which both are 1.1×10^{-5} . In principle, these small CP asymmetries are understandable by, for example, an inclusion of s_{low} or s_{high} dependence on δ . What important is, our division of the phase space $f_0(500)$ and $\rho^0(770)$ enlarge the effect of CP violations caused by the interference, consequently, can be used to determine the strong phase.

5 Conclusion

In this paper, we study the localized CP violation and differential decay width of the decay channel $D^\pm \rightarrow \pi^+\pi^-\pi^+$. We focus our attention on phase space where the invariant mass of $\pi^+\pi^-$ are around the vicinities of $\rho^0(770)$ and $f_0(500)$. We consider a mechanism which can generate large CP asymmetries on three-body decays of B meson, that is localized CP asymmetries caused by the interference of amplitudes corresponding to resonances with different spins. We found that by properly choosing a relative strong phase δ , the interference of amplitude corresponding to resonances $f_0(500)$ and $\rho^0(770)$ gives predictions that are consistent with experimental data both on CP asymmetries and differential decay widths.

¹Strictly speaking, since there are two identical particles in the final state for $D^\pm \rightarrow \pi^\pm\pi^+\pi^-$, the terms of amplitudes are doubled by $[s_{\text{low}} \leftrightarrow s_{\text{high}}]$, as is shown in Eq. 9. As a result, the contribution of resonance ω in ΔA_{CP} cannot be cancelled exactly. Besides, the interference between the amplitudes corresponding to $f_0(500)$ and ω is also small due to the smallness of ω 's width.

Our results generate no CP asymmetry differences ($\Delta A_{CP} = 0$) when the strong phase $\delta = 4.50$. In the same time, the behaviour of event distribution around the vicinity of $\rho^0(770)$ and $f_0(500)$ is also understandable.

Acknowledgments

This work was partially supported by National Natural Science Foundation of China (No. 11447021), the construct program of the key discipline in Hunan province, and the Innovation Team of Nuclear and Particle Physics of USC.

References

- [1] J.H. Christenson, J.W. Cronin, V.L. Fitch, and R. Turlay. Evidence for the 2π Decay of the K_2^0 Meson. *Phys. Rev. Lett.*, 13:138–140, 1964.
- [2] Makoto Kobayashi and Toshihide Maskawa. CP Violation in the Renormalizable Theory of Weak Interaction. *Prog. Theor. Phys.*, 49:652–657, 1973.
- [3] Ashton B. Carter and A.I. Sanda. CP Violation in Cascade Decays of B Mesons. *Phys.Rev.Lett.*, 45:952, 1980.
- [4] Ikaros I.Y. Bigi and A.I. Sanda. Notes on the Observability of CP Violations in B Decays. *Nucl.Phys.*, B193:85, 1981.
- [5] Ashton B. Carter and A.I. Sanda. CP Violation in B Meson Decays. *Phys. Rev.*, D23:1567, 1981.
- [6] R. Aaij et al. Observation of CP violation in $B^+ \rightarrow DK^+$ decays. *Phys.Lett.*, B712:203–212, 2012.
- [7] Bernard Aubert et al. Evidence for Direct CP Violation from Dalitz-plot analysis of $B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm$. *Phys.Rev.*, D78:012004, 2008.
- [8] P. del Amo Sanchez et al. Measurements of branching fractions, polarizations, and direct CP -violation asymmetries in $B^+ \rightarrow \rho^0 K^{*+}$ and $B^+ \rightarrow f_0(980) K^{*+}$ decays. *Phys.Rev.*, D83:051101, 2011.
- [9] Bernard Aubert et al. Dalitz Plot Analysis of $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$ Decays. *Phys.Rev.*, D79:072006, 2009.
- [10] J.P. Lees et al. Amplitude Analysis of $B^0 \rightarrow K^+ \pi^- \pi^0$ and Evidence of Direct CP Violation in $B \rightarrow K^* \pi$ decays. *Phys.Rev.*, D83:112010, 2011.
- [11] T. Aaltonen et al. Measurements of Direct CP Violating Asymmetries in Charmless Decays of Strange Bottom Mesons and Bottom Baryons. *Phys.Rev.Lett.*, 106:181802, 2011.

- [12] R Aaij et al. First observation of CP violation in the decays of B_s^0 mesons. *Phys.Rev.Lett.*, 110(22):221601, 2013.
- [13] R Aaij et al. Measurement of CP violation in the phase space of $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ and $B^\pm \rightarrow K^\pm K^+ K^-$ decays. *Phys.Rev.Lett.*, 111:101801, 2013.
- [14] Roel Aaij et al. Measurement of CP violation in the phase space of $B^\pm \rightarrow K^+ K^- \pi^\pm$ and $B^\pm \rightarrow \pi^+ \pi^- \pi^\pm$ decays. *Phys.Rev.Lett.*, 112:011801, 2014.
- [15] Zhen-Hua Zhang, Xin-Heng Guo, and Ya-Dong Yang. CP violation in $B^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ in the region with low invariant mass of one $\pi^+ \pi^-$ pair. *Phys.Rev.*, D87(7):076007, 2013.
- [16] Zhen-Hua Zhang, Xin-Heng Guo, and Ya-Dong Yang. CP violation induced by the interference of scalar and vector resonances in three-body decays of bottom mesons. 2013.
- [17] Bhuvanajyoti Bhattacharya, Michael Gronau, and Jonathan L. Rosner. CP asymmetries in three-body B^\pm decays to charged pions and kaons. *Phys.Lett.*, B726:337–343, 2013.
- [18] I. Bediaga, T. Frederico, and O. Loureno. CP violation and CPT invariance in B^\pm decays with final state interactions. *Phys.Rev.*, D89(9):094013, 2014.
- [19] Joachim Brod, Yuval Grossman, Alexander L. Kagan, and Jure Zupan. A Consistent Picture for Large Penguins in $D \rightarrow \pi^+ \pi^-, K^+ K^-$. *JHEP*, 1210:161, 2012.
- [20] Joachim Brod, Alexander L. Kagan, and Jure Zupan. Size of direct CP violation in singly Cabibbo-suppressed D decays. *Phys.Rev.*, D86:014023, 2012.
- [21] Bhuvanajyoti Bhattacharya, Michael Gronau, and Jonathan L. Rosner. CP asymmetries in singly-Cabibbo-suppressed D decays to two pseudoscalar mesons. *Phys.Rev.*, D85:054014, 2012.
- [22] R Aaij et al. Search for direct CP violation in $D^0 \rightarrow h^- h^+$ modes using semileptonic B decays. *Phys.Lett.*, B723:33–43, 2013.
- [23] T. Aaltonen et al. Measurement of the difference of CP -violating asymmetries in $D^0 \rightarrow K^+ K^-$ and $D^0 \rightarrow \pi^+ \pi^-$ decays at CDF. *Phys.Rev.Lett.*, 109:111801, 2012.
- [24] Roel Aaij et al. Measurement of CP asymmetry in $D^0 \rightarrow K^- K^+$ and $D^0 \rightarrow \pi^- \pi^+$ decays. *JHEP*, 1407:041, 2014.
- [25] R Aaij et al. Search for CP violation in the decay $D^+ \rightarrow \pi^- \pi^+ \pi^+$. *Phys.Lett.*, B728:585–595, 2014.
- [26] Gerhard Buchalla, Andrzej J. Buras, and Markus E. Lautenbacher. Weak decays beyond leading logarithms. *Rev. Mod. Phys.*, 68:1125–1144, 1996.
- [27] Xin-Heng Guo and Anthony William Thomas. Direct CP violation in charmed hadron decays via $\rho - \omega$ mixing. *Phys.Rev.*, D61:116009, 2000.

- [28] Andrzej J. Buras, Markus E. Lautenbacher, and Gaby Ostermaier. Waiting for the top quark mass, $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $B_s^0 - \bar{B}_s^0$ mixing and CP asymmetries in B decays. *Phys.Rev.*, D50:3433–3446, 1994.
- [29] J. Charles et al. CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories. *Eur.Phys.J.*, C41:1–131, 2005.
- [30] Hai-Yang Cheng, Chun-Khiang Chua, and Chien-Wen Hwang. Covariant light front approach for s wave and p wave mesons: Its application to decay constants and form-factors. *Phys. Rev.*, D 69:074025, 2004.
- [31] G. Bonvicini et al. Dalitz plot analysis of the $D^+ \rightarrow \pi^- \pi^+ \pi^+$ decay. *Phys.Rev.*, D76:012001, 2007.
- [32] K.A. Olive et al. Review of Particle Physics. *Chin.Phys.*, C38:090001, 2014.
- [33] Pieter Maris and Peter C. Tandy. Bethe-Salpeter study of vector meson masses and decay constants. *Phys.Rev.*, C60:055214, 1999.
- [34] J. Charles, O. Deschamps, S. Descotes-Genon, H. Lacker, A. Menzel, et al. Current status of the Standard Model CKM fit and constraints on $\Delta F = 2$ New Physics. *Phys.Rev.*, D91(7):073007, 2015.
- [35] Hsiang-nan Li. PQCD factorization of two-body B decays.
- [36] M. Beneke, G. Buchalla, M. Neubert, and Christopher T. Sachrajda. QCD factorization for $B \rightarrow \pi\pi$ decays: Strong phases and CP violation in the heavy quark limit. *Phys. Rev. Lett.*, 83:1914–1917, 1999.
- [37] Christian W. Bauer, Dan Pirjol, and Iain W. Stewart. A proof of factorization for $B \rightarrow D\pi$. *Phys. Rev. Lett.*, 87:201806, 2001.




Figure 1: The CP asymmetry difference ΔA_{CP} (solid line), the CP asymmetry of region Ω^- (dashed line) and Ω^+ (dash-dotted line) as a function of the strong phase δ .




Figure 2: Differential branching ratio (in GeV^{-4}) of $D^+ \rightarrow \pi^+ \pi^- \pi^+$ when $\delta = 4.50$.




Figure 3: Differential branching ratio (in GeV^{-4}) of $D^+ \rightarrow \pi^+\pi^-\pi^+$ when $\delta = 1.06$.




Figure 4: Differential branching ratio (in GeV^{-4}) of $D^+ \rightarrow \pi^+ \pi^- \pi^+$ when $\delta = 0$.




Figure 5: Differential branching ratio (in GeV^{-4}) of $D^+ \rightarrow \pi^+ \pi^- \pi^+$, where only the amplitudes corresponding to $\rho^0(770)$ are included.