

Path Integral for Off-Shell Supersymmetry

Chiu Man Ho^{1,*}

*¹Department of Physics and Astronomy,
Michigan State University, East Lansing, MI 48824, USA*

(Dated: October 13, 2018)

Abstract

Off-shell supersymmetry, which restricts sparticles to appear only off-shell, solves the gauge hierarchy problem and unifies the gauge couplings in the usual way. Without introducing any new interactions or exacerbating the naturalness, this idea has been proposed to evade the direct detection bounds on sparticles. The original realization of off-shell supersymmetry relies on applying a generalized quantization scheme solely to sparticles. In this paper, we formulate off-shell supersymmetry from the path-integral approach.

arXiv:1506.00319v1 [hep-ph] 1 Jun 2015

*Electronic address: cmho@msu.edu

I. INTRODUCTION

Supersymmetry (SUSY) offers an elegant solution to the gauge hierarchy problem. In the Minimal Supersymmetric Standard Model (MSSM), the three gauge couplings are unified at $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV, which is desirable for the Grand Unified Theory (GUT). This makes supersymmetry a promising idea. Since SUSY is most naturally realized at the TeV scale, we had expected to discover sparticles in the first run of the Large Hadron Collider (LHC). However, no signal of sparticles has been observed so far and the hope is now on the LHC Run II with a collider energy 13 – 14 TeV. For an overview of sparticle search at the LHC, see [1, 2]. The null search results for colored sparticles such as the gluino \tilde{g} and squarks \tilde{q} imply lower bounds on their masses with roughly $m_{\tilde{g}} \gtrsim 1$ TeV and $m_{\tilde{q}} \gtrsim 600$ GeV. The lower bounds on non-colored sparticles such as sleptons, charginos and neutralinos are not so severe, with roughly a few hundred GeV.

It is possible that sparticles might have actually been produced at the LHC but have escaped from the detection because their signals are too difficult to be distinguished from the Standard Model (SM) background. This situation occurs when the energy of hadronic/leptonic jets and the missing transverse energy associated with the sparticle cascade decay are significantly reduced in some SUSY models, such as the so-called Compressed SUSY [3], Stealth SUSY [4] and R -parity violating SUSY models [5]. For these models, the lower bounds on sparticle masses can be relaxed up to a few hundred GeV. However, it is not easy to construct such models in a natural way [6, 7]. These models cannot completely hide the direct signal of sparticles forever. With the substantially improved sensitivity at the LHC Run II, it is conceivable that they will be probed and constrained soon.

Contrary to the SUSY models which can tentatively hide the sparticles from direct detection, we would like to provide an R -parity preserving MSSM scenario that can survive even if sparticles may never be directly observed at the LHC. Our guiding principle is to retain simplicity and naturalness as much as possible.

Supersymmetry at the Lagrangian does not dictate the way we should quantize the fields in the supermultiplets, and so we have the freedom to quantize the SM particles and their superpartners differently. In [8], we proposed a generalized quantization scheme under which a particle can only appear off-shell, while its contributions to quantum corrections behave exactly the same as in the usual quantum field theory (QFT). We applied this quantization

scheme solely to the sparticles in the R -parity preserving MSSM. Thus sparticles can only appear off-shell.¹ Without introducing any new interactions or exacerbating the naturalness, the sparticles could be light but would completely escape the direct detection at any experiments such as the LHC. However, our theory still retains the same desirable features of the usual MSSM at the quantum level. For instance, the gauge hierarchy problem is solved and the three MSSM gauge couplings are unified in the usual way. Although direct detection of sparticles is impossible, their existence can be revealed by precise measurements of some observables (such as the running QCD coupling) that may receive quantum corrections from them and have sizable deviations from the SM predictions. The experimental constraints from the indirect sparticle search are still applicable and the same as in the usual MSSM.

Of course, the lightest sparticle (e.g. neutralino) would no longer be a viable dark matter candidate if it can only appear off-shell. However, this should not be considered as a serious deficiency of our MSSM scenario. Providing a viable dark matter candidate is a just bonus of the usual MSSM, and it is easy enough to construct other models that give a promising dark matter candidate. Supersymmetry is most crucial for solving the gauge hierarchy problem and unification of the three MSSM gauge couplings.

In principle, one could apply our generalized quantization scheme to any other theories so as to evade the corresponding bounds from direct detection. Nevertheless, sparticles are particularly well-suited for the generalized quantization. The reason is that the most important merit of sparticles is due to their off-shell quantum contributions. For instance, we do not need on-shell sparticles in order to solve the gauge hierarchy problem and unify the three MSSM gauge couplings.

The path-integral approach is perhaps the most consistent and convenient way to quantize the non-abelian gauge fields. It also facilitates the study of non-perturbative effects in quantum field theory. A consistent formulation from the path-integral approach makes a theory more promising. The original realization of off-shell supersymmetry has relied on applying the generalized quantization scheme solely to sparticles. In order to better justify its consistency, we formulate off-shell supersymmetry from the path-integral approach in this paper.

¹ After [8] appeared on arXiv, we were informed that this possibility was wished in [9].

II. PATH-INTEGRAL QUANTIZATION IN QUANTUM FIELD THEORY

In this section, we briefly review the path-integral quantization for scalars and fermions in the usual QFT. We refer the readers to [10] for more details. In the next section, we will demonstrate how should these path-integrals be modified in order to restrict scalars and fermions to appear only off-shell.

For a non-interacting real scalar ϕ with mass m , the generating functional in the presence of a source J is given by

$$Z[J] = \int \mathcal{D}\phi \exp \left\{ i \int d^4x (\mathcal{L}(\phi) + J(x) \phi(x)) \right\}, \quad (2.1)$$

where $\mathcal{L}(\phi)$ is the free Lagrangian for the ϕ field. $Z[J]$ can be evaluated and normalized to give

$$Z_0[J] = \exp \left(-\frac{i}{2} \int d^4x d^4y J(x) \Delta(x-y) J(y) \right). \quad (2.2)$$

The Green's function $\Delta(x-y)$ obeys the inhomogeneous equation $(\square^2 + m^2 - i\epsilon) \Delta(x-y) = -\delta^{(4)}(x-y)$ and it reads

$$\Delta(x-y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{1}{p^2 - m^2 + i\epsilon}. \quad (2.3)$$

The n -point correlation function can be obtained from

$$\langle 0|T \{ \phi(x_1) \cdots \phi(x_n) \} |0\rangle = \frac{1}{i^n} \frac{\delta^n Z_0[J]}{\delta J(x_1) \cdots \delta J(x_n)} \Big|_{J=0}. \quad (2.4)$$

For a non-interacting complex scalar, $\mathcal{D}\phi$ in Eq. (2.1) is replaced by $\mathcal{D}\phi^\dagger \mathcal{D}\phi$ and $J(x) \phi(x)$ by $J^\dagger(x) \phi(x) + \phi^\dagger(x) J(x)$. In Eq. (2.2), $J(x)$ is replaced by $J^\dagger(x)$ and the overall factor of 1/2 is absent. The $2n$ -point correlation function is

$$\langle 0|T \{ \phi(x_1) \phi^\dagger(x_2) \cdots \phi(x_{2n-1}) \phi^\dagger(x_{2n}) \} |0\rangle = \frac{1}{i^{2n}} \frac{\delta^{2n} Z_0[J, J^\dagger]}{\delta J^\dagger(x_1) J(x_2) \cdots J^\dagger(x_{2n-1}) \delta J(x_{2n})} \Big|_{J=J^\dagger=0}. \quad (2.5)$$

For a non-interacting fermion ψ with mass m , the generating functional in the presence of Grassmannian sources η and $\bar{\eta}$ is given by

$$Z[\eta, \bar{\eta}] = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left\{ i \int d^4x (\mathcal{L}(\bar{\psi}, \psi) + \bar{\eta}(x) \psi(x) + \bar{\psi}(x) \eta(x)) \right\}, \quad (2.6)$$

where $\mathcal{L}(\bar{\psi}, \psi)$ is the free Lagrangian for the ψ field. $Z[\eta, \bar{\eta}]$ can be evaluated and normalized to give

$$Z_0[\eta, \bar{\eta}] = \exp\left(-i \int d^4x d^4y \bar{\eta}(x) \mathcal{S}(x-y) \eta(y)\right). \quad (2.7)$$

The Green's function $\mathcal{S}(x-y)$ obeys the inhomogeneous equation $(\square^2 + m^2 - i\epsilon) \mathcal{S}(x-y) = -\delta^{(4)}(x-y)$ and it reads

$$\mathcal{S}(x-y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{\not{p} + m}{p^2 - m^2 + i\epsilon}. \quad (2.8)$$

The $2n$ -point correlation function can be obtained from

$$\langle 0|T\{\psi(x_1)\bar{\psi}(x_2)\cdots\psi(x_{2n-1})\bar{\psi}(x_{2n})\}|0\rangle = \frac{\delta^{2n} Z_0[\eta, \bar{\eta}]}{\delta\bar{\eta}(x_1)\delta\eta(x_2)\cdots\delta\bar{\eta}(x_{2n-1})\delta\eta(x_{2n})}\Bigg|_{\eta=\bar{\eta}=0}. \quad (2.9)$$

When interactions are present, the most general generating functional involving a scalar ϕ , a fermion ψ , an abelian vector boson A_μ and a nonabelian vector boson A_ν^a is given by

$$Z[J, J^\dagger, \eta, \bar{\eta}, J_\mu, J_\nu^a] = \exp\left(i \int d^4x \mathcal{L}_I\right) Z_0, \quad (2.10)$$

where Z_0 is the product of the normalized generating functionals for the scalar, fermion, abelian and nonabelian vector bosons. \mathcal{L}_I is the Lagrangian that contains the interaction operators under consideration, but with the fields for the scalar, fermion, abelian and non-abelian vector bosons substituted by:

$$\phi(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J(x)}, \quad \text{for real } \phi \quad (2.11)$$

$$\phi(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J^\dagger(x)}, \quad \text{for complex } \phi \quad (2.12)$$

$$\phi^\dagger(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J(x)}, \quad \text{for complex } \phi \quad (2.13)$$

$$\psi(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta \bar{\eta}}, \quad \bar{\psi}(x) \rightarrow -\frac{1}{i} \frac{\delta}{\delta \eta}, \quad (2.14)$$

$$A_\mu(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J_\mu}, \quad A_\mu^a(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J_\mu^a}. \quad (2.15)$$

III. PATH-INTEGRAL APPROACH TO OFF-SHELL SUPERSYMMETRY

In [8], we proposed a generalized quantization scheme which restricts particles to appear only off-shell. We applied this quantization scheme solely to the sparticles in the R -parity

preserving MSSM. Thus sparticles can only appear off-shell. (In the MSSM, we do not have spin-1 sparticles, so the application of our generalized quantization scheme to vector bosons is not relevant.) Under the generalized quantization scheme, we derived the propagators for scalars and fermions [8]. The propagator for scalars is given by

$$\langle 0|T\{\phi(x)\phi(y)\}|0\rangle = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{1}{2} \left(\frac{i}{p^2 - m^2 + i\epsilon} + \frac{i}{p^2 - m^2 - i\epsilon} \right), \quad (3.1)$$

while that for Dirac fermions is given by

$$\begin{aligned} \langle 0|T\{\psi_\alpha(x)\bar{\psi}_\beta(y)\}|0\rangle &\equiv i\mathcal{S}_{\alpha\beta}(x-y) \\ &= \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} (\not{p} + m)_{\alpha\beta} \frac{1}{2} \left(\frac{i}{p^2 - m^2 + i\epsilon} + \frac{i}{p^2 - m^2 - i\epsilon} \right). \end{aligned} \quad (3.2)$$

For Majorana fermions with $\psi = \psi^c \equiv C\bar{\psi}^T$ where C is the charge conjugation matrix, there are two additional propagators:

$$\langle 0|T\{\psi_\alpha(x)\psi_\beta(y)\}|0\rangle = [C^{-1}\mathcal{S}(x-y)]_{\alpha\beta}, \quad (3.3)$$

$$\langle 0|T\{\bar{\psi}_\alpha(x)\bar{\psi}_\beta(y)\}|0\rangle = [-\mathcal{S}(x-y)C]_{\alpha\beta}. \quad (3.4)$$

As shown in [8], the propagators for both scalars and fermions turn out to be half-retarded and half-advanced. Under the generalized quantization scheme, a particle simultaneously propagates positive and negative energy modes both forward and backward in time with equal amplitudes. Therefore, the particle behaves like a “standing wave” (in time) — there is *no* net energy flux being transferred. Since an on-shell particle carries a nonzero net energy flux, a particle propagated by the half-retarded and half-advanced propagator *cannot* be on-shell.

In this section, we would like to demonstrate how should the usual path-integrals for scalars and fermions be modified in order to restrict them to appear only off-shell. Under the generalized quantization scheme proposed in [8], a scalar or fermionic field, *as a single field*, acquires both of the propagators $\frac{1}{p^2 - m^2 + i\epsilon}$ and $\frac{1}{p^2 - m^2 - i\epsilon}$. This suggests that in the path-integral formulation, a single J or a single pair of η and $\bar{\eta}$ is not sufficient. We need separate (Grassmannian) sources, which can account for each of $\frac{1}{p^2 - m^2 + i\epsilon}$ and $\frac{1}{p^2 - m^2 - i\epsilon}$, for the same field. A natural way of doing this is to introduce $\{J_+, J_-\}$ for scalars while $\{\eta_+, \bar{\eta}_+\}$ and $\{\eta_-, \bar{\eta}_-\}$ for fermions. The (Grassmannian) sources with subscripts + and - are required to generate $\frac{1}{p^2 - m^2 + i\epsilon}$ and $\frac{1}{p^2 - m^2 - i\epsilon}$ respectively. We will carry out this strategy in what follows.

A. Scalars

For a non-interacting real scalar ϕ with mass m , we propose that the generating functional in the presence of sources $\{J_+, J_-\}$ is given by

$$Z[J_+, J_-] = \int \mathcal{D}\phi \exp \left\{ i \int d^4x \left(\mathcal{L}(\phi) + \frac{1}{2} J_+(x) \phi(x) + \frac{1}{2} J_-(x) \phi(x) \right) \right\} \quad (3.5)$$

$$= \int \mathcal{D}\phi \exp \left\{ \frac{i}{2} \int d^4x \left(\mathcal{L}(\phi) + J_+(x) \phi(x) + \frac{i}{2} \epsilon \phi^2(x) \right) \right\} \\ \exp \left\{ \frac{i}{2} \int d^4x \left(\mathcal{L}(\phi) + J_-(x) \phi(x) - \frac{i}{2} \epsilon \phi^2(x) \right) \right\}, \quad (3.6)$$

with $\epsilon \rightarrow 0^+$. At first glance, it is not obvious that the path integral above is convergent as the factors $+\frac{i}{2} \epsilon \phi^2(x)$ and $-\frac{i}{2} \epsilon \phi^2(x)$ cancel each other. One way to justify the convergence is to think of these two factors as $+\frac{i}{2} \epsilon_+ \phi^2(x)$ and $-\frac{i}{2} \epsilon_- \phi^2(x)$ with $\epsilon_{\pm} \rightarrow 0^+$. The convergence of the path integral can then be guaranteed by assuming $\epsilon_+ - \epsilon_- \rightarrow 0^+$. For simplicity, we will set $\epsilon_+ = \epsilon_- = \epsilon$ from now on.

Except from an extra overall factor of $1/2$, the first term in Eq. (3.6) is the usual path integral for a non-interacting real scalar ϕ . For the second term in Eq. (3.6), one simply needs to replace $m^2 - i\epsilon$ by $m^2 + i\epsilon$ in the usual path integral for a non-interacting real scalar ϕ (apart from the extra overall factor of $1/2$). Thus, following the similar derivations as in the usual path-integral formulation, the normalized generating functional will be given by

$$Z_0[J_+, J_-] = Z_0[J_+] Z_0[J_-], \quad (3.7)$$

where

$$Z_0[J_+] = \exp \left(-\frac{i}{4} \int d^4x d^4y J_+(x) \Delta_+(x-y) J_+(y) \right), \quad (3.8)$$

$$Z_0[J_-] = \exp \left(-\frac{i}{4} \int d^4x d^4y J_-(x) \Delta_-(x-y) J_-(y) \right). \quad (3.9)$$

The Green's functions $\Delta_+(x-y)$ and $\Delta_-(x-y)$:

$$\Delta_+(x-y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{1}{p^2 - m^2 + i\epsilon}, \quad (3.10)$$

$$\Delta_-(x-y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{1}{p^2 - m^2 - i\epsilon}, \quad (3.11)$$

obey the inhomogeneous equations $(\square^2 + m^2 - i\epsilon) \Delta_+(x-y) = -\delta^{(4)}(x-y)$ and $(\square^2 + m^2 + i\epsilon) \Delta_-(x-y) = -\delta^{(4)}(x-y)$ respectively.

The two-point correlation function can be obtained from

$$\begin{aligned} \langle 0|T\{\phi(x)\phi(y)\}|0\rangle &= \frac{1}{i} \left(\frac{\delta}{\delta J_+(x)} + \frac{\delta}{\delta J_-(x)} \right) \frac{1}{i} \left(\frac{\delta}{\delta J_+(y)} + \frac{\delta}{\delta J_-(y)} \right) Z_0[J_+, J_-] \Big|_{J_{\pm}=0} \quad (3.12) \\ &= \frac{1}{2} [i\Delta_+(x-y) + i\Delta_-(x-y)], \end{aligned}$$

which is consistent with Eq. (3.1). This can be generalized to the n -point correlation function:

$$\begin{aligned} &\langle 0|T\{\phi(x_1)\cdots\phi(x_n)\}|0\rangle \\ &= \frac{1}{i^n} \left(\frac{\delta}{\delta J_+(x_1)} + \frac{\delta}{\delta J_-(x_1)} \right) \cdots \left(\frac{\delta}{\delta J_+(x_n)} + \frac{\delta}{\delta J_-(x_n)} \right) Z_0[J_+, J_-] \Big|_{J_{\pm}=0}. \quad (3.13) \end{aligned}$$

For a non-interacting complex scalar, $\mathcal{D}\phi$ in Eq. (3.5) is replaced by $\mathcal{D}\phi^\dagger\mathcal{D}\phi$ and $J_{\pm}(x)\phi(x)$ by $J_{\pm}^\dagger(x)\phi(x) + \phi^\dagger(x)J_{\pm}(x)$. In Eq. (3.6), $\pm\frac{i}{2}\epsilon\phi^2(x)$ is replaced by $\pm i\epsilon\phi^\dagger(x)\phi(x)$. In Eq. (3.8) and Eq. (3.9), $J_{\pm}(x)$ is replaced by $J_{\pm}^\dagger(x)$ and the overall factor is $-i/2$ instead of $-i/4$. The $2n$ -point correlation function is

$$\begin{aligned} &\langle 0|T\{\phi(x_1)\phi^\dagger(x_2)\cdots\phi(x_{2n-1})\phi^\dagger(x_{2n})\}|0\rangle \\ &= \frac{1}{i^{2n}} \left(\frac{\delta}{\delta J_+^\dagger(x_1)} + \frac{\delta}{\delta J_-^\dagger(x_1)} \right) \left(\frac{\delta}{\delta J_+(x_2)} + \frac{\delta}{\delta J_-(x_2)} \right) \cdots \\ &\quad \left(\frac{\delta}{\delta J_+^\dagger(x_{2n-1})} + \frac{\delta}{\delta J_-^\dagger(x_{2n-1})} \right) \left(\frac{\delta}{\delta J_+(x_{2n})} + \frac{\delta}{\delta J_-(x_{2n})} \right) Z_0[J_+, J_+^\dagger, J_-, J_-^\dagger] \Big|_{J_{\pm}=J_{\pm}^\dagger=0}. \quad (3.14) \end{aligned}$$

B. Fermions

For a non-interacting fermion ψ with mass m , we propose that the generating functional in the presence of sources $\{\eta_+, \bar{\eta}_+\}$ and $\{\eta_-, \bar{\eta}_-\}$ is given by

$$\begin{aligned} &Z[\eta_+, \bar{\eta}_+, \eta_-, \bar{\eta}_-] \\ &= \int \mathcal{D}\bar{\psi}\mathcal{D}\psi \exp \left\{ i \int d^4x \left[\mathcal{L}(\bar{\psi}, \psi) + \frac{1}{2} (\bar{\eta}_+(x)\psi(x) + \bar{\psi}(x)\eta_+(x)) \right. \right. \\ &\quad \left. \left. + \frac{1}{2} (\bar{\eta}_-(x)\psi(x) + \bar{\psi}(x)\eta_-(x)) \right] \right\} \quad (3.15) \end{aligned}$$

$$\begin{aligned} &= \int \mathcal{D}\bar{\psi}\mathcal{D}\psi \exp \left\{ \frac{i}{2} \int d^4x (\mathcal{L}(\bar{\psi}, \psi) + \bar{\eta}_+(x)\psi(x) + \bar{\psi}(x)\eta_+(x) + i\epsilon\bar{\psi}\psi) \right\} \\ &\quad \exp \left\{ \frac{i}{2} \int d^4x (\mathcal{L}(\bar{\psi}, \psi) + \bar{\eta}_-(x)\psi(x) + \bar{\psi}(x)\eta_-(x) - i\epsilon\bar{\psi}\psi) \right\}, \quad (3.16) \end{aligned}$$

with $\epsilon \rightarrow 0^+$. Similar to the scalar case, it is not obvious that the path integral above is convergent as the factors $+i\epsilon\bar{\psi}\psi$ and $-i\epsilon\bar{\psi}\psi$ cancel each other. One way to justify the

convergence is to think of these two factors as $+i\epsilon_+ \bar{\psi}\psi$ and $-i\epsilon_- \bar{\psi}\psi$ with $\epsilon_{\pm} \rightarrow 0^+$. The convergence of the path integral can then be guaranteed by assuming $\epsilon_+ - \epsilon_- \rightarrow 0^+$. For simplicity, we will set $\epsilon_+ = \epsilon_- = \epsilon$ from now on.

Except from an extra overall factor of $1/2$, the first term in Eq. (3.16) is the usual path integral for a non-interacting fermion ψ . For the second term in Eq. (3.16), one simply needs to replace $m^2 - i\epsilon$ by $m^2 + i\epsilon$ in the usual path integral for a non-interacting fermion ψ (apart from the extra overall factor of $1/2$). Thus, following the similar derivations as in the usual path-integral formulation, the normalized generating functional will be given by

$$Z_0[\eta_+, \bar{\eta}_+, \eta_-, \bar{\eta}_-] = Z_0[\eta_+, \bar{\eta}_+] Z_0[\eta_-, \bar{\eta}_-], \quad (3.17)$$

where

$$\begin{aligned} Z_0[\eta_+, \bar{\eta}_+] &= \exp\left(-\frac{i}{2} \int d^4x d^4y \bar{\eta}_+(x) \mathcal{S}_+(x-y) \eta_+(y)\right), \\ Z_0[\eta_-, \bar{\eta}_-] &= \exp\left(-\frac{i}{2} \int d^4x d^4y \bar{\eta}_-(x) \mathcal{S}_-(x-y) \eta_-(y)\right). \end{aligned} \quad (3.18)$$

The Green's functions $\mathcal{S}_+(x-y)$ and $\mathcal{S}_-(x-y)$

$$\begin{aligned} \mathcal{S}_+(x-y) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{\not{p} + m}{p^2 - m^2 + i\epsilon}, \\ \mathcal{S}_-(x-y) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{\not{p} + m}{p^2 - m^2 - i\epsilon}. \end{aligned} \quad (3.19)$$

obey the inhomogeneous equations $(\square^2 + m^2 - i\epsilon) \mathcal{S}_+(x-y) = -\delta^{(4)}(x-y)$ and $(\square^2 + m^2 + i\epsilon) \mathcal{S}_-(x-y) = -\delta^{(4)}(x-y)$ respectively.

The two-point correlation function can be obtained from

$$\begin{aligned} &\langle 0|T\{\psi(x)\bar{\psi}(y)\}|0\rangle \\ &= \frac{1}{i} \left(\frac{\delta}{\delta\bar{\eta}_+(x)} + \frac{\delta}{\delta\bar{\eta}_-(x)} \right) \left(-\frac{1}{i} \right) \left(\frac{\delta}{\delta\eta_+(y)} + \frac{\delta}{\delta\eta_-(y)} \right) Z_0[\eta_+, \bar{\eta}_+, \eta_-, \bar{\eta}_-] \Bigg|_{\eta_{\pm}=\bar{\eta}_{\pm}=0} \end{aligned} \quad (3.20)$$

$$= \frac{1}{2} [i\mathcal{S}_+(x-y) + i\mathcal{S}_-(x-y)], \quad (3.21)$$

which is consistent with Eq. (3.2). This can be generalized to the $2n$ -point correlation function:

$$\begin{aligned} &\langle 0|T\{\psi(x_1)\bar{\psi}(x_2)\cdots\psi(x_{2n-1})\bar{\psi}(x_{2n})\}|0\rangle \\ &= \left(\frac{\delta}{\delta\bar{\eta}_+(x_1)} + \frac{\delta}{\delta\bar{\eta}_-(x_1)} \right) \left(\frac{\delta}{\delta\eta_+(x_2)} + \frac{\delta}{\delta\eta_-(x_2)} \right) \cdots \\ &\quad \left(\frac{\delta}{\delta\bar{\eta}_+(x_{2n-1})} + \frac{\delta}{\delta\bar{\eta}_-(x_{2n-1})} \right) \left(\frac{\delta}{\delta\eta_+(x_{2n})} + \frac{\delta}{\delta\eta_-(x_{2n})} \right) Z_0[\eta_+, \bar{\eta}_+, \eta_-, \bar{\eta}_-] \Bigg|_{\eta_{\pm}=\bar{\eta}_{\pm}=0}. \end{aligned} \quad (3.22)$$

C. Interacting Theory

When interactions are present, the most general generating functional involving a scalar ϕ , a fermion ψ , an abelian vector boson A_μ and a nonabelian vector boson A_ν^a is given by

$$Z[J_+, J_+^\dagger, J_-, J_-^\dagger, \eta_+, \bar{\eta}_+, \eta_-, \bar{\eta}_-, J_\mu, J_\nu^a] = \exp\left(i \int d^4x \tilde{\mathcal{L}}_I\right) \tilde{Z}_0, \quad (3.23)$$

where \tilde{Z}_0 is the product of the normalized generating functionals for the scalar, fermion, abelian and nonabelian vector bosons. $\tilde{\mathcal{L}}_I$ is the Lagrangian that contains the supersymmetric interaction operators under consideration, but with the fields for the scalar, fermion, abelian and nonabelian vector bosons substituted by:

$$\phi(x) \rightarrow \frac{1}{i} \left(\frac{\delta}{\delta J_+(x)} + \frac{\delta}{\delta J_-(x)} \right), \quad \text{for real } \phi \quad (3.24)$$

$$\phi(x) \rightarrow \frac{1}{i} \left(\frac{\delta}{\delta J_+^\dagger(x)} + \frac{\delta}{\delta J_-^\dagger(x)} \right), \quad \text{for complex } \phi \quad (3.25)$$

$$\phi^\dagger(x) \rightarrow \frac{1}{i} \left(\frac{\delta}{\delta J_+(x)} + \frac{\delta}{\delta J_-(x)} \right), \quad \text{for complex } \phi \quad (3.26)$$

$$\psi(x) \rightarrow \frac{1}{i} \left(\frac{\delta}{\delta \bar{\eta}_+(x)} + \frac{\delta}{\delta \bar{\eta}_-(x)} \right), \quad (3.27)$$

$$\bar{\psi}(x) \rightarrow -\frac{1}{i} \left(\frac{\delta}{\delta \eta_+(x)} + \frac{\delta}{\delta \eta_-(x)} \right), \quad (3.28)$$

$$A_\mu(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J^\mu}, \quad A_\mu^a(x) \rightarrow \frac{1}{i} \frac{\delta}{\delta J_\mu^a}. \quad (3.29)$$

IV. CONCLUSIONS

The idea of off-shell supersymmetry evades the direct detection bounds on sparticles. On the other hand, it solves the gauge hierarchy problem and unifies the gauge couplings in the usual way. This idea fundamentally changes our understanding about supersymmetry. It explains why we have not directly observed the sparticles at the LHC so far and suggests a new way of searching for them.

The original realization of off-shell supersymmetry relies on a generalization of the way we quantize the quantum fields and its sole application to sparticles. A natural way to reinforce its consistency is through the path-integral approach. In this paper, we show that off-shell supersymmetry can be formulated from the path-integral approach.

Acknowledgments. We thank Pierre Ramond for useful conversations. CMH was supported by the Office of the Vice-President for Research and Graduate Studies at Michigan State University.

- [1] **ATLAS** Collaboration, <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>
- [2] **CMS** Collaboration, <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>
- [3] T. J. LeCompte and S. P. Martin, *Phys. Rev. D* **84**, 015004 (2011); T. J. LeCompte and S. P. Martin, *Phys. Rev. D* **85**, 035023 (2012).
- [4] J. Fan, M. Reece and J. T. Ruderman, *JHEP* **1111**, 012 (2011); J. Fan, M. Reece and J. T. Ruderman, *JHEP* **1207**, 196 (2012).
- [5] R. Barbier, C. Berat, M. Besancon, M. Chemtob, A. Deandrea, E. Dudas, P. Fayet and S. Lavignac *et al.*, *Phys. Rept.* **420**, 1 (2005).
- [6] J. L. Feng, *Ann. Rev. Nucl. Part. Sci.* **63**, 351 (2013).
- [7] N. Craig, arXiv:1309.0528 [hep-ph].
- [8] C. M. Ho and N. Okada, arXiv:1412.2734 [hep-ph].
- [9] D. M. Ghilencea, *Nucl. Phys. B* **876**, 16 (2013).
- [10] L. H. Ryder, *Quantum Field Theory*, Cambridge University Press, Cambridge, United Kingdom, 1985; M. E. Peskin and D. V. Schroeder, *An Introduction To Quantum Field Theory*, Westview Press, Boulder, Colorado, 1995.