Journal of the Korean Astronomical Society
00: 1 ∼ 4, 2015 June
The Korean Astronomical Society (2015)

preprint - no DOI assigned
pISSN: 1225-4614 · eISSN: 2288-890X
http://jkas.kas.org

arXiv:1506.00730v1 [astro-ph.CO] 2 Jun 2015

Milgrom's Law and Λ's Shadow: How Massive Gravity Connects Galactic and Cosmic Dynamics

Sascha Trippe
Department of Physics and Astronomy, Seoul National University, Seoul 151-742, Korea; trippe@astro.snu.ac.kr

Received March 11, 2015; accepted June 2, 2015

Abstract: Massive gravity provides a natural solution for the dark energy problem of cosmology and is also a candidate for resolving the dark matter problem. I demonstrate that, assuming reasonable scaling relations, massive gravity can provide for Milgrom's law of gravity (or "modified Newtonian dynamics") which is known to remove the need for particle dark matter from galactic dynamics. Milgrom's law comes with a characteristic acceleration, Milgrom's constant, which is observationally constrained to $a_0 \approx 1.1 \times 10^{-10}\,\mathrm{m\,s^{-2}}$. In the derivation presented here, this constant arises naturally from the cosmologically required mass of gravitons like $a_0 \propto c\sqrt{\Lambda} \propto cH_0\sqrt{3\Omega_\Lambda}$, with Λ, H_0, and Ω_Λ being the cosmological constant, the Hubble constant, and the third cosmological parameter, respectively. My derivation suggests that massive gravity could be the mechanism behind both, dark matter and dark energy.

Key words: gravitation — cosmology — dark matter — dark energy

1. Introduction

Modern standard cosmology suffers from two critical issues: the *dark matter* problem and the *dark energy* problem. Canonical ΛCDM cosmology (e.g., Bahcall et al. 1999) requires about 95% (e.g., Ade et al. 2014) of the mass/energy content of the universe to be provided by exotic dark components for which physical counterparts have not been identified. Dark matter and dark energy are commonly assumed to be unrelated: dark matter is identified with as yet undiscovered new exotic elementary particles, whereas dark energy is linked to a *cosmological constant*, Λ, which is inserted into Einstein's field equations and which triggers an accelerated expansion of the universe on its largest scales.

ΛCDM cosmology is based on the assumption that gravitation is Einsteinian on all scales. Modified theories of gravity might be able to describe the universe without dark components; this idea has initiated a vast number of works exploring multiple models of gravitation (see Clifton et al. 2012 for an exhaustive review). A quantum-field theoretical modified theory of gravity, first proposed by Fierz & Pauli (1939), is *massive gravity* in which gravitation is mediated by virtual bosons, *gravitons*, that have a very small (compared to all other elementary particles) non-zero mass. Massive gravitons, being virtual exchange particles, have a limited life time governed by Heisenberg's uncertainty relation for energy and time. If the mass of the graviton is such that its life time is $\sqrt{1/\Lambda}$, the decay of gravity induces an accelerated expansion of the universe like the one actually observed – thus providing an elegant resolution of the dark energy problem (recently, e.g., Cardone et al. 2012; Clifton et al. 2012; Hinterbichler 2012; Volkov 2012; Tasinato et al. 2013; De Felice et al. 2013; de Rham 2014).

Corresponding author: S. Trippe

On smaller scales, the dynamics of galaxies is in excellent agreement with a modification of Newtonian dynamics (the *MOND paradigm*) in the limit of small acceleration (or gravitational field strength) g, expressed in *Milgrom's law*

$$g_\mathrm{N} = \mu(g/a_0)\, g \tag{1}$$

(Milgrom 1983a) with g_N being the Newtonian field strength, a_0 denoting *Milgrom's constant*, and μ being a transition function with $\mu = 1$ for $g/a_0 \gg 1$ and $\mu = g/a_0$ for $g/a_0 \ll 1$. Such a modification implies that the ratio of dynamical and luminous masses of a stellar system, the *mass discrepancy* M_dyn/M_0, exceeds unity and becomes a function of acceleration (or field strength) when assuming Newton's law of gravity. In the "deep MOND" limit $g/a_0 \ll 1$, the circular speed v_c of stars in rotation-supported dynamical systems (especially disk galaxies) is

$$v_\mathrm{c}^4 = G\, M_0\, a_0 \tag{2}$$

with G being Newton's constant (Milgrom 1983a,b,c); note that $g = v_\mathrm{c}^2/r = GM_\mathrm{dyn}/r^2$ and $g_\mathrm{N} = GM_0/r^2$ for circular orbits with radius r. For pressure-supported systems (especially elliptical galaxies and galaxy clusters) with 3D velocity dispersion σ, the mass–velocity relation takes the form

$$\sigma^4 = \frac{4}{9}\, G\, M_0\, a_0 \tag{3}$$

(Milgrom 1984, 1994). Equations (1, 2, 3), with $a_0 \approx 1.1 \times 10^{-10}\,\mathrm{m\,s^{-2}}$, naturally provide for the fundamental scaling laws of galactic kinematics, specifically the baryonic Tully–Fisher, baryonic Faber–Jackson, mass discrepancy–acceleration (MDA), and surface mass density–acceleration

relations, plus the asymptotic flattening of rotation curves and the occurrence of "dark rings" in galaxy clusters (Sanders 1994, 2010; Rhee 2004a,b; McGaugh 2004, 2005a,b; Milgrom & Sanders 2008; McGaugh 2011; Gentile et al. 2011; Cardone et al. 2011; Famaey & McGaugh 2012; Trippe 2014; Walker & Loeb 2014; Wu & Kroupa 2015; Milgrom 2015; Chae & Gong 2015). Milgrom's law eliminates particle dark matter from galactic dynamics together with its substantial difficulties (cf., e.g., Kroupa 2012, 2015).

As pointed out by Trippe (2013a,b,c, 2015), massive gravity can, at least in principle, provide Milgrom's law together with an expression for the function μ in agreement with observations. Following up on this ansatz, I investigate the link of galactic dynamics to the cosmic expansion history. A physical connection is established by the graviton mass $m_\mathrm{g} \propto \sqrt{\Lambda}$. Milgrom's constant is given by $a_0 \propto c\sqrt{\Lambda} \propto cH_0\sqrt{3\Omega_\Lambda}$, with c, Λ, H_0, and Ω_Λ being the speed of light, the cosmological constant, the (present-day) Hubble constant, and the (present-day) third cosmological parameter, respectively. Accordingly, massive gravity might offer a natural alternative explanation for the phenomena conventionally associated with dark matter and dark energy.

2. CALCULATIONS

Massive gravity implies (Trippe 2013a,b,c, 2015) that any luminous ("baryonic") mass M_0 is the source of a spherical graviton halo with mass density

$$\rho_\mathrm{g}(R) = m_\mathrm{g}\, n_\mathrm{g}(R) = m_\mathrm{g}\, \eta\, M_0\, R^{-2} \tag{4}$$

where m_g is the graviton mass, n_g is the graviton particle density, R is the radial coordinate, and η is a scaling factor. The proportionality $\rho_\mathrm{g} \propto M_0 R^{-2}$ follows from (i) consistency with the classical force law in the Newtonian limit, and (ii) the inverse-square-of-distance law of flux conservation (but see also the discussion in Section 3). For a circular orbit of radius r around M_0, the total, dynamical mass experienced by a test particle follows from integrating $\rho_\mathrm{g}(R)$ from 0 to r,

$$M_\mathrm{dyn} = M_0 + M_\mathrm{g} = M_0\,(1 + 4\,\pi\,\eta\, m_\mathrm{g}\, r) \tag{5}$$

with M_g being the mass contributed by the graviton halo. One can bring this expression into the more intuitive form

$$\frac{M_\mathrm{dyn}}{M_0} = 1 + 4\,\pi\, f\, \frac{m_\mathrm{g}}{x}\, \frac{r}{y} \tag{6}$$

where f is a dimensionless number, x is a mass scale, and y is a length scale (with $\eta \equiv f/(xy)$).

With gravitons being virtual exchange particles, their effective mass can be estimated from (cf., e.g., Griffiths 2008) Heisenberg's uncertainty relation for energy and time,

$$m_\mathrm{g} c^2 \times T_\mathrm{g} \approx \hbar \tag{7}$$

with T_g being the life time of gravitons and $\hbar$ being the reduced form of Planck's constant. Consistency with the cosmic expansion history requires $T_\mathrm{g} \approx \sqrt{1/\Lambda}$, resulting in

$$m_\mathrm{g} \approx \frac{\hbar}{c^2}\,\sqrt{\Lambda}\;. \tag{8}$$

According to the standard "cosmic triangle" formalism (e.g., Bahcall et al. 1999), $\Lambda = 3H_0^2\Omega_\Lambda$, resulting in

$$m_\mathrm{g} \approx \frac{\hbar}{c^2}\, H_0\, \sqrt{3\Omega_\Lambda}\;; \tag{9}$$

for $H_0 \approx 70\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$ and $\Omega_\Lambda \approx 0.7$ (e.g., Ade et al. 2014), $\Lambda \approx 1.1 \times 10^{-35}\,\mathrm{s^{-2}}$ and thus $m_\mathrm{g} \approx 4 \times 10^{-69}\,\mathrm{kg} \approx 2 \times 10^{-33}\,\mathrm{eV}\,c^{-2}$.

A natural choice for a quantum-physical mass scale, x in our case, is the *Planck mass*

$$x \equiv m_\mathrm{P} = \sqrt{\frac{\hbar\, c}{G}} \tag{10}$$

with G being Newton's constant; combination of Equations (9) and (10) leads to $m_\mathrm{g}/m_\mathrm{P} \approx 2 \times 10^{-61}$. The choice for the scale y is less obvious; for our purpose, we require a scale characteristic for gravitationally bound dynamical systems. A good choice (to be discussed in Section 3) is the magnitude of the gravitational potential,

$$\epsilon = \frac{G\, M_\mathrm{dyn}}{r\, c^2} \tag{11}$$

(cf. also Baker et al. 2015). As ϵ is dimensionless, we can supply y with the unit of a length by multiplying ϵ and the *Planck length*

$$l_\mathrm{P} = \sqrt{\frac{\hbar\, G}{c^3}} \tag{12}$$

so that $y \equiv \epsilon\, l_\mathrm{P}$.

Inserting Equations (8, 10, 11, 12) into Equation (6) leads to

$$\frac{M_\mathrm{dyn}}{M_0} = 1 + 4\,\pi\, f\, c\sqrt{\Lambda}\;\frac{r^2}{G\, M_\mathrm{dyn}}\;. \tag{13}$$

Using the classical expression for gravitational field strength, $g = GM_\mathrm{dyn}/r^2$, one finds

$$\frac{M_\mathrm{dyn}}{M_0} = 1 + 4\,\pi\, f\;\frac{c\sqrt{\Lambda}}{g} \tag{14}$$

or, using the expression given by Equation (9),

$$\frac{M_\mathrm{dyn}}{M_0} = 1 + 4\,\pi\, f\;\frac{c\, H_0\, \sqrt{3\,\Omega_\Lambda}}{g}\;. \tag{15}$$

Comparison of Equations (14, 15) to the expression following from the "simple μ function" (e.g., Famaey & McGaugh 2012) of MOND,

$$\frac{M_\mathrm{dyn}}{M_0} = 1 + \frac{a_0}{g}\;, \tag{16}$$

shows that the expressions are equivalent, with Milgrom's constant being given by $a_0 = 4\pi f c\sqrt{\Lambda}$. Matching this expression with the empirical value $a_0 \approx 1.1 \times 10^{-10}\,\mathrm{m\,s^{-2}}$ requires a (universal) value of $f \approx 0.0088$ (or $4\pi f \approx 1/9$). It is straightforward to see that in the limit $g \ll a_0$, Equation (16) leads to Equation (2).

3. Discussion

As demonstrated in Section 2, massive gravity can be used to link the cosmological constant to a specific version of Milgrom's law, namely the one comprising the "simple μ function". The transition function given by Equation (16), with $a_0 = (1.06 \pm 0.05) \times 10^{-10}\,\mathrm{m\,s^{-2}}$, is in excellent agreement with the observed mass discrepancy–acceleration relation of disk galaxies (Trippe 2013c). I thus present a physical mechanism that provides astrophysically meaningful expressions for both μ and Milgrom's constant. My derivation implies that dark matter and dark energy could be understood as phenomena arising from the same effect – the non-zero mass and thus finite life time of gravitons. This mechanism also explains why $a_0 \sim cH_0$ as already noted by Milgrom (1983a). I also note that a similar approach has been discussed by van Putten (2014) in the specific context of de-Sitter space (but see also van Putten 2015).

The result expressed in Equation (14) critically depends on the choice of the scales x and y. Identifying x with the Planck mass is a natural choice. Identification of y with the magnitude of the gravitational potential, ϵ, (multiplied with the Planck length) is suggested by the properties of gravity on astronomical scales. As pointed out recently by Baker et al. (2015), astrophysical gravitating systems are characterized completely by their location in a two-dimensional plane spanned by (i) the parameter ϵ and (ii) the *Kretschmann scalar* which corresponds to the magnitude of the Riemann curvature tensor (thus quantifying the local strength of spacetime curvature in the frame of general relativity) and which takes the form $\xi = \sqrt{48}\,GM_{\mathrm{dyn}}/(r^3c^2)$ for spherical systems. However, ξ can be expressed in units of the cosmic curvature, Λ/c^2; thus the curvature (or at least a specific value thereof) already appears in Equation (8), which suggests the introduction of ϵ (multiplied with the Planck length for dimensional reasons) as a new scaling parameter. Conveniently, my choice of x and y cancels out the constant $\hbar$ coming with the graviton mass (Equation (8)) because $m_{\mathrm{P}}\,l_{\mathrm{P}} = \hbar/c$. The dimensionless factor f arises from the need to provide a normalization factor for the graviton halo density in Equation (4) of the form $\eta \equiv f/(xy)$; I emphasize that f is the only free parameter left in the derivation because neither m_{P} nor l_{P} nor ϵ are tunable. For mass density distributions, scaling factors $0 < f \leq 1$ – in our case, $f \approx 0.9\%$ – can be interpreted generically as filling or efficiency factors (in the absence of further information, as is the case here). I also note that Equation (4) follows explicitly from the limiting case of Newtonian gravity. Obviously, the inverse-square-of-distance law of gravity does not hold globally in massive gravity but only in the limit $g \gg a_0$. The choice of Equation (4) is motivated by the need to connect Newtonian and modified Newtonian gravity in the strong field limit.

Even though they are quite intuitive already, the relations presented in Section 2 should eventually follow from, or be included in, a complete theory of massive gravity – which has not been found yet. Especially, it will be necessary to connect the classical derivation of Equation (16) with general relativity and the resulting deviations from Newtonian gravity in the strong-field limit ($g \gg a_0$). A complete theory should also provide for the basic properties of gravitons, especially the absence of graviton–graviton interactions which is required for the validity of the classical force law (as used in Equation (4); cf., Trippe 2013a – but see also Vainshtein 1972; Babichev & Deffayet 2013; Avilez-Lopez et al. 2015). Last but not least, a complete theory of massive gravity should also comprise non-dynamical effects that are currently unexplained in the frame of general relativity; possible candidates are the "radio flux ratio anomaly" affecting the images of multiply gravitationally lensed quasars (e.g., Xu et al. 2015, and references therein) and the discrepancy between mass estimates based on galactic dynamics and those based on gravitational lensing found recently for filaments of the Virgo cluster of galaxies (Lee et al. 2015).

4. Summary and Conclusions

I explore a possible physical connection between dark matter and dark energy. Assuming that gravitation is mediated by virtual gravitons with non-zero mass (massive gravity), the resulting limited life time of gravitons provides for a decay of gravity on cosmological scales and thus an accelerated expansion of the universe ("dark energy"); the graviton mass follows from the cosmological constant like $m_{\mathrm{g}} \approx \hbar\sqrt{\Lambda}/c^2$ via Heisenberg's uncertainty relation. Massive gravity also implies that any luminous ("baryonic") mass is surrounded by a (electromagnetically invisible) halo of gravitons that contributes additional mass ("dark matter"). Assuming reasonable scaling relations for such graviton halos, one recovers Milgrom's law of gravity and finds that Milgrom's constant is $a_0 = 4\pi f c\sqrt{\Lambda} = 4\pi f c H_0\sqrt{3\Omega_\Lambda}$ with $f \approx 0.9\%$ (or $4\pi f \approx 1/9$).

My derivation suggests that dark matter and dark energy could be interpreted as two effects arising from the same physical mechanism: massive gravity. This would imply a natural connection between the dynamics of galaxies and the dynamics of the universe.

Acknowledgments

I am grateful to Benoit Famaey and Christian Boily (both: Observatoire Astronomique de Strasbourg) for inspiring discussion, and to an anonymous referee for valuable comments. I acknowledge financial support from the Korean National Research Foundation (NRF) via Basic Research Grant 2012-R1A1A-2041387.

References

Ade, P. A. R., et al. 2014, Planck 2013 Results. XVI. Cosmological Parameters, A&A, 571, A16

Avilez-Lopez, A., Padilla, A., Saffin, P. M., & Skordis, C. 2015, The Parameterized Post-Newtonian-Vainshteinian Formalism, arXiv:1501.01985

Bahcall, N. A., Ostriker, J. P., Perlmutter, S., & Steinhardt, P. J. 1999, The Cosmic Triangle: Revealing the State of the Universe, Science, 284, 1481

Babichev, E. & Deffayet, C. 2013, An Introduction to the Vainshtein Mechanism, Class. Quantum Grav., 30, 184001

Baker, T., Psaltis, D., & Skordis, C. 2015, Linking Tests of Gravity on All Scales: From the Strong-Field Regime to Cosmology, ApJ, 802, 63

Cardone, V. F., Angus, G., Diaferio, A., et al. 2011, The Modified Newtonian Dynamics Fundamental Plane, MNRAS, 412, 2617

Cardone, V. F., Radicella, N., & Parisi, L. 2012, Constraining Massive Gravity with Recent Cosmological Data, Phys. Rev. D, 85, 124005

Chae, K.-H. & Gong, I.-T. 2015, Testing Modified Newtonian Dynamics through Statistics of Velocity Dispersion Profiles in the Inner Regions of Elliptical Galaxies, arXiv:1505.02936

Clifton, T., Ferreira, P. G., Padilla, A., & Skordis, C. 2012, Modified Gravity and Cosmology, Phys. Rep., 513, 1

De Felice, A., Gümrükcüoglu, A. E., Lin, C., & Mukohyama, S. 2013, On the Cosmology of Massive Gravity, Class. Quantum Grav., 30, 184004

de Rham, C. 2014, Massive Gravity, Living Rev. Relativ., 17, 7

Famaey, B. & McGaugh, S. S. 2012, Modified Newtonian Dynamics (MOND): Observational Phenomenology and Relativistic Extensions, Living Rev. Relativ., 15, 10

Fierz, M. & Pauli, W. 1939, On Relativistic Wave Equations for Arbitrary Spin in an Electromagnetic Field, Proc. R. Soc. London A, 173, 211

Gentile, G., Famaey, B., & de Blok, W. J. G. 2011, THINGS about MOND, A&A, 527, A76

Griffiths, D. 2008, Introduction to Elementary Particles, 2nd edn. (Weinheim: Wiley-VCH)

Hinterbichler, K. 2012, Theoretical Aspects of Massive Gravity, Rev. Mod. Phys., 84, 671

Kroupa, P. 2012, The Dark Matter Crisis: Falsification of the Current Standard Model of Cosmology, PASA, 29, 395

Kroupa, P. 2015, Galaxies as Simple Dynamical Systems: Observational Data Disfavor Dark Matter and Stochastic Star Formation, Can. J. Phys., 93, 169

Lee, J., Kim, S., & Rey, S.-C. 2015, A New Dynamical Mass Estimate for the Virgo Cluster Using the Radial Velocity Profile of the Filament Galaxies, arXiv:1501.07064

McGaugh, S. S. 2004, The Mass Discrepancy–Acceleration Relation: Disk Mass and the Dark Matter Distribution, ApJ, 609, 652

McGaugh, S. S. 2005, The Baryonic Tully–Fisher Relation of Galaxies with Extended Rotation Curves and the Stellar Mass of Rotating Galaxies, ApJ, 632, 859

McGaugh, S. S. 2005, Balance of Dark and Luminous Mass in Rotating Galaxies, Phys. Rev. Lett., 95, 171302

McGaugh, S. S. 2011, Novel Test of Modified Newtonian Dynamics with Gas Rich Galaxies, Phys. Rev. Lett., 106, 121303

Milgrom, M. 1983, A Modification of the Newtonian Dynamics as a Possible Alternative to the Hidden Mass Hypothesis, ApJ, 270, 365

Milgrom, M. 1983, A Modification of the Newtonian Dynamics: Implications for Galaxies, ApJ, 270, 371

Milgrom, M. 1983, A Modification of the Newtonian Dynamics: Implications for Galaxy Systems, ApJ, 270, 384

Milgrom, M. 1984, Isothermal Spheres in the Modified Dynamics, ApJ, 287, 571

Milgrom, M. 1994, Modified Dynamics Predictions Agree with Observations of the H I Kinematics in Faint Dwarf Galaxies Contrary to the Conclusions of Lo, Sargent, and Young, ApJ, 429, 540

Milgrom, M. 2015, MOND Theory, Can. J. Phys. 93, 107

Milgrom, M. & Sanders, R. H. 2008, Rings and Shells of "Dark Matter" as MOND Artifacts, ApJ, 678, 131

Rhee, M.-H. 2004, On the Physical Basis of the Tully–Fisher Relation, JKAS, 37, 15

Rhee, M.-H. 2004, Mass-to-Light Ratio and the Tully–Fisher Relation, JKAS, 37, 91

Sanders, R. H. 1994, A Faber–Jackson Relation for Clusters of Galaxies: Implications for Modified Dynamics, A&A, 284, L31

Sanders, R. H. 2010, The Universal Faber–Jackson Relation, MNRAS, 407, 1128

Tasinato, G., Koyama, K., & Niz, G. 2013, Exact Solutions in Massive Gravity, Class. Quantum Grav., 30, 184002

Trippe, S. 2013, A Simplified Treatment of Gravitational Interaction on Galactic Scales, JKAS, 46, 41

Trippe, S. 2013, A Derivation of Modified Newtonian Dynamics, JKAS, 46, 93

Trippe, S. 2013, Can Massive Gravity Explain the Mass Discrepancy–Acceleration Relation of Disk Galaxies?, JKAS, 46, 133

Trippe, S. 2014, The 'Missing Mass Problem' in Astronomy and the Need for a Modified Law of Gravity, Z. Naturforsch. A, 69, 173

Trippe, S. 2015, The "Graviton Picture": a Bohr Model for Gravitation on Galactic Scales?, Can. J. Phys., 93, 213

Vainshtein, A. I. 1972, To the Problem of Non-Vanishing Gravitation Mass, Phys. Lett. B, 39, 393

van Putten, M. H. P. M. 2014, Galaxy Rotation Curves in de Sitter Space, arXiv:1411.2665

van Putten, M. H. P. M. 2015, Accelerated Expansion from Cosmological Holography, MNRAS, 450, L48

Volkov, M. S. 2012, Cosmological Solutions with Massive Gravitons in the Bigravity Theory, J. High Energy Phys., 2012, 35

Walker, M. G. & Loeb, A. 2014, Is the Universe Simpler than ΛCDM?, Contemp. Phys., 55, 198

Wu, X. & Kroupa, P. 2015, Galactic Rotation Curves, the Baryon-to-Dark-Halo-Mass Relation and Space-Time Scale Invariance, MNRAS, 446, 330

Xu, D., Sluse, D., Gao, L., et al. 2015, How Well Can Dark-Matter Substructures Account for the Observed Radio Flux-Ratio Anomalies?, MNRAS, 447, 3189