Sensitivity of High-Scale SUSY in Low Energy Hadronic FCNC

Morimitsu Tanimoto^{a,1} and Kei Yamamoto^{b,2}

^bDepartment of Physics, Niigata University, Niigata 950-2181, Japan ^bInstitute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan

Abstract

We discuss the sensitivity of the high-scale SUSY at 10-1000 TeV in B^0 , B_s , K^0 and D meson systems together with the neutron EDM and the mercury EDM. In order to estimate the contribution of the squark flavor mixing to these FCNCs, we calculate the squark mass spectrum, which is consistent with the recent Higgs discovery. The SUSY contribution in ϵ_K could be large, around 40% in the region of the SUSY scale 10-100 TeV. The neutron EDM and the mercury EDM are also sensitive to the SUSY contribution induced by the gluino-squark interaction. The predicted EDMs are roughly proportional to $|\epsilon_K^{\text{SUSY}}|$. If the SUSY contribution is the level of $\mathcal{O}(10\%)$ for ϵ_K , the neutron EDM is expected to be discovered in the region of 10^{-28} - 10^{-26} ecm. The mercury EDM also gives a strong constraint for the gluino-squark interaction. The SUSY contribution. The SUSY contribution of ΔM_D is also discussed.

 $^{^1\}mathrm{E}\text{-}\mathrm{mail}$ address: tanimoto@muse.sc.niigata-u.ac.jp

²E-mail address: kei.yamamoto@kek.jp

1 Introduction

The supersymmetry (SUSY) is one of the most attractive theories beyond the standard model (SM). Therefore, the SUSY has been expected to be observed at the LHC experiments. However, no signals of the SUSY have been discovered yet. The present searches for the SUSY particles give us important constraints for the SUSY. Since the lower bounds of the superparticle masses increase gradually, the squark and the gluino masses are supposed to be at the higher scale than 1 TeV [1, 2, 3]. On the other hand, the SUSY model has been seriously constrained by the Higgs discovery, in which the Higgs mass is 125 GeV [4]. Based on this theoretical and experimental situations, we consider the high-scale SUSY models, which have been widely discussed with a lot of attention [5]-[20].

If the squark and slepton masses are at the high-scale $\mathcal{O}(10\text{-}1000)$ TeV, the lightest Higgs mass can be pushed up to 125 GeV, whereas SUSY particles are out of the reach of the LHC experiment. Therefore, the indirect search of the SUSY particles becomes important in the low energy flavor physics [21, 22, 23].

The flavor physics is also on the new stage in the light of LHCb data. The LHCb collaboration has reported new data of the CP violation of the B_s meson and the branching ratios of rare B_s decays [24]-[36]. For many years the CP violation in the K and B^0 mesons has been successfully understood within the framework of the standard model (SM), so called Kobayashi-Maskawa (KM) model [37], where the source of the CP violation is the KM phase in the quark sector with three families. However, the new physics has been expected to be indirectly discovered in the precise data of B^0 and B_s meson decays at the LHCb experiment and the further coming experiment, Belle-II.

There are new sources of the CP violation if the SM is extended to the SUSY models. The soft squark mass matrices contain the CP violating phases, which contribute to the flavor changing neutral current (FCNC) with the CP violation [38]. Therefore, we can expect the SUSY effect in the CP violating phenomena. However, the clear deviation from the SM prediction has not been observed yet in the LHCb experiment [24]-[36]. Actually, we have found that the CP violation of B^0 and B_s meson systems are suppressed if the SUSY scale is above 10 TeV [39]. On the other hand, the CKMfitter group presented the current limits on new physics contributions of $\mathcal{O}(10\%)$ in B^0 , B_s and K^0 systems [40]. They have also estimated the sensitivity to new physics in B^0 and B_s mixing achievable with $50ab^{-1}$ of Belle-II and $50fb^{-1}$ of LHCb data. Therefore, we should carefully study the sensitivity of the high-scale SUSY to the hadronic FCNC.

In this work, we discuss the high-scale SUSY contribution to the B^0 , B_s and K^0 meson systems. Furthermore, we also discuss the sensitivity to the D meson and the electric dipole moment (EDM) of the neutron and the mercury. For these modes, the most important process of the SUSY contribution is the gluino-squark mediated flavor changing process [41]-[56]. The CP violation of K meson, ϵ_K , provides a severe constraint to the gluino-squark mediated FCNC [57, 58]. In addition, the recent work have found that the chromo-electric dipole moment (cEDM) is sensitive to the high-scale SUSY [59]. It is noted that the upperbound of the neutron EDM (nEDM) [60] gives a severe constraint for the gluino-squark interaction through the cEDM [61]-[66]. It is also remarked that the upper bound of the mercury EDM (HgEDM) [67] can give an important constraint [68]. In order to estimate the gluino-squark mediated FCNC of the K, B^0 , B_s and D mesons, we work in the basis of the squark mass eigenstate with the non-minimal squark (slepton) flavor mixing. There are three reasons why the SUSY contribution to the FCNC considerably depends on the squark mass spectrum. The first one is that the GIM mechanism works in the squark flavor mixing, and the second one is that the loop functions depend on the mass ratio of squark and gluino. The last one is that we need the mixing angle between the left-handed sbottom and right-handed sbottom, which dominates the $\Delta B = 1$ decay processes. Therefore, we discuss the squark mass spectrum, which is consistent with the recent Higgs discovery. Taking the universal soft parameters at the SUSY breaking scale, we obtain the squark mass spectrum at the matching scale where the SM emerges, by using the Renormalization Group Equations (RGEs) of the soft masses. On the other hand, the 6 × 6 mixing matrix between squarks and quarks is taken to be free at the low energy.

In section 2, we discuss the squark and gluino mass spectrum and the squark mixing. In section 3, we present the formulation of the FCNC with $\Delta F = 2$ in K, B^0 , B_s and D meson systems together with nEDM and HgEDM. We present numerical results and discussions in section 4. Section 5 is devoted to the summary. The relevant formulations are presented in Appendices A, B, C and D.

2 SUSY Spectrum and Squark mixing

The low energy FCNCs depend significantly on the spectrum of the SUSY particles, which depend on the model. As well known, the lightest Higgs mass can be pushed up to 125 GeV if the squark masses are expected to be $\mathcal{O}(10)$ TeV. Therefore, let us consider the heavy SUSY particle mass spectrum in the framework of the minimal supersymmetric standard model (MSSM), which is consistent with the observed Higgs mass. The discussion how to obtain the SUSY spectrum have been given in Refs. [69, 70].

We outline how to obtain the SUSY spectrum in our work. The details are presented in Appendix A. At the SUSY breaking scale Λ , we write the quadratic terms in the MSSM potential as

$$V_2 = m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_3^2 (H_1 \cdot H_2 + h.c.) .$$
(1)

Then, the Higgs mass parameter m^2 is expressed in terms of m_1^2 , m_2^2 and $\tan \beta$ as:

$$m^{2} = \frac{m_{1}^{2} - m_{2}^{2} \tan^{2} \beta}{\tan^{2} \beta - 1} .$$
⁽²⁾

After running down to the Q_0 scale, in which the SM emerges, by the one-loop SUSY Renormalization Group Equations (RGEs) [71], the scalar potential is the SM one as follows:

$$V_{SM} = -m^2 |H|^2 + \frac{\lambda}{2} |H|^4 .$$
(3)

Here, the Higgs coupling λ is given in terms of the SUSY parameters at the leading order as

$$\lambda(Q_0) = \frac{1}{4} (g^2 + g'^2) \cos^2 2\beta + \frac{3h_t^2}{8\pi^2} X_t^2 \left(1 - \frac{X_t^2}{12} \right) , \qquad X_t = \frac{A_t(Q_0) - \mu(Q_0) \cot \beta}{Q_0} , \quad (4)$$

and h_t is the top Yukawa coupling of the SM. The parameters m_2 and λ run with the two-loop SM RGEs with $\overline{\text{MS}}$ scheme [72] down to the electroweak scale $Q_{EW} = m_H$, and then give

$$m_H^2 = 2m^2(m_H) = \lambda(m_H)v^2$$
 (5)

When $m_H = 125$ GeV is put, $\lambda(Q_0)$ and $m^2(Q_0)$ are obtained. This input constrains the SUSY mass spectrum of the MSSM. In our work, we take the universal soft breaking parameters at the SUSY breaking scale Λ as follows:

$$m_{\tilde{Q}_{i}}(\Lambda) = m_{\tilde{U}_{i}^{c}}(\Lambda) = m_{\tilde{D}_{i}^{c}}(\Lambda) = m_{\tilde{L}_{i}}(\Lambda) = m_{\tilde{E}_{i}^{c}}(\Lambda) = m_{0}^{2} \ (i = 1, 2, 3) ,$$

$$M_{1}(\Lambda) = M_{2}(\Lambda) = M_{3}(\Lambda) = m_{1/2} , \qquad m_{1}^{2}(\Lambda) = m_{2}^{2}(\Lambda) = m_{0}^{2} ,$$

$$A_{U}(\Lambda) = A_{0}y_{U}(\Lambda) , \qquad A_{D}(\Lambda) = A_{0}y_{D}(\Lambda) , \qquad A_{E}(\Lambda) = A_{0}y_{E}(\Lambda) .$$
(6)

By inputting $m_H = 125$ GeV and taking the heavy scalar mass $m_H \simeq Q_0$ (see Appendix A), we can obtain the SUSY spectrum for the fixed Q_0 and $\tan \beta$. The details and numerical results are presented in Appendix A.

Let us consider the squark flavor mixing. As discussed above, there is no flavor mixing at Λ in the MSSM. However, in order to consider the non-minimal flavor mixing framework, we allow the off diagonal components of the squark mass matrices at the 10% level, which leads to the flavor mixing of order 0.1. We take these flavor mixing angles as free parameters at low energies. Now we consider the 6×6 squark mass matrix $M_{\tilde{q}}$ in the super-CKM basis. In order to move the mass eigenstate basis of squark masses, we should diagonalize the mass matrix by rotation matrix $\Gamma_{G}^{(q)}$ as

$$m_{\tilde{q}}^2 = \Gamma_G^{(q)} M_{\tilde{q}}^2 \Gamma_G^{(q)\dagger} , \qquad (7)$$

where $\Gamma_G^{(q)}$ is the 6 × 6 unitary matrix, and we decompose it into the 3 × 6 matrices as $\Gamma_G^{(q)} = (\Gamma_{GL}^{(q)}, \Gamma_{GR}^{(q)})^T$ in the following expressions:

$$\Gamma_{GL}^{(d)} = \begin{pmatrix} c_{13}^L & 0 & s_{13}^L e^{-i\phi_{13}^L} c_{\theta} & 0 & 0 & -s_{13}^L e^{-i\phi_{13}^L} s_{\theta} e^{i\phi} \\ -s_{23}^L s_{13}^L e^{i(\phi_{13}^L - \phi_{23}^L)} & c_{23}^L & s_{23}^L c_{13}^L e^{-i\phi_{23}^L} c_{\theta} & 0 & 0 & -s_{23}^L c_{13}^L e^{-i\phi_{23}^L} s_{\theta} e^{i\phi} \\ -s_{13}^L c_{23}^L e^{i\phi_{13}^L} & -s_{23}^L e^{i\phi_{23}^L} & c_{13}^L c_{23}^L c_{\theta} & 0 & 0 & -c_{13}^L c_{23}^L s_{\theta} e^{i\phi} \end{pmatrix},$$

$$\Gamma_{GR}^{(d)} = \begin{pmatrix} 0 & 0 & s_{13}^R s_{\theta} e^{-i\phi_{13}^R} e^{-i\phi} & c_{13}^R & 0 & s_{13}^R e^{-i\phi_{13}^R} c_{\theta} \\ 0 & 0 & s_{23}^R c_{13}^R s_{\theta} e^{-i\phi_{23}^R} e^{-i\phi} & -s_{13}^R s_{23}^R e^{i(\phi_{13}^R - \phi_{23}^R)} & c_{23}^R & s_{23}^R c_{13}^R e^{-i\phi_{23}^R} c_{\theta} \\ 0 & 0 & c_{13}^R c_{23}^R s_{\theta} e^{-i\phi} & -s_{13}^R c_{23}^R e^{i\phi_{13}^R} & -s_{23}^R e^{i\phi_{23}^R} & c_{13}^R c_{23}^R c_{\theta} \end{pmatrix}, \quad (8)$$

where we use abbreviations $c_{ij}^{L,R} = \cos \theta_{ij}^{L,R}$, $s_{ij}^{L,R} = \sin \theta_{ij}^{L,R}$, $c_{\theta} = \cos \theta$ and $s_{\theta} = \sin \theta$. Here θ is the left-right mixing angle between \tilde{b}_L and \tilde{b}_R , which is discussed in Appendix A. It is remarked that we take $s_{12}^{L,R} = 0$ due to the degenerate squark masses of the first and second families as discussed in Appendix A.

The gluino-squark-quark interaction is given as

$$\mathcal{L}_{\text{int}}(\tilde{g}q\tilde{q}) = -i\sqrt{2}g_s \sum_{\{q\}} \tilde{q}_i^*(T^a) \overline{\tilde{G}^a} \left[(\Gamma_{GL}^{(q)})_{ij}L + (\Gamma_{GR}^{(q)})_{ij}R \right] q_j + \text{h.c.} , \qquad (9)$$

where $L = (1-\gamma_5)/2$, $R = (1+\gamma_5)/2$, and \tilde{G}^a denotes the gluino field, q^i are three left-handed (i=1,2,3) and three right-handed quarks (i=4,5,6). This interaction leads to the gluino-squark mediated flavor changing processes with $\Delta F = 2$ and $\Delta F = 1$ through the box and penguin diagrams.

The chargino (neutralino)-squark-quark interaction can be also discussed in the similar way.

3 FCNC of $\Delta F = 2$

In our previous work [39], we have probed the high-scale SUSY, which is at 10-50 TeV scale, in the CP violations of K, B^0 and B_s mesons. It is found that ϵ_K is most sensitive to the SUSY even if the SUSY scale is at 50 TeV. The SUSY contributions for the time dependent CP asymmetries of B^0 and B_s with $\Delta B = 1$ are suppresses at the SUSY scale of 10 TeV. Furthermore, the SUSY contribution for the $b \to s\gamma$ process is also suppressed since the left-right mixing angle, which induces the chiral enhancement, is very small as discussed in Appendix A. Therefore, we discuss the neutral meson mixing $P^0 - \bar{P}^0(P^0 = K, B^0, B_s, D)$, which are FCNCs with $\Delta F = 2$.

In those FCNCs, the dominant SUSY contribution is given through the gluino-squark interaction. Then, the dispersive part of meson mixing $M_{12}^{P^0}(P^0 = K, B^0, B_s)$ are written as

$$M_{12}^{P^0} = M_{12}^{P^0,\rm SM} + M_{12}^{P^0,\rm SUSY},\tag{10}$$

where $M_{12}^{q,\text{SUSY}}$ are given by the squark mixing parameters in Eq.(8) and its explicit formulation is given in Appendices B and C.

At first, we discuss the $\Delta B = 2$ process, that is, the mass differences ΔM_{B^0} and ΔM_{B_s} , and the CP-violating phases ϕ_d and ϕ_s . In general, the contribution of the new physics (NP) to the dispersive part M_{12}^q is parameterized as

$$M_{12}^{B_q} = M_{12}^{q,\text{SM}} + M_{12}^{q,\text{NP}} = M_{12}^{q,\text{SM}} (1 + h_q e^{2i\sigma_q}) , \quad (q = B^0, B_s)$$
(11)

where $M_{12}^{q,\text{NP}}$ are the NP contributions. The generic fits for B^0 and B_s mixing have given the constraints on (h_q, σ_q) [40], where it is assumed that the NP does not significantly affect the SM tree-level charged-current interaction, that is, the absorptive part Γ_{12}^q is dominated by the decay $b \to c\bar{c}s$. At present, the NP contribution h_q are 10-35% and 15-25% depending on σ_q for B^0 and B_s , respectively. Thus, we can expect the sizable NP contribution of $\mathcal{O}(20\%)$. We will discuss whether the high-scale SUSY can fill in the magnitude of the present NP contribution of $\mathcal{O}(20\%)$.

Next, we discuss $\Delta S = 2$ process, ΔM_{K^0} and the CP-violating parameter in the K meson, ϵ_K . By the similar parametrization in Eq.(11), the allowed region of (h_K, σ_K) has been estimated in Ref.[40]. The NP contribution is at least 50% although there is the strong σ_K dependence. Therefore, it is important to examine carefully the CP violating parameter ϵ_K , which is given as follows:

$$\epsilon_K = e^{i\phi_\epsilon} \sin \phi_\epsilon \left(\frac{\operatorname{Im}(M_{12}^K)}{\Delta M_K} + \xi \right), \qquad \xi = \frac{\operatorname{Im} A_0^K}{\operatorname{Re} A_0^K}, \qquad \phi_\epsilon = \tan^{-1} \left(\frac{2\Delta M_K}{\Delta \Gamma_K} \right), \tag{12}$$

with A_0^K being the isospin zero amplitude in $K \to \pi \pi$ decays. Here, M_{12}^K is the dispersive part of the K^0 - \bar{K}^0 mixing, and ΔM_K is the mass difference in the neutral K meson. The effects of $\xi \neq 0$ and $\phi_{\epsilon} < \pi/4$ give suppression effect in ϵ_K , and it is parameterized as κ_{ϵ} and estimated by Buras and Guadagnoli [73] as:

$$\kappa_{\epsilon} = 0.92 \pm 0.02 \quad . \tag{13}$$

In the SM, the dispersive part M_{12}^K is given as follows,

$$M_{K}^{12} = \langle K | \mathcal{H}_{\Delta F=2} | K \rangle$$

= $-\frac{4}{3} \left(\frac{G_F}{4\pi} \right)^2 M_W^2 \hat{B}_K F_K^2 M_K \left(\eta_{cc} \lambda_c^2 E(x_c) + \eta_{tt} \lambda_t^2 E(x_t) + 2\eta_{ct} \lambda_c \lambda_t E(x_c, x_t) \right), \quad (14)$

where $\lambda_c = V_{cs}V_{cd}^*$, $\lambda_t = V_{ts}V_{td}^*$. The E(x)'s are the one-loop functions [74] and $\eta_{cc,tt,ct}$ are the QCD corrections [73]. Then, $|\epsilon_K^{\text{SM}}|$ is given in terms of the Wolfenstein parameters λ , $\overline{\rho}$ and $\overline{\eta}$ as follows:

$$|\epsilon_K^{\rm SM}| = \kappa_\epsilon C_\epsilon \hat{B}_K |V_{cb}|^2 \lambda^2 \bar{\eta} \left(|V_{cb}|^2 (1 - \bar{\rho}) \eta_{tt} E(x_t) - \eta_{cc} E(x_c) + \eta_{ct} E(x_c, x_t) \right) , \qquad (15)$$

with

$$C_{\epsilon} = \frac{G_F^2 F_K^2 m_K M_W^2}{6\sqrt{2}\pi^2 \Delta M_K}.$$
(16)

Note that $|\epsilon_K^{\text{SM}}|$ depends on the non-perturbative parameter \hat{B}_K in Eq.(15). Recently, the error of this parameter shrank dramatically in the lattice calculations [75]. In our calculation we use the updated value by the Flavor Lattice Averaging Group [76]:

$$\ddot{B}_K = 0.766 \pm 0.010$$
 . (17)

Let us write down ϵ_K as:

$$\epsilon_K = \epsilon_K^{\rm SM} + \epsilon_K^{\rm SUSY},\tag{18}$$

where $\epsilon_K^{\rm SUSY}$ is induced by the imaginary part of the gluino-squark box diagram, which is presented in Appendices B and C. Since $s_{12}^{L(R)}$ vanishes in our scheme, ϵ_K^{SUSY} is given in the second order of the squark mixing $s_{13}^{L(R)} \times s_{23}^{L(R)}$.

In addition to the above FCNC processes, the neutron EDM, d_n arises through the cEDM of the quarks, d_q^C due to the gluino-squark mixing [61]-[66]. By using the QCD sum rules, d_n is given as

$$d_n = (0.79d_d - 0.20d_u) + e(0.3d_u^C + 0.59d_d^C) .$$
⁽¹⁹⁾

where d_q and d_q^C denote the EDM and cEDM of quarks d_q^C defined in Appendix D. On the other hand, by using the chiral perturbation theory

$$d_n = e(3.0d_u^C + 2.5d_d^C + 0.5d_s^C) . (20)$$

Therefore, the experimental upper bound [60]

$$|d_n| < 0.29 \times 10^{-25} \text{ecm} , \qquad (21)$$

provides us a strong constraint to the gluino-squark mixing.

The HgEDM can also probe the gluino-squark mixing [68]. The QCD sum rule approach gives [77]

$$d_{Hg} = e(d_u^C - d_d^C + 0.012d_s^C) \times 3.2 \times 10^{-2} , \qquad (22)$$

and the chiral Lagrangian method gives [78]

$$d_{Hg} = e(d_u^C - d_d^C + 0.0051d_s^C) \times 8.7 \times 10^{-3} .$$
⁽²³⁾

The experimental upper bound [67]

$$|d_{Hg}| < 3.1 \times 10^{-29} \text{ecm} , \qquad (24)$$

constrains the gluino-squark mixing.

At the last step, we discuss the charm sector, which is a promising field to probe for the new physics beyond the SM. The $D^0 - \overline{D}^0$ mixing is now well established [79] as follows:

$$x_D = \frac{\Delta M_D}{\Gamma_D} = (3.6 \pm 1.6) \times 10^{-3} , \qquad y_D = \frac{\Delta \Gamma_D}{2\Gamma_D} = (6.1 \pm 0.7) \times 10^{-3} , \qquad (25)$$

where ΔM_D and $\Delta \Gamma_D$ are the differences of the masses and the decay widths between the mass eigenstates of the D meson, respectively, and Γ_D is the averaged decay width of the D meson. Since the SM prediction of ΔM_D at the short distance is much suppressed compared with the experimental value due to the bottom quark loop, the SUSY contribution may be enhanced.

4 Results and Discussions

Let us estimate the SUSY contribution of the low energy FCNC. We calculate the SUSY mass spectrum at $Q_0 = 10, 50, 100, 1000$ TeV and interpolate the each mass of the SUSY particle in the region of $Q_0 = 10\text{-}1000$ TeV. This approximation is satisfied within $\mathcal{O}(10\%)$. Therefore, our numerical results should be taken with the ambiguity of $\mathcal{O}(10\%)$. The mass spectrum at $Q_0 = 10$ TeV is presented in Appendix A. See Refs.[39], [58] for the mass spectrum at $Q_0 = 50$ TeV.

Then, we have four mixing angles $\theta_{13}^{L(R)}$ and $\theta_{23}^{L(R)}$, five phase $\phi_{13}^{L(R)}$, $\phi_{23}^{L(R)}$, ϕ . We reduce the number of parameters by taking $\sin \theta_{ij}^{L} = \sin \theta_{ij}^{R} \equiv s_{ij}$ for simplicity. In the numerical calculations, we scan the phases of Eq. (8) in the region of $0 \sim 2\pi$ for fixed s_{ij} , where the Cabibbo angle 0.22 and the large angle 0.5 are taken as the typical mixing. Other relevant input parameters such as quark masses m_c , m_b , the CKM parameters V_{us} , V_{cb} , $\bar{\rho}$, $\bar{\eta}$ and f_B , f_K , etc. have been presented in our previous paper Ref. [55], which are referred from UTfit Collaboration [80] and PDG [60].



Figure 1: The SUSY components of (a) ΔM_{B^0} and (b) ΔM_{B_s} versus $m_{\tilde{Q}}$ for $s_{13} = s_{23} = 0.22$ (cyan) and 0.5 (blue). The horizontal red line denotes the experimental central value.

4.1 B^0 and B_s meson systems

At first, we examine the SUSY contribution in the $\Delta B = 2$ process. We show the SUSY scale $m_{\tilde{Q}} \equiv Q_0$ dependence of the SUSY contributions of ΔM_{B^0} and ΔM_{B_s} in Figure 1(a) and (b), where the experimental central value is shown by the red line. The experimental error-bars are 1% and 0.1% levels for ΔM_{B^0} and ΔM_{B_s} , respectively. We take $s_{13} = s_{23} = 0.22, 0.5$. There is no phase dependence in our predictions. It is found that the SUSY contributions in ΔM_{B^0} and ΔM_{B_s} are at most 1.5% and 0.1% at $m_{\tilde{Q}} = 10$ TeV, respectively. Namely, the high-scale SUSY cannot explain the NP contributions of $h_d = 0.1$ -0.35 and $h_s = 0.15$ -0.25, which have been discussed in Eq.(11). As $m_{\tilde{Q}}$ increases, the SUSY contributions of both ΔM_{B^0} and ΔM_{B_s} decrease approximately with the power of $1/m_{\tilde{Q}}^2$. Thus, there is no hope to observe the SUSY contribution in the $\Delta B = 2$ process for the high-scale SUSY. It should be noted that the SM predictions are comparable to these experimental data.

The related phenomena are the CP violation of the non-leptonic decays $B^0 \to J/\psi K_S$ and $B_s \to J/\psi \phi$. The recent experimental data of these phases are [27, 34, 35, 36]

$$\sin \phi_d = 0.679 \pm 0.020$$
, $\phi_s = 0.07 \pm 0.09 \pm 0.01$, (26)

in which the contribution of the gluino-squark-quark interaction may be included. The NP contributions in ϕ_d and ϕ_s are expressed in terms of the parameters of Eq.(11) as [55]:

$$\phi_d = 2\beta_d + \arg(1 + h_d e^{2i\sigma_d}) , \qquad \phi_s = -2\beta_s + \arg(1 + h_s e^{2i\sigma_s}) , \qquad (27)$$

where $\beta_d(\beta_s)$ is the one angle of the unitarity triangle giving by the CKM matrix elements of the SM. However, h_d and h_s in the high-scale SUSY are much suppressed compared with $h_d = 0.1$ -0.35 and $h_s = 0.15$ -0.25 of Eq.(11), one cannot find signals of the high-scale SUSY in the CP violating decays $B^0 \to J/\psi K_S$ and $B_s \to J/\psi \phi$.

4.2 Neutral K meson system

At the second step, we examine the neutral K meson. We show the SUSY contributions of ΔM_{K^0} and ϵ_K versus $m_{\tilde{Q}} \equiv Q_0$ in Figure 2(a) and (b), where the experimental central value



Figure 2: The SUSY components of (a) ΔM_{K^0} and (b) $|\epsilon_K|$ versus $m_{\tilde{Q}}$ for $s_{13} = s_{23} = 0.22$ (cyan) and 0.5 (blue). The horizontal red line denotes the experimental central value.

is shown by the horizontal red line. The experimental error-bars are 0.2% and 0.5% levels for ΔM_{K^0} and ϵ_K , respectively. Since $\theta_{12}^{L,R} = 0$, the SUSY flavor mixing arise from the second order of $s_{13} \times s_{23}$, where $s_{13} = s_{23} = 0.22, 0.5$ are put.

It is found in Figure 2(a) that the SUSY contribution in ΔM_{K^0} can be comparable to the experimental value in the case of $s_{13} = s_{23} = 0.5$ whereas it is suppressed in the case of $s_{13} = s_{23} = 0.22$ at $m_{\tilde{Q}} = 10$ TeV. Thus, ΔM_{K^0} constrains the squark mixing of s_{13} and s_{23} around $m_{\tilde{Q}} = 10$ TeV. When the SUSY scale increases to more than 20 TeV, no SUSY contribution is expected.

On the other hand, ϵ_K is very sensitive to the SUSY contribution up to 100TeV as seen in Figure 2(b). The plot is scattered due to the random phases of the squark mixing. The experimental data of ϵ_K constrains the squark mixing and phases considerably. Actually, we have already pointed out that the SUSY contribution in ϵ_K could be 40% and 35% at $m_{\tilde{Q}} = 10,50$ TeV, respectively [39]. It is found that this seizable SUSY contribution still exist up to 100 TeV in this work.

In the SM, there is only one CP violating phase. Therefore, the observed value of ϕ_d in Eq.(27), should be correlated with ϵ_K in the SM. According to the recent experimental results, it is found that the consistency between the SM prediction and the experimental data of $\sin \phi_d$ and ϵ_K is marginal. This fact was pointed out by Buras and Guadagnoli [73] and called as the tension between ϵ_K and $\sin \phi_d$. Considering the effect of the SUSY contribution $\mathcal{O}(10\%)$ in ϵ_K , this tension can be relaxed even if $m_{\tilde{Q}} = 100$ TeV. The precise determination of the unitarity triangle of B^0 is required in order to find the SUSY contribution of this level.

It is noted that the SUSY contribution of both ΔM_{K^0} and ϵ_K also decrease approximately with the power of $1/m_{\tilde{Q}}^2$ as $m_{\tilde{Q}}$ increases up to 1000 TeV.

4.3 The nEDM and HgEDM with ϵ_K

The nEDM and HgEDM are also sensitive to the SUSY contribution [59, 68]. The gluinosquark interaction leads to the cEDM of quarks, which give the nEDM as shown in Eqs.(19) and (20). We show the predicted nEDM versus $m_{\tilde{O}}$ for the case of the QCD sum rules of



Figure 3: The the neutron EDM versus (a) $m_{\tilde{Q}}$ and (b) versus $|\epsilon_K^{\text{SUSY}}|$ for $s_{13} = s_{23} = 0.22$ (cyan) and 0.5 (blue) for the case of the QCD sum rule. The horizontal red line denotes the experimental upper bound of $|d_n|$ and the vertical one is the experimental central value of $|\epsilon_K|$.



Figure 4: The the mercury EDM versus (a) $m_{\tilde{Q}}$ and (b) versus $|\epsilon_K^{\text{SUSY}}|$ for $s_{13} = s_{23} = 0.22$ (cyan) and 0.5 (blue) for the case of the QCD sum rule. The horizontal red line denotes the experimental upper bound of $|d_{Hg}|$ and the vertical one is the experimental central value of $|\epsilon_K|$.

Eqs.(19) in Figure 3(a), where the upper bound of $|d_n|$ is shown by the red line. The plot is scattered due to the random phases of the squark mixing as well as in the case of ϵ_K . We find that the contribution of EDM, d_d and d_u occupy around 25% of the neutron EDM. The SUSY contribution is close to the experimental upper bound up to 50TeV. Since the predicted nEDM depends on the phases of the squark mixing matrix significantly, we plot the nEDM versus $|\epsilon_K^{\text{SUSY}}|$ in Figure 3(b). It is found that the predicted nEDM is roughly proportional to $|\epsilon_K^{\text{SUSY}}|$. If the SUSY contribution is the level of $\mathcal{O}(10\%)$ for ϵ_K , the nEDM is expected to be discovered in the region of 10^{-27} - 10^{-26} ecm. On the other hand, if the nEDM is not observed above 10^{-28} ecm, the SUSY contribution of ϵ_K is below a few %. Thus, there is the correlation between d_n and ϵ_K^{SUSY} . We also show the predicted HgEDM versus $m_{\tilde{Q}}$ for the case of the QCD sum rules of Eq.(22) in Figure 4(a), where the upper bound of $|d_{Hg}|$ is shown by the red line. The SUSY contribution is close to the experimental upper bound up to 200TeV, which is much higher than the one of the nEDM. In Figure 4(b), we plot the HgEDM versus $|\epsilon_K^{\text{SUSY}}|$. It is found that the experimental upper bound of the HgEDM excludes completely $|\epsilon_K^{\text{SUSY}}|$ which is inconsistent with the experimental data. If the SUSY contribution is the level of $\mathcal{O}(10\%)$ for ϵ_K , the nEDM is expected to be discovered in the region of 10^{-27} - 10^{-26} ecm. If the HgEDM is not observed above 10^{-29} ecm, the SUSY contribution of ϵ_K is below a few %. Thus, the mercury EDM gives more significant information for the gluino-squark interaction compared with the neutron EDM.

However, these correlations strongly depend on the assumptions of $\theta_{23}^L = \theta_{13}^L$ and $\theta_{ij}^L = \theta_{ij}^R$. The deviation from these relations destroys these correlations. For instance, for the case of $\theta_{23}^L \gg \theta_{13}^L$ with $\theta_{ij}^L = \theta_{ij}^R$, ϵ_K^{SUSY} is much suppressed whereas the nEDM and HgEDM are still sizable. On the other hand, if $\theta_{ij}^L \gg \theta_{ij}^R$ or $\theta_{ij}^L \gg \theta_{ij}^R$ is realized, the cEDMs are suppressed because they require the chirality flipping. In conclusion, the careful studies of the mixing angle relations are required to test the correlations between EDMs and ϵ_K^{SUSY} .

We should comment on the hadronic model dependence of our numerical result. For both nEDM and HgEDM, we show the numerical result by using the hadronic model of the QCD sum rules in Eqs.(19) and (22). We have also calculated the EDMs by using the hadronic model of the chiral perturbation theory in Eqs.(20) and (23). For the neutron EDM, the prediction of the chiral perturbation theory is larger than the one of the QCD sum rule at most of factor two. However, for the mercury EDM, the prediction of the QCD sum rule is more than three times larger compared with the one of the chiral perturbation theory. Thus, predicted EDMs have the ambiguity with the factor 2 - 3 from the hadronic model.

4.4 *D*-*D* mixing

Since the SM prediction of ΔM_D at the short distance is $\mathcal{O}(10^{-18})$ GeV, which is very small compared with the experimental value due to the bottom quark loop, it is important to estimate the SUSY contribution of ΔM_D . The mixing angle $\theta_{ij}^{L(R)}$ also appears in the uptype squark mixing matrix whereas the down-type squark mixing matrix contributes to K^0 , B^0 and B_s meson systems induced by the gluino-squark-quark interaction.

We show the SUSY component of ΔM_D and x_D versus $m_{\tilde{Q}}$ for $s_{13} = s_{23} = 0.22, 0.5$ in Figure 5. At the SUSY scale of 10 TeV, the SUSY component may be comparable to the observed value. Although the accurate estimate of the long-distance effect is difficult, Cheng and Chiang estimated x_D of order 10^{-3} from the two body hadronic modes [81]. This obtained value is consistent with the experimental one. Therefore, we should take into account the long-distance effect properly in order to constrain the SUSY contribution from ΔM_D .

Before closing the presentation of the numerical results, we add a comment on the other gaugino contribution. There are additional contributions to the FCNC induced by chargino exchanging diagrams. The chargino contribution to the gluino one is approximately 10% in the above numerical study of $\Delta F = 2$. Thus, the chargino contributions are the sub-leading ones.



Figure 5: The SUSY component of (a) ΔM_D and (b) x_D versus $m_{\tilde{Q}}$ for $s_{13} = s_{23} = 0.22$ (cyan) and 0.5 (blue). The horizontal red line denotes the experimental central value.

5 Summary

We discussed the sensitivity of the high-scale SUSY at 10-1000 TeV in the B^0 , B_s and K^0 meson systems. Furthermore, we have also discussed the sensitivity to the $D-\bar{D}$ mixing, the neutron EDM and the mercury EDM. In order to estimate the contribution of the squark flavor mixing to these FCNC, we calculate the squark mass spectrum, which is consistent with the recent Higgs discovery.

The SUSY contributions in ΔM_{B^0} and ΔM_{B_s} are at most 1.5% and 0.1% at $m_{\tilde{Q}} = 10$ TeV, respectively. As $m_{\tilde{Q}}$ increases, the SUSY contributions of both ΔM_{B^0} and ΔM_{B_s} decrease approximately with the power of $1/m_{\tilde{Q}}^2$. Therefore, the SUSY scale increases to more than 10 TeV, no signal of the SUSY is expected. On the other hand, the SUSY contribution in ΔM_{K^0} can be comparable to the experimental value in the case of $s_{13} = s_{23} = 0.5$ whereas it is suppressed in the case of $s_{13} = s_{23} = 0.22$ at $m_{\tilde{Q}} = 10$ TeV. Furthermore, the SUSY contribution in ϵ_K could be large, around 40% in the region of the SUSY scale 10-100 TeV. By considering the effect of the SUSY contribution $\mathcal{O}(10\%)$ in ϵ_K , the tension between ϵ_K and $\sin \phi_d$ can be relaxed even if the SUSY scale is 100 TeV.

The neutron EDM and the mercury EDM are also sensitive to the SUSY contribution induced by the gluino-squark interaction. The $|d_n|$ is expected to be close to the experimental upper bound even if the SUSY scale is 50 TeV. The predicted nEDM is roughly proportional to $|\epsilon_K^{\text{SUSY}}|$. If the SUSY contribution is the level of $\mathcal{O}(10\%)$ for ϵ_K , the $|d_n|$ is expected to be discovered in the region of 10^{-27} - 10^{-26} cm. For the $|d_{Hg}|$, the SUSY contribution is close to the experimental upper bound up to 200 TeV, which is much higher than the one of the nEDM. If the HgEDM is not observed above 10^{-29} cm, the SUSY contribution of ϵ_K is below a few %. Thus, the mercury EDM gives more significant information for the gluinosquark interaction compared with the neutron EDM. It may be important to give a comment that these predictions depend strongly on the assumptions of $\theta_{23}^L = \theta_{13}^L$ and $\theta_{ij}^L = \theta_{ij}^R$. The deviation from these relations destroys these correlations. In conclusion, the careful studies of the mixing angle relations are required to test the correlations between EDMs and ϵ_K^{SUSY} . The predicted EDMs have also the ambiguity with the factor 2-3 from the hadronic model. Since the SM prediction of ΔM_D at the short distance is $\mathcal{O}(10^{-18})$ GeV, which is very small compared with the experimental value, it is important to estimate the SUSY contribution of ΔM_D .

In conclusion, the more detailed studies of K^0 meson system, the EDMs of the neutron and mercury are required in order to probe the high-scale SUSY at 10-1000 TeV.

Acknowledgment

This work is supported by JSPS Grand-in-Aid for Scientific Research, 24654062 and 25-5222.

Appendix A : Running of SUSY particle masses

In the framework of the MSSM, one obtains the SUSY particle spectrum which is consistent with the observed Higgs mass. The numerical analyses have been given in Refs. [69, 70]. At the SUSY breaking scale Λ , the quadratic terms in the MSSM potential is given as

$$V_2 = m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_3^2 (H_1 \cdot H_2 + h.c.) .$$
⁽²⁸⁾

The mass eigenvalues at the H_1 and $\tilde{H}_2 \equiv \epsilon H_2^*$ system are given

$$m_{\mp}^2 = \frac{m_1^2 + m_2^2}{2} \mp \sqrt{\left(\frac{m_1^2 - m_2^2}{2}\right)^2 + m_3^4} .$$
 (29)

Suppose that the MSSM matches with the SM at the SUSY mass scale $Q_0 \equiv m_0$. Then, the smaller one m_-^2 is identified to be the mass squared of the SM Higgs H with the tachyonic mass. The larger one m_+^2 is the mass squared of the orthogonal combination \mathcal{H} , which is decoupled from the SM at Q_0 , that is, $m_{\mathcal{H}} \simeq Q_0$. Therefore, we have

$$m_{-}^2 = -m^2(Q_0) , \qquad m_{+}^2 = m_{\mathcal{H}}^2(Q_0) = m_1^2 + m_2^2 + m^2 , \qquad (30)$$

with

$$m_3^4 = (m_1^2 + m^2)(m_2^2 + m^2) , \qquad (31)$$

which leads to the mixing angle between H_1 and H_2 , β as follows:

$$\tan^2 \beta = \frac{m_1^2 + m^2}{m_2^2 + m^2} , \qquad H = \cos \beta H_1 + \sin \beta \tilde{H}_2 , \qquad \mathcal{H} = -\sin \beta H_1 + \cos \beta \tilde{H}_2 .$$
(32)

Thus, the Higgs mass parameter m^2 is expressed in terms of m_1^2 , m_2^2 and $\tan \beta$:

$$m^{2} = \frac{m_{1}^{2} - m_{2}^{2} \tan^{2} \beta}{\tan^{2} \beta - 1} .$$
(33)

Below the Q_0 scale, in which the SM emerges, the scalar potential is the SM one as follows:

$$V_{SM} = -m^2 |H|^2 + \frac{\lambda}{2} |H|^4 .$$
(34)

Here, the Higgs coupling λ is given in terms of the SUSY parameters at the leading order as

$$\lambda(Q_0) = \frac{1}{4} (g^2 + g'^2) \cos^2 2\beta + \frac{3h_t^2}{8\pi^2} X_t^2 \left(1 - \frac{X_t^2}{12} \right) , \qquad X_t = \frac{A_t(Q_0) - \mu(Q_0) \cot \beta}{Q_0} , \quad (35)$$

and h_t is the top Yukawa coupling of the SM. The parameters m_2 and λ run with the SM Renormalization Group Equation down to the electroweak scale $Q_{EW} = m_H$, and then give

$$m_H^2 = 2m^2(m_H) = \lambda(m_H)v^2$$
 (36)

It is easily seen that the VEV of Higgs, $\langle H \rangle$ is v, and $\langle \mathcal{H} \rangle = 0$, taking account of $\langle H_1 \rangle = v \cos \beta$ and $\langle H_2 \rangle = v \sin \beta$, where v = 246 GeV.

Let us fix $m_H = 125$ GeV, which gives $\lambda(Q_0)$ and $m^2(Q_0)$. This experimental input constrains the SUSY mass spectrum of the MSSM. We consider the some universal soft breaking parameters at the SUSY breaking scale Λ as follows:

$$m_{\tilde{Q}_{i}}(\Lambda) = m_{\tilde{U}_{i}^{c}}(\Lambda) = m_{\tilde{D}_{i}^{c}}(\Lambda) = m_{\tilde{L}_{i}}(\Lambda) = m_{\tilde{E}_{i}^{c}}(\Lambda) = m_{0}^{2} \ (i = 1, 2, 3) ,$$

$$M_{1}(\Lambda) = M_{2}(\Lambda) = M_{3}(\Lambda) = m_{1/2} , \qquad m_{1}^{2}(\Lambda) = m_{2}^{2}(\Lambda) = m_{0}^{2} ,$$

$$A_{U}(\Lambda) = A_{0}y_{U}(\Lambda) , \qquad A_{D}(\Lambda) = A_{0}y_{D}(\Lambda) , \qquad A_{E}(\Lambda) = A_{0}y_{E}(\Lambda) .$$
(37)

Therefore, there is no flavor mixing at Λ in the MSSM. However, in order to consider the non-minimal flavor mixing framework, we allow the off diagonal components of the squark mass matrices at the 10% level, which leads to the flavor mixing of order 0.1. We take these flavor mixing angles as free parameters at low energies.

Now, we have the SUSY five parameters, Λ , $\tan \beta$, m_0 , $m_{1/2}$, A_0 , where $Q_0 = m_0$. In addition to these parameters, we take $\mu = Q_0$. Inputing $m_H = 125$ GeV and taking $m_H \simeq Q_0$, we can obtain the SUSY spectrum for the fixed Q_0 and $\tan \beta$.

We present the SUSY mass spectrum at $Q_0 = 10$ TeV. The input parameter set and the obtained SUSY mass spectra at Q_0 are summarized in Table 1, where we use $\overline{m}_t(m_t) =$ 163.5 ± 2 GeV [60, 80]. These parameter sets are easily found from the work in Ref.[69].

Input at Λ and Q_0	Output at Q_0
at $\Lambda = 10^{17}$ GeV,	$m_{\tilde{g}} = 12.8 \text{ TeV}, \ m_{\tilde{W}} = 5.2 \text{ TeV}, \ m_{\tilde{B}} = 2.9 \text{ TeV}$
$m_0 = 10$ TeV,	$m_{\tilde{b}_L} = m_{\tilde{t}_L} = 12.2 \text{ TeV}$
$m_{1/2} = 6.2 \text{ TeV},$	$m_{\tilde{b}_R} = 14.1 \text{ TeV}, \ m_{\tilde{t}_R} = 8.4 \text{ TeV}$
$A_0 = 25.803 \text{ TeV};$	$m_{\tilde{s}_L,\tilde{d}_L} = m_{\tilde{c}_L,\tilde{u}_L} = 15.1 \text{ TeV}$
at $Q_0 = 10$ TeV,	$m_{\tilde{s}_R,\tilde{d}_R} \simeq m_{\tilde{c}_R,\tilde{u}_R} = 14.6 \text{ TeV}, \ m_{\mathcal{H}} = 13.7 \text{ TeV}$
$\mu = 10$ TeV,	$m_{\tilde{\tau}_L} = m_{\tilde{\nu}_{\tau_L}} = 10.4 \text{ TeV}, \ m_{\tilde{\tau}_R} = 9.3 \text{ TeV}$
$\tan\beta=10$	$m_{\tilde{\mu}_L, \tilde{e}_L} = m_{\tilde{\nu}_{\mu_L}, \tilde{\nu}_{e_L}} = 10.8 \text{ TeV}, \ m_{\tilde{\mu}_R, \tilde{e}_R} = 10.3 \text{ TeV}$
	$X_t = -0.22, \lambda_H = 0.126$

Table 1: Input parameters at Λ and the obtained SUSY spectra at $Q_0 = 10$ TeV.

As seen in Table 1, the first and second family squarks are degenerate in their masses, on the other hand, the third ones split due to the large RGE's effect. Therefore, the mixing angle between the first and second family squarks vanishes, but the mixing angles between the first-third and the second-third family squarks are produced at the Q_0 scale. The left-right mixing angle between \tilde{b}_L and \tilde{b}_R is given as

$$\theta \simeq \frac{m_b (A_b(Q_0) - \mu \tan \beta)}{m_{\tilde{b}_L}^2 - m_{\tilde{b}_R}^2} , \qquad (38)$$

which is very small, $\mathcal{O}(0.01)$ at 10 TeV. The lightest squark is the right-handed stop and the lightest gaugino is the Bino.

Appendix B : Squark contribution in $\Delta F = 2$ process

The $\Delta F = 2$ effective Lagrangian from the gluino-sbottom-quark interaction is given as [82]:

$$\mathcal{L}_{\text{eff}}^{\Delta F=2} = -\frac{1}{2} \left[C_{VLL} O_{VLL} + C_{VRR} O_{VRR} \right] - \frac{1}{2} \sum_{i=1}^{2} \left[C_{SLL}^{(i)} O_{SLL}^{(i)} + C_{SRR}^{(i)} O_{SRR}^{(i)} + C_{SLR}^{(i)} O_{SLR}^{(i)} \right],$$
(39)

where

$$O_{VLL} = (\bar{q}_a \gamma^{\mu} L Q^a) (\bar{q}_b \gamma^{\mu} L Q^b), \qquad O_{VRR} = (\bar{q}_a \gamma^{\mu} R Q^a) (\bar{q}_b \gamma^{\mu} R Q^b),$$

$$O_{SLL}^{(1)} = (\bar{q}_a L Q^a) (\bar{q}_b L Q^b), \qquad O_{SLL}^{(2)} = (\bar{q}_a L Q^b) (\bar{q}_b L Q^a),$$

$$O_{SRR}^{(1)} = (\bar{q}_a R Q^a) (\bar{q}_b R Q^b), \qquad O_{SRR}^{(2)} = (\bar{q}_a R Q^b) (\bar{q}_b R Q^a),$$

$$O_{SLR}^{(1)} = (\bar{q}_a L Q^a) (\bar{q}_b R Q^b), \qquad O_{SLR}^{(2)} = (\bar{q}_a L Q^b) (\bar{q}_b R Q^a),$$

$$O_{SLR}^{(1)} = (\bar{q}_a L Q^a) (\bar{q}_b R Q^b), \qquad O_{SLR}^{(2)} = (\bar{q}_a L Q^b) (\bar{q}_b R Q^a),$$

(40)

with $(P, Q, q) = (B^0, b, d)$, (B_s, b, s) , (K^0, s, d) . The L, R denote $(1 \pm \gamma_5)/2$, and a, b are color indices. Then, the $P^0 - \bar{P}^0$ mixing, M_{12} , is written as:

$$M_{12} = -\frac{1}{2m_P} \langle P^0 | \mathcal{L}_{\text{eff}}^{\Delta F=2} | \bar{P}^0 \rangle .$$

$$\tag{41}$$

The hadronic matrix elements are given in terms of the non-perturbative parameters B_i as:

$$\langle P^{0} | \mathcal{O}_{VLL} | \bar{P}^{0} \rangle = \frac{2}{3} m_{P}^{2} f_{P}^{2} B_{1}, \quad \langle P^{0} | \mathcal{O}_{VRR} | \bar{P}^{0} \rangle = \langle P^{0} | \mathcal{O}_{VLL} | \bar{P}^{0} \rangle,$$

$$\langle P^{0} | \mathcal{O}_{SLL}^{(1)} | \bar{P}^{0} \rangle = -\frac{5}{12} m_{P}^{2} f_{P}^{2} R_{P} B_{2}, \quad \langle P^{0} | \mathcal{O}_{SRR}^{(1)} | \bar{P}^{0} \rangle = \langle P^{0} | \mathcal{O}_{SLL}^{(1)} | \bar{P}^{0} \rangle,$$

$$\langle P^{0} | \mathcal{O}_{SLL}^{(2)} | \bar{P}^{0} \rangle = \frac{1}{12} m_{P}^{2} f_{P}^{2} R_{P} B_{3}, \quad \langle P^{0} | \mathcal{O}_{SRR}^{(2)} | \bar{P}^{0} \rangle = \langle P^{0} | \mathcal{O}_{SLL}^{(2)} | \bar{P}^{0} \rangle,$$

$$\langle P^{0} | \mathcal{O}_{SLR}^{(1)} | \bar{P}^{0} \rangle = \frac{1}{2} m_{P}^{2} f_{P}^{2} R_{P} B_{4}, \quad \langle P^{0} | \mathcal{O}_{SLR}^{(2)} | \bar{P}^{0} \rangle = \frac{1}{6} m_{P}^{2} f_{P}^{2} R_{P} B_{5},$$

$$(42)$$

where

$$R_P = \left(\frac{m_P}{m_Q + m_q}\right)^2. \tag{43}$$

The Wilson coefficients for the gluino contribution in Eq. (39) are written as [82]:

$$\begin{split} C_{VLL}(m_{\tilde{g}}) &= \frac{\alpha_s^2}{m_{\tilde{g}}^2} \sum_{I,J=1}^6 (\lambda_{GLL}^{(d)})_I^{ij} (\lambda_{GLL}^{(d)})_J^{ij} \left[\frac{11}{18} g_{2[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) + \frac{2}{9} g_{1[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) \right], \\ C_{VRR}(m_{\tilde{g}}) &= C_{VLL}(m_{\tilde{g}}) (L \leftrightarrow R), \\ C_{SRR}^{(1)}(m_{\tilde{g}}) &= \frac{\alpha_s^2}{m_{\tilde{g}}^2} \sum_{I,J=1}^6 (\lambda_{GLR}^{(d)})_I^{ij} (\lambda_{GLR}^{(d)})_J^{ij} \frac{17}{9} g_{1[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}), \\ C_{SLL}^{(1)}(m_{\tilde{g}}) &= C_{SRR}^{(1)}(m_{\tilde{g}}) (L \leftrightarrow R), \\ C_{SRR}^{(2)}(m_{\tilde{g}}) &= \frac{\alpha_s^2}{m_{\tilde{g}}^2} \sum_{I,J=1}^6 (\lambda_{GLR}^{(d)})_I^{ij} (\lambda_{GLR}^{(d)})_J^{ij} \left(-\frac{1}{3} \right) g_{1[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}), \\ C_{SLL}^{(2)}(m_{\tilde{g}}) &= C_{SRR}^{(2)}(m_{\tilde{g}}) (L \leftrightarrow R), \\ \end{array}$$

$$C_{SLR}^{(1)}(m_{\tilde{g}}) = \frac{\alpha_s^2}{m_{\tilde{g}}^2} \sum_{I,J=1}^6 \left\{ (\lambda_{GLR}^{(d)})_I^{ij} (\lambda_{GRL}^{(d)})_J^{ij} \left(-\frac{11}{9} \right) g_{2[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) + (\lambda_{GLL}^{(d)})_I^{ij} (\lambda_{GRR}^{(d)})_J^{ij} \left[\frac{14}{3} g_{1[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) - \frac{2}{3} g_{2[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) \right] \right\},$$

$$C_{SLR}^{(2)}(m_{\tilde{g}}) = \frac{\alpha_s^2}{m_{\tilde{g}}^2} \sum_{I,J=1}^6 \left\{ (\lambda_{GLR}^{(d)})_I^{ij} (\lambda_{GRL}^{(d)})_J^{ij} \left(-\frac{5}{3} \right) g_{2[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) + (\lambda_{GLL}^{(d)})_I^{ij} (\lambda_{GRR}^{(d)})_J^{ij} \left[\frac{2}{9} g_{1[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) + \frac{10}{9} g_{2[1]}(x_I^{\tilde{g}}, x_J^{\tilde{g}}) \right] \right\},$$

$$(44)$$

where

$$(\lambda_{GLL}^{(d)})_{K}^{ij} = (\Gamma_{GL}^{(d)\dagger})_{i}^{K} (\Gamma_{GL}^{(d)})_{K}^{j} , \quad (\lambda_{GRR}^{(d)})_{K}^{ij} = (\Gamma_{GR}^{(d)\dagger})_{i}^{K} (\Gamma_{GR}^{(d)})_{K}^{j} , (\lambda_{GLR}^{(d)})_{K}^{ij} = (\Gamma_{GL}^{(d)\dagger})_{i}^{K} (\Gamma_{GR}^{(d)})_{K}^{j} , \quad (\lambda_{GRL}^{(d)})_{K}^{ij} = (\Gamma_{GR}^{(d)\dagger})_{i}^{K} (\Gamma_{GL}^{(d)})_{K}^{j} .$$

$$(45)$$

Here we take (i, j) = (1, 3), (2, 3), (1, 2) which correspond to B^0 , B_s , and K^0 mesons, respectively. The loop functions are given as follows:

• If
$$x_{I}^{\tilde{g}} \neq x_{J}^{\tilde{g}} (x_{I,J}^{\tilde{g}} = m_{\tilde{d}_{I,J}}^{2}/m_{\tilde{g}}^{2}),$$

 $g_{1[1]}(x_{I}^{\tilde{g}}, x_{J}^{\tilde{g}}) = \frac{1}{x_{I}^{\tilde{g}} - x_{J}^{\tilde{g}}} \left(\frac{x_{I}^{\tilde{g}} \log x_{I}^{\tilde{g}}}{(x_{I}^{\tilde{g}} - 1)^{2}} - \frac{1}{x_{I}^{\tilde{g}} - 1} - \frac{x_{J}^{\tilde{g}} \log x_{J}^{\tilde{g}}}{(x_{J}^{\tilde{g}} - 1)^{2}} + \frac{1}{x_{J}^{\tilde{g}} - 1} \right),$
 $g_{2[1]}(x_{I}^{\tilde{g}}, x_{J}^{\tilde{g}}) = \frac{1}{x_{I}^{\tilde{g}} - x_{J}^{\tilde{g}}} \left(\frac{(x_{I}^{\tilde{g}})^{2} \log x_{I}^{\tilde{g}}}{(x_{I}^{\tilde{g}} - 1)^{2}} - \frac{1}{x_{I}^{\tilde{g}} - 1} - \frac{(x_{J}^{\tilde{g}})^{2} \log x_{J}^{\tilde{g}}}{(x_{J}^{\tilde{g}} - 1)^{2}} + \frac{1}{x_{J}^{\tilde{g}} - 1} \right).$ (46)

• If $x_I^{\tilde{g}} = x_J^{\tilde{g}}$,

$$g_{1[1]}(x_{I}^{\tilde{g}}, x_{I}^{\tilde{g}}) = -\frac{(x_{I}^{\tilde{g}} + 1)\log x_{I}^{\tilde{g}}}{(x_{I}^{\tilde{g}} - 1)^{3}} + \frac{2}{(x_{I}^{\tilde{g}} - 1)^{2}} ,$$

$$g_{2[1]}(x_{I}^{\tilde{g}}, x_{I}^{\tilde{g}}) = -\frac{2x_{I}^{\tilde{g}}\log x_{I}^{\tilde{g}}}{(x_{I}^{\tilde{g}} - 1)^{3}} + \frac{x_{I}^{\tilde{g}} + 1}{(x_{I}^{\tilde{g}} - 1)^{2}} .$$
(47)

Taking account of the case that the gluino mass is much smaller than the squark mass scale Q_0 , the effective Wilson coefficients are given by using the RGEs for higher-dimensional operators in Eq.(39) at the leading order of QCD as follows:

$$C_{VLL}(m_b(\Lambda = 2 \text{ GeV})) = \eta_{VLL}^{B(K)} C_{VLL}(Q_0), \quad C_{VRR}(m_b(\Lambda = 2 \text{ GeV})) = \eta_{VRR}^{B(K)} C_{VLL}(Q_0),$$

$$\begin{pmatrix} C_{SLL}^{(1)}(m_b(\Lambda = 2 \text{ GeV})) \\ C_{SLL}^{(2)}(m_b(\Lambda = 2 \text{ GeV})) \end{pmatrix} = \begin{pmatrix} C_{SLL}^{(1)}(Q_0) \\ C_{SLL}^{(2)}(Q_0) \end{pmatrix} X_{LL}^{-1} \eta_{LL}^{B(K)} X_{LL},$$

$$\begin{pmatrix} C_{SRR}^{(1)}(m_b(\Lambda = 2 \text{ GeV})) \\ C_{SRR}^{(2)}(m_b(\Lambda = 2 \text{ GeV})) \end{pmatrix} = \begin{pmatrix} C_{SRR}^{(1)}(Q_0) \\ C_{SRR}^{(2)}(Q_0) \end{pmatrix} X_{RR}^{-1} \eta_{RR}^{B(K)} X_{RR},$$

$$\begin{pmatrix} C_{SLR}^{(1)}(m_b(\Lambda = 2 \text{ GeV})) \\ C_{SLR}^{(2)}(m_b(\Lambda = 2 \text{ GeV})) \end{pmatrix} = \begin{pmatrix} C_{SLR}^{(1)}(Q_0) \\ C_{SLR}^{(2)}(Q_0) \end{pmatrix} X_{LR}^{-1} \eta_{LR}^{B(K)} X_{LR},$$
(48)

where

$$\begin{split} \eta^{B}_{VLL} &= \eta^{B}_{VRR} = \left(\frac{\alpha_{s}(Q_{0})}{\alpha_{s}(\tilde{g})}\right)^{\frac{6}{15}} \left(\frac{\alpha_{s}(m_{\tilde{g}})}{\alpha_{s}(m_{t})}\right)^{\frac{6}{21}} \left(\frac{\alpha_{s}(m_{t})}{\alpha_{s}(m_{b})}\right)^{\frac{6}{23}}, \\ \eta^{B}_{LL} &= \eta^{B}_{RR} = S_{LL} \begin{pmatrix} \eta^{d^{1}_{LL}}_{b\tilde{g}} & 0\\ 0 & \eta^{d^{2}_{LL}}_{b\tilde{g}} \end{pmatrix} S^{-1}_{LL}, \quad \eta^{B}_{LR} = S_{LR} \begin{pmatrix} \eta^{d^{1}_{LR}}_{b\tilde{g}} & 0\\ 0 & \eta^{d^{2}_{LR}}_{b\tilde{g}} \end{pmatrix} S^{-1}_{LR}, \\ \eta_{b\tilde{g}} &= \left(\frac{\alpha_{s}(Q_{0})}{\alpha_{s}(m_{\tilde{g}})}\right)^{\frac{1}{10}} \left(\frac{\alpha_{s}(m_{\tilde{g}})}{\alpha_{s}(m_{t})}\right)^{\frac{1}{14}} \left(\frac{\alpha_{s}(m_{t})}{\alpha_{s}(m_{b})}\right)^{\frac{3}{46}}, \\ \eta^{K}_{VLL} &= \eta^{K}_{VRR} = \left(\frac{\alpha_{s}(Q_{0})}{\alpha_{s}(m_{\tilde{g}})}\right)^{\frac{6}{15}} \left(\frac{\alpha_{s}(m_{\tilde{g}})}{\alpha_{s}(m_{t})}\right)^{\frac{6}{21}} \left(\frac{\alpha_{s}(m_{t})}{\alpha_{s}(m_{b})}\right)^{\frac{6}{23}} \left(\frac{\alpha_{s}(m_{b})}{\alpha_{s}(\Lambda = 2 \text{ GeV})}\right)^{\frac{6}{25}}, \end{split}$$

$$\eta_{LL}^{K} = \eta_{RR}^{K} = S_{LL} \begin{pmatrix} \eta_{\Lambda\tilde{g}}^{d_{LL}} & 0\\ 0 & \eta_{\Lambda\tilde{g}}^{d_{LL}} \end{pmatrix} S_{LL}^{-1}, \qquad \eta_{LR}^{K} = S_{LR} \begin{pmatrix} \eta_{\Lambda\tilde{g}}^{d_{LR}} & 0\\ 0 & \eta_{\Lambda\tilde{g}}^{d_{LR}} \end{pmatrix} S_{LR}^{-1},$$

$$\eta_{\Lambda\tilde{g}} = \begin{pmatrix} \alpha_s(Q_0)\\ \alpha_s(m_{\tilde{g}}) \end{pmatrix}^{\frac{1}{10}} \begin{pmatrix} \alpha_s(m_{\tilde{g}})\\ \alpha_s(m_t) \end{pmatrix}^{\frac{1}{14}} \begin{pmatrix} \alpha_s(m_t)\\ \alpha_s(m_b) \end{pmatrix}^{\frac{3}{46}} \begin{pmatrix} \alpha_s(m_b)\\ \alpha_s(\Lambda = 2 \text{ GeV}) \end{pmatrix}^{\frac{3}{50}},$$

$$d_{LL}^1 = \frac{2}{3}(1 - \sqrt{241}), \qquad d_{LL}^2 = \frac{2}{3}(1 + \sqrt{241}), \qquad d_{LR}^1 = -16, \qquad d_{LR}^2 = 2,$$

$$S_{LL} = \begin{pmatrix} \frac{16 + \sqrt{241}}{60} & \frac{16 - \sqrt{241}}{60}\\ 1 & 1 \end{pmatrix}, \qquad S_{LR} = \begin{pmatrix} -2 & 1\\ 3 & 0 \end{pmatrix},$$

$$X_{LL} = X_{RR} = \begin{pmatrix} 1 & 0\\ 4 & 8 \end{pmatrix}, \qquad X_{LR} = \begin{pmatrix} 0 & -2\\ 1 & 0 \end{pmatrix}.$$

(49)

For the parameters $B_i^{(d)}(i=2-5)$ of B mesons, we use values in [83] as follows:

$$B_{2}^{(B_{d})}(m_{b}) = 0.79(2)(4), \qquad B_{3}^{(B_{d})}(m_{b}) = 0.92(2)(4), B_{4}^{(B_{d})}(m_{b}) = 1.15(3)\binom{+5}{-7}, \qquad B_{5}^{(B_{d})}(m_{b}) = 1.72(4)\binom{+20}{-6}, B_{2}^{(B_{s})}(m_{b}) = 0.80(1)(4), \qquad B_{3}^{(B_{s})}(m_{b}) = 0.93(3)(8), B_{4}^{(B_{s})}(m_{b}) = 1.16(2)\binom{+5}{-7}, \qquad B_{5}^{(B_{s})}(m_{b}) = 1.75(3)\binom{+21}{-6}.$$
(50)

On the other hand, we use the most updated values for $\hat{B}_1^{(d)}$ and $\hat{B}_1^{(s)}$ as [80]:

$$\hat{B}_1^{(B_s)} = 1.33 \pm 0.06 , \qquad \hat{B}_1^{(B_s)} / \hat{B}_1^{(B_d)} = 1.05 \pm 0.07 .$$
 (51)

For the paremeters $B_i^K (i = 2 - 5)$, we use following values [84],

$$B_2^{(K)}(2\text{GeV}) = 0.66 \pm 0.04, \qquad B_3^{(K)}(2\text{GeV}) = 1.05 \pm 0.12, B_4^{(K)}(2\text{GeV}) = 1.03 \pm 0.06, \qquad B_5^{(K)}(2\text{GeV}) = 0.73 \pm 0.10,$$
(52)

and we take recent value of Eq.(17) for deriving $B_1^{(K)}(2\text{GeV})$. For the paremeters $B_i^D(i = 1 - 5)$, we use following values [85, 86],

$$B_1^{(D)}(3\text{GeV}) = 0.75 \pm 0.02, \quad B_2^{(D)}(3\text{GeV}) = 0.66 \pm 0.02, \quad B_3^{(D)}(3\text{GeV}) = 0.96 \pm 0.05, \\ B_4^{(D)}(3\text{GeV}) = 0.91 \pm 0.04, \quad B_5^{(D)}(3\text{GeV}) = 1.10 \pm 0.05.$$
(53)

Appendix C : The loop functions F_i

The loop functions $F_i(x_{\tilde{g}}^I)$ are given in terms of $x_{\tilde{g}}^I = m_{\tilde{g}}^2/m_{\tilde{d}_I}^2$ (I = 3, 6) as follows:

$$F_{1}(x_{\tilde{g}}^{I}) = \frac{x_{\tilde{g}}^{I}\log x_{\tilde{g}}^{I}}{2(x_{\tilde{g}}^{I}-1)^{4}} + \frac{(x_{\tilde{g}}^{I})^{2} - 5x_{\tilde{g}}^{I} - 2}{12(x_{\tilde{g}}^{I}-1)^{3}} , \quad F_{2}(x_{\tilde{g}}^{I}) = -\frac{(x_{\tilde{g}}^{I})^{2}\log x_{\tilde{g}}^{I}}{2(x_{\tilde{g}}^{I}-1)^{4}} + \frac{2(x_{\tilde{g}}^{I})^{2} + 5x_{\tilde{g}}^{I} - 1}{12(x_{\tilde{g}}^{I}-1)^{3}} , \\ F_{3}(x_{\tilde{g}}^{I}) = \frac{\log x_{\tilde{g}}^{I}}{(x_{\tilde{g}}^{I}-1)^{3}} + \frac{x_{\tilde{g}}^{I} - 3}{2(x_{\tilde{g}}^{I}-1)^{2}} , \quad F_{4}(x_{\tilde{g}}^{I}) = -\frac{x_{\tilde{g}}^{I}\log x_{\tilde{g}}^{I}}{(x_{\tilde{g}}^{I}-1)^{3}} + \frac{x_{\tilde{g}}^{I} + 1}{2(x_{\tilde{g}}^{I}-1)^{2}} = \frac{1}{2}g_{2[1]}(x_{\tilde{g}}^{I}, x_{\tilde{g}}^{I}) .$$

$$(54)$$

Appendix D : EDM and Chromo-EDM of quarks

We present the EDM of the strange quark from the gluino contribution as the typical example [82]:

$$d_s(Q_0) = -2\sqrt{4\pi\alpha(m_{\tilde{g}})} \text{Im}[A_s^{\gamma 22}(Q_0)],$$
(55)

where

$$A_{s}^{\gamma 22}(Q_{0}) = \frac{Q_{s}\alpha_{s}(m_{\tilde{g}})}{4\pi} \frac{8}{3} \sum_{I=1}^{6} \frac{1}{2m_{\tilde{d}_{I}}^{2}} \left\{ \left(m_{s}(\lambda_{GLL}^{(d)})_{3}^{22} + m_{s}(\lambda_{GRR}^{(d)})_{I}^{22} \right) \left(F_{2}(x_{\tilde{g}}^{I}) \right) + m_{\tilde{g}}(\lambda_{GLR}^{(d)})_{I}^{22} \left(F_{4}(x_{\tilde{g}}^{I}) \right) \right\}.$$
(56)

On the other hand, the chromo-EDM (cEDM) of the strange quark from gluino contribution is given as:

$$d_s^C(Q_0) = -2\sqrt{4\pi\alpha_s(m_{\tilde{g}})} \operatorname{Im}[A_s^{g22}(Q_0)],$$
(57)

where

$$A_{s}^{g22}(Q_{0}) = -\frac{\alpha_{s}(m_{\tilde{g}})}{4\pi} \frac{1}{3} \sum_{I=1}^{6} \frac{1}{2m_{\tilde{d}_{I}}^{2}} \left\{ \left(m_{s}(\lambda_{GLL}^{(d)})_{I}^{22} + m_{s}(\lambda_{GRR}^{(d)})_{I}^{22} \right) \left(9F_{1}(x_{\tilde{g}}^{I}) + F_{2}(x_{\tilde{g}}^{I}) \right) + m_{\tilde{g}}(\lambda_{GLR}^{(d)})_{I}^{22} \left(9F_{3}(x_{\tilde{g}}^{I}) + F_{4}(x_{\tilde{g}}^{I}) \right) \right\}$$

$$(58)$$

(59)

Including the RGE effect of QCD [87], the cEDM of the strange quark is given as

$$d_s^C(2\text{GeV}) = d_s^C(Q_0) \left(\frac{\alpha_s(Q_0)}{\alpha_s(m_t)}\right)^{\frac{14}{21}} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_b)}\right)^{\frac{14}{23}} \left(\frac{\alpha_s(m_b)}{\alpha_s(2\text{GeV})}\right)^{\frac{14}{25}} .$$
 (60)

On the other hand, the EDM operator is mixied with the cEDM operator during RGE evolution. Then, one obtains

$$d_s(2\text{GeV}) = d_s(Q_0) \left(\frac{\alpha_s(Q_0)}{\alpha_s(m_t)}\right)^{\frac{16}{21}} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_b)}\right)^{\frac{16}{23}} \left(\frac{\alpha_s(m_b)}{\alpha_s(2\text{GeV})}\right)^{\frac{16}{25}} + \frac{8}{g_s} d_s^C(Q_0) \times$$
(61)

$$\left[\left(\frac{\alpha_s(Q_0)}{\alpha_s(m_t)}\right)^{\frac{16}{21}} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_b)}\right)^{\frac{16}{23}} \left(\frac{\alpha_s(m_b)}{\alpha_s(2\text{GeV})}\right)^{\frac{16}{25}} - \left(\frac{\alpha_s(Q_0)}{\alpha_s(m_t)}\right)^{\frac{14}{21}} \left(\frac{\alpha_s(m_t)}{\alpha_s(m_b)}\right)^{\frac{14}{23}} \left(\frac{\alpha_s(m_b)}{\alpha_s(2\text{GeV})}\right)^{\frac{14}{25}} \right].$$

The EDMs and cEDMs of the down- and up-quarks induced by the gluino interaction are also given by the similar formulas.

References

- G. Aad *et al.* [ATLAS Collaboration], JHEP **1409** (2014) 176 [arXiv:1405.7875 [hepex]].
- [2] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1406** (2014) 055 [arXiv:1402.4770 [hep-ex]].
- [3] G. Aad et al. [ATLAS Collaboration], JHEP 1411 (2014) 118 [arXiv:1407.0583 [hepex]].
- [4] G. Aad et al. [ATLAS Collaboration], Phys. Lett. B 716 (2012) 1 [arXiv:1207.7214];
 S. Chatrchyan et al. [CMS Collaboration], Phys. Lett. B 716 (2012) 30 [arXiv:1207.7235];
 G. Aad et al. [ATLAS and CMS Collaborations], arXiv:1503.07589 [hep-ex].
- [5] G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP 9812, 027 (1998) [hep-ph/9810442].
- [6] J. D. Wells, hep-ph/0306127.
- [7] N. Arkani-Hamed and S. Dimopoulos, JHEP **0506**, 073 (2005) [hep-th/0405159].
- [8] G. F. Giudice and A. Romanino, Nucl. Phys. B 699, 65 (2004) [Erratum-ibid. B 706, 65 (2005)] [hep-ph/0406088].
- [9] N. Arkani-Hamed, S. Dimopoulos, G. F. Giudice and A. Romanino, Nucl. Phys. B 709, 3 (2005) [hep-ph/0409232].
- [10] J. D. Wells, Phys. Rev. D 71, 015013 (2005) [hep-ph/0411041].
- [11] M. Ibe, T. Moroi and T. T. Yanagida, Phys. Lett. B 644 (2007) 355 [hep-ph/0610277].
- [12] L. J. Hall and Y. Nomura, JHEP **1003**, 076 (2010) [arXiv:0910.2235 [hep-ph]].
- [13] L. J. Hall and Y. Nomura, JHEP **1201**, 082 (2012) [arXiv:1111.4519 [hep-ph]].
- [14] M. Ibe and T. T. Yanagida, Phys. Lett. B **709** (2012) 374 [arXiv:1112.2462 [hep-ph]].
- [15] M. Ibe, S. Matsumoto and T. T. Yanagida, Phys. Rev. D 85 (2012) 095011 [arXiv:1202.2253 [hep-ph]].
- [16] A. Arvanitaki, N. Craig, S. Dimopoulos and G. Villadoro, JHEP **1302** (2013) 126 [arXiv:1210.0555 [hep-ph]].
- [17] N. Arkani-Hamed, A. Gupta, D. E. Kaplan, N. Weiner and T. Zorawski, arXiv:1212.6971 [hep-ph].

- [18] J. L. Evans, M. Ibe, K. A. Olive and T. T. Yanagida, Eur. Phys. J. C 73 (2013) 2468 [arXiv:1302.5346 [hep-ph]].
- [19] J. Hisano, T. Kuwahara and N. Nagata, Phys. Lett. B 723 (2013) 324 [arXiv:1304.0343 [hep-ph]].
- [20] N. Nagata, H. Otono and S. Shirai, arXiv:1504.00504 [hep-ph].
- [21] W. Altmannshofer, R. Harnik and J. Zupan, JHEP 1311 (2013) 202 [arXiv:1308.3653 [hep-ph]].
- [22] T. Moroi and M. Nagai, Phys. Lett. B **723** (2013) 107 [arXiv:1303.0668 [hep-ph]].
- [23] D. McKeen, M. Pospelov and A. Ritz, Phys. Rev. D 87 (2013) 11, 113002 [arXiv:1303.1172 [hep-ph]].
- [24] R. Aaij *et al.* [LHCb Collaboration], Eur. Phys. J. C **73** (2013) 4, 2373 [arXiv:1208.3355].
- [25] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110** (2013) 24, 241802 [arXiv:1303.7125].
- [26] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110** (2013) 22, 221601 [arXiv:1304.6173].
- [27] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. D 87 (2013) 11, 112010 [arXiv:1304.2600].
- [28] R. Aaij et al. [LHCb Collaboration], JHEP 1307 (2013) 084 [arXiv:1305.2168].
- [29] R. Aaij *et al.* [LHCb Collaboration], JHEP **1308** (2013) 131 [arXiv:1304.6325, arXiv:1304.6325].
- [30] R. Aaij et al. [LHCb Collaboration], New J. Phys. 15, 053021 (2013) [arXiv:1304.4741].
- [31] M. Vesterinen [LHCb Collaboration], PoS Beauty **2013** (2013) 005 [arXiv:1306.0092].
- [32] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110** (2013) 021801 [arXiv:1211.2674].
- [33] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **110** (2013) 031801 [arXiv:1210.4492].
- [34] Y. Amhis *et al.* [Heavy Flavor Averaging Group], arXiv:1207.1158 [hep-ex].
- [35] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108** (2012) 101803 [arXiv:1112.3183].
- [36] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett. B **707** (2012) 497 [arXiv:1112.3056].

- [37] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49 (1973) 652.
- [38] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nucl. Phys. B 477 (1996) 321 [hep-ph/9604387].
- [39] M. Tanimoto and K. Yamamoto, Phys. Lett. B 735 (2014) 426 [arXiv:1404.0520 [hepph]].
- [40] J. Charles, S. Descotes-Genon, Z. Ligeti, S. Monteil, M. Papucci and K. Trabelsi, Phys. Rev. D 89 (2014) 3, 033016 [arXiv:1309.2293 [hep-ph]].
- [41] S. F. King, JHEP **1009** (2010) 114 [arXiv:1006.5895 [hep-ph]].
- [42] M. Endo, S. Shirai and T. T. Yanagida, Prog. Theor. Phys. 125 (2011) 921 [arXiv:1009.3366 [hep-ph]].
- [43] M. Endo and N. Yokozaki, JHEP **1103** (2011) 130 [arXiv:1012.5501 [hep-ph]].
- [44] J. Kubo and A. Lenz, Phys. Rev. D 82 (2010) 075001 [arXiv:1007.0680].
- [45] Y. Kaburaki, K. Konya, J. Kubo and A. Lenz, Phys. Rev. D 84 (2011) 016007 [arXiv:1012.2435].
- [46] J. Virto, JHEP **0911** (2009) 055 [arXiv:0907.5376 [hep-ph]].
- [47] J. Virto, JHEP **1201** (2012) 120 [arXiv:1111.0940 [hep-ph]].
- [48] P. Ko and J. -h. Park, Phys. Rev. D 80 (2009) 035019 [arXiv:0809.0705 [hep-ph]].
- [49] P. Ko and J. -h. Park, Phys. Rev. D 82 (2010) 117701 [arXiv:1006.5821 [hep-ph]].
- [50] R. -M. Wang, Y. -G. Xu, Q. Chang and Y. -D. Yang, Phys. Rev. D 83 (2011) 095010 [arXiv:1102.2031 [hep-ph]].
- [51] J. K. Parry, Phys. Lett. B **694** (2011) 363 [arXiv:1006.5331 [hep-ph]].
- [52] A. Hayakawa, Y. Shimizu, M. Tanimoto and K. Yamamoto, Phys. Lett. B 710 (2012) 446 [arXiv:1202.0486 [hep-ph]].
- [53] Y. Shimizu, M. Tanimoto and K. Yamamoto, Prog. Theor. Phys. 128 (2012) 273 [arXiv:1205.1705].
- [54] Y. Shimizu, M. Tanimoto and K. Yamamoto, Phys. Rev. D 87 (2013) 056004 [arXiv:1212.6486].
- [55] Y. Shimizu, M. Tanimoto and K. Yamamoto, Phys. Lett. B 726 (2013) 747 [arXiv:1307.0374].
- [56] A. Hayakawa, Y. Shimizu, M. Tanimoto and K. Yamamoto, PTEP 2014 (2014) 023B04, arXiv:1311.5974 [hep-ph].

- [57] F. Mescia and J. Virto, Phys. Rev. D 86 (2012) 095004 [arXiv:1208.0534 [hep-ph]].
- [58] M. Tanimoto and K. Yamamoto, arXiv:1503.06270 [hep-ph].
- [59] K. Fuyuto, J. Hisano, N. Nagata and K. Tsumura, JHEP **1312** (2013) 010 [arXiv:1308.6493].
- [60] K.A. Olive et al. (Particle Data Group), Chin. Phys. C, 38, 090001 (2014).
- [61] M. Pospelov and A. Ritz, Phys. Rev. D 63 (2001) 073015 [hep-ph/0010037].
- [62] J. Hisano and Y. Shimizu, Phys. Lett. B 581 (2004) 224 [hep-ph/0308255].
- [63] J. Hisano and Y. Shimizu, Phys. Rev. D 70 (2004) 093001 [hep-ph/0406091].
- [64] J. Hisano, M. Nagai and P. Paradisi, Phys. Rev. D 80 (2009) 095014 [arXiv:0812.4283].
- [65] J. Hisano, J. Y. Lee, N. Nagata and Y. Shimizu, Phys. Rev. D 85 (2012) 114044 [arXiv:1204.2653 [hep-ph]].
- [66] K. Fuyuto, J. Hisano and N. Nagata, Phys. Rev. D 87 (2013) 054018 [arXiv:1211.5228].
- [67] W. C. Griffith, M. D. Swallows, T. H. Loftus, M. V. Romalis, B. R. Heckel and E. N. Fortson, Phys. Rev. Lett. **102** (2009) 101601.
- [68] C. C. Chiou, O. C. W. Kong and R. D. Vaidya, Phys. Rev. D 76 (2007) 013003 [arXiv:0705.3939 [hep-ph]].
- [69] A. Delgado, M. Garcia and M. Quiros, Phys. Rev. D 90 (2014) 1, 015016 [arXiv:1312.3235].
- [70] G. F. Giudice and R. Rattazzi, Nucl. Phys. B **757** (2006) 19 [hep-ph/0606105].
- [71] For example, see S. P. Martin, "A Supersymmetry primer," in *Kane, G.L. (ed.): Perspectives on supersymmetry II* 1-153 [hep-ph/9709356].
- [72] For example, see S. Iso, arXiv:1304.0293 [hep-ph];
 S. Iso and Y. Orikasa, PTEP **2013** (2013) 023B08 [arXiv:1210.2848 [hep-ph]];
 L. g. Bian, arXiv:1303.2402 [hep-ph].
- [73] A. J. Buras and D. Guadagnoli, Phys. Rev. D 78 (2008) 033005 [arXiv:0805.3887].
- [74] T. Inami and C. S. Lim, Prog. Theor. Phys. 65 (1981) 297 [Erratum-ibid. 65 (1981) 1772].
- [75] T. Bae, Y. -C. Jang, H. Jeong, J. Kim, J. Kim, K. Kim, S. Kim and W. Lee *et al.*, PoS LATTICE **2013**, 476 (2013) [arXiv:1310.7319 [hep-lat]].
- [76] S. Aoki, Y. Aoki, C. Bernard, T. Blum, G. Colangelo, M. Della Morte, S. Durr and A. X. E. Khadra *et al.*, arXiv:1310.8555 [hep-lat].

- [77] T. Falk, K. A. Olive, M. Pospelov and R. Roiban, Nucl. Phys. B 560 (1999) 3 [hep-ph/9904393].
- [78] J. Hisano, M. Kakizaki, M. Nagai and Y. Shimizu, Phys. Lett. B 604 (2004) 216 [hep-ph/0407169].
- [79] A. J. Bevan et al. [UTfit Collaboration], JHEP 1403 (2014) 123 [arXiv:1402.1664 [hep-ph]].
- [80] UTfit Collaboration (http://www.utfit.org).
- [81] H. Y. Cheng and C. W. Chiang, Phys. Rev. D 81 (2010) 114020 [arXiv:1005.1106 [hep-ph]].
- [82] T. Goto, http://research.kek.jp/people/tgoto/ .
- [83] D. Becirevic, V. Gimenez, G. Martinelli, M. Papinutto and J. Reyes, JHEP 0204 (2002) 025 [hep-lat/0110091].
- [84] C. R. Allton, L. Conti, A. Donini, V. Gimenez, L. Giusti, G. Martinelli, M. Talevi and A. Vladikas, Phys. Lett. B 453 (1999) 30 [hep-lat/9806016].
- [85] A. J. Buras, M. Misiak and J. Urban, Nucl. Phys. B 586 (2000) 397 [hep-ph/0005183].
- [86] N. Carrasco, M. Ciuchini, P. Dimopoulos, R. Frezzotti, V. Gimenez, V. Lubicz, G. C. Rossi and F. Sanfilippo *et al.*, Phys. Rev. D **90** (2014) 1, 014502 [arXiv:1403.7302 [hep-lat]].
- [87] G. Degrassi, E. Franco, S. Marchetti and L. Silvestrini, JHEP 0511 (2005) 044 [hep-ph/0510137].