

Cluster automorphism groups and automorphism groups of exchange graphs

Wen Chang¹ and Bin Zhu²

Abstract

For a coefficient free cluster algebra \mathcal{A} , we study the cluster automorphism group $Aut(\mathcal{A})$ and the automorphism group $Aut(E_{\mathcal{A}})$ of its exchange graph $E_{\mathcal{A}}$. We show that these two groups are isomorphic with each other, if \mathcal{A} is of finite type excepting types of rank two and type F_4 , or if \mathcal{A} is of skew-symmetric finite mutation type.

Key words. Cluster algebras; Exchange graphs; Cluster automorphism groups.

Mathematics Subject Classification. 13F60; 05E40

1 Introduction

Cluster algebras are introduced by Sergey Fomin and Andrei Zelevinsky in [10]. In this paper we consider cluster algebras with trivial coefficients, which can be defined through a skew-symmetrizable square matrix. Such a cluster algebra is a \mathbb{Z} -subalgebra of rational function field with n indeterminates. More precisely, a *seed* is a pair consisting of a set (*cluster*) of n indeterminates (*cluster variables*) in the field and a skew-symmetrizable square matrix (*exchange matrix*) of size n . Starting from an initial seed, we get a new seed by an operate so called mutation. Then the cluster algebra is algebraic-generated by all the cluster variables obtained by iterated mutations. The cluster algebra has nice combinatorial structures which are (in some sense) given by mutations, and these structures are captured by its exchange graph, which is a graph with seeds as vertices and with mutations as edges.

We focus in this paper on two special types of cluster algebras: the *finite type* and the *finite mutation type*. Cluster algebras of finite type are those algebras with finite number of clusters. They are classified in [11], which corresponds to the Killing-Cartan classification of complex semisimple Lie algebras, equivalently, corresponds to the classification of root systems in Euclidean space. If there are finite many matrix classes in the seeds of a cluster algebra, then we say it is of finite mutation type, where two matrices are in the same class if one of them can be obtained from the other by simultaneous relabeling of the rows and columns. The cluster algebras of finite mutation type with skew-symmetric exchange matrices are classified in [9], a large class of them arises from marked Riemann surfaces (possibly with boundary) [7], and there are 11 exceptional ones. The classification of skew-symmetrizable cluster algebras of finite mutation type is given in [8] via operations so called unfoldings upon the skew-symmetric cluster algebras of finite mutation type.

We consider the relations in this paper between two groups associated to the cluster algebras. One is the cluster automorphism group consisting of *cluster automorphisms*, which are permutations of the clusters that commute with mutations. This group is introduced in [1] for a coefficient free

¹Supported by Shaanxi Province, Shaanxi Normal University and the NSF of China (Grant 11601295)

²Supported by the NSF of China (Grant 11671221)

cluster algebra, and in [4] for a cluster algebra with coefficients, it reveals the combinatorial and algebraic symmetries of the cluster algebra. Another is the automorphism group of the exchange graph, which consists of graph automorphism of the exchange graph. This group describes the symmetries of the exchange graph, in other words, describes combinatorial symmetries of the cluster algebra. The problem that consider the relations between these two groups is stated in [16].

The exchange graph is a fairly coarse invariant of a cluster algebra, e.g. all infinite type cluster algebras of rank 2 have the same exchange graph. This article suggests that, nonetheless, the exchange graph is already rich enough to capture most of the symmetries of the cluster algebra.

For a coefficient free cluster algebra \mathcal{A} with exchange graph $E_{\mathcal{A}}$, we write the cluster automorphism group of \mathcal{A} and the automorphism group of $E_{\mathcal{A}}$ as $Aut(\mathcal{A})$ and $Aut(E_{\mathcal{A}})$ respectively. In general, $Aut(\mathcal{A})$ is a subgroup of $Aut(E_{\mathcal{A}})$, and may be a proper subgroup, see Example 2.4. The main result of this paper is that these two groups are isomorphic with each other, if \mathcal{A} is of finite type, excepting types of rank two and type F_4 (Theorem 3.8), or \mathcal{A} is of skew-symmetric finite mutation type (Theorem 3.10). Therefore in some degree, for these cluster algebras, the algebraic symmetries are also captured by the exchange graphs. In particular, we compute the automorphism group of the exchange graph of a finite type cluster algebra in table 1, see Remark 3.9.

Dynkin type	Automorphism group $Aut(E_{\mathcal{A}})$
$A_n(n \geq 2)$	\mathbb{D}_{n+3}
B_2	\mathbb{D}_6
$B_n(n \geq 3)$	\mathbb{D}_{n+1}
C_2	\mathbb{D}_6
$C_n(n \geq 3)$	\mathbb{D}_{n+1}
D_4	$\mathbb{D}_4 \times S_3$
$D_n(n \geq 5)$	\mathbb{Z}_2
E_6	\mathbb{D}_{14}
E_7	\mathbb{D}_{10}
E_8	\mathbb{D}_{16}
F_4	$\mathbb{D}_7 \rtimes \mathbb{Z}_2$
G_2	\mathbb{D}_8

Table 1: Automorphism groups of exchange graphs of cluster algebras of finite type

To prove these results, we describe $E_{\mathcal{A}}$ more precisely. In Section 3.1, we define layers of geodesic loops of $E_{\mathcal{A}}$ by using the distance of a vertex to a fixed vertex on $E_{\mathcal{A}}$. An easy observation is that an isomorphism of exchange graphs should maintain the combinatorial numbers of the layers of geodesic loops based on the corresponding vertices see Remark 3.2(4). By this observation, we directly show in Examples 5, 6, 8, 9 that for a cluster algebra of type A_3 , B_3 , C_3 , \tilde{A}_2 or T_3 (the cluster algebra from an once punctured torus), we have $Aut(\mathcal{A}) \cong Aut(E_{\mathcal{A}})$. For the general cases we reduce them to above five cases (Theorems 3.8, 3.10).

The paper is organized as follows: we recall preliminaries on cluster algebras, cluster algebras of finite mutation type and cluster automorphisms in section 2, then we prove the main Theorems in section 3.

2 Preliminaries

2.1 Cluster algebras

Definition 2.1. [10](Labeled seeds). A labeled seed is a pair $\Sigma = (\mathbf{x}, B)$, where

- $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ is an ordered set of n indeterminates;
- $B = (b_{x_j, x_i})_{n \times n} \in M_{n \times n}(\mathbb{Z})$ is a skew-symmetrizable matrix labeled by $\mathbf{x} \times \mathbf{x}$, that is, there exists a diagonal matrix D with positive integer entries such that DB is skew-symmetric.

The set \mathbf{x} is called the *cluster* with elements the *cluster variables*, and B is called the *exchange matrix*. An element b_{x_j, x_i} in B is also written as b_{ji} for brevity. We always assume through the paper that B is indecomposable, that is, for any $1 \leq i, j \leq n$, there is a sequence $i_0 = i, i_1, \dots, i_m, i_{m+1} = j$, such that $b_{i_k, i_{k+1}} \neq 0$ for any $0 \leq k \leq m$. We also assume that $n > 1$ for convenience. One may produce a new labeled seed by a mutation at direction k for any cluster variable x_k .

Definition 2.2. [10](Seed mutations). The labeled seed $\mu_k(\Sigma) = (\mu_k(\mathbf{x}), \mu_k(B))$ obtained by the mutation of Σ in the direction k is given by:

- $\mu_k(\mathbf{x}) = (\mathbf{x} \setminus \{x_k\}) \sqcup \{\mu_{x_k, \mathbf{x}}(x_k)\}$ where

$$x_k \mu_{x_k, \mathbf{x}}(x_k) = \prod_{\substack{1 \leq j \leq n \\ b_{jk} > 0}} x_j^{b_{jk}} + \prod_{\substack{1 \leq j \leq n \\ b_{jk} < 0}} x_j^{-b_{jk}}.$$

- $\mu_k(B) = (b'_{ji})_{n \times n} \in M_{n \times n}(\mathbb{Z})$ is given by

$$b'_{ji} = \begin{cases} -b_{ji} & \text{if } i = k \text{ or } j = k ; \\ b_{ji} + \frac{1}{2}(|b_{ji}|b_{ik} + b_{ji}|b_{ik}|) & \text{otherwise.} \end{cases}$$

It is easy to check that a mutation is an involution, that is $\mu_k \mu_k(\Sigma) = \Sigma$.

Definition 2.3. [13](n -cluster patterns). An n -regular tree \mathbb{T}_n is a diagram, whose edges are labeled by $1, 2, \dots, n$, such that the n edges emanating from each vertex receive different labels. A n -cluster pattern is an assignment of a labeled seed $\Sigma_t = (\mathbf{x}_t, B_t)$ to every vertex $t \in \mathbb{T}_n$, so that the labeled seeds assigned to the endpoints of any edge labeled by k are obtained from each other by the seed mutation in direction k . The elements of Σ_t are written as follows:

$$\mathbf{x}_t = (x_{1;t}, \dots, x_{n;t}), \quad B_t = (b_{ij}^t). \quad (1)$$

Note that \mathbb{T}_n is in fact determined by any fixed labeled seed on it. Now we are ready to define cluster algebras.

Definition 2.4. [13](Cluster algebras). Given a seed Σ and a cluster pattern \mathbb{T}_n associated to it, we denote

$$\mathcal{X} = \bigcup_{t \in \mathbb{T}_n} \mathbf{x}_t = \{x_{i,t} : t \in \mathbb{T}_n, 1 \leq i \leq n\}, \quad (2)$$

the union of clusters of all the seeds in the pattern. We call the elements $x_{i,t} \in \mathcal{X}$ the cluster variables. The cluster algebra \mathcal{A} associated with Σ is the \mathbb{Z} -subalgebra of the rational function field $\mathcal{F} = \mathbb{Q}(x_1, x_2, \dots, x_n)$, generated by all cluster variables: $\mathcal{A} = \mathbb{Z}[\mathcal{X}]$.

For a skew-symmetrizable matrix $B = (b_{ji})_{n \times n}$, one can associate it to a *valued quiver* (quiver for brevity) $Q = (Q_0, Q_1, \nu)$ as follows: $Q_0 = \{1, 2, \dots, n\}$ is a set of vertices; for any two vertices j and i , if $b_{ji} > 0$, then there is an arrow α from j to i , these arrows form the set Q_1 ; for an arrow α from j to i , we assign it a pair of values: $(\nu_1(\alpha), \nu_2(\alpha)) = (b_{ji}, -b_{ij})$. Since B is an indecomposable skew-symmetrizable matrix, the defined valued quiver Q is connected and there is no loops nor 2-cycles in Q . Then we can define a mutation of the valued quiver by the mutation of the matrix, we refer to [10, 15] for details. We say two quivers Q and Q' *mutation equivalent*, if the corresponding matrices are mutation equivalent, that is, one of them can be obtained from the other one by a finite sequence of mutations. We also write (\mathbf{x}, Q) for the labeled seed (\mathbf{x}, B) , and write \mathcal{A}_Q to the cluster algebra defined by Σ . The quiver and the defined cluster algebra are called skew-symmetric, if the corresponding matrix is skew-symmetric. If the cluster algebra is of finite type [11] or of skew-symmetric type, then the cluster determines the quiver [14], and we denote the quiver of a cluster \mathbf{x} by $Q(\mathbf{x})$.

Example 1. Let B be the following matrix, it is a skew-symmetrizable matrix with skew-symmetrizer $D = \text{diag}\{2, 2, 1, 1\}$. The quiver corresponding to B is Q , where we always delete the trivial pairs of values $(1, 1)$, and replace a arrow assigning pair (m, m) by m arrows.

$$B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & -1 & 0 \\ 0 & 2 & 0 & 2 \\ 0 & 0 & -2 & 0 \end{pmatrix}$$

$$Q : \quad 1 \longrightarrow 2 \xleftarrow{(2,1)} 3 \Longrightarrow 4$$

Definition 2.5. [13](Seeds) Given two labeled seeds $\Sigma = (\mathbf{x}, B)$ and $\Sigma' = (\mathbf{x}', B')$, we say that they define the same seed if Σ' is obtained from Σ by simultaneous relabeling of the sets \mathbf{x} and the corresponding relabeling of the rows and columns of B .

We denote by $[\Sigma]$ the seed represented by a labeled seed Σ . The cluster \mathbf{x} of a seed $[\Sigma]$ is a unordered n -element set. For any $x \in \mathbf{x}$, there is a well-defined mutation $\mu_x([\Sigma]) = [\mu_x(\Sigma)]$ of $[\Sigma]$ at direction x , where $x = x_k$. For two same rank skew-symmetrizable matrices B and B' , we say $B \cong B'$, if B' is obtained from B by simultaneous relabeling of the rows and columns of B . Then the exchange matrices in any two labeled seeds representing a same seed are isomorphic. The isomorphism of two exchange matrices induces an isomorphism of corresponding quivers. For convenience, in the rest of the paper, we also denote by Σ the seed $[\Sigma]$ represented by Σ .

Definition 2.6. [13](Exchange graphs) The exchange graph of a cluster algebra is the n -regular graph whose vertices are the seeds of the cluster algebra and whose edges connect the seeds related by a single mutation. We denote by $E_{\mathcal{A}}$ the exchange graph of a cluster algebra \mathcal{A} .




Figure 1: Quivers of finite type

Clearly, the exchange graph of a cluster algebra is a quotient graph of the n -regular tree, its vertices are equivalent classes of labeled seeds. The exchange graph not necessary be a finite graph, if it is finite, then we say the corresponding cluster algebra (and its cluster pattern) is of *finite type*.

Definition 2.7. [11, page 70](Cluster complexes) A cluster complex Δ of \mathcal{A} is a simplicial complex on the ground set \mathcal{X} with the clusters as the maximal simplices.

Then Δ is an n -dimensional complex. In particular, if \mathcal{A} is of finite type or skew-symmetric, then the vertices of $E_{\mathcal{A}}$ are clusters, thus the dual graph of Δ is $E_{\mathcal{A}}$.

2.2 Finite types and finite mutation types

By the classification of cluster algebras of finite type [11], a cluster algebra is of finite type if and only if there is a seed whose quiver is one of quivers depicted in Figure 1. Note that the underlying graphs of quivers in Figure 1 are trees, thus any two quivers with the same underlying graph are mutation-equivalent.

Definition 2.8. [7, 9] A block is a quiver isomorphic to one of the quivers with black / white colored vertices shown on Figure 2. Vertices marked in white are called outlets. A connected quiver




Figure 2: Blocks. Outlets are colored white, dead ends are black.

Q is called *block-decomposable* (decomposable for brevity) if it can be obtained from a collection of blocks by identifying outlets of different blocks along some partial matching (matching of outlets of the same block is not allowed), where two arrows with same endpoints and opposite directions cancel out. If Q is not block-decomposable then we call Q non-decomposable.

Then it is proved in [7, Theorem 13.3] that a quiver is decomposable if and only if it is a quiver of a triangulation of an oriented marked Riemann surface, and thus a quiver mutation equivalent to a decomposable quiver is also decomposable. Note that all arrow multiplicities of a decomposable quiver are 1 or 2. Therefore decomposable quivers are mutation finite. It is clear that a quiver of rank two, that is, a quiver with two vertices, is mutation finite. Besides these two kinds of quivers, there are exactly 11 exceptional skew-symmetric quivers of finite mutation type, see Theorem 6.1 in [9]. We list the exceptional quivers in Figure 3.

2.3 Automorphism groups

In this section, we recall the cluster automorphism group [1] of a cluster algebra, and the automorphism group of the corresponding exchange graph [4].

Definition 2.9. [1](Cluster automorphisms) For a cluster algebra \mathcal{A} and a \mathbb{Z} -algebra automorphism $f : \mathcal{A} \rightarrow \mathcal{A}$, we call f a cluster automorphism, if there exists a labeled seed (\mathbf{x}, B) of \mathcal{A} such that the following conditions are satisfied:

1. $f(\mathbf{x})$ is a cluster;
2. f is compatible with mutations, that is, for every $x \in \mathbf{x}$ and $y \in \mathbf{x}$, we have

$$f(\mu_{x, \mathbf{x}}(y)) = \mu_{f(x), f(\mathbf{x})}(f(y)).$$

Then a cluster automorphism maps a labeled seed $\Sigma = (\mathbf{x}, B)$ to a labeled seed $\Sigma' = (\mathbf{x}', B')$. Under our assumption that B is indecomposable, we have the following

Lemma 2.10. [1] A \mathbb{Z} -algebra automorphism $f : \mathcal{A} \rightarrow \mathcal{A}$ is a cluster automorphism if and only if there exists a labeled seed $\Sigma = (\mathbf{x}, B)$ of \mathcal{A} , such that $f(\mathbf{x})$ is the cluster in a labeled seed $\Sigma' = (\mathbf{x}', B')$ of \mathcal{A} with $B' = B$ or $B' = -B$.

We call the cluster automorphism such that $B = B'$ ($B = -B'$ respectively) a *direct cluster automorphism* (an *inverse cluster automorphism* respectively). Clearly, all the cluster automorphism of a cluster algebra \mathcal{A} form a group with homomorphism composition as multiplication. We call this group the *cluster automorphism group* of \mathcal{A} , and denote it by $\text{Aut}(\mathcal{A})$. We call the group




Figure 3: Representatives of non-decomposable quivers of finite mutation type

$\text{Aut}^+(\mathcal{A})$ consisting of the direct cluster automorphisms of \mathcal{A} the *direct cluster automorphism group* of \mathcal{A} , which is a subgroup of $\text{Aut}(\mathcal{A})$ with index at most two, see [1].

Definition 2.11. (*Automorphism of exchange graphs*) [16, 4] An automorphism of the exchange graph $E_{\mathcal{A}}$ of a cluster algebra \mathcal{A} is an automorphism of $E_{\mathcal{A}}$ as a graph, that is, a permutation σ of the vertex set, such that the pair of vertices (u, v) forms an edge if and only if the pair $(\sigma(u), \sigma(v))$ also forms an edge.

Clearly, the natural composition of two automorphisms of $E_{\mathcal{A}}$ is again an automorphism. We define an *automorphism group* $\text{Aut}(E_{\mathcal{A}})$ of $E_{\mathcal{A}}$ as a group consisting of automorphisms of $E_{\mathcal{A}}$. It is clearly that a cluster automorphism induces a unique automorphism of the exchange graph. Thus $\text{Aut}(\mathcal{A})$ is a subgroup of $\text{Aut}(E_{\mathcal{A}})$, see [4]. By the definition, an automorphism σ of an exchange graph maps clusters to clusters, and induces an automorphism of its dual graph: cluster complex Δ , we denote this automorphism by σ_{Δ} . Then σ_{Δ} is a permutation of cluster variables in \mathcal{X} , which maps a maximal simplex to a maximal simplex, but the map may not be compatible with the algebra relations among cluster variables in \mathcal{A} , thus it is not necessarily a cluster automorphism. In fact, $\text{Aut}(\mathcal{A})$ maybe a proper subgroup of $\text{Aut}(E_{\mathcal{A}})$, see Example 2.4. The following Lemma can be viewed as a description of $\text{Aut}(\mathcal{A})$ as a subgroup of $\text{Aut}(E_{\mathcal{A}})$, as those exchange graph automorphisms which happen to preserve B-matrices (perhaps up to global reversal of sign) up to simultaneously relabeling of the rows and columns. In this point of view, the main thrust of this paper is to show that, typically for the cluster algebras we consider, any graph automorphism has the property of preserving B-matrices.

Lemma 2.12. Let $\Phi : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}}$ be an automorphism which maps a seed $\Sigma = (\mathbf{x}, B)$ to a seed $\Sigma' = (\mathbf{x}', B')$. If $B \cong B'$ or $B \cong -B'$ under the correspondence $\mathbf{x} \rightarrow \mathbf{x}'$, then the map $\mathbf{x} \rightarrow \mathbf{x}'$ induces a cluster automorphism Ψ of \mathcal{A} and the induced automorphism $\Psi_E : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}}$ coincides with Φ .

Proof. Since $B \cong B'$ or $B \cong -B'$, the map $\mathbf{x} \rightarrow \mathbf{x}'$ induces a cluster automorphism Ψ of \mathcal{A} by Lemma 2.10. Noticing that $\Phi(\mathbf{x}) = \Psi(\mathbf{x})$, then by inductions on the mutations, we have $\Phi = \Psi$ on each clusters of $E_{\mathcal{A}}$, so $\Phi = \Psi_E$ as automorphisms of the exchange graph $E_{\mathcal{A}}$. \square

3 Automorphism groups of exchange graphs

In this section we consider relations between the groups $\text{Aut}(\mathcal{A})$ and $\text{Aut}(E_{\mathcal{A}})$ for a cluster algebra \mathcal{A} of finite type or of skew-symmetric finite mutation type. For this, it is needed to describe $E_{\mathcal{A}}$ more precisely. In the following we will recall the basic structures of $E_{\mathcal{A}}$ from [10, 11], and then introduce layers of geodesic loops on $E_{\mathcal{A}}$.

3.1 Layers of geodesic loops

Let $\Sigma = (\mathbf{x}, B)$ be a labeled seed on the cluster pattern of \mathcal{A} . Let \mathbf{x}' be a proper subset of \mathbf{x} , then \mathbf{x}' is a non-maximal simplex in the cluster complex Δ . We denote by $\Delta_{\mathbf{x}'}$ the *link* of $\mathbf{x} \setminus \mathbf{x}'$, which is the simplicial complex on the ground set $\mathcal{X}_{\mathbf{x}'} = \{\alpha \in \mathcal{X} - (\mathbf{x} \setminus \mathbf{x}') : (\mathbf{x} \setminus \mathbf{x}') \cup \{\alpha\} \in \Delta\}$, such that \mathbf{x}'' is a simplex in $\Delta_{\mathbf{x}'}$ if and only if $\mathbf{x} \setminus \mathbf{x}' \cup \mathbf{x}''$ is a simplex in Δ . Let $\Gamma_{\mathbf{x}'}$ be the dual graph of $\Delta_{\mathbf{x}'}$. We view $\Gamma_{\mathbf{x}'}$ as a subgraph of $E_{\mathcal{A}}$ whose vertices are the maximal simplices in

Δ that contain $\mathbf{x} \setminus \mathbf{x}'$. In fact, as we explain now, $\Gamma_{\mathbf{x}'}$ is the exchange graph of a cluster algebra \mathcal{A}_f defined by a frozen seed $\Sigma_f = (\mathbf{x}', \mathbf{x} \setminus \mathbf{x}', B_f)$, which is the freezing of Σ at $\mathbf{x} \setminus \mathbf{x}'$ (see [3, Definition 2.25]), where B_f is obtained from B by deleting the columns labeled by variables in $\mathbf{x} \setminus \mathbf{x}'$. Then elements in $\mathbf{x} \setminus \mathbf{x}'$ are coefficients of \mathcal{A}_f (we refer to [10, 13] for a cluster algebra with coefficients). Let \mathcal{A}' be cluster algebra defined by a seed $\Sigma' = (\mathbf{x}', B')$, where B' is obtained from B by deleting rows and columns labeled by variables in $\mathbf{x} \setminus \mathbf{x}'$. In our settings, that is, cluster algebras are of finite type or of skew-symmetric finite type, the exchange graph of a cluster algebra (with coefficients) only depends on the principal part of the exchange matrix (see [11, 2]) which is the submatrix labeled by $\mathbf{x} \setminus \mathbf{x}' \times \mathbf{x} \setminus \mathbf{x}'$, thus the graph $\Gamma_{\mathbf{x}'}$ coincides with the exchange graph $E_{\mathcal{A}'}$.

For an 2-dimensional subcomplex \mathbf{x}' of Δ , we call the dual graph $\Gamma_{\mathbf{x}'}$ a *geodesic loop* of $E_{\mathcal{A}}$. We mention that the definition of geodesic loop is slightly different with the definition used in [11], where a line is not a geodesic loop. If \mathcal{A} is of finite type, then $E_{\mathcal{A}}$ is a finite graph, and $\Gamma_{\mathbf{x}'}$ is a polygon. Notice that in the seed $\Sigma' = (\mathbf{x}', B')$ constructed above, B' is of Dynkin type, that is, one of types A_2, B_2, C_2 or G_2 . Therefore $\Gamma_{\mathbf{x}'}$ is a $h + 2$ -polygon, where h is the Coxeter number of the corresponding Dynkin type, see [11]. If \mathcal{A} is of finite mutation type, then $\Gamma_{\mathbf{x}'}$ may be a line. We fix a basepoint $\Sigma = (\mathbf{x}, B)$ and introduce the following concept.

- Definition 3.1.**
1. Let Σ' be a point of $E_{\mathcal{A}}$, the distance $\ell(\Sigma, \Sigma')$ between Σ and Σ' is the minimal length of paths between Σ and Σ' ;
 2. Let L be a geodesic loop of $E_{\mathcal{A}}$, the distance $\ell_{\Sigma}(L)$ between Σ and L is the minimal length $\min\{\ell(\Sigma, \Sigma'), \Sigma' \in L\}$;
 3. Let $m \in \mathbb{Z}_{\geq 0}$ be a non-negative integer, denote by ℓ_{Σ}^m the set of geodesic loop whose distance to Σ is m . We call it the m -layer of geodesic loops of $E_{\mathcal{A}}$ based on Σ ;
 4. For any $m \in \mathbb{Z}_{\geq 0}$, denote $N(\ell_{\Sigma}^m)$ the set of amounts of edges belonging to geodesic loops in the m -layer ℓ_{Σ}^m .

Remark 3.2. The following observations are directly derived from the definitions:

1. The elements in ℓ_{Σ}^0 are those geodesic loops $\Gamma_{\mathbf{x}'}$ for the 2-dimensional subcomplex \mathbf{x}' of Δ , where \mathbf{x}' is a subset of the cluster \mathbf{x} in Σ ;
2. For $m_1 \neq m_2$, $\ell_{\Sigma}^{m_1} \cap \ell_{\Sigma}^{m_2} = \emptyset$;
3. The disjoint union $\sqcup_{m \geq 0} \ell_{\Sigma}^m$ is the set of all the geodesic loops of $E_{\mathcal{A}}$;
4. If $\sigma : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}'}$ is an isomorphism of graphs, such that the image of Σ is Σ' , then for every $m \in \mathbb{Z}_{\geq 0}$, $N(\ell_{\Sigma}^m) = N(\ell_{\Sigma'}^m)$ as sets.

3.2 Cases of rank two and rank three

In this subsection, we consider the relations between $\text{Aut}(\mathcal{A})$ and $\text{Aut}(E_{\mathcal{A}})$ for a cluster algebra \mathcal{A} of rank two or rank three.

Example 2. For a finite type cluster algebra \mathcal{A} of rank 2, that is, one of types A_2, B_2, C_2 or G_2 , its exchange graph $E_{\mathcal{A}}$ is a $(h+2)$ -polygon, thus $\text{Aut}(E_{\mathcal{A}})$ is isomorphic to the dihedral group \mathbb{D}_{h+2} , where h is the Coxeter number. If \mathcal{A} is of type A_2 , then $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_5$ [1], thus $\text{Aut}(\mathcal{A}) \cong \text{Aut}(E_{\mathcal{A}})$. If \mathcal{A} is of type B_2, C_2 or G_2 , [5, Theorem 3.5] shows that $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_{(h+2)/2}$, thus $\text{Aut}(\mathcal{A}) \subsetneq \text{Aut}(E_{\mathcal{A}})$.

Example 3. For an infinite type skew-symmetric cluster algebra \mathcal{A} of rank 2, its exchange graph $E_{\mathcal{A}}$ is a line, thus $\text{Aut}(E_{\mathcal{A}}) = \langle s \rangle \times \langle r \rangle \cong \mathbb{Z} \rtimes \mathbb{Z}_2 = \mathbb{D}_{\infty}$, where s is a left shift of $E_{\mathcal{A}}$ which maps a cluster to the left adjacent cluster and r is a reflection with respect to a fixed cluster. Then s corresponds to a direct cluster automorphism of \mathcal{A} and r corresponds to an inverse cluster automorphism of \mathcal{A} , thus by Lemma 2.12 $\text{Aut}(E_{\mathcal{A}}) \subseteq \text{Aut}(\mathcal{A})$. Therefore $\text{Aut}(E_{\mathcal{A}}) \cong \text{Aut}(\mathcal{A}) \cong \mathbb{D}_{\infty}$.

Example 4. For an infinite type non-skew-symmetric cluster algebra \mathcal{A} of rank 2, its exchange graph $E_{\mathcal{A}}$ is also a line, thus as showed in Example 3, $\text{Aut}(E_{\mathcal{A}}) = \langle s \rangle \times \langle r \rangle \cong \mathbb{Z} \rtimes \mathbb{Z}_2$, where s corresponds to a direct cluster automorphism of \mathcal{A} , while r dose not correspond to any cluster automorphism of \mathcal{A} , since there is no non-trivial symmetry of the quiver in any seed of \mathcal{A} . Thus $\text{Aut}(\mathcal{A}) \cong \mathbb{Z} \subsetneq \text{Aut}(E_{\mathcal{A}})$.

Example 5. We consider the cluster algebra \mathcal{A} of type A_3 with an initial labeled seed $\Sigma_0 = (\{x_1, x_2, x_3\}, Q)$, where Q is $1 \longrightarrow 2 \longleftarrow 3$. Then its exchange graph $E_{\mathcal{A}}$ is depicted in Figure 4. Note that there are three quadrilaterals and six pentagons in $E_{\mathcal{A}}$. Then as shown in [5, Example 3], $\text{Aut}(\mathcal{A}) = \langle f_-, f_+ \rangle \cong \mathbb{D}_6$, where f_- is defined by:

$$f_- : \begin{cases} x_1 \mapsto x_1 \\ x_2 \mapsto \mu_2(x_2) \\ x_3 \mapsto x_3 \end{cases} \quad (3)$$

It maps Σ_0 to Σ_1 , and induces a reflection with respect to the horizontal central axis of $E_{\mathcal{A}}$. The cluster automorphism f_+ is defined by:

$$f_+ : \begin{cases} x_1 \mapsto \mu_1(x_1) \\ x_2 \mapsto x_2 \\ x_3 \mapsto \mu_3(x_3) \end{cases} \quad (4)$$

It gives a reflection on $E_{\mathcal{A}}$, which maps Σ_0 to Σ_5 . In fact as shown in [5], a direct cluster automorphism of \mathcal{A} is of the form $(f_+ f_-)^m, 0 \leq m \leq 5$, which induces a rotation of seeds in $\{\Sigma_0, \Sigma_1, \Sigma_2, \Sigma_3, \Sigma_4, \Sigma_5\}$, thus $\text{Aut}^+(\mathcal{A})$ can be viewed as the symmetry group of the bipartite belt consisting of seeds in $\{\Sigma_0, \Sigma_1, \Sigma_2, \Sigma_3, \Sigma_4, \Sigma_5\}$, where the quivers in these seeds are the bipartite quivers isomorphic to Q .

We will prove that any automorphisms of $E_{\mathcal{A}}$ is induced from an element in $\text{Aut}(\mathcal{A})$, and thus $\text{Aut}(\mathcal{A}) \cong \text{Aut}(E_{\mathcal{A}})$. For this purpose, we should show the following claims:

- (1) there exists no automorphism of $E_{\mathcal{A}}$ which maps Σ_0 to a vertex excepting for $\Sigma_i, 0 \leq i \leq 5$;
- (2) if an automorphism of $E_{\mathcal{A}}$ maps Σ_0 to $\Sigma_i, 0 \leq i \leq 5$, then it is induced from a cluster automorphism of \mathcal{A} .




Figure 4: The exchange graph of a cluster algebra of type A_3

Let σ be an automorphism of $E_{\mathcal{A}}$, due to symmetries of $E_{\mathcal{A}}$, we only show that $\sigma(\Sigma) \neq O_i$, $i = 1, 2, 3$. By a direct computation, the sets of numbers for the layers of geodesic loops based on these vertices are as follows:

$$\begin{aligned}
 N(\ell_{\Sigma}^0) &= \{4, 5, 5\}, N(\ell_{\Sigma}^1) = \{4, 5, 5\}, N(\ell_{\Sigma}^2) = \{5, 5\}, N(\ell_{\Sigma}^3) = \{4\}; \\
 N(\ell_{O_1}^0) &= \{4, 5, 5\}, N(\ell_{O_1}^1) = \{5, 5, 5\}, N(\ell_{O_1}^2) = \{4, 4\}, N(\ell_{O_1}^3) = \{5\}; \\
 N(\ell_{O_2}^0) &= \{5, 5, 5\}, N(\ell_{O_2}^1) = \{4, 4, 4\}, N(\ell_{O_2}^2) = \{5, 5, 5\}; \\
 N(\ell_{O_3}^0) &= \{4, 5, 5\}, N(\ell_{O_3}^1) = \{5, 5, 5\}, N(\ell_{O_3}^2) = \{4, 4\}, N(\ell_{O_3}^3) = \{5\};
 \end{aligned}$$

Then by Remark 3.2 (4), $\sigma(\Sigma_0) \neq O_i$, $i = 1, 2, 3$. So the claim one is affirmed.

Now we consider the claim two. Still due to the symmetries of the graph, we may assume that $\sigma(\Sigma_0) = \Sigma_0$. Since σ is a graph automorphism, it can be seen that there are two possibilities of σ , one is the identity, another is the reflection f_0 with respect to the vertical central axis of $E_{\mathcal{A}}$, as depicted in Figure 4. Note that the identity graph automorphism is induced from the identity automorphism of the cluster algebra, while the graph automorphism f_0 is induced from the cluster automorphism $(f_+ f_-)^3$ by a direct computation. Therefore the claim two is true and we have $\text{Aut}(E_{\mathcal{A}}) \cong \text{Aut}(\mathcal{A}) \cong \mathbb{D}_6$.

Example 6. It is known from a result in [12] that the cluster algebras of type B_n and type C_n have the same exchange graph. Based on a seed Σ_0 , the exchange graph of a cluster algebra \mathcal{A}

of type B_3 or type C_3 is depicted in Figure 5. For the cluster algebra of type B_3 , the quiver of the initial seed Σ_0 is

$$1 \longrightarrow 2 \xleftarrow{(2,1)} 3.$$

For the cluster algebra of type C_3 , the quiver of the initial seed Σ_0 is

$$1 \longrightarrow 2 \xleftarrow{(1,2)} 3.$$

Let σ be an automorphism of $E_{\mathcal{A}}$. As showed by Example 4 in [5], we have $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_4$ and $\{\Sigma_0, \Sigma_1, \Sigma_2, \Sigma_3, \Sigma_4, \Sigma_5, \Sigma_6, \Sigma_7\}$ are all the seeds whose quivers are isomorphic to Q . Similarly, to get $\text{Aut}(E_{\mathcal{A}}) \cong \text{Aut}(\mathcal{A})$, we should prove the two claims stated in Example 5. For the first claim, we only need to prove that $\sigma(\Sigma_0) \neq O_i (i = 1, 2, 3, 4)$ in Figure 5, and this can be obtained by the fact that these seeds have different combinatorial numbers of layers of geodesic loops:

$$\begin{aligned} N(\ell_{\Sigma_0}^0) &= \{4, 5, 6\}, \quad N(\ell_{\Sigma_0}^1) = \{4, 5, 6\}; \\ N(\ell_{O_1}^0) &= \{5, 6, 6\}; \\ N(\ell_{O_2}^0) &= \{4, 5, 6\}, \quad N(\ell_{O_2}^1) = \{5, 6, 6\}; \\ N(\ell_{O_3}^0) &= \{4, 5, 6\}, \quad N(\ell_{O_3}^1) = \{5, 6, 6\}; \\ N(\ell_{O_4}^0) &= \{5, 6, 6\}. \end{aligned}$$

For the second claim, we may also assume $\sigma(\Sigma_0) = \Sigma_0$. Since $N(\ell_{\Sigma_0}^0) = \{4, 5, 6\}$, there are neither rotation symmetries nor reflection symmetries of $E_{\mathcal{A}}$ at Σ_0 . So σ must be the identity automorphism of $E_{\mathcal{A}}$, which is induced from the identity automorphism of the cluster algebra. Noticing that there are eight elements in $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_4$, where each one corresponds a graph automorphism which maps Σ_0 to $\Sigma_i, 0 \leq i \leq 7$.

Example 7. For cluster algebra of type F_4 , let the quiver Q of a seed Σ be

$$1 \longrightarrow 2 \xleftarrow{(2,1)} 3 \longrightarrow 4.$$

Then $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_7$ [5]. The variables x_1, x_2, x_3 and the corresponding full subquiver of Q form a seed Σ_1 of type B_3 , while x_2, x_3, x_4 and the corresponding full subquiver of Q form a seed Σ_2 of type C_3 . By pinning down Σ , rotating the graph $E_{\mathcal{A}}$ induces an automorphism σ of $E_{\mathcal{A}}$, which exchanges the graph $E_{\mathcal{A}_{\Sigma_1}}$ and the graph $E_{\mathcal{A}_{\Sigma_2}}$. However σ does not induces a cluster automorphism of \mathcal{A} , and $\text{Aut}(\mathcal{A}) \cong \mathbb{D}_7 \not\subseteq \mathbb{D}_7 \rtimes \mathbb{Z}_2 \cong \text{Aut}(E_{\mathcal{A}})$.

Proposition 3.3. Let Q be a connected quiver with three vertices, which is finite type. Let $\Sigma = (\mathbf{x}, Q)$ and $\Sigma' = (\mathbf{x}', Q')$ be two seeds (not necessarily mutation equivalent with each other). If there is an isomorphism $\sigma : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}'}$ such that $\sigma(\Sigma) = \Sigma'$, then $\Sigma' = (\mathbf{x}', Q')$ is a finite type seed with Q' connected, and

1. if Σ is of type A_3 , then $Q' \cong Q$ (or Q^{op});
2. if Σ is of type B_3 and Σ' is not of type C_3 , then $Q' \cong Q$ (or Q^{op});




Figure 5: The exchange graph of a cluster algebra of type B_3 or type C_3

3. if Σ is of type C_3 and Σ' is not of type B_3 , then $Q' \cong Q$ (or Q^{op}).

Proof. Clearly, since $E'_{\mathcal{A}} \cong E_{\mathcal{A}}$ is of finite, Q' is a Dynkin type quiver with three vertices. If Q is of type A_3 , then by Example 5,

$$N(\ell_{\Sigma}^0) = \{4, 5, 5\} \text{ or } \{5, 5, 5\}.$$

If Q is of type B_3 (or C_3), then from Example 6,

$$N(\ell_{\Sigma}^0) = \{4, 5, 6\} \text{ or } \{5, 6, 6\}.$$

If Q' is a union of a quiver of type A_2 and a point, then from Example 2,

$$N(\ell_{\Sigma}^0) = \{4, 4, 5\}.$$

If Q' is a union of a quiver of type B_2 (or C_2) and a point, then from Example 2,

$$N(\ell_{\Sigma}^0) = \{4, 4, 6\}.$$

If Q' is a union of a quiver of type G_2 and a point, then from Example 2,

$$N(\ell_{\Sigma}^0) = \{4, 4, 8\}.$$

Thus we get the proof by Remark 3.2. \square

Example 8. Let Q be the quiver in Figure 6, we call it of type \tilde{A}_2 . Then it is not hard to see that if a quiver in the mutation class of Q is not isomorphic to Q , then it must be isomorphic to the quiver Q' in Figure 6. Let \mathcal{A} be a cluster algebra with an initial seed $\Sigma = (\{x_1, x_2, x_3\}, Q)$, similarly to above examples, to show that $\text{Aut}(\mathcal{A}) \cong \text{Aut}(E_{\mathcal{A}})$, we only need to notice that:

$$N(\ell_{\Sigma}^0) = \{5, 5, \infty\},$$

$$N(\ell_{\Sigma'}^0) = \{5, 5, 5\},$$

where Σ' is a seed of \mathcal{A} with quiver isomorphic to Q' . In fact, from section 3.3 in [1], $\text{Aut}(\mathcal{A}) = \langle r_1, r_2 | r_1 r_2 = r_2 r_1, r_1^2 = r_2 > \times < \sigma | \sigma^2 = 1 \rangle \cong \mathbb{H}_{2,1} \rtimes \mathbb{Z}_2$, where

$$r_1 : \begin{cases} x_1 \mapsto x_3 \\ x_2 \mapsto \mu_1(x_1) \\ x_3 \mapsto x_2 \end{cases} \quad (5)$$

$$r_2 : \begin{cases} x_1 \mapsto x_2 \\ x_2 \mapsto \mu_3 \mu_1(x_3) \\ x_3 \mapsto \mu_1(x_1) \end{cases} \quad (6)$$

$$\sigma : \begin{cases} x_1 \mapsto x_2 \\ x_2 \mapsto x_1 \\ x_3 \mapsto x_3 \end{cases} \quad (7)$$

Thus $\text{Aut}(E_{\mathcal{A}}) \cong \mathbb{H}_{2,1} \rtimes \mathbb{Z}_2$




Figure 6: quivers of type \tilde{A}_2




Figure 7: quiver of type T_3

Example 9. Let \mathcal{A} be a cluster algebra from an once punctured torus, we call it a cluster algebra of type T_3 , then it is of finite mutation type with quiver always isomorphic to the quiver in Figure 7. Then by Lemma 2.12, we have $\text{Aut}(\mathcal{A}) \cong \text{Aut}(E_{\mathcal{A}})$.

Corollary 3.4. Let \mathcal{A} and \mathcal{A}' be two cluster algebras of finite type, or of skew-symmetric finite mutation type, with rank equal to 2 or 3. Let $\Sigma = (\mathbf{x}, B)$ and $\Sigma' = (\mathbf{x}', B')$ be two seeds of \mathcal{A} and \mathcal{A}' respectively. If $N(\ell_{\Sigma}^k) = N(\ell_{\Sigma'}^k)$ for any $k \in \mathbb{Z}_{\geq 0}$, then there exists an isomorphism $\Phi : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}'}$ such that $\Phi(\mathbf{x}) = \mathbf{x}'$.

Proof. This follows from Example 5, Example 6, Example 8 and Example 9 □

We expect the result in the Corollary is true for any finite type cluster algebras and finite mutation type cluster algebras. It means that for any seed Σ , the set $N(\ell_{\Sigma}^k)$ characterizes the exchange graph.

Lemma 3.5. Let Q be a connected skew-symmetric quiver of finite mutation type.

1. If there are 3 vertices in Q , then Q is one of the following types:
 - (1) A_3 type;
 - (2) \tilde{A}_2 type;
 - (3) T_3 type.
2. If there are at least 4 vertices in Q , then any full subquiver of Q with three vertices is of type A_3 or of type \tilde{A}_2 .

Proof. 1. From the classification of cluster algebras of finite mutation type, Q must be block-decomposable, then the proof is a straightforward check by gluing the blocks in 2.

2. We only need to notice that a quiver of type T_3 is obtained by gluing two blocks of type II in Figure 2, and thus one can not further glue it with a block to obtain a connected quiver of finite mutation type.

□

It is clear that if for any quiver in the mutation equivalent class of Q , the number of arrows between any two vertices is at most 2, then Q is of finite mutation type. The above lemma shows that the inverse statement is also true for the cases when there are at least 3 vertices, that is, we have the following corollary, which has been stated in [6, Corollary 8].

Corollary 3.6. *A connected quiver Q with at least 3 vertices is of finite mutation type if and only if for any quiver in its mutation class the number of arrows between any two vertices is at most 2.*

Proposition 3.7. *Let Q be a connected skew-symmetric quiver with three vertices, which is of finite mutation type. Let $\Sigma = (\mathbf{x}, Q)$ and $\Sigma' = (\mathbf{x}', Q')$ be two seeds. If there is an isomorphism $\sigma : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}'}$ such that $\sigma(\Sigma) = \Sigma'$, then $Q' \cong Q$ or $Q' \cong Q^{op}$.*

Proof. Similar to Proposition 3.3, this follows from Lemma 3.5, Example 9, Example 5 and Example 8. □

3.3 General cases

Theorem 3.8. *Let \mathcal{A} be a cluster algebra of finite type. Assume that it is not of type F_4 , let $\Sigma = (\mathbf{x}, Q)$ be a labeled seed of \mathcal{A} , where Q is a connected quiver with at least three vertices. Then we have $\text{Aut}(\mathcal{A}) \cong \text{Aut}(E_{\mathcal{A}})$.*

Proof. We need to show that $\text{Aut}(E_{\mathcal{A}}) \subseteq \text{Aut}(\mathcal{A})$. Let Φ be any automorphism of $E_{\mathcal{A}}$. Then it induces an automorphism ϕ of the complex Δ , in particular, which gives a permutation on the cluster variable set \mathcal{X} . Let $\mathbf{x}' \subseteq \mathbf{x}$ be a 3-dimensional complex, and let $Q(\mathbf{x}')$ be the full subquiver of $Q(\mathbf{x})$ with vertices indexed by the variables in \mathbf{x}' . Let \mathcal{A}' be the cluster algebra defined by the seed $\Sigma' = (\mathbf{x}', Q(\mathbf{x}'))$.

Noticing that since ϕ is an automorphism of a complex, it maps a simplex to a simplex, and thus induces a bijection from $\mathcal{X}_{\mathbf{x}'} = \{\alpha \in \mathcal{X} - (\mathbf{x} \setminus \mathbf{x}') : \mathbf{x} \setminus \mathbf{x}' \cup \{\alpha\} \in \Delta\}$ to $\mathcal{X}_{\phi(\mathbf{x}')} = \{\alpha \in \mathcal{X} - \phi(\mathbf{x} \setminus \mathbf{x}') : \phi(\mathbf{x} \setminus \mathbf{x}') \cup \{\alpha\} \in \Delta\}$, and also induces an isomorphism $\phi_{\mathbf{x}'}$ from the link $\Delta_{\mathbf{x}'}$ to the link $\Delta_{\phi(\mathbf{x}')}$. Moreover, the duality of the isomorphism $\phi_{\mathbf{x}'}$ gives an isomorphism between the dual graphs of the complexes, that is we have an isomorphism

$$\Phi_{\mathbf{x}'} : \Gamma_{\mathbf{x}'} \rightarrow \Gamma_{\phi(\mathbf{x}')}.$$
 (8)

Let $\bar{\Sigma}' = (\phi(\mathbf{x}'), Q(\phi(\mathbf{x}')))$ be a seed, where $Q(\phi(\mathbf{x}'))$ is the full subquiver of $Q(\Phi(\mathbf{x}))$ whose vertices are those labeled by elements in $\phi(\mathbf{x}')$. Let $\bar{\mathcal{A}}'$ be the cluster algebra defined by $\bar{\Sigma}'$. As showed in the beginning of subsection 3.1, there are isomorphisms $\Gamma_{\mathbf{x}'} \cong E_{\mathcal{A}'}$ and $\Gamma_{\phi(\mathbf{x}')} \cong E_{\bar{\mathcal{A}}'}$. Then combining with the isomorphism (8), we have $E_{\mathcal{A}'} \cong E_{\bar{\mathcal{A}}'}$. Since \mathcal{A} is not of type F_4 , if $Q(\mathbf{x}')$ is of type B_3 (type C_3 respectively), then $Q(\phi(\mathbf{x}'))$ is not of type C_3 (type B_3 respectively). Thus by Proposition 3.3, $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))$ or $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))^{op}$.

Let $\mathbf{x}' = \{x_1, x_2, x_3\} \subseteq \mathbf{x}$ and $\mathbf{x}'' = \{x_2, x_3, x_4\} \subseteq \mathbf{x}$ be two 3-dimensional complexes with exactly two common elements. By above discussion, we have $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))$ or $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))^{op}$, and $Q(\mathbf{x}'') \cong Q(\phi(\mathbf{x}''))$ or $Q(\mathbf{x}'') \cong Q(\phi(\mathbf{x}''))^{op}$. Now assume $b_{x_2 x_3} \neq 0$, that is there exists at

least one arrow in $Q(\mathbf{x})$ between the vertices labeled by x_2 and x_3 , then simultaneously we have $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))$ and $Q(\mathbf{x}'') \cong Q(\phi(\mathbf{x}''))$, or $Q(\mathbf{x}') \cong Q(\phi(\mathbf{x}'))^{op}$ and $Q(\mathbf{x}'') \cong Q(\phi(\mathbf{x}''))^{op}$. Finally, due to the arbitrariness of the choice of \mathbf{x}' and the connectness of the quiver, one may show that $Q(\Phi(\mathbf{x})) \cong Q$ or $Q(\Phi(\mathbf{x})) \cong Q^{op}$. See the inductive process in the following picture:

$$\begin{array}{ccccccccccc} & & & & & \mathbf{x}'' & & & & & \\ & & & & & \text{-----} & & & & & \\ \cdots & x_0 & \text{---} & x_1 & \text{---} & x_2 & \text{---} & x_3 & \text{---} & x_4 & \text{---} & x_5 & \text{---} & x_6 & \cdots \\ & & & & & \mathbf{x}' & & & & \mathbf{x}''' & & & & \end{array}$$

Therefore $\Phi : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}}$ induces an cluster automorphism of \mathcal{A} by Lemma 2.12. Thus $Aut(E_{\mathcal{A}}) \subseteq Aut(\mathcal{A})$ and we have $Aut(E_{\mathcal{A}}) \cong Aut(\mathcal{A})$. \square

Remark 3.9. By combining the above theorem, Table 1 in [1] and Theorem 3.5 in [5], we may compute the automorphism groups of the exchange graphs of cluster algebras of finite type, see the table 1. The cases of rank two and type F_4 is computed in Example 2 and Example 7 respectively.

Theorem 3.10. Let \mathcal{A} be a connected skew-symmetric cluster algebra of finite mutation type, then $Aut(\mathcal{A}) \cong Aut(E_{\mathcal{A}})$.

Proof. If \mathcal{A} is of finite type of rank 2, that is, of type A_2 , then the result follows from Example 2. If \mathcal{A} is of infinite type of rank 2, then the result follows from Example 3. When the rank of \mathcal{A} is at least 3, the proof is similar to the proof of Theorem 3.8 by using the connectness of the cluster algebra and Proposition 3.7. \square

Corollary 3.11. Let \mathcal{A} be a connected cluster algebra of finite type or of skew-symmetric finite mutation type, then an automorphism of $E_{\mathcal{A}}$ is determined by the image of any fixed seed Σ and the images of the seeds adjacent to Σ . More precisely, let $\Sigma = (\mathbf{x}, B)$ be a seed on $E_{\mathcal{A}}$, then an automorphism $\Phi : E_{\mathcal{A}} \rightarrow E_{\mathcal{A}}$ is determined by a pair (Σ', ϕ) , where $\Sigma' = (\mathbf{x}', B')$ is a seed on $E_{\mathcal{A}}$ and $\phi : \mathbf{x} \rightarrow \mathbf{x}'$ is a bijection such that $\Phi(\Sigma) = \Sigma'$ and $\Phi(\mu_x(\mathbf{x})) = \mu_{\phi(x)}(\mathbf{x}')$ for any $x \in \mathbf{x}$.

Proof. If \mathcal{A} is of finite type of rank 2 and type of F_4 , then the conclusion is clearly. Otherwise, note that a cluster automorphism is determined by such a pair (Σ', ϕ) , thus the proof follows from Theorem 3.8 and Theorem 3.10. \square

Acknowledgements

Both authors thank for anonymous reviewer's careful reading and many valuable suggestions. In particular, the reference [6] is pointed out by the reviewer.

References

- [1] Assem I, Schiffler R, Shramchenko V. Cluster automorphisms. Proceedings of the London Mathematical Society, 2012, 104(6):1271-1302.
- [2] Irelli C G, Keller B, Labardini-Fragoso D, Plamondon P G. Linear independence of cluster monomials for skew-symmetric cluster algebras. Compositio Mathematica, 2013, 149(10):1753-1764.

- [3] Chang W, Zhu B. On rooted cluster morphisms and cluster structures in 2-Calabi-Yau triangulated categories. *Journal of Algebra*, 2016, 458:387-421.
- [4] Chang W, Zhu B. Cluster automorphism groups of cluster algebras with coefficients. *Science China Mathematics*, 2016, 59(10): 1919-1936.
- [5] Chang W, Zhu B. Cluster automorphism groups of cluster algebras of finite type. *Journal of Algebra*, 2016, 447:490-515.
- [6] Derksen H, Owen T. New graphs of finite mutation type. *Electronic Journal of Combinatorics*. 15 (2008), no. 1, Research Paper 139, 15 pp.
- [7] Fomin S, Shapiro M, Thurston D. Cluster algebras and triangulated surfaces. Part I: Cluster complexes. *Acta Mathematica*, 2008, 201:83-146.
- [8] Felikson A, Shapiro M, Tumarkin P. Cluster algebras of finite mutation type via unfoldings. *International Mathematics Research Notices*, 2011: rnr072.
- [9] Felikson A, Shapiro M, Tumarkin P. Skew-symmetric cluster algebras of finite mutation type. *Journal of the European Mathematical Society*, 2012, 14(4): 1135-1180.
- [10] Fomin S, Zelevinsky A. Cluster algebras. I. Foundations. *Journal of the American Mathematical Society*, 2002, 15(2), 497-529.
- [11] Fomin S, Zelevinsky A. Cluster algebras. II. Finite type classification. *Inventiones Mathematicae*, 2003, 154(1):63-121.
- [12] Fomin S, Zelevinsky A. Y-systems and generalized associahedra. *Annals of Mathematics. Second Series*, 2003, 158(3):977-1018.
- [13] Fomin S, Zelevinsky A. Cluster algebras IV: Coefficients. *Compositio Mathematica*, 2007, 143:112-164.
- [14] Gekhtman M, Shapiro M, Vainshtein A. On the properties of the exchange graph of a cluster algebra, *Mathematical Research Letters*, 2008, 15(2):321-330.
- [15] Keller B. Cluster algebras and derived categories. arXiv:1202.4161. 60 pages.
- [16] Saleh I. Exchange maps on cluster algebras, *International Electronic J. Algebra*, 2014, 16: 1-15.

Wen Chang

School of Mathematics and Information Science, Shaanxi Normal University, Xi'an 710062, China &
 Department of Mathematical Sciences, Tsinghua University, Beijing 10084, China
 Email: changwen161@163.com

Bin Zhu

Department of Mathematical Sciences, Tsinghua University, Beijing 10084, China
 Email: zhu-b@mail.tsinghua.edu.cn