

# Superstatistical cluster decay

Grzegorz Wilk<sup>a</sup>, Zbigniew Włodarczyk<sup>b</sup>

<sup>a</sup>National Centre for Nuclear Research, Department of Fundamental Research, Hoża 69, 00-681 Warsaw, Poland

<sup>b</sup>Institute of Physics, Jan Kochanowski University, Świętokrzyska 15, 25-406 Kielce, Poland

---

## Abstract

We provide an overview of Tsallis statistics, presented as a special case of superstatistics and applied to the multiparticle processes described by the statistical cluster model. This model combines Boltzman statistics applied to hadronization of clusters and superstatistics induced by fluctuations of their Lorentz factors. It results in a Tsallis-like distribution of transverse momenta with some scale, the origin of which is discussed. We show that this distribution becomes a Tsallis distribution in a special case, namely when one combines fluctuations of the Lorentz factor and temperature inside the cluster, given by beta and gamma distributions, respectively.

**Keywords:** nonextensive statistics, statistical cluster model, relativistic temperature fluctuations

**PACS:** 05.90.+m 12.40.Ee 13.85.Hd

---

## 1. Introduction

Quasi-power like distributions, exemplified by the Tsallis distribution [1],

$$h(E) = \frac{2-q}{T} \left[ 1 - (1-q) \frac{E}{T} \right]^{\frac{1}{1-q}} \quad (1)$$

are nowadays recognized as a legitimate tool for use in describing different distributions of this type (usually with  $E$  replaced by transverse momentum  $p_T$ ). This is the more so when it was realized that, albeit based on nonextensive statistical considerations, it is equivalent to the so called Hagedorn distribution [2] (known in the field of particle physics)<sup>1</sup>,

$$h(E) = \frac{n-1}{nT} \left( 1 + \frac{E}{nT} \right)^{-n}, \quad (2)$$

where for  $n = \frac{1}{q-1}$  both distribution coincide. The most recent examples of different phenomenological applications of one of these formulas can be found in [4, 5, 6, 7, 8]

(and references therein<sup>2</sup>. It is remarkable that such a distribution also fits recent LHC data on transverse momentum distributions extending up to  $p_T = 200$  GeV, with cross sections dropping through  $\sim 14$  orders of magnitude, cf. for example [11]. There are numerous ways besides nonextensive statistics which lead to distributions (1) or (2), like, for example, fluctuations of the scale parameter (especially the so called superstatistics), preferential attachment or multiplicative noise - to name only a few (see [12, 13] for recent reviews)<sup>3</sup>.

## 2. Scale parameter fluctuations

Let us concentrate on the case of superstatistics (understood as a superposition of two different statistics relevant

---

Email addresses: wilk@fuw.edu.pl (Grzegorz Wilk), zbigniew.wlodarczyk@ujk.edu.pl (Zbigniew Włodarczyk)

<sup>1</sup>Actually, it was originally proposed in [3].

<sup>2</sup>One should mention that the first papers applying  $q$ -statistics to scattering data in high energy physics dates back to [9] and [10], the second also pointing out the connection to Hagedorn's theory.

<sup>3</sup>Actually, the LHC results mentioned above were also interpreted in the spirit of a Tsallis-like distribution, emerging in this case, under some approximations, from the hard collisions of partons described by QCD methods and followed by showering/hadronization processes into hadrons [14].

to driven nonequilibrium systems with a stationary state and intensive parameter fluctuations) [15]. It is based on the observation that the Tsallis distribution (1) appears in a quite natural way as a result of  $T'$  scale fluctuations in the BG distribution,

$$f(E) = \frac{1}{T'} \exp\left(-\frac{E}{T'}\right), \quad (3)$$

when these fluctuations are described by a gamma distribution in the variable  $(1/T')$  [16, 17, 18, 13],

$$g(T') = \frac{1}{nT\Gamma(n)} \left(\frac{nT}{T'}\right)^{n+1} \exp\left(-\frac{nT}{T'}\right), \quad n = \frac{1}{q-1}, \quad (4)$$

where the parameter  $q$  defines the strength of the  $T'$  fluctuations,

$$q = 1 + \frac{\text{Var}(1/T')}{\langle 1/T' \rangle^2}. \quad (5)$$

It should be mentioned at this point that depending on the statistical properties of the fluctuations, one obtains different effective statistical mechanical descriptions. Tsallis statistics follow from the above gamma distribution of an intensive variable, but other classes of generalized statistics can be obtained as well and, for small variance of the fluctuations, they all behave in a universal way. From the thermal perspective, this corresponds to a situation in which the system considered, for example a decaying cluster, is not homogeneous, but has different temperatures in different parts, which are fluctuating around some mean temperature  $T$  (an essentially identical result is obtained when the temperature fluctuates from one cluster to another remaining constant in a given cluster). It must therefore be described by two parameters: a mean temperature  $T$  and the mean strength of the fluctuations defined by  $q$ . The gamma distribution (4) itself emerges in a quite natural way, either from thermal considerations corresponding to the purely multiplicative noise [16], or in a stochastic approach without correlation between the assumed additive and multiplicative noises [17]<sup>4</sup>.

We shall now discuss this situation in more detail. Note that the distribution  $g(T')$  in Eq. 4 is in fact the product

of two distributions:

$$g(T') \sim g_1(T') \cdot g_2(T'), \quad (6)$$

where

$$g_1(T') = \left(\frac{1}{T'}\right)^\kappa \quad (7)$$

is a scale free power law, and

$$g_2(T') = \exp\left(-\frac{nT}{T'}\right), \quad (8)$$

is an exponential, which for small values of  $T'$  cuts off the distribution  $g(T')$ , with the scale parameter  $T$  determining how fast this cut-off is.

Suppose now that we fluctuate the BG distribution (3) using only the  $g_1(T')$  part of the distribution  $g(T')$ . As a result we obtain the following scale free distribution:

$$h_1(E) = \int_0^\infty dT' g_1(T') \exp\left(-\frac{E}{T'}\right) \propto E^{-\kappa+1}. \quad (9)$$

The scale appears when one cuts-off somehow the small values of  $T'$ . For example, a sharp cut-off of the small  $T'$ , i.e., limiting  $T'$  to  $T' > T$  only, results in the following distribution:

$$h_2(E) = \int_T^\infty dT' g_1(T') \exp\left(-\frac{E}{T'}\right) \propto E^{-\kappa+1} \left[ \Gamma(\kappa-1) - \Gamma\left(\kappa-1, \frac{E}{T}\right) \right], \quad (10)$$

where  $\Gamma(x, y)$  is an incomplete gamma function. The factor  $\Gamma(\kappa-1) - \Gamma(\kappa-1, E/T)$  now suppresses the power distribution (9) for small values of  $E$ . However, the form of this suppression is not the same as in the Tsallis distribution. This can be seen by comparing the expansion

$$\left(\frac{E}{T}\right)^{-\kappa} \cdot \left[ \Gamma(\kappa-1) - \Gamma\left(\kappa-1, \frac{E}{T}\right) \right] = \frac{1}{\kappa} + \sum_{i=1}^\infty \frac{\Gamma(i+\kappa)}{\Gamma(i+\kappa+1)\Gamma(i+1)} \left(-\frac{E}{T}\right)^i \quad (11)$$

with the corresponding expansion of the Tsallis factor:

$$\left(1 + \frac{E}{\kappa T}\right)^{-\kappa} = \frac{1}{\kappa} + \sum_{i=1}^\infty \frac{\Gamma(i+\kappa)}{\Gamma(1+\kappa)\Gamma(i+1)} \left(-\frac{E}{\kappa T}\right)^i. \quad (12)$$

However, if we smooth out this suppression by replacing the previous sharp limitation of the integrand by some smooth suppression factor, provided, for example, by the exponential function  $g_2(T')$  from Eq. (8), we get, as result, the Tsallis distribution defined in Eq. (1).

<sup>4</sup>The first identification of local temperature fluctuations as the source of Tsallis distributions observed in high energy physics data was done in [16], later on it was also used in analysis of data from cosmic rays events in [19].

### 3. Approximate description of cluster decay

Recently, a new statistical cluster decay model of hadronization has been analyzed numerically, showing that even without resorting to approaches of the kind mentioned above, the resulting distribution of transverse momenta follows rather closely a Tsallis distribution [20], albeit not identically so. In this work the hadronic clusters produced were supposed to decay purely thermally (i.e. by following the usual exponential Boltzmann-Gibbs (BG) statistics) but, at the same time, were supposed to move in the transverse direction with a fluctuating (transverse) Lorentz factor distributed according to the assumed power law. It turns out that the combination of both distributions follows (at least numerically) a quasi-power like distribution, closely resembling a Tsallis distribution. Note that, according to what was said before regarding superstatistics, production and decay of such clusters can be regarded as an example of superstatistics at work (and not necessarily resulting in a Tsallis distribution).

In this work we shall investigate this phenomenon in more detail, aiming to obtain its analytical justification and a deeper understanding from the nonextensive statistical point of view. We start with statistical cluster decay, discussed recently in [20]. The distribution of transverse momenta proposed there is<sup>5</sup>,

$$f(p_T) \propto \int d\gamma_T K_0\left(\gamma_T \frac{m_T}{T}\right) I_0\left(\sqrt{\gamma_T^2 - 1} \frac{p_T}{T}\right) g(\gamma_T), \quad (13)$$

where  $m_T$  is the transverse mass and  $\gamma_T$  denotes the transverse Lorentz factor ( $K_0$  and  $I_0$  are modified Bessel functions of the, respectively, second and first kind). This factor is then supposed to fluctuate according to some power-like distribution,

$$g(\gamma_T) \sim \gamma_T^{-\kappa}, \quad (14)$$

i.e. according to the  $g_1$  function from Eq. (7) with  $T'$  replaced by  $\gamma_T$ . Numerical calculations in [20] show that whereas for large  $p_T$  one has

$$f(p_T) \sim p_T^{-\kappa-1}, \quad (15)$$

there is a suppression for small values of  $p_T$ . To describe its origin note that the  $p_T$  distribution given by the product

of two Bessel functions from Eq. (13),

$$\tilde{f}(p_T) \sim K_0\left(\gamma_T \frac{m_T}{T}\right) I_0\left(\sqrt{\gamma_T^2 - 1} \frac{p_T}{T}\right), \quad (16)$$

depends strongly on the transverse Lorentz factor of the decaying cluster,  $\gamma_T$ ,

$$\tilde{f}(p_T) \sim \begin{cases} \exp(-p_T) & \text{for } \gamma_T = 1, \\ \frac{1}{p_T} & \text{for large } \gamma_T. \end{cases} \quad (17)$$

Because:

- $K_0(x)I_0(y) \simeq \frac{\exp(-x+y)}{2\sqrt{xy}}$  for  $x, y \gg \frac{1}{4}$ ,
- $\gamma_T - \sqrt{\gamma_T^2 - 1} \simeq \frac{1}{2\gamma_T}$  for large  $\gamma_T$ <sup>6</sup>,
- $m_T \simeq p_T$  for  $p_T \gg m$ ,

therefore we can rewrite Eq. (13) in the following way:

$$\begin{aligned} f(p_T) &\propto \int_1^\infty d\gamma_T \exp\left(-\frac{p_T}{T\gamma_T}\right) p_T^{-1} \gamma_T^{-\kappa-1} = \\ &= p_T^{-\kappa-1} \left[ \Gamma(\kappa) - \Gamma\left(\kappa, \frac{p_T}{T}\right) \right]. \end{aligned} \quad (18)$$

Note that because  $\gamma_T$  is always limited to values  $\gamma_T \geq 1$  only, there is a natural cut-off in Eq. (13), which makes it similar and comparable to Eq. (10) discussed above. As a result, here the power-like distribution of  $p_T$  is also suppressed in the region of small values of transverse momenta. The factor  $[\Gamma(\kappa) - \Gamma(\kappa, p_T/T)]$  introduces the behavior  $\sim p_T^\kappa$  for small  $p_T$  and remains constant for large values of  $p_T$ . To summarize, such an approximation must therefore lead to a result which is very near to the numerical results presented in [20], and not coinciding with a Tsallis distribution.

Because of the important role played by fluctuations of the Lorentz factor  $\gamma_T$  described by the distribution  $g(\gamma_T)$  in Eq. (13), it is interesting to speculate on its possible origin emerging from the quark structure of the colliding nucleons. Let us denote by  $x = p_T/p_h$  the fraction of transverse momentum of the parton,  $p_T$ , with respect to

<sup>5</sup>It is worth noting that cluster models have a very long tradition, a similar formula was already derived in [21].

<sup>6</sup>For  $\gamma_T > 5$  the difference is less than 1%. We expand in series in  $z = 1/\gamma_T$ :  $1/z - \sqrt{1/z^2 - 1} = 1/z - \sqrt{1 - z^2}/z \approx z/2$ .

the momentum of hadron,  $p_h$ , and let us assume that the density of the parton distribution is

$$w(x) = Ax^a(1-x)^b. \quad (19)$$

In the center of mass system the Lorentz factor of the cluster formed by the collision of partons with fractions of momenta equal to, respectively,  $x_1$  and  $x_2$ , is given by

$$\gamma_T = \frac{x_1 + x_2}{2\sqrt{x_1 x_2}}. \quad (20)$$

Notice that if  $x_1 = x_2$  then  $\gamma_T = 1$ , it becomes greater than unity only for nonsymmetric values of  $x_{1,2}$ . It is convenient to change variables to

$$x_1 = \rho \cos \phi, \quad x_2 = \rho \sin \phi \quad (21)$$

in which

$$\frac{x_2}{x_1} = \tan \phi, \quad x_1^2 + x_2^2 = \rho^2 \quad \text{and} \quad \gamma_T = \frac{1 + \tan \phi}{2\sqrt{\tan \phi}}. \quad (22)$$

The Jacobian of this transformation is equal  $|J| = \rho$ . The distribution of the Lorentz factor  $\gamma_T$  is now given by

$$\begin{aligned} \frac{1}{\sigma} \frac{d\sigma}{d\gamma_T} &= A^2 \int d\rho \rho^{1+2a} \cdot \int d\phi \left( \frac{\tan \phi}{1 + \tan^2 \phi} \right)^a \cdot \\ &\cdot \left( 1 - \rho \frac{1 + \tan \phi}{\sqrt{1 + \tan^2 \phi}} + \rho^2 \frac{\tan \phi}{1 + \tan^2 \phi} \right)^b \cdot \\ &\cdot \delta \left( \gamma_T - \frac{1 + \tan \phi}{2\sqrt{\tan \phi}} \right) \end{aligned} \quad (23)$$

Denoting

$$y = \gamma_T - \frac{1 + \tan \phi}{2\sqrt{\tan \phi}} \quad (24)$$

we can write

$$\int d\phi (\dots) \delta(y) = \int dy (\dots) \sum_i \frac{\delta(y)}{\left| \frac{dy}{d\phi} \right|_{\phi_{0i}}} \quad (25)$$

where  $\phi_{0i}$  are solutions of Eq. (24) for  $y = 0$ . Now,

$$\frac{dy}{d\phi} = \frac{1 - \tan \phi}{4 \tan \phi \sqrt{\tan \phi}} (1 + \tan^2 \phi) \quad (26)$$

and there are two solutions,  $\phi_{0(1,2)}$ , for which we have

$$\tan \phi_{0(1,2)} = 2\gamma_T^2 - 1 \pm 2\gamma_T \sqrt{\gamma_T^2 - 1}, \quad (27)$$

$$\text{where } \tan \phi_{01} \geq 1 \quad \text{and} \quad \tan \phi_{02} \leq 1. \quad (28)$$

It means therefore that Eq. (23) has now two terms with different limits of integrations over  $\rho$  in each of them,

$$\rho \in (0, \rho_1); \quad \rho_1 = \frac{1}{\sin \phi_{01}} = \frac{\sqrt{1 + \tan^2 \phi_{01}}}{\tan \phi_{01}} \quad (29)$$

(corresponding to  $\phi \in (\pi/4, \pi/2)$ )

$$\rho \in (0, \rho_2); \quad \rho_2 = \frac{1}{\cos \phi_{02}} = \sqrt{1 + \tan^2 \phi_{02}}. \quad (30)$$

(corresponding to  $\phi \in (0, \pi/4)$ ). However, in both cases we cover the whole range of  $x_{1,2}$ . For the choice (29) it is done preserving all the time inequality  $x_2 \geq x_1$ , whereas for the choice (30) it is done by keeping all the time  $x_2 \leq x_1$ . In both cases the final result must be the same. Therefore, choosing for example, Eq. (30,) we have that

$$\begin{aligned} \frac{1}{\sigma} \frac{d\sigma}{d\gamma_T} &= 2A^2 \frac{1}{\left| \frac{dy}{d\phi} \right|_{\phi_{02}}} \left( \frac{\tan \phi_{02}}{1 + \tan^2 \phi_{02}} \right)^a \int_0^{\rho_2} d\rho \rho^{1+2a} \cdot \\ &\cdot \left( 1 - \rho \frac{1 + \tan \phi_{02}}{\sqrt{1 + \tan^2 \phi_{02}}} + \rho^2 \frac{\tan \phi_{02}}{1 + \tan^2 \phi_{02}} \right)^b. \end{aligned} \quad (31)$$

For  $\gamma_T \gg 1$  it can be numerically approximated by

$$\frac{1}{\sigma} \frac{d\sigma}{d\gamma_T} \propto \frac{1}{\gamma_T^\kappa - 1} \simeq \gamma_T^{-\kappa}, \quad (32)$$

where for  $a = 1$  and  $b = 3$  one has  $\kappa = 5$ . These value seem to be a reasonable first guess in the case of nucleon-nucleon collisions ( $b = 2m - 1$ , where  $m$  denotes the number of spectator quarks, whereas the parameter  $a$  is given by the expected mean value  $\langle x \rangle = 1/3$ ). The exact form of Eq. (32) for  $a = 1$  and  $b = 3$  would be

$$\frac{1}{\sigma} \frac{d\sigma}{d\gamma_T} = 400 \frac{r^2}{\sqrt{\gamma_T^2 - 1}} \left( \frac{1}{35} - \frac{3r}{70} + \frac{r^2}{42} - \frac{r^3}{210} \right) \quad (33)$$

with

$$r = \tan \phi_{02} = 2\gamma_T^2 - 1 - 2\gamma_T \sqrt{\gamma_T^2 - 1}. \quad (34)$$

#### 4. Fluctuations of relativistic temperature

We close these remarks with the following observation. In Eq. (18) one encounters the scale factor  $T\gamma_T$ , not  $\gamma_T$

alone. Some time ago such a quantity was proposed as a relativistic temperature [22],

$$T^* = \gamma_T T, \quad (35)$$

(albeit in this case it would rather be a relativistic transverse temperature, whatever that means). From the discussion following Eq. (10) it follows that to get a proper Tsallis distribution when fluctuating the scale parameter, one has to use the full gamma function, Eq. (4). Therefore, fluctuating  $T\gamma_T$  should bring the distribution of the transverse momenta to the desired final Tsallis form. This, however, would mean fluctuation of the relativistic temperature,  $T^*$ , but so far its proper form is still under discussion [23]. Nevertheless, keeping this reservation in mind let us elaborate on such a possibility in more detail.

Assuming a beta distribution for fluctuations of  $1/\gamma_T$ ,

$$b\left(\frac{1}{\gamma_T}\right) = \frac{1}{\Gamma(\kappa+1)\Gamma(\alpha+1)} \left(\frac{1}{\gamma_T}\right)^\kappa \left(1 - \frac{1}{\gamma_T}\right)^\alpha, \quad (36)$$

and a gamma distribution for fluctuations of  $1/T$ ,

$$g\left(\frac{1}{T}\right) = \frac{nT_0}{\Gamma(n)} \left(\frac{nT_0}{T}\right)^{n-1} \exp\left(-\frac{nT_0}{T}\right), \quad (37)$$

one gets the joint distribution

$$g'\left(\frac{1}{T}, \frac{1}{\gamma_T}\right) = b\left(\frac{1}{\gamma_T}\right) \cdot g\left(\frac{1}{T}\right), \quad (38)$$

and can define the function

$$g'\left(\frac{1}{T^*}, \frac{1}{\gamma_T}\right) = \frac{(nT_0)^n}{\Gamma(n)\Gamma(\kappa+1)\Gamma(\alpha)} \cdot \left(\frac{1}{T^*}\right)^n \left(\frac{1}{T} - \frac{1}{T^*}\right)^\alpha \exp\left(-\frac{nT_0}{T}\right). \quad (39)$$

In the case when the parameters of the two components of this joint distribution are related in a certain way, for example if  $\alpha = n - 1 - \kappa$ , fluctuations of the relativistic temperature  $T^*$  are again given by a gamma distribution (but this time with a changed shape parameter):

$$\begin{aligned} g'\left(\frac{1}{T^*}\right) &= \int_{1/T^*}^{\infty} g'\left(\frac{1}{T^*}, \frac{1}{T}\right) d\left(\frac{1}{T}\right) = \\ &= \frac{n-1-\kappa}{\Gamma(\kappa+1)\Gamma(n+1)} \left(\frac{nT_0}{T^*}\right)^\kappa \exp\left(-\frac{nT_0}{T^*}\right). \end{aligned} \quad (40)$$

This means that one could therefore consider the existence of fluctuations of the relativistic temperature  $1/T^*$ , which are again given by a gamma distribution with parameter  $\kappa$  defining the size of these fluctuations. Relative fluctuations, defined as

$$\omega(z) = \frac{\text{Var}(1/z)}{\langle 1/z \rangle^2}, \quad (41)$$

satisfy the relation

$$\omega(T^*) = \frac{\omega(T) + \omega(\gamma_T) + 2\omega(T)\omega(\gamma_T)}{1 + 3\omega(T)} \quad (42)$$

which connects fluctuations of  $T$  and  $\gamma_T$ <sup>7</sup>. The corresponding nonextensivity parameter is

$$q^* - 1 = \frac{1}{n^*} = \omega(T^*). \quad (43)$$

Note that the parameter  $n$ , which defines fluctuations of temperature in the decaying cluster, cf. Eq. (4), is proportional to heat capacity of the system under constant volume,  $n \propto C_V$ . In the case considered here it is much greater than  $\kappa$ ,  $n \gg \kappa$ . On the other hand, as shown in [6, 12, 25], the observed power index in the corresponding Tsallis distribution is smaller than  $n$ . The above consideration provides a simple and natural explanation of this fact, the power index is defined by both  $n$  and  $\kappa$  (i.e. by fluctuations of the relativistic temperature).

## 5. Concluding remarks

To summarize, the statistical cluster decay mechanism proposed in [20] and discussed above, is yet another example of supertstatistics (which is not necessarily connected with a Tsallis distribution). Fluctuations of the Lorentz factor given by the distribution  $g(\gamma_T)$  fully specify the slope parameter of the transverse momentum distribution in the region of large values of  $p_T$ . If they are given by a scale free power-like distribution, as in Eq. (14), the resultant distribution of  $p_T$  is also a scale-free, power-like one. Its behavior for small values of  $p_T$  is dictated by the fact that the Lorentz factor is defined only

<sup>7</sup>This is another example of the generalized thermodynamic fluctuation relations discussed previously in [24].

for  $\gamma_T \geq 1$ , therefore there is always a natural cut-off in  $g(\gamma_T)$ , eliminating  $\gamma_T < 1$ . As a result one gets a distribution which is not a Tsallis distribution, remaining, however, quite close to it numerically.

One can therefore invent a more general distribution, which is a product of a beta distribution in  $1/\gamma_T$  and a gamma distribution for the parameter  $1/T$ . If the parameters defining these distribution are related in some specific way, the resultant distribution is again a gamma distribution with a modified shape. In this case one gets the true Tsallis distribution for the spectrum of the transverse momenta with the slope parameter determined by  $n^*$  from Eq. (43). Note that in this case it can be argued that it is determined by fluctuations of the relativistic temperature  $T^* = \gamma_T T$  (modulo reservations concerning its proper form as mentioned before). Combining this result with the connection between fluctuations of temperature and specific heat,  $\omega(T) \simeq 1/C_V$ , one realizes that fluctuations  $\omega(\gamma_T)$  provide a natural source of fluctuations resulting in the observed values of the slope parameter  $n^*$ , which is smaller than  $n \simeq C_V$ . In such a way the puzzle discussed in [6, 12, 25] could find its natural solution.

We close by concluding that, as shown here, the realistic superstatistics for high energy physics could have deviations from the standard Gamma distribution and hence from Tsallis statistics, although in an approximate sense Tsallis statistics is still recovered. This provides yet another support for the idea expressed in [26] that general superstatistics, based on more general distributions than Gamma distribution, might be relevant in high energy physics.

## Acknowledgments

This research was supported in part by the National Science Center (NCN) under contract Nr 2013/08/M/ST2/00598. We would like to thank warmly Dr Nicholas Keely for reading this manuscript.

## References

- [1] C. Tsallis, J. Statist. Phys. **52** (1988) 479; Eur. Phys. J. A **40** (2009) 257; Contemporary Phys. **55** (2014) 179; Acta Phys. Pol. B **46** (2015) 1089 and *Introduction to Nonextensive Statistical Mechanics* (Springer, 2009). For an updated bibliography on this subject, see <http://tsallis.cat.cbpf.br/biblio.htm>.
- [2] G. Arnison et al (UA1 Collab.), Phys. Lett. B **118** (1982) 167 and Nucl. Phys. B **309** (1988) 405; R. Hagedorn, Riv. Nuovo Cimento **6** (1984) 1.
- [3] C. Michael and L. Vanryckeghem, J. Phys. G **3** (1977) L151; C. Michael, Prog. Part. Nucl. Phys. **2** (1979) 1.
- [4] M. Rybczyński and Z. Włodarczyk, Eur. Phys. J. C **74** (2014) 2785.
- [5] L. Marques, J. Cleymans and A. Deppman, Phys. Rev. D **91** (2015) 054025.
- [6] T. S. Biró, P. Ván, G. G. Barnaföldi and K. Ürmösy, Entropy **16** (2014) 6497.
- [7] P. K. Khandai, P. Sett, P. Shukla and V. Singh, Int. J. Mod. Phys. A **28** (2013) 1350066 and J. Phys. G **41** (2014) 025105.
- [8] Bao-Chun Li., Ya-Zhou Wang and Fu-Hu Liu, Phys. Lett. B **725** (2013) 352.
- [9] I. Bediaga, E.M.F. Curado and J. Miranda, Physica A **286** (2000) 156.
- [10] C. Beck, Physica A **286** (2000) 164.
- [11] V. Khachatryan et al. (CMS Collaboration), JHEP **02** (2010) 041 and JHEP **08** (2011) 086; Phys. Rev. Lett. **105** (2010) 022002; G. Aad et al. (ATLAS Collaboration), New J. Phys. **3** (2011) 053033; B. Abelev et al. (ALICE Collaboration), Phys. Lett. B **722** (2013) 262.
- [12] G. Wilk and Z. Włodarczyk, Entropy **17** (2015) 384 and Acta Phys. Pol. B **46** (2015) 1103.
- [13] G. Wilk and Z. Włodarczyk, *Quasi-power laws in multiparticle production processes*, arXiv:1503.08704v1[hep-ph], in press in Chaos Solitons and Fractals (2015), available online 14 May 2015; DOI:10.1016/j.chaos.2015.04.016).

- [14] C.-Y. Wong and G. Wilk, *Acta Phys. Pol. B* **43** (2012) 2047 and *Phys. Rev. D* **87** (2013) 114007;  
C.-Y. Wong, G. Wilk, L.J.L. Cirto and C. Tsallis, *Rev. D* **91** (2015) 114027.
- [15] C. Beck and E. G. D. Cohen, *Physica A* **322** (2003) 267;  
F. Sattin, *Eur. Phys. J. B* **49** (2006) 219.
- [16] G. Wilk and, Z. Włodarczyk, *Phys. Rev. Lett.* **84** (2000) 2770 and *Chaos, Solitons, Fractals* **13** (2002) 581.
- [17] T. S. Biró and A. Jakovác, *Phys. Rev. Lett.* **94** (2005) 132302.
- [18] C. Beck, *Phys. Rev. Lett.* **87** (2001) 180601.
- [19] C. Beck, *Physica A* **331** (2004) 173.
- [20] A. Białas, *Phys. Lett. B* **747** (2015) 150.
- [21] F. Hayot, F. S. Henyey and M. Lebellac, *Nucl. Phys. B* **80** (1974) 77.
- [22] H. Ott, *Z. Phys.* **175** (1963) 70.  
H. Arzelies, *Nuovo Cim.* **41B** (1966) 81.
- [23] T. S. Biró and P. Ván, *Eur. Phys. Lett.* **89** (2010) 30001.
- [24] G. Wilk and, Z. Włodarczyk, *Physica A* **390** (2011) 3566.
- [25] T. S. Biró, G. G. Barnaföldi and P. Ván, *Physica A* **417** (2015) 215.
- [26] C. Beck, *Eur. Phys. J. A* **267** (2009) 267.