

Relativistic and nonrelativistic annihilation of dark matter: a sanity check using an effective field theory approach

Mirco Cannoni

Departamento de Física Aplicada, Facultad de Ciencias Experimentales, Universidad de Huelva, 21071 Huelva, Spain

We find an exact formula for the thermally averaged cross section times the relative velocity $\langle\sigma v_{\text{rel}}\rangle$ with relativistic Maxwell-Boltzmann statistics. The formula is valid in the effective field theory approach when the masses of the annihilation products can be neglected compared with the dark matter mass and cut-off scale. The expansion at $x = m/T \gg 1$ directly gives the nonrelativistic limit of $\langle\sigma v_{\text{rel}}\rangle$ which is usually used to compute the relic abundance for heavy particles that decouple when they are nonrelativistic. We compare this expansion with the one obtained by expanding the total cross section $\sigma(s)$ in powers of the nonrelativistic relative velocity v_r . We show the correct invariant procedure that gives the nonrelativistic average $\langle\sigma_{nr} v_r\rangle_{nr}$ coinciding with the large x expansion of $\langle\sigma v_{\text{rel}}\rangle$ in the comoving frame. We explicitly formulate flux, cross section, thermal average, collision integral of the Boltzmann equation in an invariant way using the true relativistic relative v_{rel} , showing the uselessness of the Møller velocity and further elucidating the conceptual and numerical inconsistencies related with its use.

I. INTRODUCTION

While there are compelling evidences in astrophysics and cosmology that most of the mass of the Universe is composed by a new form of non baryonic dark matter (DM), there is a lack of evidence for the existence of new physics at LHC and other particle physics experiments. On the theory side, many specific models with new particles and interactions beyond the standard model have been proposed to account for DM.

Under these circumstances where no clear indications in favour of a particular model are at our disposal, the phenomenology of DM as been studied in a model independent way using an effective field theory approach, see for example [1–23].

Measurements of the parameters of standard model of cosmology [24, 25] furnish the present day mass density of DM, the relic abundance, $\Omega h^2 \sim 0.11$ with an uncertainty at the level of 1%. Any model that pretends to account for DM must reproduce this number, which, on the other hand, sets strong constraints on the free parameters of the model.

When the DM particles are weakly interacting massive particles that decouple from the primordial plasma at a temperature when they are nonrelativistic, the relativistic averaged annihilation rate $\langle\sigma v_{\text{rel}}\rangle$ can be well approximated by taking the nonrelativistic average of the first two terms of the expansion of σ in powers of the nonrelativistic relative velocity. With v_{rel} we indicate the *relativistic relative velocity* and with v_r the *nonrelativistic relative velocity*, as defined in B. To describe collisions in a gas, and in particular in the primordial plasma, the reference frame that matters is the comoving frame (COF) where the observer sees the gas at rest as a whole and the colliding particles have general velocities $\mathbf{v}_{1,2}$ without any further specification of the kinematics.

It is thus desirable to formulate cross sections and rates in a relativistic invariant way, such that all the formulas and nonrelativistic expansions are valid automatically in

the COF. Obviously, invariant formulas give the same results in the lab frame (LF), the frame where one massive particle is at rest, and in the center of mass frame (CMF) where the total momentum is zero. We will see that the key for the invariant formulation is v_{rel} .

On the contrary, in DM literature [26] instead of v_{rel} it is used the so-called Møller velocity \bar{v} , see B. That this is incorrect was already discussed in Ref. [27] but papers using \bar{v} continue to appear. The problem with \bar{v} , which is not the relative velocity, is its non invariant and nonphysical nature, for it can take values larger than c .

In this paper we first find an exact formula for $\langle\sigma v_{\text{rel}}\rangle$ as a function of $x = m/T$ calculated with the relativistic Maxwell-Boltzmann statistics. The formula is valid in the effective field theory framework such that the masses of the annihilation products can be neglected compared with the DM and the cut-off scale. For concreteness we work with fermion DM. We find the thermal functions corresponding to various interactions and in particular those corresponding to s and p wave scattering in the nonrelativistic limit which is given by the expansion at $x \gg 1$. This is done in Section II, and A contains some mathematical results needed for the derivation of the exact formula and its asymptotic expansions.

Then, in Section III, we present the correct invariant method for obtaining the same expansion by expanding the total annihilation cross section $\sigma(s)$ in powers of v_r .

We then discuss in Section IV the problems with the use \bar{v} , while the numerical impact on the relic abundance of some incorrect methods employed in literature is evaluated in Section V.

B is preparatory for the whole paper: we remind how relativistic flux, cross section, rate, collision term of the Boltzmann equation and thermal averaged rate can be defined in the invariant way in terms of v_{rel} showing the uselessness of the Møller velocity.




Figure 1. s and t channel annihilation diagrams reducing to the effective vertex corresponding to the lagrangian Eq. (1).

II. EXACT FORMULA FOR THE THERMAL AVERAGE IN THE EFFECTIVE APPROACH

We consider a DM fermion field χ that couples to other fermion fields ψ through an effective dimension-6 operator of the type

$$\mathcal{L}_\Lambda = \frac{\lambda_a \lambda_b}{\Lambda^2} (\bar{\chi} \Gamma_a \chi) (\bar{\psi} \Gamma_b \psi). \quad (1)$$

The DM particles can be of Dirac or Majorana nature and have mass m , while ψ are the standard model fermions or new ones. Here $\lambda_{a,b}$ are dimensionless coupling associated with the interactions described by combination of Dirac matrices $\Gamma_{a,b}$. Λ is the energy scale below which the effective field theory is valid. In the exact theory Λ corresponds to the mass of a heavy scalar or vector boson mediator that appears in the propagators. The ψ masses can be neglected compared to Λ and m . The exchange of a heavy mediator with mass Λ may take place in the s -channel and/or in t -channel, as depicted in Figure 1, depending on the specific model.

A. Exact formula for $\langle \sigma v_{\text{rel}} \rangle$

In all generality, for $2 \rightarrow 2$ processes, the matrix elements depend only on two independent Mandelstam variables, for example s and t , and the squared matrix element is dimensionless. After integrating over the CMF angle, for example, the only remaining dependence is on s and m . Any amplitude related to the operator (1) gives an integrated squared matrix element $|\overline{\mathcal{M}}|^2$ summed over the final spins and averaged over the initial spins that is a simple polynomial of the type

$$w = \int |\overline{\mathcal{M}}|^2 d\cos\theta = p_2 s^2 + p_1 m^2 s + p_0 m^4, \quad (2)$$

with p_0, \dots, p_2 depending on Λ and $\lambda_{a,b}$. To get the formula for $\langle \sigma v_{\text{rel}} \rangle$ in a useful form, it is convenient to define the *reduced cross section*

$$\sigma_0 = \frac{1}{2m^2} \frac{1}{32\pi} w, \quad (3)$$

and the *effective cross section*

$$\sigma_\Lambda = \frac{\lambda_a^2 \lambda_b^2}{4\pi} \frac{m^2}{\Lambda^4}, \quad (4)$$

which contains all the couplings. In terms of the effective cross section (4), and of the dimensionless variable $y =$

$s/(4m^2)$, the reduced cross section Eq. (3) becomes

$$\sigma_0 = \sigma_\Lambda \left(a_2 y^2 + \frac{a_1}{4} y + \frac{a_0}{16} \right), \quad (5)$$

where now a_2, \dots, a_0 are pure numbers. The total unpolarized cross section then is

$$\sigma = \frac{2m^2}{s} \frac{\sqrt{\lambda(s, m_3^2, m_4^2)}}{\sqrt{\lambda(s, m_1^2, m_2^2)}} \sigma_0. \quad (6)$$

We now set $m_1 = m_2 = m$, $m_3 = m_4 = 0$ in Eq. (6) and in Eq. (B20), and change variable to y . Thus Eq. (B20) becomes

$$\langle \sigma v_{\text{rel}} \rangle = \frac{2x}{K_2^2(x)} \int_1^\infty dy \sqrt{y-1} K_1(2x\sqrt{y}) \sigma_0(y). \quad (7)$$

Using the integrals of A, we find

$$\begin{aligned} \langle \sigma v_{\text{rel}} \rangle = \sigma_\Lambda \frac{1}{16} [8a_2 + 2a_1 + (5a_2 + 2a_1 + a_0) \frac{K_1^2(x)}{K_2^2(x)} \\ + 3a_2 \frac{K_3^2(x)}{K_2^2(x)}]. \end{aligned} \quad (8)$$

In the case $m_3 = m_4 = 0$ the pure mass terms do not appear in the cross sections, thus $a_0 = 0$. Furthermore, we can relate a_2 and a_1 each other by an appropriate multiplicative factor,

$$a_1 = k a_2, \quad (9)$$

and express the cross sections as a function of a_2 only. The general formula (8) thus finally becomes

$$\langle \sigma v_{\text{rel}} \rangle = \sigma_\Lambda a_2 \mathcal{F}_k(x), \quad (10)$$

with

$$\mathcal{F}_k(x) = \frac{1}{16} \left(8 + 2k + (5 + 2k) \frac{K_1^2(x)}{K_2^2(x)} + 3 \frac{K_3^2(x)}{K_2^2(x)} \right) \quad (11)$$

the factored out thermal function.

The nonrelativistic thermal average is given by the expansion at $x \gg 1$. Using the asymptotic expansions Eq. (A2) we find

$$\langle \sigma_{nr} v_r \rangle_{nr} = \sigma_\Lambda a_2 \left(1 + \frac{k}{4} - \frac{3k}{8x} \right) + \mathcal{O}(x^{-2}). \quad (12)$$

In the ultrarelativistic limit, $x \ll 1$, using the expansions (A3), the thermal functions behave as $3/x^2$, thus

$$\langle \sigma v_{\text{rel}} \rangle_{ur} \sim \sigma_\Lambda a_2 \frac{3}{x^2} = \frac{\lambda_a^2 \lambda_b^2}{4\pi \Lambda^4} 3a_2 T^2, \quad (13)$$

which is the expected result for massless particles.

The exact integration is possible because the effective operator removes the momentum dependence in the propagators that are reduced to a multiplicative constant

and the assumption $m_3 = m_4 = 0$ allows to simplify the square root $\sqrt{\lambda(s, m_3^2, m_4^2)} = s$ in the cross section (6). For example, with $m_3 = m_4 = m_\psi$, equation (7) becomes

$$\langle \sigma v_{\text{rel}} \rangle = \frac{2x}{K_2^2(x)} \int_{y_0}^{\infty} dy \sqrt{y - \rho} \sqrt{y - 1} K_1(2x\sqrt{y}) \sigma_0(y, \rho).$$

with $\rho = m_\psi^2/m^2$ and $y_0 = 1$ if $m \geq m_\psi$, $y_0 = \rho$ if $m < m_\psi$. In this case the exact integration is not possible but nonrelativistic expansions exist also in the case $\rho = 1$ and $\rho \gg 1$ as we have shown in Ref. [27].

B. Applications

In order to show the thermal behaviour of different interactions, we calculate the cross sections for various operators of the type (1), both for s and t channel annihilation. We list the quantity $\varpi = \Lambda^4/(\lambda_a^2 \lambda_b^2) w$ and the resulting average Eq. (10).

For the s -channel annihilation we find:

1) Scalar: $(\bar{\chi}\chi)(\bar{\psi}\psi)$, $(\bar{\chi}\chi)(\bar{\psi}\gamma^5\psi)$.

$$\varpi = 2s(s - 4m^2), \quad \langle \sigma_S v_{\text{rel}} \rangle = \sigma_\Lambda 2\mathcal{F}_{-4}(x). \quad (14)$$

2) Pseudo-scalar: $(\bar{\chi}\gamma^5\chi)(\bar{\psi}\gamma^5\psi)$, $(\bar{\chi}\gamma^5\chi)(\bar{\psi}\psi)$:

$$\varpi = 2s^2, \quad \langle \sigma_{PS} v_{\text{rel}} \rangle = \sigma_\Lambda 2\mathcal{F}_0(x). \quad (15)$$

3) Chiral: $(\bar{\chi}P_{L,R}\chi)(\bar{\psi}P_{L,R}\psi)$.

$$\varpi = \frac{1}{2}s(s - 2m^2), \quad \langle \sigma_C v_{\text{rel}} \rangle = \sigma_\Lambda \frac{1}{2}\mathcal{F}_{-2}(x). \quad (16)$$

4) Pseudo-vector: $(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{\psi}\gamma_\mu\gamma_5\psi)$, $(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{\psi}\gamma_\mu\psi)$.

$$\varpi = \frac{8}{3}s(s - 4m^2), \quad \langle \sigma_{PV} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{8}{3}\mathcal{F}_{-4}(x) \quad (17)$$

5) Vector: $(\bar{\chi}\gamma^\mu\chi)(\bar{\psi}\gamma_\mu\psi)$, $(\bar{\chi}\gamma^\mu\chi)(\bar{\psi}\gamma_\mu\gamma^5\psi)$.

$$\varpi = \frac{8}{3}s(s + 2m^2), \quad \langle \sigma_V v_{\text{rel}} \rangle = \sigma_\Lambda \frac{8}{3}\mathcal{F}_2(x). \quad (18)$$

6) Vector-chiral: $(\bar{\chi}\gamma^\mu P_{L,R}\chi)(\bar{\psi}\gamma_\mu P_{L,R}\psi)$.

$$\varpi = \frac{8}{3}s(s - m^2), \quad \langle \sigma_{VC} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{8}{3}\mathcal{F}_{-1}(x). \quad (19)$$

The tensor interaction $\sigma^{\mu\nu}$ gives the same function as the vector case and is not reported. In the case of a Majorana χ clearly the vector and tensor interactions are absent, and the inclusion of a factor 1/2 in the operator (1) cancels the factor 4 due to the presence of the exchange diagram of the initial identical particles.

Now we consider some examples of t -channel annihilation for operators common to Dirac and Majorana DM annihilation:




Figure 2. The thermal function (11) for the interactions and annihilation cross sections considered in the text.

1) Scalar, pseudo-scalar: $(\bar{\chi}\chi)(\bar{\psi}\psi)$, $(\bar{\chi}\chi)(\bar{\psi}\gamma^5\psi)$, $(\bar{\chi}\gamma^5\chi)(\bar{\psi}\gamma^5\psi)$, $(\bar{\chi}\gamma^5\chi)(\bar{\psi}\psi)$.

$$\varpi_D = \frac{2}{3}s(s - m^2), \quad \langle \sigma_{S,PS}^{D,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{2}{3}\mathcal{F}_{-1}(x). \quad (20)$$

$$\varpi_M = \frac{1}{3}s(5s - 14m^2), \quad \langle \sigma_{S,PS}^{M,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{1}{3}\mathcal{F}_{-\frac{14}{5}}(x). \quad (21)$$

2) Chiral: $(\bar{\chi}P_{L,R}\chi)(\bar{\psi}P_{L,R}\psi)$.

$$\varpi_D = \frac{1}{6}s(s - m^2), \quad \langle \sigma_C^{D,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{1}{6}\mathcal{F}_{-1}(x). \quad (22)$$

$$\varpi_M = \frac{1}{3}s(s - 4m^2), \quad \langle \sigma_C^{M,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{1}{3}\mathcal{F}_{-4}(x). \quad (23)$$

3) Pseudo-vector: $(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{\psi}\gamma_\mu\gamma_5\psi)$, $(\bar{\chi}\gamma^\mu\gamma_5\chi)(\bar{\psi}\gamma_\mu\psi)$.

$$\varpi_D = \frac{4}{3}s(4s - 7m^2), \quad \langle \sigma_{PV}^{D,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{4}{3}\mathcal{F}_{-\frac{7}{4}}(x). \quad (24)$$

$$\varpi_M = \frac{8}{3}s(7s - 16m^2), \quad \langle \sigma_{PV}^{M,t} v_{\text{rel}} \rangle = \sigma_\Lambda \frac{8}{3}\mathcal{F}_{-\frac{16}{7}}(x). \quad (25)$$

The thermal functions corresponding to the previous cases are shown in Figure 2 where the asymptotic behaviours are clearly seen. In particular we note that

$$\mathcal{F}_0(x) = \frac{1}{16} \left(8 + 5 \frac{K_1^2(x)}{K_2^2(x)} + 3 \frac{K_3^2(x)}{K_2^2(x)} \right), \quad (26)$$

$$\mathcal{F}_{-4}(x) = \frac{3}{16} \left(-\frac{K_1^2(x)}{K_2^2(x)} + \frac{K_3^2(x)}{K_2^2(x)} \right), \quad (27)$$

behave in the nonrelativistic limit as

$$\mathcal{F}_0(x) \sim 1 + \mathcal{O}(x^{-2}), \quad \mathcal{F}_{-4}(x) \sim \frac{3}{2x} + \mathcal{O}(x^{-2}).$$

The function $\mathcal{F}_0(x)$, which appears in the s -channel annihilation through a pseudoscalar interaction, is the only

case where the term of order $\mathcal{O}(x^{-1})$ is absent, while $\mathcal{F}_{-4}(x)$, which appears in the scalar and axial-vector s -channel annihilation and in the chiral t -channel Majorana fermion annihilation, is the only case where the constant $\mathcal{O}(x^0)$ term is zero. These are the exact temperature dependent factors that correspond to the phenomenological interpolating functions proposed in Ref. [28] to model the s -wave and p -wave behaviour in the nonrelativistic limit. For all other interactions both s -wave and p -wave contribution are present. The function $\mathcal{F}_{-4}(x)$ can be also read off from the formulas of Ref. [29] where the t -channel annihilation of Majorana fermions with the exchange of a scalar with chiral couplings was considered.

We note that although we have concentrated on the case of fermion DM, the formula is valid for DM scalar and vector candidates as well, with the necessary redefinition of σ_Λ .

III. EXPANSION OF THE CROSS SECTION IN POWERS OF THE RELATIVE VELOCITY

In the general case $m_3 = m_4 = m_\psi \neq 0$ the exact integration is not possible. If the relative velocity of the annihilating particles is small compared with the velocity of light we can work directly with nonrelativistic formulas. The exothermic annihilation cross section in the nonrelativistic limit, to the lowest orders in v_r , is usually expanded as $\sigma_{nr} \sim a/v_r + bv_r$, and multiplying by v_r ,

$$\sigma_{nr}v_r \sim a + bv_r^2. \quad (28)$$

Then, using Eq. (B18) and (B19), the nonrelativistic thermal average of Eq. (28) is

$$\langle \sigma_{nr}v_r \rangle_{nr} \sim a + 6\frac{b}{x}. \quad (29)$$

In the case of our cross sections, comparing Eq. (29) with Eq. (12), the coefficients are thus ¹

$$a = \sigma_\Lambda a_2 \left(1 + \frac{k}{4}\right), \quad b = -\sigma_\Lambda a_2 \frac{k}{16}. \quad (30)$$

We now ask, given $\sigma(s)$, how to perform the expansion in terms of the relative velocity to find the coefficients a and b that correspond to the large x expansion

¹ This result must coincide with the expansion of Ref. [27, 30]. With our notation the expansion is

$$\langle \sigma_{nr}v_r \rangle_{nr} \sim \sigma_0|_{y=1} + \frac{3}{x} \left(-\sigma_0|_{y=1} + \frac{1}{2}\sigma'_0|_{y=1} \right),$$

where the prime indicate derivative respect to the variable y . Comparison with the expansion (29) requires to identify

$$a \equiv \sigma_0|_{y=1}, \quad b \equiv \frac{1}{2} \left(\frac{1}{2}\sigma'_0|_{y=1} - \sigma_0|_{y=1} \right).$$

Using Eq. (5) with $a_0 = 0$ and $a_2 = ka_1$, it is easy to verify that one obtains again Eq. (12).

of the relativistic thermal average in the COF. Combining equations (5), (6), (9), the general total annihilation cross section reads

$$\sigma = \sigma_\Lambda \frac{a_2}{2} \frac{\sqrt{s}}{\sqrt{s - 4m^2}} \left(\frac{s}{4m^2} + \frac{k}{4} \right). \quad (31)$$

The correct way to proceed is to use the invariant relation Eq. (B5) with $m_1 = m_2 = m$ and to solve it for s as a function of v_{rel} :

$$s = 2m^2 \left(1 + \frac{1}{\sqrt{1 - v_{\text{rel}}^2}} \right). \quad (32)$$

This formula is valid in every frame and substituted in Eq. (31) gives the exact dependence of the cross section on the relativistic relative velocity, $\sigma(v_{\text{rel}})$. Then, if $v_{\text{rel}} \sim v_r \ll 1$, we can expand the obtained expression to the desired order in v_r and the nonrelativistic average taken using Eq. (B19) will coincide with the expansion of Eq. (10) for $x \gg 1$, that is the expansion (12).

Equivalently, in order to find the expansion (28), we note that the squared roots in the annihilation cross section (31) imply that a term of order v_r^4 in s will contribute to the order v_r^2 in σ . Thus we need to expand s , formula (32), at least to order v_r^4 ,

$$s \sim 4m^2 + m^2v_r^2 + \frac{3}{4}m^2v_r^4. \quad (33)$$

Substituting Eq. (33) in Eq. (31) and performing the expansion in powers of v_r it easy to find

$$\sigma_{nr}v_r \sim \sigma_\Lambda a_2 \left(1 + \frac{k}{4} - \frac{k}{16}v_r^2 \right), \quad (34)$$

in agreement with (30).

In the case of coannihilations [31], for example when a DM particles scatter off another particle with different mass, the Mandelstam invariant takes the form

$$s = (m_1 - m_2)^2 + 2m_1m_2 \left(1 + \frac{1}{\sqrt{1 - v_{\text{rel}}^2}} \right), \quad (35)$$

with the expansion

$$s \sim (m_1 - m_2)^2 + m_1m_2v_r^2 + \frac{3}{4}m_1m_2v_r^4. \quad (36)$$

This procedure gives the correct expansion in the COF where the velocities $\mathbf{v}_{1,2}$ of the colliding particles are specified in this frame. Clearly, the same expansion with the same coefficients is obtained in the LF and in the CMF.

IV. THE PROBLEMS WITH THE MØLLER VELOCITY

The simple outlined procedure has not been recognized in DM literature where, incorrectly, the Møller velocity

\bar{v} , Eq. (B22), instead of v_{rel} is considered. As reminded in B, \bar{v} is a non-invariant, non-physical velocity. The expression of \bar{v} in terms of s is thus different in different frames and the expansion of σ takes different values in different frames.

Before discussing the problems with the Møller velocity we note that if we take the limit $m_f \rightarrow 0$ in the analogous expansions published many papers [4–11], we do not reproduce the expansion (34). The reason is that in these papers the expansion of s is truncated to the lowest order in v_r^2 ,

$$s \sim 4m^2 + m^2 v_r^2. \quad (37)$$

If we substitute this in Eq. (31) and expand, we find

$$\sigma_{nr} v_r \sim \sigma_{\Lambda} a_2 \left(1 + \frac{k}{4} + \frac{12+k}{32} v_r^2 \right), \quad (38)$$

with an incorrect coefficient b . Clearly the same wrong result is obtained truncating (33) to order v_r^2 , whatever the frame in which v_r is specified, CMF, LF or COF.

We now go back to the Møller velocity (B22). Evaluated in the CMF taking $m_1 = m_2 = m$ reads

$$\bar{v}_* = \frac{2}{\sqrt{s_*}} \sqrt{s_* - 4m^2}. \quad (39)$$

We indicate the quantities evaluated in the CMF with a “*”. By inverting Eq. (39) we find

$$s_* = \frac{4m^2}{1 - \frac{\bar{v}_*^2}{4}}. \quad (40)$$

This relation is different from (32) and is often incorrectly identified as the relation between s and the relative velocity in the CMF, see for example [31], [10]. In facts, the expansion to order $\mathcal{O}(v_{r,*}^4)$ reads

$$s_* \sim 4m^2 + m^2 v_{r,*}^2 + \frac{m^2}{4} v_{r,*}^4. \quad (41)$$

When used in (31), it gives the following nonrelativistic expansion of the cross section

$$\sigma_{nr} v_{r,*} \sim \sigma_{\Lambda} a_2 \left(1 + \frac{k}{4} + \frac{1}{4} v_{r,*}^2 \right), \quad (42)$$

which is different from the correct expansion (34).

Other authors, [26] and [20–23], perform the expansion with the Møller velocity evaluated in the rest frame of one particle. Indicating with “ ℓ ” the quantities in this frame, Eq. (B22) becomes

$$\bar{v}_{\ell} = \frac{\sqrt{s_{\ell}} \sqrt{s_{\ell} - 4m^2}}{s_{\ell} - 2m^2}, \quad (43)$$

and by inverting Eq. (43) we obtain

$$s_{\ell} = 2m^2 \left(1 + \frac{1}{\sqrt{1 - \bar{v}_{\ell}^2}} \right). \quad (44)$$

This expression is formally identical to Eq. (32), thus when $\bar{v}_{\ell} \sim v_{r,\ell}$ and s_{ℓ} is expanded up to the order $v_{r,\ell}^4$ we obtain the expansion $\sigma_{nr} v_{r,\ell}$ which formally coincides with Eq. (34), with $v_{r,\ell}$ in place of v_r .

It should be clear that this is just a mathematical coincidence due to the fact that \bar{v} reduces to v_{rel} only when one of the two velocities $\mathbf{v}_{1,2}$ is zero as it is evident from the definitions Eq. (B2) and Eq. (B22). In other words, the expansion found in Refs. [20–23] are correct because the authors have implicitly used the relative velocity, Eq. (B5) and (33).

We thus emphasize some common statements found in DM literature and why they do not subsist:

1) *In the relativistic Boltzmann equation the v in σv is \bar{v} and $\langle \sigma v \rangle$ must be calculated in the LF frame.*

This is not true, as shown in details in Ref. [27] and in B. Using v_{rel} and recognizing the nonphysical nature of \bar{v} , one works always with invariant quantities and the consistency of the relativistic and nonrelativistic formulas and expansions is obtained in the comoving frame without any further specification of the kinematics. The LF, also called Møller frame in Ref. [23], cannot be a privileged frame for the relic abundances calculation also because for massless particles the rest frame does not exist.

2) *The Møller velocity coincides with relative velocity in a frame where the velocities are collinear.*

This is not true because, for example, in the CMF where the particles have velocities v_* , the Møller velocity is $2v_*$ while the relative velocity is $2v_*/(1 + v_*^2)$. Note that the true relative velocity is never superluminal.

V. IMPACT ON THE RELIC ABUNDANCE

Only in the case $k = -4$ the incorrect expansions (38) and (42) coincide, incidentally, with the expansion (34). While the lowest order coefficient a turns out to be always the same, the coefficient b is different in any other case. To illustrate the impact of b on the value of the relic abundance we consider the case of the s -channel annihilation with vector interaction, Eq. (18), and the s -channel annihilation with a pseudoscalar exchange, Eq. (15). In the first case $k = 2$, $a_2 = 8/3$, and the correct coefficients a and b are

$$a_V = 4\sigma_{\Lambda}, \quad b_V = -\frac{\sigma_{\Lambda}}{3}, \quad (45)$$

while the incorrect coefficient b in (38) and (42) is

$$b_{V_1} = \frac{7}{6}\sigma_{\Lambda}, \quad b_{V_2} = \frac{2}{3}\sigma_{\Lambda}. \quad (46)$$

In the second case, $k = 0$ and $a_2 = 2$, thus

$$a_{PS} = 2\sigma_{\Lambda}, \quad b_{PS} = 0, \quad (47)$$

and the wrong b coefficients are

$$b_{PS_1} = \frac{3}{4}\sigma_{\Lambda}, \quad b_{PS_2} = \frac{1}{2}\sigma_{\Lambda}. \quad (48)$$

We calculate the relic abundance following the exact theory of freeze out presented in Ref. [32]. We briefly recall the main points. Let $Y_0 = 45/(4\pi^4)(g_\chi/g_s)x^2 K_2(x)$ be the initial equilibrium abundance (number density over the entropy density), with $g_\chi = 2$ for spin 1/2 fermions and g_s the relativistic degrees of freedom associated with the entropy density. The function $Y_1(x)$ that gives the abundance up to the point x_* where $Y_1(x) - Y_0(x)$ is maximal is

$$Y_1(x) = (1 + \delta(x))Y_0(x), \quad (49)$$

$$\delta(x) = \sqrt{1 - \frac{x^2}{C\langle\sigma v_{\text{rel}}\rangle Y_0} \frac{1}{Y_0} \frac{dY_0}{dx}} - 1, \quad (50)$$

with x_* given by the condition

$$-\frac{1}{Y_0(x)} \frac{dY_0(x)}{dx} = \frac{1}{\delta(x)} \frac{d\delta(x)}{dx} \text{ at } x = x_*. \quad (51)$$

The abundance at $x > x_*$ is found by integrating numerically the usual equation

$$\frac{dY}{dx} = \frac{C}{x^2} \langle\sigma v_{\text{rel}}\rangle (Y_0^2 - Y^2), \quad (52)$$

with the initial condition $(x_*, Y(x_*) = Y_1(x_*))$. The factor C is defined by $C = \sqrt{\frac{\pi}{45}} M_P m_\chi \sqrt{g_*}$, where M_P is the Plank mass and $\sqrt{g_*} = g_s/\sqrt{g_\rho}(1 + T/3 d(\ln g_s)/dT)$ accounts for the temperature dependence of the relativistic degrees of freedom associated with the energy density, g_ρ , and g_s [26, 30]. For WIMP masses larger than 10 GeV we can neglect the temperature dependence of the degrees of freedom [33, 34] and take $g_s = g_\rho = g = 100$, $\sqrt{g_*} = \sqrt{g}$. In solving numerically (52) and (51) with the exposed method, we use the exact formula for $\langle\sigma v_{\text{rel}}\rangle$, Eq. (10).

We compare the previous numerical solution with the one obtained using the nonrelativistic freeze out approximation (FOA) that is commonly employed in literature. The FOA consists in integrating equation (52) with an initial condition $(x_f, Y(x_f))$ such that the equilibrium term proportional Y_0^2 can be neglected. We choose the freeze out point at the point x_2 where $Y(x_2) \simeq Y_1(x_2) = 2Y_0(x_2)$. As shown in Ref. [32], $Y_1(x)$ well approximates the true abundance also in the interval $x_* < x < x_2$. x_2 is the optimal point for the FOA and corresponds to the temperature where the extent of the inverse creation reaction $\psi\bar{\psi} \rightarrow \chi\chi$ is maximal. The solution in the freeze out approximation is then

$$Y_{\text{FOA}} = \frac{2Y_0(x_2)}{1 + 2Y_0(x_2) \frac{C}{x_2} (a + 3\frac{b}{x_2})}. \quad (53)$$

The freeze out point x_2 is given by the condition $-\frac{1}{Y_0} \frac{dY_0}{dx} = 3\frac{C}{x^2} \langle\sigma v_{\text{rel}}\rangle Y_0$, which, in terms of the method of Ref. [35] corresponds to $c(c+2) = 3$, that is $c = 1$. Using the nonrelativistic form of Y_0 ,

$$Y_0 = \frac{45}{4\pi^4} \frac{g_\chi}{g_s} \sqrt{\frac{\pi}{2}} x^{3/2} e^{-x}, \quad (54)$$




Figure 3. Ratio of the relic abundance obtained by solving numerically equation (52) over the value given by the freeze out approximation, for the pseudoscalar and vector interactions. In the bottom blue curves for the FOA the correct coefficients (45) and (47) are used. The red and the black curves show the effect of the wrong coefficients (46) and (48), respectively.

x_2 is given by the root of

$$3C \left(a + 6\frac{b}{x} \right) \sqrt{\frac{\pi}{2}} x^{-1/2} e^{-x} = 1. \quad (55)$$

Calling $\alpha = 3aC\sqrt{\pi/2}$, an accurate analytical approximate solution of Eq. (55) is given by

$$x_2 = \ln \alpha - \frac{1}{2} \ln(\ln \alpha) + \ln(1 + \frac{6b}{a} (\ln \alpha)^{-1}). \quad (56)$$

The relic abundance normalized over the critical density is $\Omega h^2 = 2.755 \times 10^8 (m/\text{GeV}) Y_\infty$ for a Majorana fermion and two times that quantity for a Dirac fermion with the same density of antiparticles. We now compare the exact relic abundance Ωh^2 with the value $(\Omega h^2)_{\text{FOA}}$ furnished by the nonrelativistic FOA calculated using the correct and the wrong expansions. We take the couplings $\lambda_{a,b} = 1$ for illustrative purposes and two values of the cut off scale, $\Lambda = 1, 10$ TeV. The value of the freeze out points x_* and x_2 varies roughly between 18 and 30 in the parameter space with $m < \Lambda$ where the effective treatment is supposed to be valid. The ratio $\Omega h^2 / (\Omega h^2)_{\text{FOA}}$ is shown in Figure 3 as a function of the DM mass for the chosen examples. The bottom blue curves show that the FOA with the correct coefficients (45) and (47) underestimates the numerical value by less than 2%, and that in most part of the parameter space the error is at the level of 1% or less. This is a test of goodness for our FOA, and confirms what shown in Ref. [32]. The red and the black curves show the effect of the wrong coefficients (46) and (48), respectively. The wrong expansions underestimate the relic abundance by a factor between 3% and 12% for both interactions for masses larger than 10 GeV as shown in the plot. The behaviour is similar

for the other interactions not shown in figure. The error becomes even larger at smaller masses and we have verified that using for example $c = 1/2$ and other values we get even worst approximations. Clearly this kind of error nowadays is not compatible with the precision with which the experimental value is known.

ACKNOWLEDGMENTS

This work was supported by a grant under the MINECO/FEDER project: SOM: Sabor y origen de la Materia (CPI-14-397). The author acknowledges Roberto Ruiz de Austri, Nuria Rius and Pilar Hernandez for hospitality and for useful discussions at the Instituto de Fisica Corpuscolar (IFIC) in Valencia where part of this work was done.

Appendix A: Integrals and expansions

Equation (7) can be written as

$$\langle \sigma v_{\text{rel}} \rangle = \sigma_{\Lambda} \frac{2x}{K_2^2(x)} \left(a_2 A_2 + \frac{a_1}{4} A_1 + \frac{a_0}{16} A_0 \right). \quad (\text{A1})$$

The integrals are evaluated with methods similar to those described in Ref. [27] in terms of Bessel functions of the second kind:

$$\begin{aligned} A_0 &= \int_1^\infty dy \sqrt{y-1} K_1(2x\sqrt{y}) = \frac{1}{2x} K_1^2(x), \\ A_1 &= \int_1^\infty dy \sqrt{y-1} y K_1(2x\sqrt{y}) = \frac{1}{2x} \frac{K_1^2(x) + K_2^2(x)}{2}, \\ A_2 &= \int_1^\infty dy \sqrt{y-1} y^2 K_1(2x\sqrt{y}) \\ &= \frac{1}{2x} \frac{1}{16} [5K_1^2(x) + 8K_2^2(x) + 3K_3^2(x)]. \end{aligned}$$

The expansions at $x \gg 1$ are

$$\frac{K_1^2(x)}{K_2^2(x)} \sim 1 - \frac{3}{x} + \mathcal{O}(x^{-2}), \quad \frac{K_3^2(x)}{K_2^2(x)} \sim 1 + \frac{5}{x} + \mathcal{O}(x^{-2}), \quad (\text{A2})$$

while for $x \ll 1$ are

$$\frac{K_1^2(x)}{K_2^2(x)} \sim \frac{x^2}{4} + \mathcal{O}(x^3), \quad \frac{K_3^2(x)}{K_2^2(x)} \sim \frac{16}{x^2} + \mathcal{O}(x^2). \quad (\text{A3})$$

Appendix B: Invariant formulation using v_{rel}

In this Appendix we remind, based on the results of Ref. [27], the main points about the relation between the relative velocity, the Møller velocity, flux and thermal average which are used in the main text.

1. Invariant relative velocity

The relativistic relative velocity that generalizes the nonrelativistic relative velocity

$$v_r = |\mathbf{v}_1 - \mathbf{v}_2|, \quad (\text{B1})$$

is given by

$$v_{\text{rel}} = \frac{\sqrt{(\mathbf{v}_1 - \mathbf{v}_2)^2 - \frac{(\mathbf{v}_1 \times \mathbf{v}_2)^2}{c^2}}}{1 - \frac{\mathbf{v}_1 \cdot \mathbf{v}_2}{c^2}}. \quad (\text{B2})$$

We have explicitly written the dependence on the velocity of light c to make manifest that v_{rel} coincide with v_r in the nonrelativistic limit because the scalar and vector products are of order $(v/c)^2$. In the following we go back to natural units.

The relative velocity v_{rel} can be written using the Mandelstam invariant $s = (p_1 + p_2)^2$, where $p_{1,2}$ are the four-momenta, and λ , the Mandelstam triangular function,

$$\lambda(s, m_1^2, m_2^2) = [s - (m_1 + m_2)^2][s - (m_1 - m_2)^2], \quad (\text{B3})$$

in a generic frame,

$$v_{\text{rel}} = \frac{\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}}{p_1 \cdot p_2} \quad (\text{B4})$$

$$= \frac{\sqrt{\lambda(s, m_1^2, m_2^2)}}{s - (m_1^2 + m_2^2)}, \quad (\text{B5})$$

showing its invariant nature.

2. Flux factor

Given two bunches of particles with number densities $n_{1,2}$ and velocities $\mathbf{v}_{1,2}$ in a generic inertial frame, in non-relativistic physics the flux is $F_{nr} = n_1 n_2 v_r$. To obtain the relativistic invariant flux that reduces to F_{nr} in the nonrelativistic limit, the easiest way is to consider the 4-currents $J_i = (n_i, n_i \mathbf{v}_i)$, thus

$$F = (J_1 \cdot J_2) v_{\text{rel}} = n_1 n_2 (1 - \mathbf{v}_1 \cdot \mathbf{v}_2) v_{\text{rel}}. \quad (\text{B6})$$

Note that the factor $(1 - \mathbf{v}_1 \cdot \mathbf{v}_2)$ that guarantees the Lorentz invariance of the product of the number densities can also be written as

$$1 - \mathbf{v}_1 \cdot \mathbf{v}_2 = \frac{\gamma_r}{\gamma_1 \gamma_2} = \frac{p_1 \cdot p_2}{E_1 E_2}, \quad (\text{B7})$$

where $\gamma_r = 1/\sqrt{1 - v_{\text{rel}}^2}$ is the Lorentz factor associated with v_{rel} and γ_i the Lorentz factors associated with \mathbf{v}_i .

If the element of Lorentz invariant phase space is defined as usual

$$d\tilde{p}_i = \frac{d^3 \mathbf{p}_i}{(2\pi)^3 2E_i}, \quad (\text{B8})$$

and one particle states for bosons and fermions are normalized to $2E_i$ such that the density per unit volume is $2E_i$, then, using (B7), the flux (B6) simplifies to

$$F = 4(p_1 \cdot p_2)v_{\text{rel}}. \quad (\text{B9})$$

Substituting the expression of v_{rel} in the momentum representation, formula (B4), in Eq. (B9), the scalar product $p_1 \cdot p_2$ cancels out and the standard explicit form is recovered

$$F = 4\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}. \quad (\text{B10})$$

3. Cross section and collision integral

The integrated collision term of the Boltzmann equation, neglecting quantum effects, can be written as,

$$\int \prod_{i=1}^4 d\tilde{p}_i [f_3 f_4 W(3, 4|1, 2) - f_1 f_2 W(1, 2|3, 4)],$$

where $W(ij|kl) = (2\pi)^4 \delta^4(P_{ij} - P_{kl}) \sum_{s_i, s_f} |\mathcal{M}_{ij \rightarrow kl}|^2$, and f_i is the phase space distribution.

Using the unitary condition $\int d\tilde{p}_3 d\tilde{p}_4 W(3, 4|1, 2) = \int d\tilde{p}_3 d\tilde{p}_4 W(1, 2|3, 4)$ to write the collision integral only in terms of the annihilation rate

$$\frac{1}{1 + \delta_{12}} \int \prod_{i=1}^4 d\tilde{p}_i (f_3 f_4 - f_1 f_2) W(1, 2|3, 4), \quad (\text{B11})$$

we keep out a statistical factor accounting for the possibility of identical particles.

By definition, the invariant cross section, using the flux in the form (B9), is

$$\sigma = \frac{1}{4(p_1 \cdot p_2)v_{\text{rel}}} \int d\tilde{p}_3 d\tilde{p}_4 \frac{W(1, 2|3, 4)}{g_1 g_2}, \quad (\text{B12})$$

being $g_i = (2s_i + 1)$ the spin degrees of freedom.

Assuming as usual that the annihilation products are described by the equilibrium phase space distribution at zero chemical potential $f_{0,i}$, we have $f_3 f_4 = f_{0,3} f_{0,4} = f_{0,1} f_{0,2}$, the last equality following from energy conservation. Hence

$$\frac{g_1 g_2}{1 + \delta_{12}} \int \prod_{i=1}^2 \frac{d^3 \mathbf{p}_i}{(2\pi)^3 E_i} (p_1 \cdot p_2) (f_{0,1} f_{0,2} - f_1 f_2) \sigma v_{\text{rel}}.$$

The equilibrium phase-space distribution $f_{0,i}$ is related to the number density n_0 and to the momentum distribution $f_{0,p}(\mathbf{p})$ by $g_i/(2\pi)^3 f_{0,i} = n_{0,i} f_{0,p}(\mathbf{p})$. Assuming further that the non-equilibrium phase-space function at finite chemical potential f_i remains proportional to the equilibrium momentum distribution by a factor given by the non-equilibrium number density n_i , $g_i/(2\pi)^3 f_i = n_i f_{0,p}(\mathbf{p})$, we obtain

$$\frac{1}{1 + \delta_{12}} (n_{0,1} n_{0,2} - n_1 n_2) \langle \sigma v_{\text{rel}} \rangle. \quad (\text{B13})$$

When the species 1 and 2 are the same, it takes the usual form $\langle \sigma v_{\text{rel}} \rangle (n_0^2 - n^2)$ with the factor $1/2$ cancelled by stoichiometric coefficient appearing in the left-hand side of the complete kinetic equation, see for example Ref. [32].

4. Averaged thermal rate

In Eq. (B13) the general definition of relativistic thermal averaged rate is

$$\langle \sigma v_{\text{rel}} \rangle = \int \prod_{i=1}^2 \frac{d^3 \mathbf{p}_i}{E_i} (p_1 \cdot p_2) f_{0,p}(\mathbf{p}_1) f_{0,p}(\mathbf{p}_2) \sigma v_{\text{rel}}. \quad (\text{B14})$$

In the case of the relativistic Maxwell-Boltzmann-Jüttner statistics, the momentum distribution is

$$f_{0,p}(\mathbf{p}) = \frac{1}{4\pi m^2 T K_2(x)} e^{-\sqrt{\mathbf{p}^2 + m^2}/T}, \quad (\text{B15})$$

and as shown in Ref. [27], the six-dimensional integral on the right-hand side of Eq. (B14) reduces to

$$\langle \sigma v_{\text{rel}} \rangle = \int_0^1 dv_{\text{rel}} \mathcal{P}(v_{\text{rel}}) \sigma v_{\text{rel}}, \quad (\text{B16})$$

where the probability distribution of v_{rel} , for example for $m_1 = m_2 = m$, is

$$\mathcal{P}(v_{\text{rel}}) = \frac{x}{\sqrt{2} K_2^2(x)} \frac{\gamma_r^3 (\gamma_r^2 - 1)}{\sqrt{\gamma_r + 1}} K_1(\sqrt{2x} \sqrt{\gamma_r + 1}). \quad (\text{B17})$$

This is completely analogous to the nonrelativistic case where the probability distribution of v_r , for $m_1 = m_2 = m$, is

$$P(v_r) = \sqrt{\frac{2}{\pi}} x^{3/2} v_r^2 e^{-x \frac{v_r^2}{4}}, \quad (\text{B18})$$

and the thermal average reads

$$\langle \sigma_{nr} v_r \rangle_{nr} = \int_0^\infty dv_r P(v_r) \sigma_{nr} v_r. \quad (\text{B19})$$

Given the total annihilation cross section σ the product σv_{rel} will reduce to the nonrelativistic limit $\sigma_{nr} v_r$ and $\langle \sigma v_{\text{rel}} \rangle$ to $\langle \sigma_{nr} v_r \rangle_{nr}$ in the COF when $v_{\text{rel}} \sim v_r \ll 1$. Expressing Eq. (B16) in terms of s using (B5), we obtain the usual integral [27, 36] useful for practical calculation

$$\langle \sigma v_{\text{rel}} \rangle = \frac{1}{8T \prod_i m_i^2 K_2(x_i)} \times \int_{M^2}^\infty ds \frac{\lambda(s, m_1^2, m_2^2)}{\sqrt{s}} K_1\left(\frac{\sqrt{s}}{T}\right) \sigma, \quad (\text{B20})$$

with $x_i = m_i/T$ and $M = (m_1 + m_2)$.

We have recently become aware of the paper [37] where, probably for the first time, the thermal average of

relativistic rates was discussed and it was realized that with the relativistic Maxwell-Boltzmann statistics formula (B14) reduces to a single integral over the distribution over the relative momentum. With some algebra and change of variables it is easy to verify that for example Eqs. (11b) and (12a) of [37] coincide with Eqs. (29) and (37) of Ref. [27]. In Ref. [37] the cases of collisions of two massive particles, two massless particles and a massive with a massless particles are treated separately as if different definitions of flux and cross sections were necessary in each case. Clearly this distinction is unnecessary for the formulation we have given is completely general and valid in any case. We finally note that an integral formula similar to (B20) was also given in Ref. [29].

5. No need for the Møller velocity.

By noting that in Eq. (B6) the factor $(1 - \mathbf{v}_1 \cdot \mathbf{v}_2)$ can cancel the same factor in the denominator of v_{rel} , the invariant flux can also be written in the form

$$F = n_1 n_2 \sqrt{(\mathbf{v}_1 - \mathbf{v}_2)^2 - (\mathbf{v}_1 \times \mathbf{v}_2)^2}. \quad (\text{B21})$$

In the textbook by Landau and Lifschits [38] this form is attributed to Pauli without giving any reference, while its origin is more generally attributed to Møller [39].

It is interesting to look at original paper by Møller [39]. With our notation, he wants to prove that the flux given (B21) is invariant. In order to do that he shows that this can be written as a product of two invariant quantities: the ratio $\frac{n_1 n_2}{E_1 E_2}$ and the quantity $B = \sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}$ and there he stops.

The flux factor written in the form (B21) has the same structure of the nonrelativistic expression $n_1 n_2 v_r$. Probably for this reason it has been later introduced in the literature the notion of Møller velocity

$$\bar{v} = \sqrt{(\mathbf{v}_1 - \mathbf{v}_2)^2 - (\mathbf{v}_1 \times \mathbf{v}_2)^2} = \frac{\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}}{E_1 E_2}. \quad (\text{B22})$$

It is worth to stress that neither Møller nor Landau and Lifschits attribute any particular meaning to Eq. (B22) and do not define it as a particular velocity, even less as relative velocity. Clearly \bar{v} is nothing but the numerator of the formula defining v_{rel} because $\bar{v} = (1 - \mathbf{v}_1 \cdot \mathbf{v}_2) v_{\text{rel}}$, where the factor $(1 - \mathbf{v}_1 \cdot \mathbf{v}_2)$ comes from the definition of the invariant flux (B6). Already this fact indicates that \bar{v} is not a fundamental physical quantity and overall, it is not the relative velocity, nor when the velocities are collinear.

On the contrary, in DM literature and in textbooks, when defining the flux factor for the relativistic invariant cross section, it is incorrectly asserted that in a frame where the velocities are collinear the quantity $|\mathbf{v}_1 - \mathbf{v}_2|$ is the relative velocity, while in a generic frame is given by (B22). The form (B21) of the flux is a simple consequence of the fundamental quantities (B2) and (B6), there is no new physics or concept in it. For these reasons, and for its noninvariant and nonphysical nature, \bar{v} should not be used.

-
- [1] A. Kurylov and M. Kamionkowski, “Generalized analysis of weakly interacting massive particle searches,” *Phys. Rev. D* **69**, 063503 (2004) [hep-ph/0307185].
 - [2] J. Fan, M. Reece and L. T. Wang, “Non-relativistic effective theory of dark matter direct detection,” *JCAP* **1011**, 042 (2010) [1008.1591].
 - [3] A. L. Fitzpatrick, W. Haxton, E. Katz, N. Lubbers and Y. Xu, “The Effective Field Theory of Dark Matter Direct Detection,” *JCAP* **1302**, 004 (2013). [1203.3542].
 - [4] M. Beltran, D. Hooper, E. W. Kolb and Z. C. Krusberg, “Deducing the nature of dark matter from direct and indirect detection experiments in the absence of collider signatures of new physics,” *Phys. Rev. D* **80**, 043509 (2009) [0808.3384].
 - [5] H. Zhang, Q. H. Cao, C. R. Chen and C. S. Li, “Effective Dark Matter Model: Relic density, CDMS II, Fermi LAT and LHC,” *JHEP* **1108**, 018 (2011) [0912.4511].
 - [6] M. R. Buckley, “Asymmetric Dark Matter and Effective Operators,” *Phys. Rev. D* **84**, 043510 (2011) [1104.1429].
 - [7] S. Matsumoto, S. Mukhopadhyay and Y. L. S. Tsai, “Singlet Majorana fermion dark matter: a comprehensive analysis in effective field theory,” *JHEP* **1410**, 155 (2014) [1407.1859].
 - [8] M. A. Fedderke, J. Y. Chen, E. W. Kolb and L. T. Wang, “The Fermionic Dark Matter Higgs Portal: an effective field theory approach,” *JHEP* **1408**, 122 (2014) [1404.2283].
 - [9] J. Y. Chen, E. W. Kolb and L. T. Wang, “Dark matter coupling to electroweak gauge and Higgs bosons: an effective field theory approach,” *Phys. Dark Univ.* **2**, 200 (2013) [1305.0021].
 - [10] A. Berlin, D. Hooper and S. D. McDermott, “Simplified Dark Matter Models for the Galactic Center Gamma-Ray Excess,” *Phys. Rev. D* **89**, 115022 (2014) [1404.0022].
 - [11] S. Chang, R. Edezhath, J. Hutchinson and M. Luty, “Effective WIMPs,” *Phys. Rev. D* **89**, 015011 (2014) [1307.8120].
 - [12] C. Balzs, T. Li and J. L. Newstead, “Thermal dark matter implies new physics not far above the weak scale,” *JHEP* **1408**, 061 (2014) [1403.5829].
 - [13] K. Cheung, P. Y. Tseng, Y. L. S. Tsai and T. C. Yuan, “Global Constraints on Effective Dark Matter Interactions: Relic Density, Direct Detection, Indirect Detection, and Collider,” *JCAP* **1205**, 001 (2012) [1201.3402].
 - [14] J. Goodman, M. Ibe, A. Rajaraman, W. Shep-

- herd, T. M. P. Tait and H. B. Yu, “Constraints on Light Majorana dark Matter from Colliders,” *Phys. Lett. B* **695** (2011) 185 [1005.1286].
- [15] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait and H. B. Yu, “Constraints on Dark Matter from Colliders,” *Phys. Rev. D* **82**, 116010 (2010) [1008.1783].
- [16] O. Buchmueller, M. J. Dolan and C. McCabe, “Beyond Effective Field Theory for Dark Matter Searches at the LHC,” *JHEP* **1401**, 025 (2014) [1308.6799].
- [17] G. Busoni, A. De Simone, E. Morgante and A. Riotto, “On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC,” *Phys. Lett. B* **728**, 412 (2014) [1307.2253].
- [18] G. Busoni, A. De Simone, J. Gramling, E. Morgante and A. Riotto, “On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC, Part II: Complete Analysis for the s -channel,” *JCAP* **1406**, 060 (2014) [1402.1275].
- [19] G. Busoni, A. De Simone, T. Jacques, E. Morgante and A. Riotto, “On the Validity of the Effective Field Theory for Dark Matter Searches at the LHC Part III: Analysis for the t -channel,” *JCAP* **1409**, 022 (2014) [1405.3101].
- [20] J. M. Zheng, Z. H. Yu, J. W. Shao, X. J. Bi, Z. Li and H. H. Zhang, “Constraining the interaction strength between dark matter and visible matter: I. fermionic dark matter,” *Nucl. Phys. B* **854**, 350 (2012) [1012.2022].
- [21] H. Dreiner, M. Huck, M. Krmer, D. Schmeier and J. Tattersall, “Illuminating Dark Matter at the ILC,” *Phys. Rev. D* **87**, 075015 (2013) [1211.2254].
- [22] J. Blumenthal, P. Gretsikov, M. Krmer and C. Wiebusch, “Effective field theory interpretation of searches for dark matter annihilation in the Sun with the IceCube Neutrino Observatory,” *Phys. Rev. D* **91**, 035002 (2015) [1411.5917].
- [23] G. Busoni, A. De Simone, T. Jacques, E. Morgante and A. Riotto, “Making the Most of the Relic Density for Dark Matter Searches at the LHC 14 TeV Run,” *JCAP* **1503**, 022 (2015) [1410.7409].
- [24] C. L. Bennett *et al.* [WMAP Collaboration], “Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results,” *Astrophys. J. Suppl.* **208**, 20 (2013) [1212.5225].
- [25] P. A. R. Ade *et al.* [Planck Collaboration], “Planck 2013 results. XVI. Cosmological parameters,” *Astron. Astrophys.* **571**, A16 (2014) [1303.5076].
- [26] P. Gondolo and G. Gelmini, “Cosmic abundances of stable particles: Improved analysis,” *Nucl. Phys. B* **360**, 145 (1991).
- [27] M. Cannoni, “Relativistic $\langle\sigma v_{\text{rel}}\rangle$ in the calculation of relics abundances: a closer look,” *Phys. Rev. D* **89**, 103533 (2014) [1311.4494] [1311.4508].
- [28] M. Drees, M. Kakizaki and S. Kulkarni, “The Thermal Abundance of Semi-Relativistic Relics,” *Phys. Rev. D* **80**, 043505 (2009) [0904.3046].
- [29] M. Claudson, L. J. Hall and I. Hinchliffe, “Cosmological Baryon Generation At Low Temperatures,” *Nucl. Phys. B* **241**, 309 (1984).
- [30] M. Srednicki, R. Watkins and K. A. Olive, “Calculations of Relic Densities in the Early Universe,” *Nucl. Phys. B* **310**, 693 (1988).
- [31] K. Griest and D. Seckel, “Three exceptions in the calculation of relic abundances,” *Phys. Rev. D* **43**, 3191 (1991).
- [32] M. Cannoni, “Exact theory of freeze out,” *Eur. Phys. J. C* **75**, 106 (2015) [1407.4108].
- [33] G. Steigman, B. Dasgupta and J. F. Beacom, “Precise Relic WIMP Abundance and its Impact on Searches for Dark Matter Annihilation,” *Phys. Rev. D* **86**, 023506 (2012) [1204.3622].
- [34] M. Drees, F. Hajkarim and E. R. Schmitz, “The Effects of QCD Equation of State on the Relic Density of WIMP Dark Matter,” *JCAP* **1506**, 025 (2015) [1503.03513].
- [35] R. J. Scherrer and M. S. Turner, “On the Relic, Cosmic Abundance of Stable Weakly Interacting Massive Particles,” *Phys. Rev. D* **33**, 1585 (1986).
- [36] J. Edsjo and P. Gondolo, “Neutralino relic density including coannihilations,” *Phys. Rev. D* **56**, 1879 (1997) [hep-ph/9704361].
- [37] T. A. Weaver, *Reaction rates in a relativistic plasma*, *Phys. Rev. A* **13**, 1563 (1976).
- [38] L. D. Landau and E. M. Lifschits, *The Classical Theory of Fields: Course of Theoretical Physics, Vol. 2*, (Pergamon Press, New York, 1975).
- [39] C. Möller, “General Properties of the Characteristic Matrix in the Theory of Elementary Particles,” *D. Kgl. Danske Vidensk. Selsk. Mat.-Fys. Medd.* **23**, No. 1 (1945). [PDF available here](#)