

Wigner time delay and related concepts Application to transport in coherent conductors [☆]

Christophe Texier^a

^a*LPTMS, CNRS, Univ. Paris Sud, Université Paris Saclay, Bât. 100, F-91405 Orsay, France*

Abstract

The concepts of Wigner time delay and Wigner-Smith matrix allow to characterize temporal aspects of a quantum scattering process. The article reviews the statistical properties of the Wigner time delay for disordered systems ; the case of disorder in 1D with a chiral symmetry is discussed and the relation with exponential functionals of the Brownian motion underlined. Another approach for the analysis of time delay statistics is the random matrix approach, from which we review few results. As a practical illustration, we briefly outline a theory of non-linear transport and AC transport developed by Büttiker and coworkers, where the concept of Wigner-Smith time delay matrix is a central piece allowing to describe screening properties in out-of-equilibrium coherent conductors.

Published online the 8 october 2015 in: *Physica E* **82**, 16–33 (2016).

▷ THIS ARXIV VERSION WAS UPDATED AND EXTENDED AFTER THE PUBLICATION IN PHYSICA. ◁

Keywords:

PACS: 73.23.-b, 73.20.Fz, 72.15.Rn

Contents

1 Introduction	2	3.3 Disorder in 1D with a chiral symmetry	9
2 Wigner time delay and other characteristic times	2	3.4 Related questions and remarks . . .	10
2.1 Scattering on the half line	2	3.5 Time delay, absorption/amplification and analyticity (Dirac case)	11
2.2 The scattering problem in one dimension : several characteristic times	3	3.6 Beyond 1D	13
2.3 Krein-Friedel relation and Virial expansion	5	4 Random matrix approach for quantum dots	14
2.4 Few remarks	6	4.1 Distribution and moments for perfect contacts	14
3 Disordered systems (localised regime)	7	4.2 Distributions and moments for non-ideal contacts	18
3.1 Exponential functionals of the Brownian motion	7	4.3 Energy correlations for perfect coupling	21
3.2 Disorder on the half line - Single parameter scaling and universality . . .	8	4.4 Other symmetry classes	22
		5 Generalised concepts : injectance, emittance, etc.	23
		5.1 Partial DoS, injectance and emittance	23
		5.2 A generalisation of the Feynman-Hellmann theorem for a continuous spectrum	24
		5.3 Statistical analysis in quantum dots	25

[☆]Contribution to a special issue “*Frontiers in quantum electronic transport – in memory of Markus Büttiker*”

Email address: christophe.texier@u-psud.fr
(Christophe Texier)

6	Non-linear transport in coherent conductors	26
6.1	Non-linear conductances	26
6.2	Characteristic potentials	26
6.3	Non-linear conductances	27
6.4	Recent developments	28
7	AC transport	28
7.1	The AC response	28
7.2	The charge relaxation resistance . .	29
7.3	Final remarks	30
8	Conclusion	31
Appendix A Calculation of $\langle R_q \rangle$ for $N \gg 1$ – Mapping to the study of absorption in the cavity		31
Appendix B Partial <i>versus</i> proper times		32
Appendix B.1	Symmetric case	33
Appendix B.2	Asymmetric case . . .	33
Bibliography		33

1. Introduction

The purpose of this article is to review several results linked to a concept which has been very influential in the work of Markus Büttiker, namely the concept of Wigner time delay, and several of its extensions – traversal time, Wigner-Smith matrix, injectance, etc. In my opinion, this topic allows one to have a good flavor of Markus Büttiker’s style as a physicist : a combination between formal developments with mathematical elegance and motivations from practical questions of fundamental condensed matter physics. Besides, the choice of this theme is also related to a more personal anecdote as it was the subject of my first scientific exchange with Markus Büttiker, which I rediscovered during the preparation of this article, finding in a box a copy of an old email, dated the 8th July 1997, addressed to me and my PhD advisor Alain Comtet. Here are few sentences extracted from this message : “*I noticed yesterday your paper on the cond-mat network [60]. I like to react to two things: (...) In your introduction you jump from the work of Fyodorov and Sommers immediately to the work of Brouwer et al. Both of these are of course fine works. But the work of Brouwer et al. as they make clear in very strong terms takes its starting point from my work*

by Gopar and Mello [96], and that alone should have been a good enough reason to not simply leave it out. (...) Therefore, it is my hope that you can give this work the place it deserves. Sincerely, Markus Büttiker”. Of course, he was absolutely right and we soon after amended our paper according to his remarks. Despite this somewhat awkward beginning, Markus Büttiker hired me as a postdoctoral assistant at the University of Geneva two years later and I enjoyed very much the pleasant and stimulating atmosphere of the Physics Department. I especially appreciated the freedom which Markus gave to his postdocs and learnt a lot from our regular exchanges.

The notion of time delay was introduced in the fifties by Eisenbud and Wigner in the context of the quantum theory of scattering. It allows to capture temporal aspects of the scattering process [72, 210, 203, 25, 132]. The most fundamental aspects of time delays have been reviewed by Carvalho and Nussenzveig [67]. Some other articles have reviewed more specific aspects : Beenakker has considered the case of wave guides in the localised regime [14]. In Ref. [117], Kottos has reviewed the random matrix approach for ergodic systems (chaotic cavities) and the case of (non ergodic) disordered systems –diffusive or strongly localised. We have emphasized in [56] the connection with the theory of exponential functional of the Brownian motion. The purpose of the present article is to review few results on time delays from different contexts and to emphasize the diversity of the physical situations where this concept has found some applications.

The outline is the following : in Section 2 we discuss several definitions of times characterizing the scattering process. Section 3 reviews the statistical analysis of time delays for disordered systems. In Section 4, we discuss few aspects of the random matrix approach. Section 5 introduces several generalized concepts (partial DoS, injectance, etc.) which will find practical applications for the analysis of non-linear transport (Section 6) and AC transport (Section 7).

2. Wigner time delay and other characteristic times

2.1. Scattering on the half line

The most simple situation allowing to introduce the concept of time delay is the scattering problem

on the half line (this also corresponds to project a rotational invariant problem in an orbital momentum channel, in higher dimensions). We consider the Schrödinger equation

$$-\psi''_\varepsilon(x) + V(x)\psi_\varepsilon(x) = \varepsilon\psi_\varepsilon(x) \quad (1)$$

for $x \in \mathbb{R}^+$ (with $\hbar^2/(2m) = 1$), describing the scattering of a particle by a potential defined on a finite interval $[0, L]$ (Fig. 1). At $x = 0$ we choose to impose a Dirichlet boundary condition, $\psi_\varepsilon(0) = 0$. In the “asymptotic” region ($x > L$), the stationary




Figure 1: A plane wave is sent on a potential living on the interval $[0, L]$.

scattering state of energy $\varepsilon = k^2$ is the superposition of an incident wave $e^{-ik(x-L)}$ and a reflected wave $r e^{+ik(x-L)}$:

$$\psi_\varepsilon(x) \underset{x>L}{=} \frac{1}{\sqrt{2\pi\hbar v}} \left(e^{-ik(x-L)} + r e^{ik(x-L)} \right). \quad (2)$$

The normalisation constant, involving the group velocity $v = (1/\hbar)d\varepsilon/dk = 2k$, corresponds to associate a measure $d\varepsilon$ to the eigenstate.¹ The reflection probability amplitude has unit modulus as a consequence of current conservation: $r = e^{i\delta_r}$. The Wigner time delay is defined as the derivative of the reflection phase

$$\tau_W(\varepsilon) = -i\hbar r^* \frac{\partial r}{\partial \varepsilon} = \hbar \frac{\partial \delta_r(\varepsilon)}{\partial \varepsilon}. \quad (4)$$

It measures the time spent by a wave packet of energy ε in the domain $[0, L]$. This can be easily understood by considering the time evolution of a wave packet

$$\Phi(x; t) = \int_0^\infty d\varepsilon \hat{\Phi}(\varepsilon) \psi_\varepsilon(x) e^{-i\varepsilon t}, \quad (5)$$

where $\hat{\Phi}(\varepsilon)$ is a narrow function centered around an energy ε_0 (from now on, we will set $\hbar = 1$ for simplicity). Normalisation of the wave

¹ This choice of normalisation ensures that orthonormalisation reads

$$\langle \psi_\varepsilon | \psi_{\varepsilon'} \rangle = \delta(\varepsilon - \varepsilon') \quad (3)$$

and closure relation $\int d\varepsilon |\psi_\varepsilon\rangle \langle \psi_\varepsilon| = 1$ [191].

function is $\int dx |\Phi(x; t)|^2 = \int_0^\infty d\varepsilon |\hat{\Phi}(\varepsilon)|^2 = 1$. We can split the wave packet into an incident part and a reflected part: $\Phi(x; t) = \Phi_{\text{inc}}(x; t) + \Phi_{\text{ref}}(x; t)$. In the free region $x > L$, neglecting the effect of dispersion, $\Phi_{\text{inc}}(x; t) \simeq (1/\sqrt{4\pi}) \int_0^\infty d\varepsilon \varepsilon^{-1/4} \hat{\Phi}(\varepsilon) e^{-i\sqrt{\varepsilon}(x-L) - i\varepsilon t}$ can be rewritten under the form

$$\Phi_{\text{inc}}(x; t) \propto f(-x + L - v_0 t), \quad (6)$$

where v_0 is the group velocity at energy ε_0 and $f(x) \approx \int dK \hat{\Phi}(\varepsilon_0 + v_0 K) e^{iKx}$. In the reflected wave packet, the expansion of the phase shift produces a shift in time and we obtain

$$\Phi_{\text{ref}}(x; t) \propto f(x - L - v_0 t + v_0 \tau_W(\varepsilon_0)) \quad (7)$$

(cf. also chapter 10 of the book [191]). The interpretation of τ_W as the delay in time relies on following the motion of a wave packet with sufficiently narrow dispersion in energy. As trivial illustrations we can consider the case of an impenetrable region when $\delta_r = \pi$, leading to $\tau_W = 0$, and a free region ($V(x) = 0$), leading to $\delta_r = \pi + 2kL$ and $\tau_W = 2L/v$.




Figure 2: The Wigner time delay $\tau_W(\varepsilon)$ measures the delay of the reflected wave packet with group velocity v_ε (bottom), with respect to the wave packet reflected on the boundary of the scattering region (top). (Dispersion is neglected here).

2.2. The scattering problem in one dimension : several characteristic times

The scattering problem on the infinite line illustrates that, although the situation is still simple, many other characteristic times can be already introduced. We consider a plane wave sent from $-\infty$ (or $+\infty$) on a potential defined on the interval $[-L/2, L/2]$ (Fig. 3). The scattering properties can be encoded in two stationary scattering states $\psi_{\varepsilon,L}(x)$ (Fig. 3) and $\psi_{\varepsilon,R}(x)$, controlled by two pairs of reflection/transmission amplitudes r, t

and r' , t' characterizing transmission from the left and from the right, respectively. A general scattering state is a linear combination

$$\Psi_\varepsilon(x) = A_L \psi_{\varepsilon,L}(x) + A_R \psi_{\varepsilon,R}(x), \quad (8)$$

where $A = (A_L, A_R)$ are the amplitudes of the two incoming plane waves. They are related to the amplitudes $B = (B_L, B_R)$ of the outgoing plane waves by the 2×2 scattering matrix $B = \mathcal{S} A$:

$$\mathcal{S} = \begin{pmatrix} r & t' \\ t & r' \end{pmatrix}. \quad (9)$$




Figure 3: The stationary scattering state $\psi_{\varepsilon,L}(x)$ describes the scattering of an incident plane wave from the left, which is encoded by a reflection amplitude r and a transmission amplitude t .

As a consequence of current conservation, the scattering matrix is a unitary matrix $\mathcal{S} \in U(2)$, parametrised by four independent real parameters, what is made clear by expressing \mathcal{S} in the polar representation:

$$\mathcal{S} = e^{i\Phi_f/2} \begin{pmatrix} \sqrt{1-\mathcal{T}} e^{i\alpha} & i\sqrt{\mathcal{T}} e^{-i\chi} \\ i\sqrt{\mathcal{T}} e^{i\chi} & \sqrt{1-\mathcal{T}} e^{-i\alpha} \end{pmatrix}. \quad (10)$$

$\mathcal{T} = |t|^2 = |t'|^2 \in [0, 1]$ denotes the transmission probability. The three independent phases are a global phase $\Phi_f = \det \mathcal{S}$, called the “Friedel phase”, the phase α controlling the left/right asymmetry and a magnetic phase χ (which can be removed by a gauge transformation in the 1D case). As the scattering process is characterised by a matrix, there is much more freedom to define characteristic times, as we now discuss.

2.2.1. Friedel phase and Wigner time delay

The most simple generalization of the definition (4) is

$$\tau_W(\varepsilon) = -\frac{i}{N} \text{Tr} \left\{ \mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} \right\} = -\frac{i}{N} \frac{\partial \ln \det \mathcal{S}}{\partial \varepsilon} \quad (11)$$

$$= \frac{1}{N} \frac{\partial \Phi_f}{\partial \varepsilon}, \quad (12)$$

where N is the number of scattering channels (here $N = 2$). Although this is an important quantity, as we will explain in the § 2.3, τ_W does not have a simple interpretation as a wave packet’s delay in time.

2.2.2. Transmission and reflection group delays

A plane wave sent on the scattering region from the left is splitted into two parts to which may be associated two phases: a reflection phase $\delta_r = \Phi_f/2 + \alpha$ and a transmission phase $\delta_t = \Phi_f/2 + \chi + \pi/2$ (we also introduce $\delta_{r'} = \Phi_f/2 - \alpha$). Extending the argument exposed in § 2.1 would lead to introduce three characteristic times: one *transmission group delay time* $\tilde{\tau}_t = \partial \delta_t / \partial \varepsilon$, and two *reflection group delay times* $\tilde{\tau}_r = \partial \delta_r / \partial \varepsilon$ and $\tilde{\tau}_{r'} = \partial \delta_{r'} / \partial \varepsilon$ (note that $\alpha = 0$ or π , and thus $\tilde{\tau}_r = \tilde{\tau}_{r'}$ for a symmetric potential).

2.2.3. Partial time delays: diagonalisation and derivation

The two scattering states $\psi_{\varepsilon,\alpha}(x)$, with $\alpha \in \{L, R\}$, describing a particle incoming from the left/right (Fig. 3), may be recombined in order to form other basis of solutions. An important case corresponds to the two partial scattering states, which behave “asymptotically” as²

$$\phi_{\varepsilon,a}(x) \underset{|x| > L/2}{=} [C_a \theta_H(-x) + D_a \theta_H(x)] \cos(k|x| + \eta_a) \quad (13)$$

with $a \in \{1, 2\}$. $\theta_H(x)$ is the Heaviside function. Decomposing $\phi_{\varepsilon,a}(x)$ over the left/right scattering states, as in Eq. (8), leads to [189, 191] $\det(\mathcal{S} - e^{2i\eta_a} \mathbf{1}_2) = 0$, i.e. the two partial waves are eigenvectors of the scattering matrix with eigenvalues $\{e^{2i\eta_a}\}$. We can write:

$$\mathcal{S} = \mathcal{U} \begin{pmatrix} e^{2i\eta_1} & 0 \\ 0 & e^{2i\eta_2} \end{pmatrix} \mathcal{U}^\dagger, \quad (14)$$

where \mathcal{U} is a unitary matrix and $\{\eta_a\}$ the scattering phase shifts. In particular, using the polar decomposition, one finds the expressions of the two phase shifts

$$e^{2i\eta_{1,2}} = e^{i\Phi_f/2} \left(\sqrt{1-\mathcal{T}} \cos \alpha \pm i\sqrt{1-(1-\mathcal{T})\cos^2 \alpha} \right), \quad (15)$$

leading to $2(\eta_1 + \eta_2) = \Phi_f$, as it should. Two other characteristic times could be introduced, which are the two *partial time delays*

$$\tilde{\tau}_a = 2 \frac{\partial \eta_a}{\partial \varepsilon}. \quad (16)$$

² For a symmetric potential $V(x) = V(-x)$, the two partial waves are the symmetric and antisymmetric solutions (i.e. $C_1 = D_1$ and $C_2 = -D_2$).

As the partial waves (13) may be combined in order to form wave packets, the partial time delays can be interpreted as delays in time for such wave packets.

2.2.4. Proper time delays : derivation and diagonalisation

The determination of the partial time delays involves a diagonalisation of the scattering matrix and a derivation of its eigenvalues. The converse of these two operations leads to introduce the *proper time delays* $\{\tau_a\}$, which are eigenvalues of the Wigner-Smith time delay matrix

$$\mathcal{Q} = -i\mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} = \mathcal{V} \begin{pmatrix} \tau_1 & 0 \\ 0 & \tau_2 \end{pmatrix} \mathcal{V}^\dagger, \quad (17)$$

where \mathcal{V} is a unitary matrix, which differs from \mathcal{U} involved in (14) in general (see Appendix B). A related matrix, which will play a role, is

$$\tilde{\mathcal{Q}} = -i \frac{\partial \mathcal{S}}{\partial \varepsilon} \mathcal{S}^\dagger = \mathcal{S} \mathcal{Q} \mathcal{S}^\dagger. \quad (18)$$

Hermiticity, $\mathcal{Q} = \mathcal{Q}^\dagger$ and $\tilde{\mathcal{Q}} = \tilde{\mathcal{Q}}^\dagger$, follows from unitarity of \mathcal{S} . Obviously, \mathcal{Q} and $\tilde{\mathcal{Q}}$ are characterised by the same spectrum of eigenvalues $\{\tau_a\}$, as they are related by a unitary transformation. Proper time delays and partial time delays have an important difference : being derivatives of the \mathcal{S} -matrix eigenvalues, the partial time delays are intrinsic properties of the scattering process, whereas the proper time delays depend on the particular choice of basis in which the \mathcal{S} -matrix is expressed (see Appendix B). Although they do not coincide in general, they satisfy the sum rule

$$-i \frac{\partial}{\partial \varepsilon} \ln \det \mathcal{S} = \sum_{a=1}^N \tilde{\tau}_a = \sum_{a=1}^N \tau_a = N \tau_W \quad (19)$$

where N is the number of scattering channels (here $N = 2$).

2.2.5. Other characteristic times

The time delay interpretation of the various characteristic times deduced from scattering phases may lead to paradoxal conclusions in the presence of tunneling barriers, like superluminal propagation. For this reason “clock approaches” have been proposed in order to provide more satisfactory answers to the question “how much time needs a wave packet to travel in a given region?”. We refer to the review

articles [109, 126, 38, 125, 44, 67]³ and Büttiker’s contributions [46, 37, 47].

2.3. Krein-Friedel relation and Virial expansion

An important aspect behind the notion of time delay is its link with the spectral properties of open systems with continuous spectra, which can be understood as follows. Let us come back to the simple situation of scattering on the semi-infinite line for simplicity (Fig. 1). The scattering states (2) satisfying the Dirichlet boundary condition $\psi_\varepsilon(0) = 0$ may be used as a basis in order to consider the spectral problem on the *finite* interval $[0, L]$, which involves a second boundary condition at $x = L$. We choose a Neumann boundary condition $\psi'_\varepsilon(L) = 0$ for instance. The eigenenergies $\{\varepsilon_n\}$ are solutions of the quantisation equation $\delta_r(\varepsilon_n) = 2n\pi$ for n integer. When two successive energies are sufficiently close, we can expand the relation $\delta_r(\varepsilon_{n+1}) - \delta_r(\varepsilon_n) = 2\pi$ as $\delta'_r(\varepsilon_n) \delta\varepsilon_n \simeq 2\pi$, where $\delta\varepsilon_n = \varepsilon_{n+1} - \varepsilon_n$ is the level spacing. We deduce the expression of the density of states $\nu(\varepsilon) \simeq 1/\delta\varepsilon_n \simeq \tau_W(\varepsilon)/(2\pi\hbar)$. A more precise connection between spectral and scattering properties was obtained by Friedel [81] and Smith [180], who established the relation

$$\int_0^L dx |\psi_\varepsilon(x)|^2 = \frac{1}{2\pi} \left(\tau_W(\varepsilon) + \frac{\sin \delta_r(\varepsilon)}{2\varepsilon} \right) \quad (20)$$

(see also Ref. [194]). The left hand side of the equation is simply related to the local density of states (DoS)⁴

$$\nu(x; \varepsilon) = \langle x | \delta(\varepsilon - H) | x \rangle = |\psi_\varepsilon(x)|^2. \quad (21)$$

Thus, Eq. (20) shows that the Wigner time delay provides a measure of the DoS of the interval $[0, L]$, i.e. of the integral of the local DoS over the support of the potential. It may be written in the more general form [193]

$$\nu(\varepsilon) = \int_0^L dx \nu(x; \varepsilon) = \frac{1}{2i\pi} \text{Tr} \left\{ \mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} + \frac{\mathcal{S} - \mathcal{S}^\dagger}{4\varepsilon} \right\}, \quad (22)$$

so that it also applies to more general situations such as the one of Fig. 3, for a potential vanishing outside $[0, L]$, or even complex structures like

³ The introduction of Ref. [18], which introduces a sojourn time from wave attenuation, provides a good overview.

⁴ The choice of the normalisation for the scattering state (2) is important.

metric graphs [189, 193, 195]. Such equalities are known as Krein-Friedel relations [80, 119, 81, 36] (or Birman-Krein formula [22]).⁵ They have a long history and first appeared in the virial expansion of the equation of state of real gases

$$p = nT [1 + B_2(T)n + B_3(T)n^2 + \dots], \quad (23)$$

where p is the pressure, n the density and T the temperature. As shown by Beth and Uhlenbeck [204, 20], the second virial coefficient can be written as

$$B_2(T) = -2^{-5/2} \Lambda_T^3 (\pm 1 + 16Z_{\text{int}}) \quad (24)$$

for bosons (+) and fermions (-), where $\Lambda_T = \sqrt{2\pi\hbar^2/(mT)}$ is the thermal length and m the mass of the particles. The first contribution to B_2 encodes the quantum correlations arising from the symmetrisation postulate, whereas the second term describes the correlations due to the interaction, Z_{int} being the partition function of the two body problem (in the relative coordinates)

$$Z_{\text{int}} = \sum_{\ell} (2\ell + 1) \int_0^{\infty} \frac{d\varepsilon}{2\pi\hbar} \tau_{\ell}(\varepsilon) e^{-\varepsilon/T}, \quad (25)$$

where summation runs over orbital momentum. $\tau_{\ell} = 2\partial\eta_{\ell}/\partial\varepsilon$ is the time delay related to the scattering phase shift⁶ η_{ℓ} characterizing the scattering in the channel of orbital momentum ℓ (cf. § 77 of [124]); in the presence of bound states, the partition function receives additional contributions. This approach was later made more systematic by Dashen, Ma and Bernstein [66]. Note that a similar analysis of real gases based on time delay can be developed within a purely classical frame, as beautifully explained by Ma [134].

⁵ The Krein-Friedel relation is sometimes written in terms of the spectral shift function

$$\Delta(\varepsilon) = \int_{-\infty}^{+\infty} dx [\nu(x; \varepsilon) - \nu_0(x; \varepsilon)],$$

which measures the variation of the DoS due to the introduction of the scattering potential, where $\nu_0(x; \varepsilon)$ is the free local DoS. The precise relationship between the two formulations has been discussed in Ref. [189] for the 1D situation. When the scattering matrix is defined like in (2), we have

$$\Delta(\varepsilon) = (2i\pi)^{-1} \partial_{\varepsilon} [\ln \det \mathcal{S} - 2i\sqrt{\varepsilon}L].$$

Note that the second term can be absorbed in a redefinition of the phases of \mathcal{S} (see Appendix of Ref. [189] for details).

⁶ Note that (25) assumes that the scattering phase shifts are measured with respect to the free eigenstates, $\psi_{\ell}(r) \sim (1/r)[e^{-ikr} + e^{ikr+2i\eta_{\ell}}]$, as it is done usually in scattering theory, instead of (2).

2.4. Few remarks

Characterisation of the spectral properties of open systems has important applications as it allows to express several physical observables in terms of scattering properties at the heart of the Landauer-Büttiker description of quantum transport. For example, (22) allows to measure the charge inside a conductor in terms of its scattering matrix, an idea which turns out to be central in many developments of Büttiker involving screening, which are reviewed below. Several remarks :

- (i) The Krein-Friedel relation was studied in metric graphs in [189, 193]. In such systems with non trivial topology, the system might exhibit so-called “*bound states in the continuum*” (BIC), i.e. a discrete spectrum *superimposed* onto the continuum spectrum (this occurs when symmetries allow some bound states to remain uncoupled to the continuum) [208, 185]. In this case, the Krein-Friedel relation only characterises the continuous part of the spectrum [193, 195, 190] (see [189, 193] for illustrations).
- (ii) A related problem concerns the role of the transmission phase. As it is clear from the polar representation (10), the Friedel phase Φ_f , which enters the relation (22) as $\det \mathcal{S} = e^{i\Phi_f}$, is related to the transmission phase $\delta_t = (\Phi_f + \pi)/2$ in 1D, as the magnetic field can always be removed by a gauge transformation on the infinite line. Thus the DoS can be as well related to δ_t in 1D [11]. If the system has a complex topology, the transmission may vanish which causes transmission phase jumps. As a consequence of these phase jumps, the Friedel phase and the transmission phase differ ; the density of states is then only related to the Friedel phase, as discussed by Taniguchi and Büttiker [187]. This question has been examined numerically in quantum dots by Lévy Yeyati and Büttiker [129].
- (iii) In Ref. [182], the Krein-Friedel relation was considered in the quasi-one-dimensional situation (wave guide) which is relevant in order to describe the electric contacts of mesoscopic structures.
- (iv) A formulation of the time dependent transport was developed in [158] (using the Keldysh formalism) and was shown to involve the time delay matrix expressed in terms of Green’s functions.

- (v) In Subsection 2.1, the Wigner time delay has been introduced by considering the delay of the position of a wave packet with narrow dispersion in energy. This corresponds to follow the *mean* position of the particle, cf. Fig. 2 (where averaging would be taken with respect to measurements in a given quantum state). The remark also holds for all times introduced in Section 2. Elaborating on Refs. [133, 23], Smilansky has introduced a distribution of time delay for a particle injected in channel i and detected in channel f [179]

$$P_{fi}(\tau) = \frac{1}{2\pi} \left| \int_0^\infty d\varepsilon \hat{\Phi}(\varepsilon) \mathcal{S}_{fi}(\varepsilon) e^{-i\varepsilon\tau} \right|^2, \quad (26)$$

where the wave packet envelope $\hat{\Phi}(\varepsilon)$ is a real and narrow normalised function. Smilansky has also discussed the classical limit of the distribution. In the simple case of scattering on the half line, we can check that the mean time is related to the Wigner time delay $\tau_W(\varepsilon) = -i\mathcal{S}(\varepsilon)^* \mathcal{S}'(\varepsilon)$:

$$\begin{aligned} \int d\tau P(\tau) \tau &= -i \int d\varepsilon [\hat{\Phi}(\varepsilon) \mathcal{S}(\varepsilon)]^* [\hat{\Phi}(\varepsilon) \mathcal{S}(\varepsilon)]' \\ &= \int d\varepsilon \hat{\Phi}(\varepsilon)^2 \tau_W(\varepsilon) \end{aligned} \quad (27)$$

where we have used unitarity $|\mathcal{S}|^2 = 1$ and normalisation $\int_0^\infty d\varepsilon \hat{\Phi}(\varepsilon)^2 = 1$. For several channels, a similar relation with the Wigner-Smith time delay matrix (17) only holds with an additional summation

$$\sum_f \int d\tau P_{fi}(\tau) \tau = \int d\varepsilon \hat{\Phi}(\varepsilon)^2 \mathcal{Q}_{ii}. \quad (28)$$

Similarly, a summation over the ingoing channel involves the matrix (18)

$$\sum_i \int d\tau P_{fi}(\tau) \tau = \int d\varepsilon \hat{\Phi}(\varepsilon)^2 \tilde{\mathcal{Q}}_{ff}. \quad (29)$$

We stress that the distribution (26) describes the fluctuations of time due to the stochasticity of the scattering process in *a given sample*. It should *not* be confused with the time delay distributions studied below in Sections 3 and 4, which characterize the *sample to sample* (mesoscopic) fluctuations of the Wigner time delay $\tau_W(\varepsilon)$ or the related quantities (Wigner-Smith matrix, proper times, partial times, etc).

3. Disordered systems (localised regime)

The review [56] has underlined the close relationship between Wigner time delay for 1D disordered systems and exponential functionals of the Brownian motion, which have been widely studied in the mathematical literature [212] (see also [58] for a brief review). Let us first recall few properties which will be useful for the following.

3.1. Exponential functionals of the Brownian motion

We introduce the random variable

$$Z_X^{(\mu)} = \int_0^X dx e^{-2(\mu x + B(x))}, \quad (30)$$

where $B(x)$ is a normalised Brownian motion starting from $B(0) = 0$ (Wiener process). The distribution of (30) was found in [147, 57] (see also [58]) :

$$\begin{aligned} \psi_X^{(\mu)}(Z) &= 2 \frac{e^{-1/(2Z)}}{(2Z)^{1+\mu}} \sum_{0 \leq n < \mu/2} \frac{(-1)^n (\mu - 2n)}{\Gamma(1 + \mu - n)} \\ &\quad \times e^{-2Xn(\mu - n)} (2Z)^n L_n^{\mu-2n} \left(\frac{1}{2Z} \right) \\ &+ \frac{e^{-1/(4Z)}}{2\pi^2 (2Z)^{(1+\mu)/2}} \int_0^\infty ds s \sinh(\pi s) \left| \Gamma \left(\frac{is - \mu}{2} \right) \right|^2 \\ &\quad \times W_{(1+\mu)/2, is/2} \left(\frac{1}{2Z} \right) e^{-(X/2)(\mu^2 + s^2)}, \end{aligned} \quad (31)$$

where $L_n^\alpha(x)$ is a Laguerre polynomial and $W_{\mu,\nu}(z)$ the Whittaker function [105]. Eq. (31) shows that the random variable admits the limit law

$$\psi_\infty^{(\mu)}(Z) = \frac{1}{2^\mu \Gamma(\mu)} Z^{1+\mu} e^{-1/(2Z)} \quad \text{for } \mu > 0. \quad (32)$$

For finite X , the functional is characterised by exponential moments

$$\langle (Z_X^{(\mu)})^n \rangle \underset{X \rightarrow \infty}{\simeq} 2^{-n} \frac{\Gamma(n - \mu)}{\Gamma(2n - \mu)} e^{2n(n - \mu)X}. \quad (33)$$

The precise expression of the moments can be found in Ref. [154] for $\mu = 0$ and Ref. [147] in the general case.⁷

⁷ Note however that Eq. 4.4 of Ref. [147] for $\mu = 0$ contains a misprint : C_{2n}^k should be replaced by C_{2n}^{n-k} .

3.2. Disorder on the half line - Single parameter scaling and universality

The interest for time delays in disordered systems has started with the work of Sulem and coworkers [83] on stochastic resonances (see also [160]). The idea that localised states could produce sharp resonances was later developed by Azbel [12]. The first statistical analysis of the Wigner time delay in a one-dimensional situation was provided by Jayanavar, Vijayagovindan and Kumar [114]. These authors have studied the time delay distribution $P_L(\tau)$ for the Schrödinger equation (1) on the half line (Fig. 1) when $V(x)$ is a Gaussian white noise. They have identified the existence of a limit law with power law tail $P_\infty(\tau) \sim \tau^{-2}$ for an infinitely long disordered region, $L \rightarrow \infty$, although their expression of $P_\infty(\tau)$ was partly incorrect (with a non vanishing distribution as $\tau \rightarrow 0$). Similar conclusions were obtained in Ref. [115] when the potential $V(x)$ is the *integral* of a white noise, i.e. a Brownian motion. As it was recognised later [169], the incorrect non vanishing $P_\infty(0)$ of Refs. [114, 115] was due to some inappropriate averaging over the fast phase variable in the weak disorder regime. Short after, Heinrichs identified that the exponential moments, $\langle \tau^n \rangle \sim \exp[2n(n-1)L/\xi]$ where ξ is the localisation length, are characteristic of a log-normal tail $P_L(\tau)$ when the length L is finite [110]. Inspired by the work of Faris and Tsay [76], we established in Ref. [60] the connection between the Wigner time delay and exponential functionals of the Brownian motion. We analysed two different disordered models exhibiting qualitatively different spectral and localisation properties, namely the Schrödinger Hamiltonian $H = -\partial_x^2 + V(x)$ and the supersymmetric Hamiltonian $H = -\partial_x^2 + m(x)^2 + m'(x)$ [i.e. the square of the Dirac Hamiltonian (41)], where the potential $V(x)$ and the mass $m(x)$ are Gaussian white noises. This analysis suggested the universality of the statistical properties, which was later demonstrated in Ref. [194]. This can be understood from the integral representation (20): in the *weak disorder* regime we may write $\tau_W \simeq 2\pi \int_0^L dx |\psi_\varepsilon(x)|^2$. Using that the logarithm of the envelope of the wave function is a Brownian motion with drift, we have related the time delay to the functional (30) [194, 56]⁸

$$\tau_W \stackrel{(\text{law})}{=} 2\tau_\xi Z_{L/\xi}^{(1)}, \quad (34)$$

⁸ An equality in law $\stackrel{(\text{law})}{=}$ relates two quantities with the same statistical properties. For example, the well-known

where ξ is the localisation length⁹ and $\tau_\xi = \xi/v = \xi/(2k)$ is the time needed for a ballistic motion on the localisation length ξ . The value of the drift $\mu = 1$ originates from the equality

$$\langle \ln |\psi_\varepsilon(x)| \rangle \simeq \text{Var}(\ln |\psi_\varepsilon(x)|) \quad \text{for } x \gg \xi, \quad (35)$$

valid in the weak disorder regime [10]. Eq. (35) is known as “*single parameter scaling*” [7, 55], as the full distribution of $\ln |\psi_\varepsilon(x)|$ (or of the conductance) is characterised by a unique length scale, the localisation length defined by $1/\xi = \lim_{x \rightarrow \infty} (1/x) \ln |\psi_\varepsilon(x)|$. This question has been re-discussed more recently for 1D disordered systems [68, 69, 177, 170]. The universality of the Wigner time delay statistical properties are thus understood as a direct consequence of the universality of the localisation properties in 1D in the weak disorder regime.¹⁰ In other terms, Eq. (34) means that the distribution of the Wigner time delay is, in the universal (weak disorder) regime,

$$P_L(\tau) = \frac{1}{2\tau_\xi} \psi_{L/\xi}^{(1)}\left(\frac{\tau}{2\tau_\xi}\right). \quad (36)$$

If the length L of the disordered region goes to infinity we get the limit law [60, 188, 194]

$$P_\infty(\tau) = \frac{\tau_\xi}{\tau^2} e^{-\tau_\xi/\tau}. \quad (37)$$

The limits $L \rightarrow \infty$ and $\tau \rightarrow \infty$ do not commute: for finite L the tail of the distribution is log-normal [110, 194], $P_L(\tau) \sim \exp[-(\xi/8L) \ln^2(\tau/\tau_\xi)]$ for $\tau \rightarrow \infty$. The leading term of the moments was also given in Ref. [60]:

$$\langle \tau \rangle = 2vL \quad (38)$$

$$\langle \tau^n \rangle \simeq \frac{(n-2)!}{(2n-2)!} \tau_\xi^n e^{2n(n-1)L/\xi} \quad \text{for } n > 1, \quad (39)$$

confirming the main exponential behaviour found earlier by Heinrichs [110] and providing the pre-exponential factor (a more precise expression of the moments was given in Ref. [194]). Note that

scaling properties of the Brownian motion may be conveniently written $B(\lambda x) \stackrel{(\text{law})}{=} \lambda^{1/2} B(x)$, where $B(x)$ is a Brownian motion.

⁹ For the model (1) with $\langle V(x)V(x') \rangle = \sigma \delta(x-x')$, the localisation length is $\xi \simeq 8\varepsilon/\sigma$ for weak disorder $\sigma^{2/3} \ll \varepsilon$.

¹⁰ The time delay distribution was also studied in the non universal (strong disorder) regime in [194] for a particular model.

the log-normal distribution is characterised by moments of the form $\exp[cn^2]$; the fact that the moments (39) increase with a lower rate with n , as $\sim \exp[2n(n-1)L/\xi]$, reflects the existence of a limit law with tail τ^{-2} (see § 3.1). The importance of Azbel resonances was emphasized as it allowed to recover the moments by some heuristic argument [194]. A lattice model was studied in Ref. [157], leading to the same conclusions.

3.3. Disorder in 1D with a chiral symmetry

The localisation properties of disordered systems are mainly controlled by their dimensionality [1] and symmetries. A first classification of symmetries was given by Wigner and Dyson in the context of random matrix theory, depending on the presence of time reversal symmetry and spin rotational symmetry, leading to the orthogonal ($\beta = 1$), unitary ($\beta = 2$) and symplectic classes ($\beta = 4$) [140]. These three symmetry classes were later completed by others, identified according to the presence or not of two other types of discrete symmetries which might occur in condensed matter physics : the chiral (or sublattice) symmetry and the particle-hole symmetry. These has led to the classification in terms of ten symmetry classes [214, 6, 75]. A one-dimensional disordered model with a chiral symmetry which has attracted a lot of attention is the Dirac equation with a random mass (see [27, 61, 196] for reviews)¹¹

$$\mathcal{H}_D \Psi(x) = \varepsilon \Psi(x) \quad (40)$$

with

$$\mathcal{H}_D = i\sigma_2 \partial_x + \sigma_1 m(x), \quad (41)$$

where σ_i are the Pauli matrices. A chiral symmetry is the anticommutation of the Hamiltonian with a unitary operator, here $\sigma_3 \mathcal{H}_D \sigma_3 = -\mathcal{H}_D$. The Wigner time delay distribution for this model was obtained in Ref. [60] in the “universal” (weak disorder) regime, leading to the same conclusions as for the Schrödinger equation, Eqs. (34,36), provided that the localisation length is modified, $\tau_\xi = \xi \simeq 2/g$ for $\varepsilon \gg g$ (in the Dirac equation, the velocity is equal to unity). The energy ε can be viewed as a

¹¹ The case of higher dimensions is also very important : besides that it appears in several contexts of condensed matter physics (graphene, etc), it is of fundamental interest as it was shown that all symmetry classes can be represented within the 2D Dirac equation with randomness [19].

parameter which tunes the chiral symmetry breaking : at the symmetry point $\varepsilon = 0$, where chiral symmetry holds, the Wigner time delay presents different properties as we now discuss.

The scattering problem on the half line for the Dirac equation is settled as follows : we consider again the situation of Fig. 1 where the mass is non zero on the interval $[0, L]$. In the free region we can decompose the spinor of energy $\varepsilon > 0$ as the combination of an incoming plane wave and a reflected plane wave :

$$\Psi_\varepsilon(x) \underset{x>L}{=} \begin{pmatrix} 1 \\ i \end{pmatrix} e^{-i\varepsilon(x-L)} + \begin{pmatrix} -1 \\ i \end{pmatrix} e^{i\varepsilon(x-L)+i\delta_r(\varepsilon)}. \quad (42)$$

Following [60, 188] we can parametrize the spinor on $[0, L]$ as $\Psi(x) = \rho(x) (\sin \theta(x), -\cos \theta(x))$, which leads to the equation for the phase

$$\frac{\partial \theta(x)}{\partial x} = \varepsilon + m(x) \sin [2\theta(x)]. \quad (43)$$

The matching with (42) shows that the phase shift is given by $\delta_r = 2\theta(L)$. We choose the initial condition $\theta(0) = 0$ or $\pi/2$, which corresponds to an infinite mass $m(x) = +\infty$ or $m(x) = -\infty$ for $x < 0$, respectively (these boundary conditions confine the particle on \mathbb{R}_+ ; they are the only ones which do not break the chiral symmetry [196]). Following [60] we introduce the variable $\mathcal{Z}(x) = 2\partial\theta(x)/\partial\varepsilon$ which obeys

$$\frac{\partial \mathcal{Z}(x)}{\partial x} = 2 + 2m(x) \mathcal{Z}(x) \cos [2\theta(x)] \quad (44)$$

with initial condition $\mathcal{Z}(0) = 0$ and provides the value of the Wigner time delay $\tau_W = \mathcal{Z}(L)$.

The analysis of the symmetry point $\varepsilon = 0$ is easy : the phase remains locked at 0 or $\pi/2$ (corresponding to $m(x) = \pm\infty$ for $x < 0$), therefore one can integrate (44). On gets

$$\tau_W = 2 \int_0^L dx e^{\pm 2 \int_x^L dx' m(x')} \quad \text{at } \varepsilon = 0. \quad (45)$$

For the sake of concreteness, we now consider the situation where the mass is a Gaussian white noise

$$\begin{cases} \langle m(x)m(x') \rangle_c = g \delta(x-x') \\ \langle m(x) \rangle = \mu g \end{cases} \quad (46)$$

where $\langle XY \rangle_c = \langle XY \rangle - \langle X \rangle \langle Y \rangle$. The representation (45) makes clear the identity

$$\tau_W \stackrel{(\text{law})}{=} \frac{2}{g} Z_{gL}^{(\mp\mu)} \quad \text{at } \varepsilon = 0, \quad (47)$$

which is distributed according to ¹²

$$P_L(\tau) = \frac{g}{2} \psi_{gL}^{(\mp\mu)} \left(\frac{g\tau}{2} \right) \quad \text{at } \varepsilon = 0, \quad (49)$$

where the dimensionless function was defined above, Eq. (31).

3.3.1. The critical case $\langle m(x) \rangle = 0$

The distribution for $\mu = 0$ was obtained in [184] (although the distribution has been already determined in another context [147], what was used in [188]). It is characterised by a log-normal tail $P_L(\tau) \sim \exp[-1/(8gL)\ln^2(\tau)]$ and exponential moments

$$\langle \tau^n \rangle \simeq \frac{(n-1)!}{(2n-1)!} g^{-n} e^{2n^2 gL} \quad \text{for } n \geq 1. \quad (50)$$

Although the distribution has no limit law, the large L limit presents the power law behaviour [147]

$$P_L(\tau) \underset{L \rightarrow \infty}{\simeq} \frac{1}{\sqrt{2\pi gL} \tau} e^{-1/(g\tau)} \quad (51)$$

cut off by the log-normal tail for $\tau \rightarrow \infty$. The absence of a limit law may be associated with the delocalisation of the model for $\varepsilon = 0$, as all time scales extracted from the distribution increase with L (see [27, 61, 196] for reviews).

3.3.2. The case $\langle m(x) \rangle = \mu g \neq 0$: time delay as a probe for zero mode

We now analyse the case $\langle m(x) \rangle = \mu g \neq 0$ which was not considered in the literature. Although the spectral properties of the model are invariant under the change of the sign of the mass, this is not the case for the scattering properties : neither for the phase distribution [104] nor for the Wigner time delay distribution, as we demonstrate here. The distribution $P_L(\tau)$ has an interesting property : the existence of a limit law is correlated with the choice of the boundary condition at $x = 0$ and the sign of the average mass $\langle m(x) \rangle = \mu g$. If we choose the boundary condition corresponding to $m(x) = \pm\infty$ for $x < 0$, we obtain the limit law only for $\mp\mu > 0$

$$P_\infty(\tau) = \frac{g}{\Gamma(|\mu|)(g\tau)^{1+|\mu|}} e^{-1/(g\tau)}. \quad (52)$$

¹² It is interesting to compare (49) for $\varepsilon = 0$ with the high energy ($\varepsilon \gg g$) distribution (36) obtained in the absence of a chiral symmetry, which takes the explicit form

$$P_L(\tau) = (g/4) \psi_{gL/2}^{(1)}(g\tau/4) \quad \text{at } \varepsilon \gg g. \quad (48)$$

Conversely, for $\pm\mu > 0$, there is no limit law as $L \rightarrow \infty$. We can correlate the existence of the limit law with the presence of a chiral zero mode located at the boundary $x = 0$, what occurs when the sign of the mass changes at the boundary :

$m(x)$ for $x < 0$	$\langle m(x) \rangle$ for $x > 0$	zero mode	limit law $P_\infty(\tau)$
$-\infty$	> 0	yes	yes
$-\infty$	< 0	no	no
$+\infty$	> 0	no	no
$+\infty$	< 0	yes	yes

The existence of zero modes in the multichannel Dirac equation with random mass was recently studied in Ref. [104], where their topological nature was underlined.

3.4. Related questions and remarks

- (i) An alternative derivation of the stationary distribution (37) was proposed in Ref. [168], by using the relation between the time delay and the reflection coefficient in the presence of a constant imaginary component in the potential (wave amplification), i.e. making use of analytic properties of the scattering matrix. This point is further discussed in the next paragraph.
- (ii) A discrete Anderson model with Cauchy disorder (Lloyd model [131]) was studied in Ref. [68], for which the relation between the two cumulants of $\ln|\psi_\varepsilon(x)|$ presents an extra factor 2 (i.e. drift $\mu = 1/2$), compared with the usual form for the single parameter scaling, Eq. (35).¹³ Some time ago, I performed some (unpublished) numerical investigations, which had shown that the distribution of the Wigner time delay is however still characterised by a power law tail τ^{-2} (and not $\tau^{-3/2}$). It would thus be interesting to clarify this observation in connection with the discussion of the section.
- (iii) Correlations in energy $\langle \tau_W(\varepsilon) \tau_W(\varepsilon') \rangle$ were analysed by Titov and Fyodorov [201].
- (iv) Time delay and resonance width are closely related (see the review [117]). The resonance width distribution was studied in [123, 106].

¹³ Localisation for potentials with power law distribution was studied in [21, 69, 131, 202].

- (v) We have focused the discussion on the Wigner time delay characterizing the reflection problem on the half line. The time delay describing the transmitted wave through a disordered medium was analysed in [26].

3.5. Time delay, absorption/amplification and analyticity (Dirac case)

We discuss here an interesting connection between time delay and the question of absorption/amplification. This relation, consequence of the analytic properties of the scattering matrix, is completely general and has been used in various contexts : 1D random Schrödinger Hamiltonians [165, 168], multichannel disordered models [17] (standard class) or chiral/BdG classes [200] and also in Ref. [84] (Appendix A also follows from this idea). The idea is simple : given the $N \times N$ scattering matrix (without absorption) $\mathcal{S}(\varepsilon)$, the case with a uniform absorption/amplification (in space) is described by the reflection matrix can be accounted for through the substitution

$$\varepsilon \longrightarrow \tilde{\varepsilon} = \varepsilon + i\Gamma/2. \quad (53)$$

Thus, the evolution is not anymore unitary and the reflection matrix

$$\begin{aligned} [\mathcal{S}(\tilde{\varepsilon})]^\dagger \mathcal{S}(\tilde{\varepsilon}) &= \mathbf{1}_N + i\Gamma \mathcal{S}^\dagger \partial_\varepsilon \mathcal{S} + \mathcal{O}(\Gamma^2) \\ &= \mathbf{1}_N - \Gamma \mathcal{Q} + \mathcal{O}(\Gamma^2) \end{aligned} \quad (54)$$

encodes useful information. In the one channel case, the weak absorption/amplification limit of the reflection coefficient

$$R = |\mathcal{S}(\tilde{\varepsilon})|^2 = 1 - \Gamma \tau_W + \mathcal{O}(\Gamma^2) \quad (55)$$

is related to the Wigner time delay at lowest order in Γ [the same holds in general if one traces Eq. (54)]. This expression shows that the absorption probability (for $\Gamma > 0$) is proportional to the time spent in the scattering region, at lowest order in Γ .

We consider below the scattering problem for the one-dimensional Dirac equation with a random mass, which is more rich than the Schrödinger case discussed previously in Refs. [165, 168].

3.5.1. The scattering problem

We start from the Dirac equation (40,41). We can restrict to $\varepsilon \geq 0$ without loss of generality. In order to introduce a reflection coefficient in a more

general setting, we consider the scattering problem on the full line : the scattering state incoming from the right is

$$\Psi_\varepsilon^{(R)}(x) = \begin{cases} t \begin{pmatrix} 1 \\ i \end{pmatrix} e^{-i\varepsilon x} & \text{for } x < 0 \\ \alpha(x) \begin{pmatrix} 1 \\ i \end{pmatrix} e^{-i\varepsilon x} - \beta(x) \begin{pmatrix} -1 \\ i \end{pmatrix} e^{i\varepsilon x} & \text{for } 0 < x < L \\ \begin{pmatrix} 1 \\ i \end{pmatrix} e^{-i\varepsilon(x-L)} + r' \begin{pmatrix} -1 \\ i \end{pmatrix} e^{i\varepsilon(x-L)} & \text{for } x > L \end{cases} \quad (56)$$

injecting this form into (40), we get two differential equations for the two functions

$$\frac{d\alpha(x)}{dx} = m(x) \beta(x) e^{+2i\varepsilon x} \quad (57)$$

$$\frac{d\beta(x)}{dx} = m(x) \alpha(x) e^{-2i\varepsilon x} \quad (58)$$

Matching at boundaries gives $\alpha(0) = t$ and $\beta(0) = 0$ while $\alpha(L) = e^{+i\varepsilon L}$ and $\beta(L) = -r' e^{-i\varepsilon L}$. The reflection coefficient r' is most naturally obtained by introducing the variable $\Upsilon(x) \stackrel{\text{def}}{=} \beta(x)/\alpha(x)$, which obeys

$$\frac{d\Upsilon(x)}{dx} = m(x) (e^{-2i\varepsilon x} - \Upsilon(x)^2 e^{+2i\varepsilon x}) \quad (59)$$

for initial condition $\Upsilon(0) = 0$. The reflection coefficient is then given by $\Upsilon(L) = -r' e^{-2i\varepsilon L}$. Knowing $\Upsilon(x)$, the transmission coefficient could be in principle deduced by dividing (57) by α and integrating : one gets $t = \exp \{ i\varepsilon L - \int_0^L dx m(x) \Upsilon(x) e^{2i\varepsilon x} \}$.

At this point, it is convenient to parametrize the complex variable $\Upsilon(x) = -\rho(x) e^{2i\theta(x) - 2i\varepsilon x}$ in terms of two real variables ρ and θ . The reflection amplitude is $r' = +\rho(L) e^{2i\theta(L)}$. Thus

$$\frac{d\theta}{dx} = \varepsilon + m(x) \frac{\rho^2 + 1}{2\rho} \sin 2\theta \quad (60)$$

$$\frac{d\rho}{dx} = 2m(x) (\rho^2 - 1) \cos 2\theta \quad (61)$$

for the initial conditions ¹⁴ $\theta(0) = \frac{\pi}{4}$ and $\rho(0) = 0$.

¹⁴ The initial condition for the phase follows from the perturbative expression of the reflection coefficient : from the Lippmann-Schwinger formula we obtain

$$\psi(x) \simeq \psi^{(0)}(x) + \int dx' G_0^R(x, x'; k^2) V(x') \psi^{(0)}(x') \quad (62)$$

If instead we impose the boundary conditions $\theta(0) = 0$ and $\rho(0) = 1$, corresponding to confinement on the positive axis (i.e. scattering on the half line), we see that the amplitude remains locked at $\rho(x) = 1 \forall x$ and we recover Eq. (43) for the phase.

3.5.2. Absorption/amplification

We now add the effect of absorption or amplification by introducing a complex term in the Hamiltonian (41)

$$\mathcal{H} = i\sigma_2 \partial_x + \sigma_1 m(x) + A_0 \quad \text{with } A_0 = -i\Gamma/2 \quad (64)$$

$\Gamma > 0$ for absorption and $\Gamma < 0$ for amplification. This is equivalent to perform the substitution (53) in the Dirac equation (40). Performing the same steps as in § 3.5.1, we obtain that Eqs. (60,61) are replaced by

$$\frac{d\theta}{dx} = \text{Re}(\tilde{\varepsilon}) + m(x) \frac{\rho^2 + 1}{2\rho} \sin 2\theta \quad (65)$$

$$\frac{d\rho}{dx} = -2 \text{Im}(\tilde{\varepsilon}) \rho + m(x) (\rho^2 - 1) \cos 2\theta \quad (66)$$

3.5.3. Dirac point ($\varepsilon = 0$)

We first consider the Dirac point ($\varepsilon = \text{Re } \tilde{\varepsilon} = 0$), where chiral symmetry holds. As we have seen in § 3.3, the phase is locked at $\theta = \pi/2$ (this ensures the existence of a zero mode and a limit law for the time delay when $\mu > 0$). Then

$$\frac{d\rho}{dx} = -\Gamma \rho - m(x) (\rho^2 - 1) \quad (67)$$

We consider $R = \rho^2$, which obeys

$$\frac{dR}{dx} = -2\Gamma R + 2m(x) \sqrt{R} (1 - R) \quad (68)$$

(at $x = L$, $R(L)$ is the reflection probability). We see on this equation that the diffusion term in Eq. (68) vanishes at $R = 1$. Only the term $-2\Gamma R$ of the drift allows the process to cross the point $R = 1$. For absorption, the drift is negative, therefore the distribution has support $[0, 1]$. For amplification it has support $[1, \infty[$. Assuming that $m(x)$ is a Gaussian white noise, Eq. (46), the (forward) generator

describing the diffusion (68) is

$$\mathcal{G}_R^\dagger = 2 \frac{\partial}{\partial R} \left(\Gamma R - \mu g \sqrt{R} (1 - R) \right) + 2g \frac{\partial}{\partial R} \sqrt{R} (1 - R) \frac{\partial}{\partial R} \sqrt{R} (1 - R) \quad (69)$$

i.e. the distribution of the process obeys the Fokker-Planck equation $\partial_x W_x(R) = \mathcal{G}_R^\dagger W_x(R)$. For $\Gamma > 0$ (absorption), we find the stationary distribution solution of $\mathcal{G}_R^\dagger W_\infty(R) = 0$:

$$W_\infty(R) = \frac{C}{\sqrt{R} (1 - \sqrt{R})^{1+\mu} (1 + \sqrt{R})^{1-\mu}} e^{-\frac{\Gamma/g}{1-R}} \quad (70)$$

for $R \in [0, 1]$, where C is a normalisation constant. For $\Gamma < 0$ (amplification), we get

$$W_\infty(R) = \frac{C}{\sqrt{R} (\sqrt{R} - 1)^{1+\mu} (\sqrt{R} + 1)^{1-\mu}} e^{-\frac{|\Gamma|/g}{R-1}} \quad (71)$$

with support $[1, \infty[$.

Using (55) for $\Gamma \rightarrow 0$ allows to map the stationary distribution $W_\infty(R)$ onto (52). Moreover, the correspondence can be established at the level of the two generators

$$\mathcal{G}_R^\dagger \xrightarrow{\Gamma \rightarrow 0} \mathcal{G}_\tau^\dagger = \frac{\partial}{\partial \tau} (2\mu g \tau - 2) + 2g \frac{\partial}{\partial \tau} \tau \frac{\partial}{\partial \tau} \tau \quad (72)$$

where \mathcal{G}_τ^\dagger describes the diffusion (44) for $\theta(x) = \pi/2$.

3.5.4. Universal (high energy) regime

Unitary evolution —. Let us first briefly discuss the case without absorption/amplification ($\Gamma = 0$). In the universal (high energy) regime $\varepsilon \gg g$, we can average over the phase (fast variable) Eqs. (65,66), assuming a uniform distribution. The distribution of $\rho(x) = e^{-w(x)}$ (reflection coefficient) is described by the diffusion equation controlling the radial diffusion on a surface of constant negative curvature [94, 10] (see also [188]) :

$$\partial_x Q(w; x) \simeq 2\gamma \partial_w^2 [\sinh^2 w Q(w; x)] \quad (73)$$

where $\gamma \simeq g/2$ is the high energy Lyapunov exponent. The fact that one gets the same result for the Dirac (here) and Schrödinger [10] equations has its origin in the universality of the high energy regime. The distribution has no limit law for $x \rightarrow \infty$ as the reflection coefficient decays exponentially with the system size L , hence the process $w(x)$ grows linearly.

hence

$$r' = -\frac{i}{2k} \int_0^L dx' V(x') e^{-2ikx'} + \mathcal{O}(V^3) \quad (63)$$

thus $r'/|r'| \rightarrow -i$ in the limit $L \rightarrow 0$.

Finite absorption/amplification rate —. For $\Gamma \neq 0$, from Eq. (66), we see that the (forward) generator receives another contribution $\mathcal{G}_w^\dagger = g\partial_w^2 \sinh^2 w \rightarrow \mathcal{G}_w^\dagger = -\Gamma\partial_w + g\partial_w^2 \sinh^2 w$. At this point it is more convenient to deal with the reflection probability and introduce the variable $R = \rho^2 = e^{-2w}$. The new generator is given by $\mathcal{G}_R^\dagger = (\partial w/\partial R)\mathcal{G}_w^\dagger(\partial R/\partial w)$. We get finally

$$\mathcal{G}_R^\dagger = 2\frac{\partial}{\partial R} \left(\Gamma R - \frac{g}{4}(1 - R^2) \right) + g\frac{\partial}{\partial R} \sqrt{R}(1 - R) \frac{\partial}{\partial R} \sqrt{R}(1 - R). \quad (74)$$

For finite absorption/amplification rate Γ , the reflection coefficient has a limit law for $L \rightarrow \infty$. For $\Gamma > 0$ (absorption), it is easy to show that the stationary distribution is

$$W_\infty(R) = \frac{\Gamma\tau_\xi}{(1 - R)^2} e^{-\frac{\Gamma\tau_\xi}{1 - R}} \quad (75)$$

with support $R \in [0, 1]$. We have introduced the parameter $\tau_\xi = \xi/v = 2/g$ (with $v = 1$ for the Dirac equation) in order to stress the universal character of this result. This is in exact correspondence with the distribution found in [168] for the disordered Schrödinger equation, as it should since we are dealing here with the universal regime. As pointed out by Ramakrishna and Kumar, in the limit $\Gamma \rightarrow 0$, Eq. (75) can be mapped onto the limit law of the time delay distribution (37) by using (55). Here, we stress that, as at the Dirac point, this mapping can be established at the level of the generators and is thus more general than the one stated in [168]: using (55), it is straightforward to check that

$$\mathcal{G}_R^\dagger \xrightarrow{\Gamma \rightarrow 0} \mathcal{G}_\tau^\dagger = \frac{\partial}{\partial \tau} (g\tau - 2) + g\frac{\partial}{\partial \tau} \tau \frac{\partial}{\partial \tau} \tau \quad (76)$$

which is indeed the generator controlling the distribution of the Wigner time delay in the high energy limit (given in [60, 188]). This is the generator of the process (30) for $\mu = 1$, according to § 3.2.

3.6. Beyond 1D

3.6.1. Multichannel disordered wires

The extension to the quasi-1D situation (a wave guide with N channels) has been considered by several authors (e.g. see [149, 206]). Using the relation between the time delays and the reflection probabilities in an absorbing or amplifying medium (i.e. the analyticity of the \mathcal{S} -matrix), Beenakker

and Brouwer obtained the distribution of the proper time delays describing the scattering on a semi-infinite disordered wave guide [14, 17]:

$$\mathcal{P}(\gamma_1, \dots, \gamma_N) \propto \prod_{i < j} |\gamma_i - \gamma_j|^\beta \prod_k e^{-\beta\gamma_k/2}, \quad (77)$$

where $\gamma_k = \tau_s/\tau_k$ is the inverse of the proper time and τ_s a characteristic time related to the disorder strength. The result relies on the isotropy assumption (among the channels), i.e. describes the weakly disordered (quasi-1D diffusive) regime; in this case the localisation length increases with the number of channels as [13] $\xi^{(\text{quasi 1D})} = \xi^{(1D)} [1 + \beta(N - 1)/2]$. The distribution (77) is a particular instance of the Laguerre ensemble of random matrix theory (RMT). For one channel $N = 1$, one recovers the limit law (37) by setting $\beta\tau_s/2 = \tau_\xi$. In the limit of large N , the marginal distribution of the rates γ_a 's is the Marčenko-Pastur law $\rho_{\text{MP}}(x) = (2\pi)^{-1} \sqrt{(4 - x)/x}$, after rescaling $x = \gamma/N$. Accordingly, the marginal distribution of the proper times is

$$w_N^{(\beta)}(\tau) \underset{N \rightarrow \infty}{\simeq} N \rho(N\tau) \text{ with } \rho(y) = \frac{\sqrt{4y - 1}}{2\pi y^2}. \quad (78)$$

Starting from (77), the distribution of the Wigner time delay $\tau_W = (1/N) \sum_i \gamma_i^{-1}$ was obtained in Ref. [103, 97]:

$$\mathcal{P}_N^{(\beta)}(\tau) \underset{N \rightarrow \infty}{\simeq} \frac{C_\beta}{\tau^2} \times \exp \left\{ -\frac{27\beta}{64\tau^2} + \left(1 - \frac{\beta}{2} \right) \frac{9(2 - \sqrt{3})}{4\tau} \right\} \quad (79)$$

where β is the Dyson index for orthogonal ($\beta = 1$) or unitary ($\beta = 2$) symmetry classes. C_β is a normalisation constant. The distribution presents the same power law tail as in the strictly one-dimensional case $\mathcal{P}_N^{(\beta)}(\tau) \sim 1/\tau^2$. This can be understood from the fact that the physics at large time (i.e. large scale) is expected to be dominated by a single channel (the less localised one).

Denoting by $\nu_L(\varepsilon_F)$ the DoS of the multichannel disordered wire of length L , (79) can be interpreted as the limit law for $\tau_W \simeq 2\pi\nu_L(\varepsilon_F)/N$ in the $L \rightarrow \infty$ limit. The fact that all moments of τ_W are infinite in this case is thus simply related to the divergence of the DoS when $L \rightarrow \infty$, as in the strictly 1D situation.

Finally, we quote [155], where the marginal distribution of the proper times was derived for the

multichannel case for finite L (it converges towards (78) for $L \rightarrow \infty$).

3.6.2. Higher dimensions

- (i) Some measurements of time delays were performed with electromagnetic waves in Ref. [178, 93, 52].
- (ii) The case of higher dimensions was considered by Ossipov and Fyodorov [156] who analysed a two dimensional situation in the weakly disordered regime. The localised regime was studied by numerical simulations in Ref. [211].
- (iii) A study of time delay at a critical point like the metal-insulator transition was performed in Ref. [118, 84, 156] (see the review [117]).

4. Random matrix approach for quantum dots

In systems with ergodic properties, like a chaotic cavity with narrow contacts (Figs. 9 and 10), or a weakly disordered cavity in the ergodic regime, it is natural to make a maximum entropy assumption [143, 82, 13, 141, 142] leading to postulate that the scattering matrix is uniformly distributed over the unitary group (for perfect couplings at the contacts). Although this approach has led to many successes in the description of several properties of coherent conductors (conductance, shot noise,...) [13, 142], it does not provide any information about the *energy dependence* of the scattering matrix, which is probed by the Wigner-Smith matrix (17). Two approaches were proposed to develop a random matrix description for the energy dependence of the scattering matrix :

- (i) the “Hamiltonian Approach” (HA) of chaotic scattering pushed forward by Fyodorov, Savin, Sommers and coworkers,
- (ii) the “Alternative Stochastic Approach” (ASA), pioneered by the work of Brouwer and Büttiker [30] and mostly developed by Brouwer, Beenakker and coworkers.

One of the first result within RMT was the calculation of the two point correlation function of the Wigner time delay [127] given below, Eq. (133). As we have seen in the previous section, one can introduce several time scales, characterised by their distributions :

- The marginal law for partial time delays

$$\tilde{w}_N^{(\beta)}(\tau) \stackrel{\text{def}}{=} (1/N) \sum_a \langle \delta(\tau - \tilde{\tau}_a) \rangle, \quad (80)$$

which was the first to be considered by Fyodorov and Sommers [87, 88] in the unitary class ($\beta = 2$), Eq. (94).¹⁵

- We also review the known results for the marginal law for the proper times

$$w_N^{(\beta)}(\tau) \stackrel{\text{def}}{=} (1/N) \sum_i \langle \delta(\tau - \tau_i) \rangle. \quad (81)$$

- The distribution of the Wigner time delay

$$\mathcal{P}_N^{(\beta)}(\tau) \stackrel{\text{def}}{=} \langle \delta(\tau - (1/N) \sum_i \tau_i) \rangle \quad (82)$$

is also of great interest.

Obviously, these distributions do not encode the full statistical information, which also require to characterise correlations (between partial/proper times, or energy correlations).

The known results corresponding to the standard (Wigner-Dyson) symmetry classes are reviewed in the next three sections. The final section reports few results obtained in other symmetry classes.

4.1. Distribution and moments for perfect contacts

In this subsection, I review the results for ideal contacts, i.e. when the transmission probability between a channel of the wave guide and the QD is equal to unity.

4.1.1. Proper time delays $\{\tau_i\}$

Gopar, Mello and Büttiker obtained the Wigner time delay distribution in the one channel case for the three symmetry classes [96] :

$$\mathcal{P}_1^{(\beta)}(\tau) = \frac{(\beta/2)^{\beta/2}}{\Gamma(\beta/2)} \tau^{-2-\beta/2} e^{-\beta/(2\tau)} \quad (83)$$

(obviously $\mathcal{P}_1^{(\beta)}(\tau) = w_1^{(\beta)}(\tau) = \tilde{w}_1^{(\beta)}(\tau)$). The proper times are measured in unit of the Heisenberg time $\tau_H = 2\pi\hbar/\Delta$, where Δ is the mean level spacing of the cavity (of more correctly the “mean resonance spacing” as we deal with a scattering problem).

¹⁵ Note that a similar route was followed in Ref. [209], however the result of the reference is partly incorrect as explained in [88].

An important advance was the work of Beenakker, Brouwer and Frahm who were able to obtain the joint distribution of the proper time delays within ASA in Ref. [31, 32]. They established the relationship with the Laguerre ensemble of random matrices : the inverse of the symmetrised Wigner-Smith matrix ¹⁶

$$\mathcal{Q}_s \stackrel{\text{def}}{=} -i\mathcal{S}^{-1/2} \frac{\partial \mathcal{S}}{\partial \varepsilon} \mathcal{S}^{-1/2} \quad (84)$$

is a Wishart matrix ; i.e. $\mathcal{Q}_s^{-1} = X^\dagger X$ where X is a $N \times (2N - 1 + 2/\beta)$ matrix with identical and independent Gaussian elements (this interpretation only holds for $\beta = 1, 2$). The joint distribution of the inverse of the proper times, $\gamma_i = \tau_H/\tau_i$ being given by [31, 32]

$$\mathcal{P}(\gamma_1, \dots, \gamma_N) \propto \prod_{i < j} |\gamma_i - \gamma_j|^\beta \prod_k \gamma_k^{\beta N/2} e^{-\beta \gamma_k/2}. \quad (85)$$

For $N = 1$, the Gamma-law $\mathcal{P}(\gamma_1) \propto \gamma_1^{\beta/2} e^{-\beta \gamma_1/2}$ corresponds to Gopar, Mello and Büttiker's result, Eq. (83).

In the large N limit, the marginal law for proper time delays takes the form

$$w_N^{(\beta)}(\tau) \underset{N \rightarrow \infty}{\simeq} N \rho(N\tau) \quad (86)$$

where [32, 197] ¹⁷

$$\rho(y) = \frac{1}{2\pi y^2} \sqrt{(x_+ - y)(y - x_-)} \quad (87)$$

with $x_\pm = (\sqrt{2} \pm 1)^2$. The marginal law is not strictly zero out of the interval $[x_-/N, x_+/N]$ but

¹⁶ The definition of \mathcal{Q}_s is intermediate between (17) and (18). The three matrices have the same spectrum of eigenvalues $\{\tau_a\}$.

¹⁷ The distribution is related to the Marčenko-Pastur law by $\rho(y) = (1/y^2) \rho_{\text{MP}}(1/y)$, where

$$\rho_{\text{MP}}(x) = [1/(2\pi x)] \sqrt{(x_+ - x)(x - x_-)}$$

is the distribution of the rescaled eigenvalues $\gamma = Nx$ for $N \times N$ matrices in the Laguerre ensemble, Eq. (85).

presents finite N corrections with asymptotics ¹⁸

$$w_N^{(\beta)}(\tau) \underset{\tau \rightarrow 0}{\sim} \tau^{-2-3\beta N/2} e^{-\beta/(2\tau)} \quad (88)$$

$$\underset{\tau \rightarrow \infty}{\sim} \tau^{-2-\beta N/2}. \quad (89)$$




Figure 4: Marginal law for the proper time delay.

We can deduce the moments :

$$\langle \tau_i \rangle = \frac{1}{N} \quad \text{and} \quad \text{Var}(\tau_i) \simeq \frac{1}{N^2} \quad \forall \beta. \quad (90)$$

The approach of Ref. [120] provides the precise expression of the variance [174] :

$$\text{Var}(\tau_i) = \frac{N[\beta(N-1)+2]+2}{N^2(N+1)(\beta N-2)}. \quad (91)$$

Because the moments of the sum of partial times are known [see below, Eq. (100)], using the symmetry between channels, it is possible to deduce the covariance. Writing $\text{Var}(N\tau_W) = N \text{Var}(\tau_i) + N(N-1) \text{Cov}(\tau_i, \tau_j)$ and using (91) provides the covariance

$$\text{Cov}(\tau_i, \tau_j) = -\frac{1}{N^2(N+1)}, \quad (92)$$

showing that the proper times are surprisingly independent of the symmetry index, and *anti-correlated*. The anti-correlations are not a surprise as the proper

¹⁸ In the general case, the Laguerre ensemble describes $N \times N$ random Hermitian matrices Γ with positive eigenvalues distributed according to $P(\Gamma) \propto (\det \Gamma)^{\alpha\beta N/2} \exp[-(\beta/2) \text{Tr} \{\Gamma\}]$ (Eq. (85) corresponds to $\alpha = 1$ and (77) to $\alpha = 0$). The Marčenko-Pastur law $p_N(\gamma) \simeq (1/N) \rho_{\text{MP}}(\gamma/N)$ for the density of eigenvalues has support $[Nx_-, Nx_+]$ where $x_\pm = (\sqrt{1+\alpha} \pm 1)^2$. Finite N corrections to the Marčenko-Pastur law for $\gamma \in [0, Nx_-] \cup [Nx_+, \infty[$ were obtained by Forrester in Ref. [79]. Asymptotic expansions of Forrester's result are $p_N(\gamma) \sim \gamma^{\alpha\beta N/2}$ for $\gamma \rightarrow 0$ and $p_N(\gamma) \sim \gamma^{(\alpha+2)\beta N/2} e^{-(\beta/2)\gamma}$ for $\gamma \rightarrow \infty$.

times are the eigenvalues of an invariant random matrix, see Eq. (85). It is interesting to rewrite the result as

$$\frac{\text{Cov}(\tau_i, \tau_j)}{\sqrt{\text{Var}(\tau_i) \text{Var}(\tau_j)}} = -\frac{\beta N - 2}{N[\beta(N-1) + 2] + 2} \simeq -\frac{1}{N}, \quad (93)$$

which show that they present the maximal anti-correlation (for identical variables).

Remark :

- The joint moments of the proper times were studied in [138].

4.1.2. Partial time delays $\{\tilde{\tau}_a\}$

The marginal law for partial times was obtained for $\beta = 2$ in [87, 88] and for $\beta = 1$ in [86]. The distribution $\forall \beta$ is given in Ref. [175] :

$$\tilde{w}_N^{(\beta)}(\tau) = \frac{(\beta/2)^{\beta N/2}}{N \Gamma(\beta N/2)} \tau^{-2-\beta N/2} e^{-\beta/(2\tau)}. \quad (94)$$

The distribution matches with (83) for $N = 1$, as it should. We deduce the moments $\langle \tilde{\tau}_a^n \rangle = (\beta/2)^n \Gamma(1-n+\beta N/2)/\Gamma(1+\beta N/2)$, i.e.

$$\langle \tilde{\tau}_a \rangle = \frac{1}{N} \quad \text{and} \quad \text{Var}(\tilde{\tau}_a) = \frac{2}{N^2(\beta N - 2)} \quad (95)$$

Thus the fluctuations of the partial time delays are reduced compared to those of the proper time delays : $\text{Var}(\tilde{\tau}_a) \simeq 2 \text{Var}(\tau_i)/(\beta N)$.

Using the same remark as for proper times, we can also deduce the covariances of the partial times :

$$\text{Cov}(\tilde{\tau}_a, \tilde{\tau}_b) = \frac{2}{N^2(N+1)(N\beta-2)}. \quad (96)$$

Hence, contrary to the proper times which presents anti-correlations, the partial times present positive correlations. We can rewrite the result as

$$\frac{\text{Cov}(\tilde{\tau}_a, \tilde{\tau}_b)}{\sqrt{\text{Var}(\tilde{\tau}_a) \text{Var}(\tilde{\tau}_b)}} = \frac{1}{N+1} \simeq +\frac{1}{N}. \quad (97)$$

Note that the joint probability distribution for two partial time delays was obtained in Ref. [175] (Eq. 23).

4.1.3. Wigner time delay τ_W

Despite the joint distribution for the proper times was explicitly known [31], the statistical properties of their sum, the Wigner time delay $\tau_W = (1/N) \sum_i \tau_i = (1/N) \sum_a \tilde{\tau}_a$, remained unknown for

a while. The distribution for $N = 2$ channels was deduced in Ref. [175] :

$$\mathcal{P}_2^{(\beta)}(\tau) = \frac{\beta^{3\beta+2} \Gamma(3(\beta+1)/2)}{\Gamma(\beta+1) \Gamma(3\beta+2)} \times \tau^{-3(\beta+1)} U\left(\frac{\beta+1}{2}, 2(\beta+1); \beta/\tau\right) e^{-\beta/\tau}, \quad (98)$$

where $U(a, b; z)$ is a Kummer function (confluent hypergeometric function) [2]. A method providing a systematic determination of the cumulants was proposed by Mezzadri and Simm in Ref. [146]. The authors gave explicitly the first four cumulants

$$\langle \tau_W \rangle = \frac{\tau_H}{N} \quad (99)$$

$$\langle \tau_W^2 \rangle_c = \frac{4 \tau_H^2}{N^2(N+1)(N\beta-2)} \quad (100)$$

$$\langle \tau_W^3 \rangle_c = \frac{96 \tau_H^3}{N^3(N+2)(N+1)(N\beta-2)(N\beta-4)} \quad (101)$$

$$\langle \tau_W^4 \rangle_c = \begin{cases} \frac{96(53N^2-68N-156)\tau_H^4}{N^4(N+3)(N+2)(N+1)^2(N-2)^2(N-4)(N-6)} & \text{for } \beta = 1 \\ \frac{12(53N^2-77)\tau_H^4}{N^4(N+3)(N+2)(N+1)^2(N-1)^2(N-2)(N-3)} & \text{for } \beta = 2 \\ \frac{12(53N^2+34N-39)\tau_H^4}{N^4(N+3)(N+2)(N+1)^2(2N-1)^2(N-1)(2N-3)} & \text{for } \beta = 4 \end{cases} \quad (102)$$

It is worth emphasizing the different scalings of the variances with N :

$$\text{Var}(N \tau_W) = \text{Var}\left(\sum_i \tau_i\right) \sim \text{Var}(\tau_i) \quad (103)$$

$$= \text{Var}\left(\sum_a \tilde{\tau}_a\right) \simeq 2N \text{Var}(\tilde{\tau}_a). \quad (104)$$

A systematic expression of the moments $\langle \tau_W^n \rangle$ of the Wigner time delay for $\beta = 2$ was given more recently by Novaes [152]. We see that the variance diverges for $N\beta \leq 2$, and the third cumulant for $N\beta \leq 4$. Writing the denominator of the fourth cumulant as $N^4(N+3)(N+2)(N+1)^2(N-2/\beta)^2(N-4/\beta)(N-6/\beta)$, we see that $\langle \tau_W^4 \rangle_c$ diverges for $N\beta \leq 6$. These observations suggest the power law tail $\mathcal{P}_N^{(\beta)}(\tau) \sim \tau^{-2-\beta N/2}$, conjectured in Ref. [88] for $\beta = 2$ on the basis of some heuristic argument involving resonances [i.e. identifying the tail of $\mathcal{P}_N^{(\beta)}(\tau)$ with the one of $\tilde{w}_N^{(\beta)}(\tau)$, Eq. (94)]. Using a Egdeworth expansion, Mezzadri and Simm




Figure 5: Wigner time delay distribution in the limit $N \gg 1$: sketch of the rescaled distribution $P_N(s) = \tau_d \mathcal{P}_N^{(\beta)}(\tau = s\tau_d)$, where $\tau_d = h/(N\Delta)$ is the dwell time. The small curves are sketches of the optimal distributions of eigenvalues of \mathcal{Q}^{-1} with the constraint that $\text{Tr}\{\mathcal{Q}\}$ is fixed. The transition between the sharp Gaussian peak and the power law tail is associated with a phase transition in the density of eigenvalues. Figure from [197].

concluded that the distribution weakly converges towards a Gaussian form as N grows. The full distribution of the Wigner time delay for $N \gg 1$ was however shown to present a richer structure in Ref. [197] where the large deviations were studied in detail, leading to the behaviours

$$\mathcal{P}_N^{(\beta)}(\tau) \underset{N \rightarrow \infty}{\sim} \begin{cases} \tau^{-3\beta N^2/4} e^{-\beta N/(2\tau)} & \text{for } \tau \rightarrow 0 \\ \exp\left\{-\frac{\beta N^2}{8}(N\tau - 1)^2\right\} & \text{for } \tau \sim 1/N \\ (N\tau - 1)^{-2-\beta N/2} & \text{for } N\tau - 1 \gg \sqrt{\frac{2}{N} \ln N} \end{cases} \quad (105)$$

The distribution is sketched in Fig. 5. In particular, the transition between the sharp Gaussian peak and the power law tail was shown to be related to a phase transition in the underlying Coulomb gas [197].

In Fig. 6, we compare the different distributions, for proper, partial and Wigner time delays.

Few remarks :

- (i) A systematic analysis of the “moments” $\langle \text{Tr}\{\mathcal{Q}^n\} \rangle$ within RMT was carried out by several authors : [144, 145, 62, 152, 64].
- (ii) The determination of the variance (189) and the covariance (190) in Ref. [102, 97] (and also [207]) has involved the correlation between the two linear statistics $\text{Tr}\{\mathcal{Q}\}$ and




Figure 6: Sketch of the distributions for perfect contacts. Figure from Ref. [101].

$\text{Tr}\{\mathcal{Q}^2\}$. A systematic analysis of the covariance $\text{Cov}(\text{Tr}\{\mathcal{Q}^n\}, \text{Tr}\{\mathcal{Q}^m\})$ was carried out by Cunden recently [62], based on the recent work [65] where a general formula for the covariance of two linear statistics was derived with the Coulomb gas method. More recently, the authors were also able to further analyse the correlations $\langle \text{Tr}\{\mathcal{Q}^{n_1}\} \text{Tr}\{\mathcal{Q}^{n_2}\} \dots \rangle_c$ in Ref. [63].

- (iii) The distribution of the truncated sum $\sum_{a=1}^K \tau_a$ of proper times, with $K < N$, was studied in Ref. [100].

The case where the sum is further restricted to the largest (or smallest) times can be analysed along the lines of Ref. [99, 97] (such an analysis allows to quantify the contribution of the fraction of eigenchannels contributing the most to the DoS).

- (iv) The distribution of the trace of a diagonal subblock of the matrix \mathcal{Q} was studied in an appendix of Ref. [100] in the unitary case, based on the result of [175]. The trace of the diagonal subblock $\sum_{a=1}^K \mathcal{Q}_{aa}$ is more physical than $\sum_{a=1}^K \tau_a$, as it corresponds with the “injectance” discussed in Section 5. The different scaling of the two quantities with N was emphasized : although they are characterized by the same average $\langle \sum_{a=1}^K \mathcal{Q}_{aa} \rangle = \langle \sum_{a=1}^K \tau_a \rangle = \kappa$, where $\kappa = K/N$, the variances scale differently with the channel num-

ber [100] (here for $\beta = 2$) :

$$\text{Var} \left(\sum_{a=1}^K \mathcal{Q}_{aa} \right) = \frac{\kappa(1+\kappa)}{N^2-1} \quad (106)$$

$$\text{Var} \left(\sum_{a=1}^K \tau_a \right) = \frac{\kappa(1-\kappa)}{N+1} + \frac{2\kappa}{N^2-1}. \quad (107)$$

For $K = N$, the two variances coincide with (100), as it should.

4.2. Distributions and moments for non-ideal contacts

Less is known about the case of non-ideal contacts, when the QD is connected to the waveguides through tunnel barriers (Fig. 7). The distribution (85) of Brouwer, Frahm and Beenakker (BFB) was recently extended in Ref. [101]. The brief review of this Subsection 4.2 mostly borrows from Ref. [101].

4.2.1. Wigner-Smith time delay matrix

The derivation is based on the idea that the scattering matrix \mathcal{S} for non-ideal contacts can be related to the scattering matrix \mathcal{S}_0 for perfect contacts

$$\mathcal{S} = r_b + t'_b \left(\mathcal{S}_0^\dagger - r'_b \right)^{-1} t_b, \quad (108)$$

where r_b , r'_b , t_b and t'_b are reflection and transmission matrices describing the tunneling barrier, here supposed fixed (non random). This is illustrated in Fig. 7 for one contact (obviously the approach is not restricted to a one contact situation).




Figure 7: Description of the model for non-ideal coupling. Figure from Ref. [101].

Assuming that the matrix \mathcal{S}_0 is uniformly distributed over the unitary group, we deduce that the coupling between the QD and the external can be encoded in the averaged (“optical”) scattering matrix $\langle \mathcal{S} \rangle = r_b$. In the following, we consider the case where all couplings are equal for simplicity, $\langle \mathcal{S} \rangle = \bar{\mathcal{S}} \mathbf{1}_N$ (the case of perfect coupling discussed in Subsection 4.1 corresponds to $\bar{\mathcal{S}} = 0$).

More precisely, we choose $r_b = -(r'_b)^\dagger = \mathbf{1}_N \bar{\mathcal{S}}$ and $t_b = t'_b = \mathbf{1}_N \sqrt{1 - |\bar{\mathcal{S}}|^2}$. This picture makes easy to recover the Poisson kernel describing the statistical properties of the \mathcal{S} -matrix for non ideal contacts [143, 28, 142]

$$D\mathcal{S} P_{\mathcal{S}}(\mathcal{S}) \propto D\mathcal{S} |\det(\mathbf{1}_N - \bar{\mathcal{S}}^* \mathcal{S})|^{-2-\beta(N-1)}. \quad (109)$$

where $D\mathcal{S}$ is the uniform measure over the circular ensemble (which corresponds with the Haar measure in the unitary case) ; since the distribution of \mathcal{S}_0 is uniform, Eq. (109) simply corresponds with the Jacobian of the transformation (108), which now takes the form $\mathcal{S} = (\bar{\mathcal{S}} \mathbf{1}_N + \mathcal{S}_0) (\mathbf{1}_N + \bar{\mathcal{S}}^* \mathcal{S}_0)^{-1}$ [101].

Introducing the notation $\Gamma = \mathcal{Q}_s^{-1}$ for the inverse of the symmetrised Wigner-Smith matrix $\mathcal{Q}_s = -i\mathcal{S}^{-1/2} \partial_\epsilon \mathcal{S} \mathcal{S}^{-1/2}$, the distribution of eigenvalues (85) for perfect contacts (superscript “(0)”) can be related to the matrix distribution

$$P_\Gamma^{(0)}(\Gamma) \propto (\det \Gamma)^{\beta N/2} \exp\{-(\beta/2) \text{Tr} \{\Gamma\}\}. \quad (110)$$

A key result of Refs. [31, 32] is the statistical independence of the scattering matrix \mathcal{S}_0 and the symmetrised Wigner-Smith matrix \mathcal{Q}_{0s} . Making use of (i) the statistical independence of \mathcal{S}_0 and \mathcal{Q}_{0s} ,

(ii) the uniform distribution of \mathcal{S}_0 ,

(iii) BFB’s distribution (110), and

(iv) assuming that the barrier coefficients are independent of the energy,

we have obtained in Ref. [101] the joint distribution

$$\begin{aligned} D\mathcal{S} D\Gamma P_{\mathcal{S},\Gamma}(\mathcal{S},\Gamma) &= D\mathcal{S} D\Gamma c_{N,\beta} \Theta(\Gamma) \quad (111) \\ &\times \left| \det(\mathbf{1}_N - \bar{\mathcal{S}}^* \mathcal{S}) \right|^{\beta N} (\det \Gamma)^{\beta N/2} \\ &\times \exp \left[-\beta \frac{\text{Tr} \left\{ (\mathbf{1}_N - \bar{\mathcal{S}}^* \mathcal{S})(\mathbf{1}_N - \bar{\mathcal{S}} \mathcal{S}^\dagger) \Gamma \right\}}{2(1 - |\bar{\mathcal{S}}|^2)} \right], \end{aligned}$$

where $c_{N,\beta}$ is a normalisation constant. $D\Gamma$ is the Lebesgue measure over the set of Hermitian matrices. $\Theta(\Gamma) = 1$ when all eigenvalues of Γ are positive and zero otherwise. The representation (111) may be regarded as an extension of the Poisson kernel (109). For $\bar{\mathcal{S}} = 0$, we recover Eq. (110). Note that a representation similar to (111) was given in Ref. [136], with extensions to the BdG symmetry classes describing scattering in an Andreev billiard.

The marginal distribution of the Wigner-Smith matrix can be obtained from (111) by a matrix integration over \mathcal{S} . We have preferred a more convenient

form in terms of a matrix integral over Hermitian matrices \mathcal{K} [101]

$$P_{\Gamma}(\Gamma) = b_{N,\beta} \Theta(\Gamma) (\det \Gamma)^{\beta N/2} \quad (112)$$

$$\times \int \mathrm{D}\mathcal{K} \frac{\det(\mathbf{1}_N + \mathcal{K}^2)^{\beta N/2}}{\det(\mathbf{1}_N + \kappa^2 \mathcal{K}^2)^{1-\frac{\beta}{2}+\beta N}} \times \exp \left[-\frac{\beta}{2} \kappa \operatorname{Tr} \left\{ \frac{\mathbf{1}_N + \mathcal{K}^2}{\mathbf{1}_N + \kappa^2 \mathcal{K}^2} \Gamma \right\} \right]$$

where $b_{N,\beta}$ is a normalisation constant and the coupling constant $\kappa > 0$ is related to the transmission probability \mathcal{T} by

$$\mathcal{T} \equiv 1 - |\bar{\mathcal{S}}|^2 = \frac{4\kappa}{(1+\kappa)^2}. \quad (113)$$

For perfect contacts ($\mathcal{T} = \kappa = 1$), we recover the Laguerre distribution (110), as it should. It is interesting to point that the Hermitian matrix \mathcal{K} in the matrix integral in Eq. (112) corresponds to the Wigner's reaction matrix characterizing the QD for perfect contacts. Hence (112) can be deduced by integrating (111) over \mathcal{S} and eventually performing the two changes of variable $\mathcal{S} = (\bar{\mathcal{S}} \mathbf{1}_N + \mathcal{S}_0) (\mathbf{1}_N + \bar{\mathcal{S}}^* \mathcal{S}_0)^{-1}$ and $\mathcal{S}_0 = (\mathbf{1}_N - i\mathcal{K}) (\mathbf{1}_N + i\mathcal{K})^{-1}$. Note that the uniform distribution of \mathcal{S}_0 corresponds to the Cauchy distribution

$$P_{\mathcal{K}}^{(0)}(\mathcal{K}) \propto [\det(\mathbf{1}_N + \mathcal{K}^2)]^{-1-\beta(N-1)/2}, \quad (114)$$

defined over the set of Hermitian matrices. We can indeed check that, removing the matrix integral in (112), we recover $P_{\Gamma}^{(0)}(\Gamma) P_{\mathcal{K}}^{(0)}(\mathcal{K})$ for $\kappa = 1$.

Remarks :

- (i) Another integral representation, equivalent to (112), was derived in Ref. [101] in terms of a matrix integral over the unitary group, which turned out to be more convenient for numerical simulations.
- (ii) The case of unequal couplings can also be analysed along the same lines : the diagonal matrix $\langle \mathcal{S} \rangle$ is not proportional to the identity matrix, and a set of N transmission coefficients $\mathcal{T}_1, \dots, \mathcal{T}_N$ (or κ_a 's) should be introduced. The form (112) was extended in Ref. [101].

4.2.2. Proper time delays

The joint distribution for the proper time delays $\tau_a = 1/\gamma_a$ can in principle be deduced from (112) by integrating over all degrees of freedom related

to the eigenvectors of the matrix. This was done in Ref. [101] in the unitary case by using Harish-Chandra-Itzykson-Zuber integral :

$$\mathcal{P}(\gamma_1, \dots, \gamma_N) \propto \Delta_N(\gamma) \prod_n \theta_H(\gamma_n) \gamma_n^N$$

$$\times \int_{\mathbb{R}^N} dk_1 \dots dk_N \frac{\Delta_N(k)^2}{\Delta_N(k^2)} \prod_n \frac{(1+k_n^2)^N}{(1+\kappa^2 k_n^2)^{N+1}}$$

$$\times \det \left[\exp \left(-\kappa \frac{1+k_i^2}{1+\kappa^2 k_i^2} \gamma_j \right) \right] \quad (115)$$

where $\Delta_N(\gamma) = \prod_{i < j} (\gamma_i - \gamma_j)$ denotes the Vandermonde determinant (here, I have slightly simplified the form given in [101]). This form reduces to (85) for $\kappa \rightarrow 1$, as we now check. In the limit $\kappa \rightarrow 1^-$ we have $\kappa \simeq 1 - 2\bar{\mathcal{S}}$. The limit of the last determinant of the integral can be analysed by using the property

$$\det(e^{\epsilon x_i y_j})_{i,j=1}^N \underset{\epsilon \rightarrow 0}{\simeq} \frac{\epsilon^{N(N-1)/2}}{G(N+1)} \Delta_N(x) \Delta_N(y) \quad (116)$$

where $G(N+1) = \prod_{n=1}^{N-1} n!$ is the Barnes- G function, we obtain

$$\det \left[\exp \left(-\kappa \frac{1+k_i^2}{1+\kappa^2 k_i^2} \gamma_j \right) \right] \quad (117)$$

$$\underset{\kappa \rightarrow 1}{\simeq} \frac{(4\bar{\mathcal{S}})^{N(N-1)/2}}{G(N+1)} \frac{\prod_n e^{-\gamma_n}}{\prod_n (1+k_n^2)^{N-1}} \Delta_N(\gamma) \Delta_N(k^2)$$

Finally, we deduce that the limit $\kappa \rightarrow 1$ of (115) is

$$\mathcal{P}(\gamma_1, \dots, \gamma_N) \propto \Delta_N(\gamma)^2 \prod_n \theta_H(\gamma_n) \gamma_n^N e^{-\gamma_n}$$

$$\times \int_{\mathbb{R}^N} dk_1 \dots dk_N \Delta_N(k)^2 \prod_n (1+k_n^2)^{-N} \quad (118)$$

where the multiple integral is a constant corresponding to the normalisation of the Cauchy ensemble (114). Eq. (118) is precisely the BFB result (85) for $\beta = 2$, as it should.

Although complicate, the form (115) was used in order to derive the distribution of the Wigner time delay [101]. Extracting the marginal distribution of the proper time or correlations seems however difficult. We now review some results obtained earlier by other means. The marginal distribution of the proper time delays was obtained in the unitary case by Sommers, Savin and Sokolov [181] : the result was given under a rather complicate form involving a sum of multiple derivative of Bessel function.

Here, we only give the limiting behaviours of the distribution for weak coupling [181, 101] :

$$\frac{\mathcal{T}}{4} w_N^{(2)} \left(\tau \simeq \frac{\mathcal{T}}{4} t \right) \underset{\mathcal{T} \rightarrow 0}{\sim} \begin{cases} c_N t^{-2N-1/2} e^{-1/t} & \text{for } t \lesssim 1/N \\ b_N t^{-3/2} & \text{for } 1/N \lesssim t \lesssim \frac{1}{N\mathcal{T}^2} \\ \frac{a_N \mathcal{T}^3}{8} \left(\frac{4}{\mathcal{T}^2 t} \right)^{N+2} & \text{for } t \gtrsim \frac{1}{N\mathcal{T}^2} \end{cases} \quad (119)$$

The two coefficients a_N and b_N are given below, Eqs. (124,125), and the last coefficient is

$$c_N = \frac{2^{2(N-1)}}{\sqrt{\pi} N(2N-1)!} . \quad (120)$$

Thus $w_N^{(\beta)}(\tau)$ and $\tilde{w}_N^{(\beta)}(\tau)$ coincide in the two regimes above the scale $\tau_{\text{lower}} \sim \mathcal{T}/N$ (Fig. 8). The form of the distribution shows that the universal power law $w_N^{(\beta)}(\tau) \sim \tau^{-3/2}$ (which holds for any β) is cut off at the scale $\tau_{\text{upper}} \simeq 8e/(N\mathcal{T})$. This cutoff controls the positive moments

$$\langle \tau_a^k \rangle \sim b_N \sqrt{\mathcal{T}} \tau_{\text{upper}}^{k-1/2} \quad (121)$$

for $1 \leq k < 1 + \beta N/2$ (and $\langle \tilde{\tau}_a^k \rangle = \langle \tau_a^k \rangle = \infty$ for $k \geq 1 + \beta N/2$).




Figure 8: Sketch of the distributions for perfect contacts ; compare with Fig. 6. Figure from Ref. [101].

4.2.3. Partial time delays

The marginal distribution of the partial time delays is known exactly for any symmetry class

[88, 86, 175]. A convenient integral representation was derived in Appendix C of [101] by extending the method of Ref. [95] to arbitrary number of channels. Here, we only give the asymptotic behaviours

$$\frac{\mathcal{T}}{4} \tilde{w}_N^{(\beta)} \left(\tau \simeq \frac{\mathcal{T}}{4} t \right) \underset{\mathcal{T} \rightarrow 0}{\sim} \begin{cases} \tilde{c}_N t^{-\beta N/2-3/2} e^{-1/t} & \text{for } t \lesssim 1/N \\ b_N t^{-3/2} & \text{for } 1/N \lesssim t \lesssim \frac{1}{N\mathcal{T}^2} \\ \frac{a_N \mathcal{T}^3}{8} \left(\frac{4}{\mathcal{T}^2 t} \right)^{2+\beta N/2} & \text{for } t \gtrsim \frac{1}{N\mathcal{T}^2} \end{cases} \quad (122)$$

involving the three coefficients

$$\tilde{c}_N = \frac{1}{\sqrt{\pi} \Gamma(1 + \beta N/2)} , \quad (123)$$

$$b_N = \frac{1}{\pi} \frac{\Gamma(1/2 + \beta N/2)}{\Gamma(1 + \beta N/2)} , \quad (124)$$

$$a_N = \frac{2^{1+\beta N}}{\sqrt{\pi}} \frac{\Gamma(1/2 + \beta N/2)}{\Gamma(1 + \beta N/2)^2} . \quad (125)$$

In Ref. [101], we have also stressed that the weak coupling limit ($\mathcal{T} \rightarrow 0$) of the marginal distribution takes the simple form

$$\lim_{\mathcal{T} \rightarrow 0} \frac{\mathcal{T}}{4} \tilde{w}_N^{(\beta)} \left(\tau \simeq \frac{\mathcal{T}}{4} t \right) = \frac{1}{\sqrt{\pi} \Gamma(1 + \beta N/2)} \frac{e^{-1/t}}{t^{2+\beta N/2}} U \left(\frac{1}{2}, \frac{\beta N + 3}{2}, \frac{1}{t} \right) , \quad (126)$$

in terms of the Kummer function $U(a, c, z)$ [2]. It is quite remarkable to obtain such a universal form in this limit.

The variance of the partial times are exactly known for $\beta = 2$ [88] :

$$\text{var}(\tilde{\tau}_a) = \frac{2N(\mathcal{T}^{-1} - 1) + 1}{N^2(N-1)} \underset{\mathcal{T} \ll 1}{\sim} \frac{2}{\mathcal{T} N^2} . \quad (127)$$

By combining (127) with the variance of the Wigner time delay given below, Eq. (130), we get the co-variance :

$$\text{cov}(\tilde{\tau}_a, \tilde{\tau}_b) = \frac{1}{\mathcal{T}^2 N(N-1)^2} \quad (128)$$

$$\underset{\mathcal{T} \ll 1}{\sim} -\frac{1}{N^2} \times \begin{cases} \frac{2}{\mathcal{T} N} & \text{for } N\mathcal{T} \gg 1 \\ 1 & \text{for } N\mathcal{T} \ll 1 \end{cases} \quad (129)$$

Contrary to the ideal contact case, where the correlations are always positive, the covariances change both in sign and scaling with N as transmission crosses over from perfect to weak coupling. At this point it is important to stress that the *weak transmission* $\mathcal{T} \ll 1$ regime and *weak coupling* $N\mathcal{T} \ll 1$ regime should not be confused.

4.2.4. Wigner time delay

The statistical properties of the Wigner time delay $\tau_W = \text{Tr} \{ \Gamma^{-1} \}$ is in principle encoded in the matrix distribution (112), however explicit calculations from the matrix integral seems a difficult task. We rather collect the known results obtained by other means. The variance of the Wigner time delay for non ideal coupling was obtained in the unitary ($\beta = 2$) case [87, 88] :

$$\text{var}(\tau_W) = \frac{2[1 - (1 - \mathcal{T})^{N+1}]}{\mathcal{T}^2 N^2 (N^2 - 1)}. \quad (130)$$

In the limit of weak transmission per channel $\mathcal{T} \ll 1$, we get from (130) two different possible behaviours, which depend on the product $N\mathcal{T}$ describing the degree of the resonance overlap (thus controlling the overall coupling to the leads) :

$$\frac{\text{var}(\tau_W)}{\langle \tau_W \rangle^2} \underset{\tau \ll 1}{\simeq} \begin{cases} \frac{4}{\beta(N\mathcal{T})^2} \ll 1 & \text{for } N\mathcal{T} \gg 1 \\ \frac{2}{N\mathcal{T}} \gg 1 & \text{for } N\mathcal{T} \ll 1 \end{cases}. \quad (131)$$

Here we have reintroduced β in order to match with the known $\beta = 1$ result [127].

Starting from (115), the distribution of the Wigner time delay was obtained in [101] in the weak coupling regime $N\mathcal{T} \ll 1$, by a careful analysis of the characteristic function :

$$\begin{aligned} \mathcal{P}_N^{(\beta)}(\tau) & \quad (132) \\ \sim & \begin{cases} \tau^{-\frac{\beta N^2}{2} - \frac{3}{2}} e^{-\beta N\mathcal{T}/(8\tau)} & \text{for } \tau \ll \mathcal{T} \\ \frac{1}{\mathcal{T}} (\mathcal{T}/\tau)^{3/2} & \text{for } \mathcal{T} \ll \tau \ll \frac{1}{N^2 \mathcal{T}} \\ \mathcal{T}^2 N^3 (\mathcal{T} N^2 \tau)^{-2 - \beta N/2} & \text{for } \tau \gg \frac{1}{N^2 \mathcal{T}} \end{cases} \end{aligned}$$

These behaviours were verified by numerical calculations. Some heuristic analysis was given in Ref. [101], relating the distribution of the Wigner time delay and that of the resonance widths. The power law $\mathcal{P}_N^{(\beta)}(\tau) \sim \tau^{-3/2}$ was then related to typical resonance with, while the power law $\mathcal{P}_N^{(\beta)}(\tau) \sim \tau^{-2 - \beta N/2}$ is due to atypically narrow resonances.

4.3. Energy correlations for perfect coupling

The knowledge of the correlation function $\langle \tau_W(\varepsilon) \tau_W(\varepsilon') \rangle_c = \langle \tau_W(\varepsilon) \tau_W(\varepsilon') \rangle - \langle \tau_W(\varepsilon) \rangle \langle \tau_W(\varepsilon') \rangle$ is of importance and has practical applications : for example, this information was used by Polianski and Büttiker in order to study the effect of thermal fluctuations on the non-linear conductance of a quantum dot [163]. The correlation function was obtained by Lehmann *et al.* [127] by a random matrix analysis in the orthogonal case ($\beta = 1$), within the HA of random matrices (see also [88]) :

$$\frac{\langle \tau_W(\varepsilon) \tau_W(\varepsilon') \rangle_c}{\langle \tau_W \rangle^2} \simeq \frac{4}{N^2} \frac{1 - (\omega \tau_d)^2}{[1 + (\omega \tau_d)^2]^2}, \quad (133)$$

where $\omega = \varepsilon - \varepsilon'$. For $\varepsilon = \varepsilon'$ we recover the cumulant (100) at leading order $1/N^4$. Eq. (133) shows that correlations occur on the scale $1/\tau_d \sim N\Delta$ much larger than the mean resonance spacing.¹⁹ We have introduced

$$\tau_d = \langle \tau_W \rangle = \frac{2\pi}{N\Delta} = \tau_H/N, \quad (134)$$

the dwell time for an electron in the cavity (Eq. (133) is valid for strongly overlapping resonances, $1/\tau_d \gg \Delta$, i.e. $N \gg 1$). The result (133) was later generalised [86] in order to include parametric correlations and describe the crossover between orthogonal and unitary cases.

Time delay correlations can also be determined within the stochastic approach : the model described in Refs. [5, 161, 33] and used by Polianski and Büttiker [162, 164, 163] provides some information about the energy and magnetic field dependence of the \mathcal{S} -matrix correlator, which reads [163]

$$\langle \mathcal{S}_{ab}(\varepsilon, \mathcal{B}) \mathcal{S}_{cd}^*(\varepsilon', \mathcal{B}') \rangle = \delta_{ac} \delta_{bd} \mathcal{D}_{\varepsilon - \varepsilon'} + \delta_{ad} \delta_{bc} \mathcal{C}_{\varepsilon - \varepsilon'}. \quad (135)$$

\mathcal{D}_ω and \mathcal{C}_ω are the (zero-dimensional) analogues of the Diffuson and the Cooperon appearing in the diagrammatic approach for weakly disordered metals [4],²⁰ given by²¹

$$\left. \begin{aligned} \mathcal{D}_\omega \\ \mathcal{C}_\omega \end{aligned} \right\} = \frac{1}{N\tau_d} \frac{1}{1/\tau_{\mathcal{D},\mathcal{C}} - i\omega}. \quad (136)$$

¹⁹ In a ballistic QD of size L , the Thouless energy, which sets the energy scale for correlations in a closed QD, is $\varepsilon_{\text{Th}} = v_F/L$, where v_F is the Fermi velocity. The open QD is characterised by the number of channels $N = k_F w/\pi$, controlled by the width w of the contacts. In this case, as we have seen, the relevant scale is a different Thouless energy [13] $\varepsilon_{\text{Th}}^{\text{open}} = 1/\tau_d$. We have $\varepsilon_{\text{Th}}^{\text{open}}/\varepsilon_{\text{Th}} \sim w/L \ll 1$.

²⁰ This can be understood from the Fisher and Lee relation [78] between scattering matrix and Green's function.

²¹ Note that (135) only satisfies unitarity at leading order

The two characteristic times

$$\frac{1}{\tau_{\mathcal{D}, \mathcal{C}}} = \frac{1}{\tau_d} + \frac{1}{\tau_{(\mathcal{B} \mp \mathcal{B}')/2}} \quad (137)$$

combines a contribution describing the escape rate $1/\tau_d$ from the cavity and dephasing due to the magnetic field. The magnetic dephasing rate is $1/\tau_{\mathcal{B}} = (v_F \ell / \text{Surf})(\Phi / \phi_0)^2$ where v_F is the Fermi velocity, $\Phi = \text{Surf } \mathcal{B}$ the magnetic flux through the cavity and $\phi_0 = h/e$ the quantum flux. ℓ is the size of the cavity in the ballistic case or the elastic mean free path in the weakly disordered (diffusive) case.

As a simple application of the formula we get $\langle \tau_W(\varepsilon) \rangle \simeq \tau_d(N^2 \mathcal{D}_0^2 + N \mathcal{C}_0^2)$, where we have used $-i\partial_\varepsilon \mathcal{D}_{\varepsilon-\varepsilon'} = \tau_H \mathcal{D}_{\varepsilon-\varepsilon'}^2$. We deduce $\langle \tau_W(\varepsilon) \rangle / \tau_d \simeq 1 + \mathcal{O}(1/N)$, whose leading order term coincides with (99).

The analysis of the correlations requires additional information : simply applying Wick's theorem with the correlator (135), we get $\langle \tau_W(\varepsilon, \mathcal{B}) \tau_W(\varepsilon', \mathcal{B}') \rangle_c^{(\text{Gaussian})} \simeq \tau_H^2 (|\mathcal{D}_{\varepsilon-\varepsilon'}|^4 + |\mathcal{C}_{\varepsilon-\varepsilon'}|^4)$, which leads, at zero magnetic field, to the incomplete expression $\langle \tau_W(\varepsilon, 0) \tau_W(\varepsilon', 0) \rangle_c^{(\text{Gaussian})} / \langle \tau_W \rangle^2 \simeq (2/N^2) [1 + (\omega \tau_d)^2]^{-2}$. This calculation, known as the “diagonal approximation” in the context of semiclassical methods, disagrees with (133) : not only the energy dependence differs but also the value at $\omega = 0$ is half the correct result [compare with the variance (100)]. This emphasizes the importance of *non Gaussian* contributions to the correlator, which are taken into account in the more precise expression of the four-point correlation function $\langle S_{ab} S_{cd}^* S_{ij} S_{kl}^* \rangle$ given in Appendix B of Ref. [33] (see also [29]).

This question has been much discussed in semiclassical approach : this discrepancy was underlined by Lewenkopf and Vallejos [130]. Kuipers and Sieber [122] identified the nature of the contributions (“trajectory quadruplets”) to be added to the diagonal approximation in order to recover the random matrix result (133) (doing so they have reconciled the two semiclassical approaches for a Wigner time delay analysis : the periodic orbit expansion based on the relation with the density of states [71], and the scattering approach involving trajectories

entering and leaving the system). The expression of the correlator, determined earlier within a semiclassical approach by Vallejos *et al.* [205] (for $\mathcal{B} = \mathcal{B}'$), is the sum of two contributions which can be identified as Diffuson and Cooperon contributions. Reintroducing the effect of the field mismatch, we get the structure obtained within the HA of random matrices in Ref. [86, 88] :

$$\frac{\langle \tau_W(\varepsilon, \mathcal{B}) \tau_W(\varepsilon', \mathcal{B}') \rangle_c}{\langle \tau_W \rangle^2} \quad (138)$$

$$\simeq \frac{2}{\tau_H^2} \left\{ \frac{1/\tau_{\mathcal{D}}^2 - \omega^2}{[1/\tau_{\mathcal{D}}^2 + \omega^2]^2} + \frac{1/\tau_{\mathcal{C}}^2 - \omega^2}{[1/\tau_{\mathcal{C}}^2 + \omega^2]^2} \right\},$$

where we recall that $\tau_H = 2\pi/\Delta$. This expression now agrees with (133) for $\tau_{\mathcal{D}} = \tau_{\mathcal{C}} = \tau_d = \tau_H/N$.

Few remarks :

- (i) The time delay may be represented in terms of as $\tau_W(\varepsilon) = (1/N) \sum_{\alpha} \Gamma_{\alpha} / [(\varepsilon - \varepsilon_{\alpha})^2 + \Gamma_{\alpha}^2/4]$, where the sum runs over the resonances [127, 88]. This establishes a connection between Wigner time delay fluctuations and Ericson fluctuations of the cross-section for strongly overlapping resonances $\Gamma \gg \Delta$ [73, 71].
- (ii) Semiclassical methods were widely used in order to analyse Wigner time delay correlations [71, 205, 130, 121, 122] ; correspondence between the random matrix and semiclassical approaches holds for strongly overlapping resonances $1/\tau_d \gg 1/\tau_H \sim \Delta$ (i.e. $N \gg 1$) [205].
- (iii) More recently, an improved semiclassical approach for the analysis of the Wigner time delay statistics was developed by Kuipers, Savin and Sieber [120], who carried out a diagrammatic calculation of the moments of the Wigner time delay. A semiclassical derivation of the moments of the time delay was also proposed by Novaes [153].

4.4. Other symmetry classes

As we have already mentioned, the three Wigner-Dyson symmetry classes (denoted AI, A and AII in the Altland-Zirnbauer classification) were completed by three chiral classes (chiral orthogonal BDI, chiral unitary AIII and chiral symplectic CII) and four Bogoliubov-de Gennes classes (C, CI, D, DIII) [214, 6, 75]. The new symmetry classes have attracted a lot of attention during the last years in relation with topological insulators [108, 167] and

in the orthogonal case : one gets $\langle (S^\dagger S)_{aa} - 1 \rangle = \mathcal{C}_0 = 1/N$ instead of zero. This can be corrected by using the expression [5] $\langle S_{ab}(\varepsilon) S_{cd}^*(\varepsilon') \rangle = [\delta_{ac} \delta_{bd} + (2/\beta - 1) \delta_{ad} \delta_{bc}] \mathcal{D}_{\varepsilon-\varepsilon'}$ with $\mathcal{D}_\omega = [N + 2/\beta - 1 - 2i\pi\omega/\Delta]^{-1}$.

topological superconductors [15, 16] (see also [172] where a classification of the various types of topological insulators was provided for the different symmetry classes and dimensions).

The analysis of the Wigner time delay distribution for different symmetry classes has been initiated by the work of Fyodorov and Ossipov [85] for the chiral-unitary class (AIII) in the $N = 1$ channel case :

$$\mathcal{P}_1^{(\text{AIII})}(\tau) = \tau^{-2} e^{-1/\tau} \quad (139)$$

in appropriate units. This should be compared with the expression (83) for the unitary case (class A) : $\mathcal{P}_1^{(2)}(\tau) = \tau^{-3} e^{-1/\tau}$. We see that the exponent of the tail is increased from 2 to 3 when the chiral symmetry is broken, as for the 1D disordered case reviewed above : compare (51) in the presence of the chiral symmetry and (37) when the chiral symmetry is broken.

More recently, the result (85) for the joint distribution of proper time delays was extended to the four BdG symmetry classes in Refs. [135, 176]. This question is reviewed in another article of this special issue [136]. The generalisation to the three chiral symmetry classes remains an open problem.

5. Generalised concepts : injectance, emittance, etc.

The concept of scattering matrix is central in the Landauer-Büttiker approach. For this reason, the Krein-Friedel relation (22) plays a very important role as it allows to express the density of states (DoS) of the *open* conductor in terms of scattering properties. In the following, we will simply write

$$\nu(\varepsilon) \simeq \frac{1}{2i\pi} \text{Tr} \left\{ \mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} \right\} = \frac{1}{2\pi\hbar} \text{Tr} \{ \mathcal{Q} \} , \quad (140)$$

where we have neglected the term $\text{Tr} \{ \mathcal{S} - \mathcal{S}^\dagger \} / \varepsilon$ in Eq. (22) (this is justified in the metallic regime). Such relation allows to characterize the amount of charge injected in an open coherent conductor, a crucial tool which has been used by Büttiker in order to describe screening properties. The out-of-equilibrium situation where the conductor is connected to several contacts (terminals) with different chemical potentials however requires to identify the contributions of each terminal to the DoS, which has led Büttiker to introduce several generalisations of (140) as the partial DoS, the injectance and the emittance. These concepts have been used

to develop a theory of non-linear transport [40, 53] and AC transport [40, 49, 50, 51, 166, 54] in coherent conductors (see the reviews [45, 41, 43, 48], or chapter 1 of [190]).

5.1. Partial DoS, injectance and emittance

Let us consider a multi-terminal structure, whose scattering properties are characterised by a basis of stationary scattering states $\psi_{\varepsilon,\alpha}(x)$ where the index α labels the terminals. We assume that the terminals support each a *single* conducting channel (i.e. contact wires are effectively one-dimensional), what simplify the analysis (the generalisation to contacts with several channels is straightforward). Furthermore, this would allow to illustrate the discussion by explicit formulae by considering the case of metric graphs for which explicit construction of the scattering matrix is possible [199, 193].

Büttiker introduced the concept of “*partial density of states*” [40, 51]

$$\nu_{\alpha\beta}(\varepsilon) \simeq \frac{1}{4i\pi} \left(\mathcal{S}_{\alpha\beta}^* \frac{\partial \mathcal{S}_{\alpha\beta}}{\partial \varepsilon} - \frac{\partial \mathcal{S}_{\alpha\beta}^*}{\partial \varepsilon} \mathcal{S}_{\alpha\beta} \right) \quad (141)$$

measuring the contribution to the DoS (140) of particles incoming from terminal β and outgoing at contact α . Another quantity which appears in Büttiker’s work is the “*injectance*”, obtained by summation of (141) over the first index

$$\bar{\nu}_\alpha(\varepsilon) = \sum_\beta \nu_{\beta\alpha}(\varepsilon) \simeq \frac{1}{2i\pi} \left(\mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} \right)_{\alpha\alpha} = \frac{\mathcal{Q}_{\alpha\alpha}}{2\pi} . \quad (142)$$

It provides the contribution to the DoS of the scattering states incoming from terminal α . Summation over the second index leads to the “*emittance*”

$$\underline{\nu}_\alpha(\varepsilon) = \sum_\beta \nu_{\alpha\beta}(\varepsilon) \simeq \frac{1}{2i\pi} \left(\frac{\partial \mathcal{S}}{\partial \varepsilon} \mathcal{S}^\dagger \right)_{\alpha\alpha} = \frac{\tilde{\mathcal{Q}}_{\alpha\alpha}}{2\pi} \quad (143)$$

which corresponds to the contribution of particles outgoing at terminal β . Injectance and emittance are related by magnetic field reversal,

$$\underline{\nu}_\alpha(\varepsilon; \mathcal{B}) = \bar{\nu}_\alpha(\varepsilon; -\mathcal{B}) , \quad (144)$$

which follows from $\mathcal{S}(-\mathcal{B}) = \mathcal{S}(\mathcal{B})^T$. Obviously

$$\nu(\varepsilon) = \sum_\alpha \bar{\nu}_\alpha(\varepsilon) = \sum_\alpha \underline{\nu}_\alpha(\varepsilon) = \sum_{\alpha,\beta} \nu_{\alpha\beta}(\varepsilon) . \quad (145)$$

Injectance and emittance allow for a preselection and a postselection, respectively, when identifying

the contribution to the density inside the conductor from particles passing through it [92].

Few remarks :

- (i) All formulae can be straightforwardly generalised to the case of multichannel contacts by adding some traces over channels.
- (ii) These concepts have been further generalised in order to deal with *local* properties, leading to the concept of partial local DoS, *injectivity* and *emissivity* [40, 92]. This can be understood from the relation

$$-\frac{1}{2i\pi} \left(\mathcal{S}^\dagger \frac{\delta \mathcal{S}}{\delta \mathcal{W}(x)} \right)_{\alpha\beta} = \psi_{\varepsilon,\alpha}^*(x) \psi_{\varepsilon,\beta}(x) \quad (146)$$

where $\mathcal{W}(x)$ is the potential (see Appendix of Ref. [42] or Refs. [189, 193, 195]). This leads to introduce the *injectivity*

$$\bar{\nu}_\alpha(x; \varepsilon) = -\frac{1}{2i\pi} \left(\mathcal{S}^\dagger \frac{\delta \mathcal{S}}{\delta \mathcal{W}(x)} \right)_{\alpha\alpha} = |\psi_{\varepsilon,\alpha}(x)|^2 \quad (147)$$

which measures the contribution of the scattering state to the local DoS $\nu(x; \varepsilon) = \sum_\alpha \bar{\nu}_\alpha(x; \varepsilon)$. Similarly, Büttiker introduced the concepts of partial *local* DoS and *emissivity* [40, 92].

- (iii) The Fisher and Lee relation [78]

$$\mathcal{S}_{\alpha\beta}(\varepsilon) = -\delta_{\alpha\beta} + i\sqrt{v_\alpha v_\beta} G^R(\alpha, \beta; \varepsilon), \quad (148)$$

where v_α is the group velocity in terminal α and where the retarded Green's function is measured at the two terminals, makes the functional derivation transparent :

$$\frac{\delta \mathcal{S}_{\alpha\beta}}{\delta \mathcal{W}(x)} = i\sqrt{v_\alpha v_\beta} G^R(\alpha, x; \varepsilon) G^R(x, \beta; \varepsilon). \quad (149)$$

Cf. also the discussion in [193].

- (iv) An illustration : the knowledge of the injectivities allows to express the density of electrons in the conductor out-of-equilibrium

$$n(x) = \sum_\alpha \int d\varepsilon f(\varepsilon - eV_\alpha) \bar{\nu}_\alpha(x; \varepsilon), \quad (150)$$

where $f(\varepsilon)$ is the Fermi function and V_α the electric potential at contact α .

- (v) The introduction of the injectivities allows to avoid the approximation made in (140) by neglecting the term $\text{Tr} \{ \mathcal{S} - \mathcal{S}^\dagger \} / \varepsilon$ of Eq. (22),

and provides a better definition for the partial DoS, injectance and emittance. Introducing a uniform potential $\mathcal{W}(x) = U$ inside the conductor (but not in the contacts), we can write

$$-\frac{1}{2i\pi} \left(\mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial U} \right)_{\alpha\beta} = \int_{\text{QD}} dx \psi_{\varepsilon,\alpha}^*(x) \psi_{\varepsilon,\beta}(x), \quad (151)$$

where $\int_{\text{QD}} \dots$ denotes integration inside the conductor [the boundaries of integration are given by the place where the scattering states are matched with plane waves in order to define scattering amplitudes, like in Eq. (2)]. In other terms [193, 195]

$$-\mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial U} = \mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial \varepsilon} + \frac{\mathcal{S} - \mathcal{S}^\dagger}{4\varepsilon}. \quad (152)$$

Therefore, instead of (142) which involves a high energy approximation, a more rigorous definition of the injectance should be

$$\bar{\nu}_\alpha(\varepsilon) = -\frac{1}{2i\pi} \left(\mathcal{S}^\dagger \frac{\partial \mathcal{S}}{\partial U} \right)_{\alpha\alpha}. \quad (153)$$

The remark also holds for partial DoS and emittances, Eqs. (141,143). The DoS takes the form $\nu(\varepsilon) = -(2i\pi)^{-1} \partial_U \ln \det \mathcal{S}$.

5.2. A generalisation of the Feynman-Hellmann theorem for a continuous spectrum

Büttiker's idea of relating the spectral properties to the scattering matrix is not limited to the DoS and the local DoS and can be generalised. This may be viewed as an extension of the famous Feynman-Hellmann theorem which applies to bounded problems with discrete energy spectra. Let us first recall this well-known theorem : consider a Hamiltonian H characterised by the discrete spectrum $\{\varepsilon_n, |\psi_n\rangle\}$. The determination of the diagonal matrix elements of an observable $X = -\partial H / \partial f$, where f is a conjugate force, does not require the knowledge of the eigenvectors but only of the eigenvalues : $\langle \psi_n | X | \psi_n \rangle = -\partial \varepsilon_n / \partial f$. In Ref. [195], we proposed that an extension of this theorem to the case of Hamiltonians with continuous spectra, characterised by scattering stationary states $\{|\psi_{\varepsilon,\alpha}\rangle\}$, is

$$\langle \psi_{\varepsilon,\alpha} | X | \psi_{\varepsilon,\beta} \rangle = \frac{1}{2i\pi} \left(\mathcal{S}^\dagger(\varepsilon) \frac{\partial \mathcal{S}(\varepsilon)}{\partial f} \right)_{\alpha\beta}. \quad (154)$$

This relation was proved in [193, 195] for various observables in metric graphs :²² for example, choosing the local density $X \rightarrow \rho(x) = |x\rangle\langle x|$ with $f \rightarrow -\mathcal{U}(x)$, we recover (146). In [195], we considered also the case of the current density I_a in a wire (a) of the graph, which involves a derivation with respect to the magnetic flux ϕ_a along the wire $\langle \psi_{\varepsilon,\alpha} | I_a | \psi_{\varepsilon,\beta} \rangle = (2i\pi)^{-1} (\mathcal{S}^\dagger \partial \mathcal{S} / \partial \phi_a)_{\alpha\beta}$. A trace over indices leads to the well-known formula obtained by Akkermans, Auerbach, Avron and Shapiro [3]²³ : $\langle I \rangle_\varepsilon = (2i\pi)^{-1} \partial_\phi \ln \det \mathcal{S}$. Note that similar considerations were used in order to analyse current fluctuations at equilibrium in Ref. [186] (a formula for current correlations out-of-equilibrium was obtained in [195]).

5.3. Statistical analysis in quantum dots

5.3.1. Mean values and covariances

In the case of chaotic quantum dots with several contacts with perfect couplings (Fig. 9), Brouwer and Büttiker [30] obtained the first two moments of the partial DoS (141) within the ASA of random matrices :²⁴

$$\langle \nu_{\mu\nu} \rangle = \frac{1}{\Delta} \left[\frac{N_\mu N_\nu}{N^2} - \frac{1}{N} \left(\frac{2}{\beta} - 1 \right) \left(\frac{N_\mu N_\nu}{N^2} - \frac{N_\mu}{N} \delta_{\mu\nu} \right) \right] \quad (155)$$

where N_μ is the number of open channels at contact μ and $N = \sum_\mu N_\mu$. We deduce $\langle \bar{\nu}_\alpha \rangle = \langle \nu_\alpha \rangle = (N_\alpha/N) \Delta^{-1}$, as expected in an ergodic device.

In the unitary case ($\beta = 2$), the covariances are [30]

$$\begin{aligned} \text{Cov}(\nu_{\mu\nu}, \nu_{\rho\sigma}) &= \frac{1}{(N\Delta)^2} \left[\frac{N_\mu N_\nu}{N^2} \left(\delta_{\mu\rho} \frac{N_\sigma}{N} + \delta_{\nu\sigma} \frac{N_\rho}{N} \right) \right. \\ &\quad \left. + \frac{3}{2} \left(\frac{N_\mu N_\rho}{N^2} - \frac{N_\mu}{N} \delta_{\mu\rho} \right) \left(\frac{N_\nu N_\sigma}{N^2} - \frac{N_\nu}{N} \delta_{\nu\sigma} \right) \right]. \end{aligned} \quad (156)$$

The correlator for the orthogonal case ($\beta = 1$) is obtained by adding similar terms obtained by permutation $\rho \leftrightarrow \sigma$ in order to fulfill the symmetry $\nu_{\rho\sigma} = \nu_{\sigma\rho}$ (for $\mathcal{B} = 0$) ; the symplectic case receives additionally a factor 1/4. Eq. (156) shows

²² Note that the expression (154) does not account for the contributions of BICs if present, as the DoS [195].

²³ See also the comment [59]

²⁴ Brouwer and Büttiker considered the dimensionless AC conductances $G_{\alpha\beta}(\omega) = G_{\alpha\beta}(0) - i\omega E_{\alpha\beta} + \mathcal{O}(\omega^2)$, which, in the unscreened limit, can be related to the partial DoS (141) : $\lim_{C \rightarrow \infty} E_{\alpha\beta} = E_{\alpha\beta}^u = 2\pi \int d\varepsilon (-\partial_\varepsilon f) \nu_{\alpha\beta}(\varepsilon)$.

that mesoscopic fluctuations are of order $\delta \nu_{\alpha\beta} \sim \langle \nu_{\alpha\beta} \rangle / N \sim 1/(N\Delta)$. Note that (156) corresponds to the leading order terms at large N (the remark also holds for the remaining expressions given in the subsection).

The correlations of injectances and emittances are also of interest. We deduce (for $\beta = 1, 2$)

$$\begin{aligned} \text{Cov}(\bar{\nu}_\mu, \bar{\nu}_\nu) &= \text{Cov}(\nu_\mu, \nu_\nu) \\ &= \frac{1}{(N\Delta)^2} \left[\frac{N_\mu N_\nu}{N^2} + \delta_{\mu\nu} \frac{N_\mu}{N} + \left(\frac{2}{\beta} - 1 \right) \frac{2N_\mu N_\nu}{N^2} \right] \end{aligned} \quad (157)$$

and

$$\begin{aligned} \text{Cov}(\bar{\nu}_\mu, \nu_\nu) &= \frac{1}{(N\Delta)^2} \\ &\times \left[\frac{2N_\mu N_\nu}{N^2} + \left(\frac{2}{\beta} - 1 \right) \left(\frac{N_\mu N_\nu}{N^2} + \delta_{\mu\nu} \frac{N_\mu}{N} \right) \right]. \end{aligned} \quad (158)$$

The interchange of the two contributions between (157) and (158) is analogous to the exchange between Diffuson and Cooperon in weakly disordered metals [198]. These expressions allow us to quantify the difference between injectance and emittance

$$\langle (\bar{\nu}_\alpha - \nu_\alpha)^2 \rangle = \left(2 - \frac{2}{\beta} \right) \frac{2}{(N\Delta)^2} \frac{N_\alpha}{N} \left(1 - \frac{N_\alpha}{N} \right). \quad (159)$$

The difference vanishes in the orthogonal case ($\beta = 1$), as a consequence of the symmetry (144). This quantity has found some application in [173].

5.3.2. Distribution of the injectance/emittance in the unitary case

As mentioned at the end of Subsection 4.1, the full distribution of the injectance has been obtained in an Appendix of Ref. [100] in the unitary case. This analysis was based on a result of Ref. [175] for the distribution of a diagonal subblock \mathcal{Q}_K of size $K \times K$ of the $N \times N$ Wigner-Smith matrix \mathcal{Q} (we recall that \mathcal{Q} and \mathcal{Q}_s have the same statistical properties only in the unitary case). Denote $\Gamma_K = \mathcal{Q}_K^{-1}$, its distribution is [175, 100] $P(\Gamma_K) \propto (\det \Gamma_K)^N \exp[-\text{Tr}\{\Gamma_K\}]$. We now introduce the dimensionless variable

$$s = \Delta \bar{\nu}_\alpha = \frac{\Delta}{2\pi} \text{Tr}\{\mathcal{Q}_K\} = \frac{1}{\tau_H} \sum_{a=1}^K \mathcal{Q}_{aa}. \quad (160)$$

The distribution was obtained in the large N limit, with $\kappa = K/N$ fixed [100] :

$$P_{N,\kappa}^{(\beta=2)}(s) \underset{N \rightarrow \infty}{\sim} \begin{cases} s^{N^2 \kappa(\kappa+2)/2} e^{-\kappa N^2/s} & \text{for } s \rightarrow 0 \\ \exp \left\{ -N^2 \frac{(s-\kappa)^2}{2\kappa(\kappa+1)} \right\} & \text{for } s \sim \kappa \\ s^{-N-2} & \text{for } s \rightarrow \infty \end{cases} \quad (161)$$

The variance $\text{Var}(s) \simeq \kappa(\kappa+1)/N^2$ corresponds with (157). In the limit $\kappa \rightarrow 1$, the distribution (161) coincides with (105).

In the unitary case, \mathcal{Q}_s , $\mathcal{Q} = \mathcal{S}^{-1/2} \mathcal{Q}_s \mathcal{S}^{1/2}$ and $\tilde{\mathcal{Q}} = \mathcal{S}^{1/2} \mathcal{Q}_s \mathcal{S}^{-1/2}$ have the same distribution,²⁵ thus (161) is also the distribution of the emittance.

6. Non-linear transport in coherent conductors

This section gives a brief presentation of the theory of non-linear transport in coherent conductors proposed by Büttiker and Christen [40, 53, 45], illustrating the interest of the quantities introduced in the previous section. We consider the case of conductors of mesoscopic dimensions such that they can be considered in the ergodic regime (like quantum dots), which slightly simplifies the presentation, although the theory was developed in a more general context in Ref. [40, 53] (see also [190]).




Figure 9: A conductor of mesoscopic dimension connected to $M = 3$ macroscopic reservoirs and capacitively coupled to a gate. When the potential at contact α is raised, current is injected in the conductor from this contact (blue arrows).

6.1. Non-linear conductances

We consider a multiterminal mesoscopic structure (Fig. 9). For small voltages $V_\alpha \rightarrow 0$, current

²⁵This follows from the fact that eigenvectors of \mathcal{Q} and \mathcal{S} are independent for $\beta = 2$ [32].

in terminal α can be written under the form of an expansion

$$I_\alpha = \frac{2_s e^2}{h} \sum_\beta g_{\alpha\beta} V_\beta + \frac{2_s e^3}{h} \sum_\beta g_{\alpha\beta\gamma} V_\beta V_\gamma + \dots \quad (162)$$

where 2_s is the spin degeneracy. $g_{\alpha\beta}$ is the dimensionless Landauer-Büttiker linear conductance [39]

$$g_{\alpha\beta} = \int d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) g_{\alpha\beta}(\varepsilon). \quad (163)$$

$f(\varepsilon)$ is the Fermi distribution and $g_{\alpha\beta}(\varepsilon) = N_\alpha \delta_{\alpha,\beta} - \text{Tr}\{\mathcal{S}_{\alpha\beta}^\dagger(\varepsilon) \mathcal{S}_{\alpha\beta}(\varepsilon)\}$ the zero temperature dimensionless conductance, N_α being the number of conducting channels in contact α .

The Landauer-Büttiker formula $I_\alpha = (2_s e/h) \sum_\beta \int d\varepsilon f(\varepsilon - eV_\beta) g_{\alpha\beta}(\varepsilon)$ already produces a contribution to the non-linear conductance

$$g_{\alpha\beta\gamma}^0 = \frac{1}{2} \delta_{\beta\gamma} \int d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) g'_{\alpha\beta}(\varepsilon). \quad (164)$$

Büttiker emphasized the importance of screening effect which produces a second contribution that we now determine. The first step is to analyse the electrostatic inside the conductor.

6.2. Characteristic potentials

Let us recall the main ingredients involved in the description of screening in good metals [213], which Büttiker adapted to deal with the non-equilibrium situation. These analysis is based on three ingredients. (i) When charges are introduced in the conductor, the total density of carriers $\delta n_{\text{tot}}(x) = \delta n_{\text{ext}}(x) + \delta n_{\text{ind}}(x)$ is the sum of the density $\delta n_{\text{ext}}(x)$ of charges injected in the conductor and the density $\delta n_{\text{ind}}(x)$ induced by screening. The concept of injectance introduced above allows us to write the number of injected charges when the conductor is out-of-equilibrium. An expansion of $\delta n_{\text{ext}}(x) = n(x) - n_{\text{equil}}(x)$ where $n(x)$ is given by (150) and $n_{\text{equil}}(x)$ is the equilibrium density, gives

$$\delta n_{\text{ext}} = \int_{\text{QD}} dx \delta n_{\text{ext}}(x) \simeq \sum_{\alpha=1}^M e V_\alpha \bar{\nu}_\alpha(\varepsilon_F), \quad (165)$$

where the summation runs over the M contacts. We assume zero temperature for simplicity ; the finite temperature formula involves an additional convolution with a Fermi function $\int d\varepsilon \left(-\partial f / \partial \varepsilon \right) \dots$.

(ii) The induced density is related to the potential energy $\mathcal{U}(x)$ by linear response theory

$$\delta n_{\text{ind}}(x) = - \int dx' \Pi(x, x') \mathcal{U}(x') \quad (166)$$

where $\Pi(x, x')$ is the (static) density-density correlation function of the non interacting electron gas (Lindhard function [35]). Assuming ergodic properties, the potential energy can be considered uniform in the conductor $\mathcal{U}(x) \simeq U$. Thomas-Fermi approximation relates the response to the DoS :

$$\delta n_{\text{ind}} = \int_{\text{QD}} dx \delta n_{\text{ind}}(x) \simeq -\nu(\varepsilon_F) U. \quad (167)$$

(iii) The potential energy and the density are related by the Poisson equation $\Delta \mathcal{U}(x) = -4\pi e^2 \delta n_{\text{tot}}(x)$ which encodes the Coulomb interaction in the conductor. The fact that (166) involves the potential $\mathcal{U}(x)$ and not the electrostatic potential related to the external density $\delta n_{\text{ext}}(x)$ makes the approach *self-consistent*. For a uniform potential, the Poisson equation is replaced by

$$C(U - eV_0) = e^2(\delta n_{\text{ext}} + \delta n_{\text{ind}}) \quad (168)$$

where C is the capacitance of the conductor and V_0 the electrostatic potential of the gate, labelled with index 0 (Fig. 9). Excess charge inside the conductor occurs when the system is driven out-of-equilibrium by the external potentials. At linear order, it is convenient to decompose the potential over contributions of the different external voltages

$$U \simeq \sum_{\alpha=0}^M u_{\alpha} eV_{\alpha} \quad (169)$$

where the “characteristic potential” u_{α} measures the response of the potential U to a shift of the external voltage V_{α} [40, 53] (one should not forget the response of the gate voltage controlled by u_0). Injecting (165,167,169) in (168) provides the characteristic potentials

$$u_{\alpha} = \frac{C_{\mu}}{C} \frac{\bar{\nu}_{\alpha}(\varepsilon_F)}{\nu(\varepsilon_F)} \quad (170)$$

for $\alpha \in \{1, \dots, M\}$ and

$$u_0 = 1 - \frac{C_{\mu}}{C} = \frac{C_{\mu}}{C_q}. \quad (171)$$

We have introduced the “mesoscopic capacitance”

$$\frac{1}{C_{\mu}} = \frac{1}{C} + \frac{1}{C_q}, \quad (172)$$

combining the “geometric capacitance” C and the “quantum capacitance”

$$C_q = e^2 \nu(\varepsilon_F). \quad (173)$$

We can check the sum rule

$$\sum_{\alpha=0}^M u_{\alpha} = 1. \quad (174)$$

The ratio $\gamma_{\text{int}} = C_{\mu}/C$ in Eq. (170) may be viewed as an interaction constant as it measures the efficiency of screening ($\gamma_{\text{int}} \simeq 1$ for efficient screening in a good metal and $\gamma_{\text{int}} \ll 1$ for weak screening). The response of the gate $u_0 = 1 - \gamma_{\text{int}}$ is thus important when screening is weak, however the presence of the gate (or the surrounding medium) can be forgotten in good metals.

6.3. Non-linear conductances

We can now determine the non-linear conductances as follows. The redefinition of the electrostatic potential inside the conductor leads to a modification of its scattering properties which depends on the potential, what contributes to the non-linear response : $g_{\alpha\beta}(\varepsilon_F) \rightarrow g_{\alpha\beta}(\varepsilon_F - U) \simeq g_{\alpha\beta}(\varepsilon_F) - g'_{\alpha\beta}(\varepsilon_F) \sum_{\gamma} u_{\gamma} eV_{\gamma}$. After symmetrisation with respect to indices $\beta \leftrightarrow \gamma$ and reintroducing the Fermi function, we get the expression of the non-linear conductance [53, 162]

$$g_{\alpha\beta\gamma} = \frac{1}{2} \int d\varepsilon \left(-\frac{\partial f}{\partial \varepsilon} \right) \times [g'_{\alpha\beta}(\varepsilon) \delta_{\beta\gamma} - g'_{\alpha\beta}(\varepsilon) u_{\gamma} - g'_{\alpha\gamma}(\varepsilon) u_{\beta}], \quad (175)$$

where the first term comes from the noninteracting theory, Eq. (164). For consistency, the $T = 0$ expression (170) should be replaced by

$$u_{\alpha} = \frac{\int d\varepsilon (-\partial_{\varepsilon} f) \bar{\nu}_{\alpha}(\varepsilon)}{C/e^2 + \int d\varepsilon (-\partial_{\varepsilon} f) \nu(\varepsilon)}. \quad (176)$$

The second term of the denominator is the finite temperature quantum capacitance $C_q = e^2 \int d\varepsilon (-\partial_{\varepsilon} f) \nu(\varepsilon)$.

Few remarks :

- (i) As emphasized above, all quantities involved in Eqs. (175,176), $g_{\alpha\beta}(\varepsilon)$, $\nu(\varepsilon)$ and $\bar{\nu}_{\alpha}(\varepsilon)$, are expressed in terms of the \mathcal{S} -matrix. The derivative of the conductance $g'_{\alpha\beta}(\varepsilon)$ can also

be related to the *sensitivies* introduced by Gasparian, Christen and Büttiker [92] :

$$\eta_{\alpha\beta}(\varepsilon) \simeq \frac{1}{4\pi} \left(\mathcal{S}_{\alpha\beta}^* \frac{\partial \mathcal{S}_{\alpha\beta}}{\partial \varepsilon} + \frac{\partial \mathcal{S}_{\alpha\beta}^*}{\partial \varepsilon} \mathcal{S}_{\alpha\beta} \right) \quad (177)$$

(note that Ref. [92] rather introduced a local version of the sensitivities).

- (ii) A statistical analysis of the non-linear conductance (175) in chaotic quantum dots requires the statistical properties of the conductance's derivative provided in Ref. [34] and those of the characteristic potentials, i.e. of the injectivities, obtained in [30], see Eqs. (155,156).

6.4. Recent developments

- (i) As mentioned, the Büttiker's scattering formalism can be extended to discuss conductors which are not in the ergodic regime [40, 53, 45]. The equivalence between this formalism and the non equilibrium Green's function approach for non-linear transport was established by Hernández and Lewenkopf [112] (see also [198]).
- (ii) An interesting aspect was identified by Sánchez and Büttiker [173] and Spivak and Zyuzin [183] : the interacting part of the non-linear conductance $g_{\alpha\beta\gamma}^{\text{int}} = -(1/2)[g'_{\alpha\beta}(\varepsilon_F)u_\gamma + g'_{\alpha\gamma}(\varepsilon_F)u_\beta]$ is not constrained to any specific symmetry under reversal of the magnetic field, contrary to the non-interacting part $g_{\alpha\beta\gamma}^0 = (1/2)g'_{\alpha\beta}(\varepsilon_F)\delta_{\beta\gamma}$. This is due to the asymmetry of the characteristic potentials u_α (i.e. of the injectance $\bar{\nu}_\alpha$). The analysis of the asymmetry of the non-linear conductance under magnetic field reversal $[g_{111}(\mathcal{B}) - g_{111}(-\mathcal{B})]/2 = -(C_\mu/C)g'_{11}(\varepsilon_F)(\bar{\nu}_1 - \nu_1)/(2\nu)$ was thus proposed as a new way for probing electronic interactions in mesoscopic structures (the parameter $\gamma_{\text{int}} = C_\mu/C$ introduced above was measured in [9] for a mesoscopic ring). Spivak and Zyuzin borrowed arguments from diagrammatic techniques in order to analyse the low magnetic field regime [183]. Sánchez and Büttiker [173] proposed a random matrix approach in the unitary case (strong magnetic field), which was later improved by Büttiker and Polianski who extended the statistical analysis in order to describe the crossover from orthogonal to unitary cases and include

thermal effects [162, 164] (for more details, cf. the excellent review article [163]). Note also the study for a ring made of strictly 1D wires [111].

- (iii) Several experimental groups have analysed the non-linear transport in mesoscopic structures and specifically the asymmetry in magnetic field [128, 137, 215, 9, 8].
- (iv) The case of disordered wires in the diffusive regime was analysed in Ref. [198].

7. AC transport

The search for fast control and manipulation of charge in coherent conductor has stimulated many developments (see the reviews [90, 24]). Büttiker, Prêtre and Thomas have proposed a theory of time-dependent response in coherent conductors [49, 50, 51] based on the scattering approach and a Hartree-Fock treatment (with Thomas-Fermi approximation) of electronic interactions (cf. chapter 1 of Ref. [190]). In order to emphasize few ideas, we will restrict ourselves here to the case of the “*quantum RC circuit*” which was studied experimentally in the integer quantum Hall regime during the PhD of Gabelli [89, 77, 91] (Fig. 10). The RC circuit was later studied from the perspective of an emitter of electronic wave packets in the PhD of Fève [77] and as such is a building block for “electron optics experiment” (see the review [24] and article [113] of this special issue). These latter developments have mostly considered the RC circuit in the integer quantum Hall regime, such that the current is carried by a single edge state along the boundary of the system, when one conducting channel is open at the constriction (quantum point contact, QPC).

Below, we do not consider the integer quantum Hall regime but we rather discuss the situation where the RC circuit of Fig. 10 is submitted to a weak magnetic field, such that the electron dynamics inside the cavity is chaotic. This justifies a random matrix approach. In this case, the opening of the constriction controls the number of channels N .

7.1. The AC response

By using arguments similar to the ones described in Section 6, Büttiker and coworkers have shown that the Coulomb interaction can be taken into account by adding the impedance of the non interacting electrons, $1/G_0(\omega)$, and the classical impedance




Figure 10: *The quantum RC circuit : a coherent conductor patterned in a 2DEG (blue) is closed by a quantum point contact (QPC) controlled by side gates. A capacitive coupling to a top gate allow to analyse its AC response. Figure from Ref. [102].*

of the capacitance $Z(\omega) \equiv 1/G(\omega) = 1/G_0(\omega) + 1/(-i\omega C)$. The admittance is [49]

$$G_0(\omega) = \frac{2_s e^2}{h} \int d\varepsilon \text{Tr} \left\{ 1 - \mathcal{S}^\dagger(\varepsilon) \mathcal{S}(\varepsilon + \omega) \right\} \times \frac{f(\varepsilon) - f(\varepsilon + \omega)}{\omega}, \quad (178)$$

where 2_s denotes the spin degeneracy. One can write the low frequency expansion of the impedance under the form given by the elementary laws of electrokinetics

$$Z(\omega) = \frac{1}{-i\omega C_\mu} + R_q + \mathcal{O}(\omega), \quad (179)$$

where the “mesoscopic capacitance” C_μ and the “charge relaxation resistance” R_q carry some information about the dynamics of electrons in a quantum coherent regime, as they control the relaxation time $\tau_{RC} = R_q C_\mu$. The presence of the mesoscopic capacitance (172) in the AC response comes from the $\omega \rightarrow 0$ expansion of $G_0(\omega)$, whose first terms involve the Wigner-Smith matrix (17). Hence, this latter controls both the quantum capacitance (173)

$$C_q = \frac{2_s e^2}{h} \text{Tr} \{ \mathcal{Q} \} \quad (180)$$

and the charge relaxation resistance

$$R_q = \frac{h}{2_s \times 2e^2} \frac{\text{Tr} \{ \mathcal{Q}^2 \}}{(\text{Tr} \{ \mathcal{Q} \})^2}. \quad (181)$$

In these equations, the Wigner-Smith matrix is taken at Fermi energy ε_F .

7.2. The charge relaxation resistance

In this section we forget about spin degeneracy factor 2_s and base the discussion on the expression

$$R_q = \frac{h}{2e^2} \frac{\text{Tr} \{ \mathcal{Q}^2 \}}{(\text{Tr} \{ \mathcal{Q} \})^2} = \frac{h}{2e^2} \frac{\sum_a \tau_a^2}{(\sum_a \tau_a)^2}. \quad (182)$$

A first important observation is that the resistance belongs to the interval :

$$\frac{h}{2Ne^2} \leq R_q \leq \frac{h}{2e^2}. \quad (183)$$

In particular the $N = 1$ channel case leads to the universal value $R_q = h/(2e^2)$ [50] (this is a property of a coherent conductor, cf. remarks closing the section).

Case $N = 2$ —. Assuming a random matrix description of the quantum dot, we can deduce the distribution of R_q starting from (85). The calculation only involves a double integral [98, 97] :

$$p_2(r_q) = A_\beta (2r_q - 1)^{(\beta-1)/2} (1 - r_q)^\beta \quad (184)$$

where $r_q = (Ne^2/h)R_q$ is the dimensionless charge relaxation resistance *per channel* ; $A_\beta = 2^{\beta+1}/B(\beta+1, (\beta+1)/2)$ is a normalisation constant. We deduce the mean value

$$\langle R_q \rangle = \frac{h}{3e^2} \quad (185)$$

which is surprisingly independent on the symmetry class.

Remark : The distribution of R_q for $N = 2$ channels was analysed by Pedersen, van Langen and Büttiker in [159], however these authors introduced a slightly different averaging procedure : following [34] the authors weighted the joint distribution of proper times by the DoS, corresponding to a “canonical” averaging. They deduced $p_2^{(\text{PvLB})}(r_q) = B_\beta (2r_q - 1)^{(\beta-1)/2} (1 - r_q)^{\beta-1}$ where the normalisation is $B_\beta = 2^\beta/B(\beta, (\beta+1)/2)$. As a result the mean resistance depends on the symmetry index $\langle R_q \rangle^{(\text{PvLB})} = (3/8) h/e^2$ for $\beta = 1$ and $\langle R_q \rangle^{(\text{PvLB})} = (5/14) h/e^2$ for $\beta = 2$.

Mean value —. Büttiker, Prêtre and Thomas [50] already pointed out that the charge relaxation resistance should scale with the number of channels as $R_q \sim 1/N$, which corresponds to the addition of resistances in parallel, although the precise prefactor was not given. The fact that the one channel case (spin polarised) leads to the universal *half* quantum of resistance $R_q = h/(2e^2)$ has produced a certain confusion for the large N case, between $R_q = h/(2Ne^2)$ (incorrect) and $R_q = h/(Ne^2)$ (correct).

The first precise calculation of the mean value $\langle R_q \rangle$ in the $N \gg 1$ limit is contained in the study of the complex admittance $G(\omega)$ by Brouwer and Büttiker [30] within the ASA of random matrices : the low frequency expansion of $\langle G(\omega) \rangle$ (Eq. 12 of Ref. [30]) leads to

$$\langle R_q \rangle = \frac{h}{Ne^2} \left[1 + \left(\frac{2}{\beta} - 1 \right) \frac{1}{N} + \mathcal{O}(N^{-2}) \right], \quad (186)$$

(without spin degeneracy 2_s). The second term can be interpreted as a weak localisation correction.²⁶ A first review paper by Büttiker [41] incorrectly quoted this result by introducing a spurious factor of $1/2$ (see Eq. 17 of Ref. [41]). In a second review [42], Büttiker gave the correct result [Eq. 9 of this reference]

$$\langle R_q \rangle \simeq \frac{h}{e^2} \frac{1}{N\mathcal{T}} \quad (187)$$

for a non perfect contact with $N\mathcal{T} \gg 1$ (without spin degeneracy), where \mathcal{T} is the transmission probability through the contact. Büttiker referred to a private communication with Carlo Beenakker and some unpublished work with myself [192], that I describe in Appendix A. This value coincides with the DC resistance of the constriction, what was expected in the large N limit.

We can summarize this discussion in the following table giving the dimensionless charge relaxation resistance *per channel* $r_q = (Ne^2/h)R_q$:

	$\langle r_q \rangle = (Ne^2/h)\langle R_q \rangle$
$N = 1$	$1/2$
$N = 2$	$2/3$
$N \gg 1$	$1 + (2/\beta - 1)N^{-1}$

Fluctuations —. As the quantum capacitance is directly proportional to the Wigner time delay, its statistical properties were determined in [197] for a contact with many channels. More recently, we have studied with Grabsch [102, 97] the statistical properties of the charge relaxation resistance, which involves the analysis of the ratio of two “linear statistics” $\sum_a \tau_a^2 / (\sum_a \tau_a)^2$. We have obtained the mean values $\langle C_q \rangle \simeq e^2/\Delta$ and $\langle R_q \rangle \simeq h/(Ne^2)$ (without spin degeneracy), in correspondence with the DC resistance of the QPC, as it

²⁶ Although the geometries are different, we can compare this expansion with those of the DC resistance of a coherent quantum dot closed by two constrictions with N_1 and N_2 channels [13] : $\langle R_{dc} \rangle = R_{class} [1 + (2/\beta - 1)N^{-1} + \dots]$ where $R_{class} = (h/e^2)(1/N_1 + 1/N_2)$ and $N = N_1 + N_2$.

should. The variance of the capacitance is given by [127, 30, 197, 146, 62]

$$\frac{\langle \delta C_q^2 \rangle}{\langle C_q \rangle^2} \simeq \frac{4}{\beta N^2} \quad (188)$$

(this coincides with the leading order term of cumulant (100) for $N \gg 1$). The variance of the resistance

$$\frac{\langle \delta R_q^2 \rangle}{\langle R_q \rangle^2} \simeq \frac{8}{\beta N^2} \quad (189)$$

and the correlations with the quantum capacitance

$$\frac{\langle \delta C_q \delta R_q \rangle}{\sqrt{\langle \delta C_q^2 \rangle \langle \delta R_q^2 \rangle}} = +1/\sqrt{2} \quad (190)$$

were found in Ref. [102], where we have also analysed the large deviations (see also [97] for further developments).

7.3. Final remarks

- (i) The multiterminal version of the general formulae reviewed in the section was provided in Refs. [50, 51].
- (ii) The finite temperature formulae are obtained by introducing convolutions with Fermi functions $\int d\varepsilon (-\partial f/\partial \varepsilon)$: the traces over channel indices in Eqs. (180,181) are then affected by additional integrations over energy.

In particular, using the representation $C_q = (e^2/h) \int d\varepsilon (-\partial_\varepsilon f) \text{Tr} \{ \mathcal{Q}(\varepsilon) \}$ with the correlator (138), the model leads to the following expression

$$\frac{\langle \delta C_q(\mathcal{B}) \delta C_q(\mathcal{B}') \rangle}{\langle C_q \rangle^2} \simeq \int d\omega \delta_T(\omega) \quad (191)$$

$$\times \frac{1}{\tau_H^2} \frac{\partial^2}{\partial \omega^2} \ln [(1 + (\omega\tau_{\mathcal{D}})^2) (1 + (\omega\tau_{\mathcal{E}})^2)]$$

where $\delta_T(\omega)$ is a (normalised) thermal function of width T which arises from the convolution of the two Fermi function's derivatives.²⁷ For $T\tau_d \gg 1$ this leads to the decay $\langle \delta C_q(\mathcal{B})^2 \rangle / \langle C_q \rangle^2 \simeq [\kappa/(\beta N^2)] (T\tau_d)^{-2}$ where $\kappa = -\int_0^\infty dx F'(x)/x \simeq 0.365$ (the function $F(x)$ is defined in the footnote). The

²⁷ $\delta_T(\omega) = F(\omega/(2T))/(2T)$ with $F(x) = (x \coth x - 1)/\sinh^2 x$.

dependence in the Dyson index β can be replaced by the precise magnetic field dependence straightforwardly.

Note that in the metallic limit ($e^2/C \gg \Delta$, i.e. $C \ll C_q$) the fluctuation of the mesoscopic capacitance is $\delta C_\mu \simeq (C/\langle C_q \rangle)^2 \delta C_q$.

- (iii) A more precise discussion of the capacitive response of quantum dots has been provided in Ref. [5], beyond mean field and also analysing the limit of a weakly transmitting constriction in the presence of Coulomb blockade.
- (iv) Nigg and Büttiker have considered in Ref. [150] the case of a single spin polarised channel. They showed that strong enough dephasing in the cavity can induce the transition from the universal value $R_q = h/(2e^2)$ to the DC resistance $R_{dc} = h/(e^2\mathcal{T})$ of the QPC, where \mathcal{T} is the transmission probability through the QPC.
- (v) The universal charge relaxation resistance $R_q = h/(2e^2)$ for a single spin non degenerate channel [50, 51, 42] has stimulated several works. This value was later obtained within more accurate treatments of electronic interactions : within a mean-field Hartree-Fock analysis [151, 171] improving the original Thomas-Fermi treatment, and beyond mean field [107, 148], within Matveev formalism [139] of QPC (see also [74]). These treatments have accounted for possible Coulomb blockade effects.

The treatment of interaction in a multichannel contact was extended more recently in [70].

8. Conclusion

Despite the huge diversity of Büttiker's work, this article has tried to show that a possible Ariane's thread is furnished by the concept of time delay and related quantities. Büttiker's interest in this matter goes back to his early fundamental study of times in quantum mechanics ; this question has led to many possible definitions, from scattering phase shifts (Wigner time delay, transmission group delays,...), clock approaches, etc. When randomness is introduced, these quantities are characterised by non-trivial distributions. I have reviewed in Section 3 the statistical analysis for the most simple of these fundamental times in the 1D situation in the presence of disorder. In particular I have emphasized the role of symmetry by considering the

effect of a chiral symmetry. In this case, a new observation was the relation between the presence of a chiral zero mode and the existence of a limit law for the Wigner time delay when the size of the disordered region goes to infinity.

One of Büttiker's major motivation for Wigner-Smith time delay matrix' analysis comes from its central role in the scattering approach for non-linear (Section 6) and AC (Section 7) coherent electronic transport. This has led to introduce several new concepts (Section 5) which have been analysed within the frame of random matrix theory : Büttiker, with Gopar and Mello, obtained one of the first important result in the Wigner time delay statistical analysis [96], motivated by the statistical analysis of the mesoscopic capacitance of a chaotic quantum dot. The first moments of the AC response were obtained with Brouwer [30] within the random matrix theory and the non-linear response with Sánchez and Polianski [173, 162]. In the present article I have provided a review of the known results on time delay statistics within the random matrix approach : joint distribution of proper times, moments of Wigner time delay and its correlation function and statistical properties of the partial DoS.

Acknowledgments

The material reviewed in this article is based on work done with several collaborators who are deeply thanked : Markus Büttiker, Alain Comtet, Pascal Degiovanni, Aurélien Grabsch, Satya Majumdar, Gilles Montambaux and Dmitry Savin. I thank Fabio Cunden for useful discussions. I am indebted to Dmitry Savin for numerous insightful comments. I am grateful to Alain Comtet for several remarks on the manuscript. Finally I thank the referee for helpful remarks.

Appendix A. Calculation of $\langle R_q \rangle$ for $N \gg 1$ – Mapping to the study of absorption in the cavity

This section reproduces some unpublished contribution due to myself and Büttiker [192] (quoted in Ref. [42] but remained unpublished so far).

Our aim is to estimate the charge relaxation resistance for a one contact conductor in the good contact limit $N\mathcal{T} \gg 1$, where N is the number of conducting channels and \mathcal{T} the transmission probability through the QPC. Our starting

point is the expression (182) (without spin degeneracy) [50]. As we mentioned in the text, the trace in the denominator is a measure of the density of states (DoS) of the cavity $\nu \simeq \text{Tr}\{\mathcal{Q}\}/h$, therefore $\langle \text{Tr}\{\mathcal{Q}\} \rangle = 2\pi/\Delta$ where Δ is the mean level spacing and the main question left is to estimate $\text{Tr}\{\mathcal{Q}^2\}$. We remark that it appears in the expansion of the dimensionless AC conductance (178)

$$\begin{aligned} G_0(\omega) &= N - \text{Tr}\left\{S^\dagger\left(\varepsilon - \frac{1}{2}\omega\right)S\left(\varepsilon + \frac{1}{2}\omega\right)\right\} \\ &= -i\omega \text{Tr}\{\mathcal{Q}\} + \frac{\omega^2}{2} \text{Tr}\{\mathcal{Q}^2\} + \dots \quad (\text{A.1}) \end{aligned}$$

Büttiker suggested to use analytic properties of the scattering matrix, $\omega \rightarrow i\Gamma$, i.e. to map the problem of AC transport onto the study of DC transport for a cavity in the presence of absorption, whose dimensionless conductance reads

$$\begin{aligned} \tilde{G} &= G_0(i\Gamma) \quad (\text{A.2}) \\ &= N - \text{Tr}\left\{\left[S\left(\varepsilon + \frac{i}{2}\Gamma\right)\right]^\dagger S\left(\varepsilon + \frac{i}{2}\Gamma\right)\right\} \\ &= \Gamma \text{Tr}\{\mathcal{Q}\} - \frac{\Gamma^2}{2} \text{Tr}\{\mathcal{Q}^2\} + \dots \end{aligned}$$

This corresponds to a situation where absorption takes place uniformly in space with absorbing rate Γ . In the ergodic regime where RMT holds, this also describes an absorbing contact (Fig. A.11).²⁸




Figure A.11: A conductor closed by a constriction with N channels (quantum point contact, QPC) and with an absorbing probe.

The idea is then to estimate \tilde{G} by noticing that a conductor with one contact and an absorbing probe is equivalent to a two terminal conductor (Fig. A.12). In this latter case, the conductance is dominated by the resistance of the constrictions, hence $1/\tilde{G} = 1/(N\mathcal{T}) + 1/(N_a\mathcal{T}_a)$, where N_a and \mathcal{T}_a characterise the second contact. We deduce

$$N - \tilde{G} = N \frac{N\mathcal{T} + N_a\mathcal{T}_a(1 - \mathcal{T})}{N\mathcal{T} + N_a\mathcal{T}_a}. \quad (\text{A.3})$$

²⁸ The relation between the dynamical problem (real ω) and the static problem with absorption (complex ω) was used in another context in Refs. [116, 168, 14, 17, 84].




Figure A.12: A two terminal conductor.

Expansion (A.2) in powers of Γ and expansion of (A.3) in powers of $N_a\mathcal{T}_a$ can be identified by setting $N_a\mathcal{T}_a = c(\Gamma/\Delta)$, where c is a dimensionless constant. Hence

$$\begin{aligned} \langle \text{Tr}\left\{\left[S\left(\varepsilon + i\frac{\Gamma}{2}\right)\right]^\dagger S\left(\varepsilon + i\frac{\Gamma}{2}\right)\right\} \rangle &= N \frac{N\mathcal{T} + (1 - \mathcal{T})c\frac{\Gamma}{\Delta}}{N\mathcal{T} + c\frac{\Gamma}{\Delta}} \\ &= N - c\frac{\Gamma}{\Delta} + \frac{1}{N\mathcal{T}} \left(c\frac{\Gamma}{\Delta}\right)^2 + \mathcal{O}(\Gamma^3). \quad (\text{A.4}) \end{aligned}$$

Comparison with (A.2) and the fact that $\langle \text{Tr}\{\mathcal{Q}\} \rangle = 2\pi/\Delta$ shows that $c = 2\pi$. We deduce

$$\langle \text{Tr}\{\mathcal{Q}^2\} \rangle = \langle \sum_i \tau_i^2 \rangle \simeq \frac{8\pi^2}{N\mathcal{T}\Delta^2}, \quad (\text{A.5})$$

leading to (187).

Appendix B. Partial versus proper times

Let us make few remarks on the relation between partial time delays and proper time delays. We consider a scattering situation with N channels. The S -matrix may be represented under the form

$$S = \mathcal{U} e^{i\Theta} \mathcal{U}^\dagger \quad (\text{B.1})$$

where $\Theta = \text{diag}(2\eta_1, \dots, 2\eta_N)$ gathers the N phase shifts. We deduce the following representation for the Wigner-Smith matrix :

$$\mathcal{Q} = \mathcal{U} (\partial_\varepsilon \Theta) \mathcal{U}^\dagger + i \left[S^\dagger (\mathcal{U} \partial_\varepsilon \mathcal{U}^\dagger) S - \mathcal{U} \partial_\varepsilon \mathcal{U}^\dagger \right] \quad (\text{B.2})$$

The first term involves the partial time delays : $\partial_\varepsilon \Theta = \text{diag}(\tilde{\tau}_1, \dots, \tilde{\tau}_N)$, which are intrinsic characteristics of the scattering matrix (independent on the basis). The second term controlled by \mathcal{U} , i.e. by the choice of basis, is the origin of the difference between partial and proper times.

We now say a little bit more on the unitary matrix \mathcal{U} in the $N = 2$ channel case. Going back to the notations introduced in § 2.2, we can rewrite the relation between partial waves (13) (eigenstates of

the \mathcal{S} -matrix) and the left/right scattering states in the matricial form

$$\begin{pmatrix} \phi_{\varepsilon,1}(x) \\ \phi_{\varepsilon,2}(x) \end{pmatrix} = \begin{pmatrix} A_{L,1} & A_{R,1} \\ A_{L,2} & A_{R,2} \end{pmatrix} \begin{pmatrix} \psi_{\varepsilon,L}(x) \\ \psi_{\varepsilon,R}(x) \end{pmatrix} \quad (\text{B.3})$$

which involves the transpose of the matrix which diagonalises the \mathcal{S} -matrix

$$\mathcal{U}(\varepsilon) = \begin{pmatrix} A_{L,1} & A_{L,2} \\ A_{R,1} & A_{R,2} \end{pmatrix} \quad (\text{B.4})$$

therefore, when eigenstates are properly normalised (associated to a measure $d\varepsilon$), we can write $\langle \phi_{\varepsilon,a} | \psi_{\varepsilon',b} \rangle = \mathcal{U}_{ba}(\varepsilon) \delta(\varepsilon - \varepsilon')$ where $a \in \{1, 2\}$ and $b \in \{L, R\}$.

Appendix B.1. Symmetric case

We first consider the symmetric case $r = r'$ and $t = t'$ [in 1D this corresponds to a symmetric potential $V(-x) = V(x)$]. The scattering states incoming from left and right are simply related by $\psi_{\varepsilon,R}(x) = \psi_{\varepsilon,L}(-x)$ and the two partial waves are the symmetric/antisymmetric eigenstates $\phi_{\varepsilon,a}(-x) = (-1)^{a+1} \phi_{\varepsilon,a}(x)$, hence the unitary matrix is independent on the energy in this case

$$\mathcal{U} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad (\text{B.5})$$

and the partial and proper times coincide

$$\tau_1 = \tilde{\tau}_1 \quad \text{and} \quad \tau_2 = \tilde{\tau}_2. \quad (\text{B.6})$$




Figure B.13: Proper time delays (continuous blue line) and partial time delays (dashed red line) as a function of $k = \sqrt{\varepsilon}$ in a particular example of 1D Hamiltonian with potential $V(x) = \lambda_1 \delta(x) + \lambda_2 \delta(x-a)$ with $\lambda_1 = 2$ and $\lambda_2 = 4$ ($a = 1$).

Appendix B.2. Asymmetric case

In general the matrix \mathcal{U} carries some energy dependence. In order to illustrate the difference between the two sets of characteristic times, we consider a specific 1D example : we compute these times for the Hamiltonian $H = -\partial_x^2 + \lambda_1 \delta(x) + \lambda_2 \delta(x-a)$. We consider the basis of left and right scattering states in which the \mathcal{S} -matrix takes the form (9). The three coefficients r , t and r' can be computed easily thanks to a transfer matrix approach. Diagonalisation of the \mathcal{S} -matrix provides the phase shifts and hence the partial time delays $\tilde{\tau}_1$ and $\tilde{\tau}_2$. We also compute the Wigner-Smith matrix in the basis of left/right scattering states (cf. § 2.2) and diagonalise it, we get the proper time delays τ_1 and τ_2 . We check that for the symmetric potential ($\lambda_1 = \lambda_2$) the two sets exactly coincide, Eq. (B.6). In the asymmetric case ($\lambda_1 \neq \lambda_2$) the four times are plotted in Fig. B.13 as a function of $\sqrt{\varepsilon}$. We see that proper and partial times are very close (they get closer as the energy grows). At the crossing points where the two partial times are equal we can however observe that partial times and proper times differ. This follows from the fact that the second term of Eq. (B.2) can be understood as a perturbation which produces anticrossing of the eigenvalues of \mathcal{Q} .

References

- [1] E. Abrahams, P. W. Anderson, D. C. Licciardello and T. V. Ramakrishnan, Scaling theory of localization: absence of quantum diffusion in two dimensions, Phys. Rev. Lett. **42**(10), 673 (1979).
- [2] M. Abramowitz and I. A. Stegun (editors), *Handbook of Mathematical functions*, Dover, New York (1964).
- [3] E. Akkermans, A. Auerbach, J. E. Avron and B. Shapiro, Relation between persistent currents and the scattering matrix, Phys. Rev. Lett. **66**(1), 76–79 (1991).
- [4] E. Akkermans and G. Montambaux, *Mesoscopic physics of electrons and photons*, Cambridge University Press, Cambridge, UK (2007).
- [5] I. L. Aleiner, P. W. Brouwer and L. I. Glazman, Quantum effects in Coulomb blockade, Phys. Rep. **358**(5–6), 309–440 (2002).
- [6] A. Altland and M. R. Zirnbauer, Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures, Phys. Rev. B **55**(2), 1142–1161 (1997).
- [7] P. Anderson, D. J. Thouless, E. Abrahams and D. S. Fisher, New method for a scaling theory of localization, Phys. Rev. B **22**(8), 3519–3526 (1980).
- [8] L. Angers, *Rectification et supraconductivité de proximité dans des anneaux mésoscopiques*, Ph.D. thesis, Université Paris-Sud (2007), <http://tel.archives-ouvertes.fr/tel-00156703>.

- [9] L. Angers, E. Zakka-Bajjani, R. Deblock, S. Guéron, H. Bouchiat, A. Cavanna, U. Gennser and M. Polianski, Magnetic-field asymmetry of mesoscopic *dc* rectification in Aharonov-Bohm rings, *Phys. Rev. B* **75**, 115309 (2007).
- [10] T. N. Antsygina, L. A. Pastur and V. A. Slyusarev, Localization of states and kinetic properties of one-dimensional disordered systems, *Sov. J. Low Temp. Phys.* **7**(1), 1–21 (1981).
- [11] Y. Avishai and Y. B. Band, One-dimensional density of states and the phase of the transmission amplitude, *Phys. Rev. B* **32**, 2674–2676 (1985).
- [12] M. Ya. Azbel, Resonance tunneling and localization spectroscopy, *Solid State Commun.* **45**(7), 527–530 (1983).
- [13] C. W. J. Beenakker, Random-matrix theory of quantum transport, *Rev. Mod. Phys.* **69**(3), 731–808 (1997).
- [14] C. W. J. Beenakker, Dynamics of localization in a waveguide, in *Photonic Crystals and Light Localization in the 21st Century*, edited by C. Soukoulis, NATO Science Series C563, pp. 489–508, Kluwer, Dordrecht (2001).
- [15] C. W. J. Beenakker, Search for Majorana Fermions in Superconductors, *Annu. Rev. Cond. Mat. Phys.* **4**, 113–136 (2013).
- [16] C. W. J. Beenakker, Random-matrix theory of Majorana fermions and topological superconductors, *Rev. Mod. Phys.* **87**, 1037–1066 (2015).
- [17] C. W. J. Beenakker and P. W. Brouwer, Distribution of the reflection eigenvalues of a weakly absorbing chaotic cavity, *Physica E* **9**, 463–466 (2001).
- [18] C. Benjamin and A. M. Jayannavar, Wave attenuation to clock sojourn time, *Solid State Commun.* **121**, 591–595 (2002).
- [19] D. Bernard and A. LeClair, A classification of 2D random Dirac fermions, *J. Phys. A: Math. Gen.* **35**, 2555–2567 (2002).
- [20] E. Beth and G. E. Uhlenbeck, The quantum theory of the non-ideal gas II. Behaviour at low temperatures, *Physica* **4**(10), 915–924 (1937).
- [21] T. Bienaimé and C. Texier, Localization for one-dimensional random potentials with large fluctuations, *J. Phys. A: Math. Theor.* **41**, 475001 (2008).
- [22] M. S. Birman and M. G. Krein, On the theory of wave operators, *Dokl. Akad. Nauk SSSR* **144**, 475–478 (1962), [*Soviet Math. Dokl.* **3**, 740–743 (1962)].
- [23] R. Blümel and U. Smilansky, Classical irregular scattering and its quantum-mechanical implications, *Phys. Rev. Lett.* **60**(6), 477–480 (1988).
- [24] E. Bocquillon, V. Freulon, F. D. Parmentier, J.-M. Berroir, B. Plaçais, C. Wahl, J. Rech, T. Jonckheere, T. Martin, C. Grenier, D. Ferraro, P. Degiovanni and G. Fève, Electron quantum optics in ballistic chiral conductors, *Ann. Phys. (Berlin)* **526**(1-2), 1–30 (2014).
- [25] D. Bollé and T. A. Osborn, Concepts of multiparticle time delay, *Phys. Rev. D* **13**, 299–311 (1976).
- [26] C. J. Bolton-Heaton, C. J. Lambert, V. I. Fal’ko, V. Prigodin and A. J. Epstein, Distribution of time constants for tunneling through a one-dimensional disordered chain, *Phys. Rev. B* **60**(15), 10569–10572 (1999).
- [27] J.-P. Bouchaud, A. Comtet, A. Georges and P. Le Doussal, Classical diffusion of a particle in a one-dimensional random force field, *Ann. Phys. (N.Y.)* **201**, 285–341 (1990).
- [28] P. W. Brouwer, Generalized circular ensemble of scattering matrices for a chaotic cavity with nonideal leads, *Phys. Rev. B* **51**, 16878–16884 (1995).
- [29] P. W. Brouwer and C. W. J. Beenakker, Diagrammatic method of integration over the unitary group, with applications to quantum transport in mesoscopic systems, *J. Math. Phys.* **37**(10), 4904–4934 (1996).
- [30] P. W. Brouwer and M. Büttiker, Charge-relaxation and dwell time in the fluctuating admittance of a chaotic cavity, *Europhys. Lett.* **37**(7), 441–446 (1997).
- [31] P. W. Brouwer, K. M. Frahm and C. W. Beenakker, Quantum mechanical time-delay matrix in chaotic scattering, *Phys. Rev. Lett.* **78**(25), 4737 (1997).
- [32] P. W. Brouwer, K. M. Frahm and C. W. Beenakker, Distribution of the quantum mechanical time-delay matrix for a chaotic cavity, *Waves Random Media* **9**, 91–104 (1999).
- [33] P. W. Brouwer, A. Lamacraft and K. Flensberg, Nonequilibrium theory of Coulomb blockade in open quantum dots, *Phys. Rev. B* **72**, 075316 (2005).
- [34] P. W. Brouwer, S. A. van Langen, K. M. Frahm, M. Büttiker and C. W. J. Beenakker, Distribution of Parametric Conductance Derivatives of a Quantum Dot, *Phys. Rev. Lett.* **79**, 913–916 (1997).
- [35] H. Bruus and K. Flensberg, *Many-body quantum theory in condensed matter physics*, Oxford University Press, Oxford (2004).
- [36] V. S. Buslaev, Trace formulas for the three-dimensional Schrödinger operator, *Dokl. Akad. Nauk SSSR* **143** (1962).
- [37] M. Büttiker, Larmor precession and the traversal time for tunneling, *Phys. Rev. B* **27**(10), 6178–6188 (1983).
- [38] M. Büttiker, Traversal, reflection and dwell time for quantum tunneling, in *Electronic properties of multilayers and low-dimensional semiconductors structures*, edited by J. M. Chamberlain et al., p. 297, Plenum Press, New York (1990).
- [39] M. Büttiker, Scattering theory of current and intensity noise correlations in conductors and wave guides, *Phys. Rev. B* **46**(19), 12485–12507 (1992).
- [40] M. Büttiker, Capacitance, admittance, and rectification properties of small conductors, *J. Phys. Cond. Matter* **5**(50), 9361–9378 (1993).
- [41] M. Büttiker, Charge relaxation resistances and charge fluctuations in mesoscopic conductors, *J. Korean Phys. Soc.* **34**, S121–130 (1999), (cond-mat/9902054).
- [42] M. Büttiker, Charge fluctuations and dephasing in Coulomb coupled conductors, in *Quantum mesoscopic phenomena and mesoscopic devices*, edited by I. O. Kulik and R. Ellialtıoglu, volume 559, p. 211, Kluwer Academic Publishers, Dordrecht (2000), (cond-mat/9911188).
- [43] M. Büttiker, Time-dependent transport in mesoscopic structures, *J. Low Temp. Phys.* **11**(5/6), 519–542 (2000).
- [44] M. Büttiker, The local Larmor clock, Partial densities of states, and mesoscopic physics, in *Time in quantum mechanics*, edited by J. G. Muga, R. S. Mayato and I. L. Egusquiza, Lecture Notes in Physics, p. 256, Springer (2002), preprint quant-ph/0103164.
- [45] M. Büttiker and T. Christen, Admittance and nonlinear transport in quantum wires, points contacts and resonant tunneling barriers, in *Mesoscopic Electron*

- Transport*, edited by L. L. Sohn, L. P. Kouwenhoven and G. Schön, pp. 259–289, Kluwer Academic Publishers, Dordrecht (1997).
- [46] M. Büttiker and R. Landauer, Traversal time for tunneling, *Phys. Rev. Lett.* **49**, 1739 (1982).
 - [47] M. Büttiker and R. Landauer, Traversal time for tunneling, *Phys. Scr.* **32**, 429–434 (1985).
 - [48] M. Büttiker and M. L. Polianski, Charge fluctuation in open chaotic cavities, *J. Phys. A: Math. Theor.* **38**, 10559–10585 (2005).
 - [49] M. Büttiker, A. Prêtre and H. Thomas, Dynamic conductance and the scattering matrix of small conductors, *Phys. Rev. Lett.* **70**(26), 4114 (1993).
 - [50] M. Büttiker, H. Thomas and A. Prêtre, Mesoscopic capacitors, *Phys. Lett. A* **180**, 364–369 (1993).
 - [51] M. Büttiker, H. Thomas and A. Prêtre, Current partition in multiprobe conductors in the presence of slowly oscillating potentials, *Z. Phys. B* **94**, 133–137 (1994).
 - [52] A. A. Chabanov and A. Z. Genack, Statistics of Dynamics of Localized Waves, *Phys. Rev. Lett.* **87**, 233903 (2001).
 - [53] T. Christen and M. Büttiker, Gauge invariant nonlinear electric transport in mesoscopic conductors, *Europhys. Lett.* **35**(7), 523–528 (1996).
 - [54] T. Christen and M. Büttiker, Low-frequency admittance of quantized Hall conductors, *Phys. Rev. B* **53**, 2064–2072 (1996).
 - [55] A. Cohen, Y. Roth and B. Shapiro, Universal distributions and scaling in disordered systems, *Phys. Rev. B* **38**(17), 12125–12132 (1988).
 - [56] A. Comtet, J. Desbois and C. Texier, Functionals of the Brownian motion, localization and metric graphs, *J. Phys. A: Math. Gen.* **38**, R341–R383 (2005).
 - [57] A. Comtet and C. Monthus, Diffusion in one-dimensional random medium and hyperbolic Brownian motion, *J. Phys. A: Math. Gen.* **29**(7), 1331–1345 (1996).
 - [58] A. Comtet, C. Monthus and M. Yor, Exponential functionals of Brownian motion and disordered systems, *J. Appl. Probab.* **35**, 255 (1998).
 - [59] A. Comtet, A. Moroz and S. Ouvry, Persistent current of free electrons in the plane, *Phys. Rev. Lett.* **74**(5), 828 (1995).
 - [60] A. Comtet and C. Texier, On the distribution of the Wigner time delay in one-dimensional disordered systems, *J. Phys. A: Math. Gen.* **30**, 8017–8025 (1997).
 - [61] A. Comtet and C. Texier, One-dimensional disordered supersymmetric quantum mechanics: a brief survey, in *Supersymmetry and Integrable Models*, edited by H. Aratyn, T. D. Imbo, W.-Y. Keung and U. Sukhatme, Lecture Notes in Physics, Vol. **502** (available as arXiv:cond-mat/9707313), pp. 313–328. Springer (1998).
 - [62] F. D. Cunden, Statistical distribution of the Wigner-Smith time-delay matrix moments for chaotic cavities, *Phys. Rev. E* **91**, 060102 (2015).
 - [63] F. D. Cunden, F. Mezzadri, N. Simm and P. Vivo, Correlators for the Wigner-Smith time-delay matrix of chaotic cavities, *J. Phys. A: Math. Theor.* **49**, 18LT01 (2016).
 - [64] F. D. Cunden, F. Mezzadri, N. Simm and P. Vivo, Large- N expansion for the time-delay matrix of ballistic chaotic cavities, *J. Math. Phys.* **57**(11), 111901 (2016).
 - [65] F. D. Cunden and P. Vivo, Universal covariance formula for linear statistics on random matrices, *Phys. Rev. Lett.* **113**, 070202 (2014).
 - [66] R. Dashen, S.-K. Ma and H. J. Bernstein, S -matrix formulation of statistical mechanics, *Phys. Rev.* **187**(1), 345–370 (1969).
 - [67] C. A. A. de Carvalho and H. M. Nussenzveig, Time delay, *Phys. Rep.* **364**, 83–174 (2002).
 - [68] L. I. Deych, A. A. Lisiansky and B. L. Altshuler, Single Parameter Scaling in One-Dimensional Localization Revisited, *Phys. Rev. Lett.* **84**(12), 2678 (2000).
 - [69] L. I. Deych, A. A. Lisiansky and B. L. Altshuler, Single Parameter Scaling in 1-D Anderson localization. Exact analytical solution, *Phys. Rev. B* **64**, 224202 (2001).
 - [70] P. Dutt, T. L. Schmidt, C. Mora and K. Le Hur, Strongly correlated dynamics in multichannel quantum RC circuits, *Phys. Rev. B* **87**, 155134 (2013).
 - [71] B. Eckhardt, Correlations in quantum time delay, *Chaos* **3**(4), 613–617 (1993).
 - [72] L. Eisenbud, *The formal properties of nuclear collisions*, Ph.D. thesis, Princeton (1948).
 - [73] T. Ericson, Fluctuations of Nuclear Cross Sections in the “Continuum” Region, *Phys. Rev. Lett.* **5**, 430–431 (1960).
 - [74] Y. Etzioni, B. Horovitz and P. Le Doussal, Rings and boxes in dissipative environments, *Phys. Rev. Lett.* **106**, 166803 (2011).
 - [75] F. Evers and A. D. Mirlin, Anderson transitions, *Rev. Mod. Phys.* **80**(4), 1355–1417 (2008).
 - [76] W. G. Faris and W. J. Tsay, Time delay in random scattering, *SIAM J. Appl. Math.* **54**(2), 443–455 (1994).
 - [77] G. Fève, *Quantification du courant alternatif : la boîte quantique comme source d’électrons uniques sub-nanoseconde*, Ph.D. thesis, Université Paris 6 (2006), <http://tel.archives-ouvertes.fr/tel-00119589>.
 - [78] D. S. Fisher and P. A. Lee, Relation between conductivity and transmission matrix, *Phys. Rev. B* **23**(12), 6851–6854 (1981).
 - [79] P. J. Forrester, Large deviation eigenvalue density for the soft edge Laguerre and Jacobi β -ensembles, *J. Phys. A: Math. Theor.* **45**, 145201 (2012).
 - [80] J. Friedel, The distribution of electrons round impurities in monovalent metals, *Phil. Mag.* **43**, 153–189 (1952).
 - [81] J. Friedel, Metallic alloys, *Nuovo Cimento Suppl.* **7**, 287–311 (1958).
 - [82] W. A. Friedman and P. A. Mello, Information theory and statistical nuclear reactions II. Many-channel case and Hauser-Feshbach formula, *Ann. Phys.* **161**(2), 276–302 (1985).
 - [83] U. Frisch, C. Froeschle, J.-P. Scheidecker and P.-L. Sulem, Stochastic resonance in one-dimensional random media, *Phys. Rev. A* **8**(3), 1416–1421 (1973).
 - [84] Y. Fyodorov, Induced vs. spontaneous breakdown of S -matrix unitarity: Probability of no return in quantum chaotic and disordered systems, *JETP Letters* **78**, 250–254 (2003).
 - [85] Y. V. Fyodorov and A. Ossipov, Distribution of the Local Density of States, Reflection Coefficient, and Wigner Delay Time in Absorbing Ergodic Systems at the Point of Chiral Symmetry, *Phys. Rev. Lett.* **92**, 084103 (2004).
 - [86] Y. V. Fyodorov, D. V. Savin and H.-J. Sommers, Parametric correlations of phase shifts and statistics of time

- delays in quantum chaotic scattering: Crossover between unitary and orthogonal symmetries, *Phys. Rev. E* **55**, R4857–R4860 (1997).
- [87] Y. V. Fyodorov and H.-J. Sommers, Parametric correlations of scattering phase shifts and fluctuations of delay times in few-channel chaotic scattering, *Phys. Rev. Lett.* **76**(25), 4709 (1996).
 - [88] Y. V. Fyodorov and H.-J. Sommers, Statistics of resonance poles, phase shifts and time delays in quantum chaotic scattering: Random matrix approach for systems with broken time-reversal invariance, *J. Math. Phys.* **38**(4), 1918–1981 (1997).
 - [89] J. Gabelli, *Mise en évidence de la cohérence quantique dans des conducteurs en régime dynamique*, Ph.D. thesis, Université Paris 6 (2006), <http://tel.archives-ouvertes.fr/tel-00011619>.
 - [90] J. Gabelli, G. Fève, J.-M. Berroir and B. Plaçais, A coherent RC circuit, *Rep. Prog. Phys.* **75**, 126504 (2012).
 - [91] J. Gabelli, G. Fève, J.-M. Berroir, B. Plaçais, A. Cavanaugh, B. Etienne, Y. Jin and C. Glattli, Violation of Kirchhoff’s laws for a coherent RC circuit, *Science* **313**(5786), 499–502 (2006).
 - [92] V. Gasparian, T. Christen and M. Büttiker, Partial densities of states, scattering matrices and Green’s functions, *Phys. Rev. A* **54**(5), 4022–4031 (1996).
 - [93] A. Z. Genack, P. Sebbah, M. Stoytchev and B. A. van Tiggelen, Statistics of wave dynamics in random media, *Phys. Rev. Lett.* **82**(4), 715 (1999).
 - [94] M. E. Gertsenshtein and V. B. Vasil’ev, Waveguides with random inhomogeneities and Brownian motion in the Lobachevsky plane, *Theory Prob. and Appl.* **4**(4), 391–398 (1959).
 - [95] V. A. Gopar and P. A. Mello, The problem of quantum chaotic scattering with direct processes reduced to the one without, *Europhys. Lett.* **42**(2), 131–136 (1998).
 - [96] V. A. Gopar, P. A. Mello and M. Büttiker, Mesoscopic capacitors: a statistical analysis, *Phys. Rev. Lett.* **77**(14), 3005 (1996).
 - [97] A. Grabsch, *Random matrices in statistical physics: quantum scattering and disordered systems*, Ph.D. thesis, Université Paris Saclay (2018), <https://tel.archives-ouvertes.fr/tel-01849097v1>.
 - [98] A. Grabsch, S. Majumdar and C. Texier, Time delays and coherent AC transport in chaotic cavities – Random matrices and Coulomb gas, in preparation (2016).
 - [99] A. Grabsch, S. N. Majumdar and C. Texier, Truncated linear statistics associated with the top eigenvalues of random matrices, *J. Stat. Phys.* **167**, 234–259 (2017).
 - [100] A. Grabsch, S. N. Majumdar and C. Texier, Truncated linear statistics associated with the eigenvalues of random matrices II. Partial sums over proper time delays for chaotic quantum dots, *J. Stat. Phys.* **167**, 1452–1488 (2017).
 - [101] A. Grabsch, D. V. Savin and C. Texier, Wigner-Smith time-delay matrix in chaotic cavities with non-ideal contacts, *J. Phys. A: Math. Theor.* **51**, 404001 (2018), special issue “*Random Matrices: the first 90 years*”.
 - [102] A. Grabsch and C. Texier, Capacitance and charge relaxation resistance of chaotic cavities – Joint distribution of two linear statistics in the Laguerre ensemble of random matrices, *Europhys. Lett.* **109**, 50004 (2015).
 - [103] A. Grabsch and C. Texier, Distribution of spectral linear statistics on random matrices beyond the large deviation function – Wigner time delay in multichannel disordered wires, *J. Phys. A: Math. Theor.* **49**, 465002 (2016).
 - [104] A. Grabsch and C. Texier, Topological phase transitions in the 1D multichannel Dirac equation with random mass and a random matrix model, *Europhys. Lett.* **116**, 17004 (2016).
 - [105] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series and products*, Academic Press, fifth edition (1994).
 - [106] E. Gurevich and B. Shapiro, Statistics of resonances in one-dimensional disordered systems, *Lithuanian Journal of Physics* **52**, 115 (2012).
 - [107] Y. Hamamoto, T. Jonckheere, T. Kato and T. Martin, Dynamic response of a mesoscopic capacitor in the presence of strong electron interactions, *Phys. Rev. B* **81**, 153305 (2010).
 - [108] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, *Rev. Mod. Phys.* **82**(4), 3045–3067 (2010).
 - [109] E. H. Hauge and J. A. Støvneng, Tunneling times: a critical review, *Rev. Mod. Phys.* **61**(4), 917–936 (1989).
 - [110] J. Heinrichs, Invariant embedding treatment of phase randomization and electrical noise at disordered surfaces, *J. Phys. Cond. Matter* **2**, 1559–1568 (1990).
 - [111] A. R. Hernández and C. H. Lewenkopf, Nonlinear Conductance in a Ballistic Aharonov-Bohm Ring, *Phys. Rev. Lett.* **103**, 166801 (2009).
 - [112] A. R. Hernández and C. H. Lewenkopf, Nonlinear electronic transport in nanoscopic devices: nonequilibrium Greens functions versus scattering approach, *Eur. Phys. J. B* **86**(4), 131 (2013).
 - [113] P. P. Hofer, D. Dasenbrook and C. Flindt, Electron waiting times for the mesoscopic capacitor, *Physica E* **82**, 3–11 (2016), *Frontiers in quantum electronic transport - In memory of Markus Büttiker*.
 - [114] A. M. Jayannavar, G. V. Vijayagovindan and N. Kumar, Energy dispersive backscattering of electrons from surface resonances of a disordered medium and $1/f$ noise, *Z. Phys. B – Condens. Matter* **75**, 77–79 (1989).
 - [115] S. K. Joshi and A. M. Jayannavar, Distribution of Wigner delay time from single channel disordered systems, *Solid State Commun.* **106**(6), 363 (1998).
 - [116] V. I. Klyatskin and A. I. Saichev, Statistical and dynamic localization of plane waves in randomly layered media, *Sov. Phys. Usp.* **35**(3), 231–247 (1992), [*Usp. Fiz. Nauk* **162**, 161 (1992)].
 - [117] T. Kottos, Statistics of resonances and delay times in random media: beyond random matrix theory, *J. Phys. A: Math. Theor.* **38**, 10761–10786 (2005).
 - [118] T. Kottos and M. Weiss, Statistics of resonances and delay times: A criterion for metal-insulator transitions, *Phys. Rev. Lett.* **89**, 056401 (2002).
 - [119] M. G. Krein, Trace formulas in perturbation theory, *Matem. Sbornik* **33**, 597 (1953).
 - [120] J. Kuipers, D. V. Savin and M. Sieber, Efficient semiclassical approach for time delays, *New J. Phys.* **16**, 123018 (2014).
 - [121] J. Kuipers and M. Sieber, Semiclassical expansion of parametric correlation functions of the quantum time delay, *Nonlinearity* **20**(4), 909–926 (2007).
 - [122] J. Kuipers and M. Sieber, Semiclassical relation between open trajectories and periodic orbits for the Wigner time delay, *Phys. Rev. E* **77**, 046219 (2008).
 - [123] H. Kunz and B. Shapiro, Statistics of resonances in a semi-infinite disordered chain, *Phys. Rev. B* **77**,

- 054203 (2008).
- [124] L. D. Landau and E. Lifchitz, *Physique statistique*, Mir (1966), tome 5.
 - [125] R. Landauer and T. Martin, Barrier interaction time in tunneling, *Rev. Mod. Phys.* **66**(1), 217–228 (1994).
 - [126] C. R. Leavens and G. C. Aers, Dwell time and phase times for transmission and reflection, *Phys. Rev. B* **39**(2), 1202–1206 (1989).
 - [127] N. Lehmann, D. V. Savin, V. V. Sokolov and H.-J. Sommers, Time delay correlations in chaotic scattering: random matrix approach, *Physica D* **86**, 572–585 (1995).
 - [128] R. Leturcq, D. Sánchez, G. Götz, T. Ihn, K. Ensslin, D. C. Driscoll and A. C. Gossard, Magnetic Field Symmetry and Phase Rigidity of the Nonlinear Conductance in a Ring, *Phys. Rev. Lett.* **96**, 126801 (2006).
 - [129] A. Levy Yeyati and M. Büttiker, Scattering phases in quantum dots: An analysis based on lattice models, *Phys. Rev. B* **62**(11), 7307–7315 (2000).
 - [130] C. H. Lewenkopf and R. O. Vallejos, Open orbits and the semiclassical dwell time, *J. Phys. A: Math. Gen.* **37**(1), 131–136 (2004).
 - [131] P. Lloyd, Exactly solvable model of electronic states in a three-dimensional disordered Hamiltonian: non-existence of localized states, *J. Phys. C: Solid St. Phys.* **2**(10), 1717–1725 (1969).
 - [132] V. L. Lyuboshitz, On collision duration in the presence of strong overlapping resonance levels, *Phys. Lett.* **72B**(1), 41–44 (1977).
 - [133] V. L. Lyuboshits, The probability distribution of the delay time of a wave packet in strong overlap of resonance levels, *Sov. J. Nucl. Phys.* **37**, 174–176 (1983), [*Yad. Fiz.* **37**, 292–297 (1983)].
 - [134] S.-K. Ma, *Statistical mechanics*, World Scientific, Singapore (1985).
 - [135] M. Marciani, P. W. Brouwer and C. W. J. Beenakker, Time-delay matrix, midgap spectral peak, and thermopower of an Andreev billiard, *Phys. Rev. B* **90**, 045403 (2014).
 - [136] M. Marciani, H. Schomerus and C. W. J. Beenakker, Effect of a tunnel barrier on the scattering from a Majorana bound state in an Andreev billiard, *Physica E* **77**, 54–64 (2016), *Frontiers in quantum electronic transport - In memory of Markus Büttiker*.
 - [137] C. A. Marlow, R. P. Taylor, M. Fairbanks, I. Shorubalko and H. Linke, Experimental Investigation of the Breakdown of the Onsager-Casimir Relations, *Phys. Rev. Lett.* **96**, 116801 (2006).
 - [138] A. M. Martínez-Argüello, M. Martínez-Mares and J. C. García, Joint moments of proper delay times, *J. Math. Phys.* **55**(8), 081901 (2014).
 - [139] K. A. Matveev, Coulomb blockade at almost perfect transmission, *Phys. Rev. B* **51**, 1743–1751 (1995).
 - [140] M. L. Mehta, *Random matrices*, Elsevier, Academic, New York, third edition (2004).
 - [141] P. A. Mello and H. U. Baranger, Interference phenomena in electronic transport through chaotic cavities: An information-theoretic approach, *Waves Random Media* **9**, 105–162 (1999).
 - [142] P. A. Mello and N. Kumar, *Quantum transport in mesoscopic systems – Complexity and statistical fluctuations*, Oxford University Press (2004).
 - [143] P. A. Mello, P. Pereyra and T. H. Seligman, Information theory and statistical nuclear reactions. I. General theory and applications to few-channel problems, *Ann. Phys.* **161**(2), 254–275 (1985).
 - [144] F. Mezzadri and N. J. Simm, Moments of the transmission eigenvalues, proper delay times, and random matrix theory. I, *J. Math. Phys.* **52**, 103511 (2011).
 - [145] F. Mezzadri and N. J. Simm, Moments of the transmission eigenvalues, proper delay times, and random matrix theory. II, *J. Math. Phys.* **53**, 053504 (2012).
 - [146] F. Mezzadri and N. J. Simm, τ -function theory of quantum chaotic transport with $\beta = 1, 2, 4$, *Commun. Math. Phys.* **324**, 465–513 (2013).
 - [147] C. Monthus and A. Comtet, On the flux distribution in a one-dimensional disordered system, *J. Phys. I (France)* **4**(6), 635–653 (1994).
 - [148] C. Mora and K. Le Hur, Universal resistances of the quantum resistance-capacitance circuit, *Nat. Phys.* **6**, 697–701 (2010).
 - [149] E. R. Mucciolo, R. A. Jalabert and J.-L. Pichard, Parametric statistics of the scattering matrix: from metallic to insulating quasi-unidimensional disordered systems, *J. Phys. I (France)* **7**, 1267 (1997).
 - [150] S. E. Nigg and M. Büttiker, Quantum to classical transition of the charge relaxation resistance of a mesoscopic capacitor, *Phys. Rev. B* **77**, 085312 (2008).
 - [151] S. E. Nigg, R. López and M. Büttiker, Mesoscopic charge relaxation, *Phys. Rev. Lett.* **97**, 206804 (2006).
 - [152] M. Novaes, Statistics of time delay and scattering correlation functions in chaotic systems. I. Random matrix theory, *J. Math. Phys.* **56**, 062110 (2015).
 - [153] M. Novaes, Statistics of time delay and scattering correlation functions in chaotic systems. II. Semiclassical approximation, *J. Math. Phys.* **56**, 062109 (2015).
 - [154] G. Oshanin, A. Mogutov and M. Moreau, Steady flux in a continuous-space Sinai chain, *J. Stat. Phys.* **73**(1/2), 379–388 (1993).
 - [155] A. Ossipov, Scattering approach to Anderson localisation, *Phys. Rev. Lett.* **121**, 076601 (2018).
 - [156] A. Ossipov and Y. V. Fyodorov, Statistics of delay times in mesoscopic systems as a manifestation of eigenfunction fluctuations, *Phys. Rev. B* **71**, 125133 (2005).
 - [157] A. Ossipov, T. Kottos and T. Geisel, Statistical properties of phases and delay times of the one-dimensional Anderson model with one open channel, *Phys. Rev. B* **61**, 11411–11415 (2000).
 - [158] H. M. Pastawski, Classical and quantum transport from generalized Landauer-Büttiker equations. II. Time-dependent resonant tunneling, *Phys. Rev. B* **46**, 4053–4070 (1992).
 - [159] M. H. Pedersen, S. A. van Langen and M. Büttiker, Charge fluctuations in quantum point contacts and chaotic cavities in the presence of transport, *Phys. Rev. B* **57**(3), 1838–1846 (1998).
 - [160] J. B. Pendry, P. D. Kirkman and E. Castano, Electrons at disordered surfaces and $1/f$ noise, *Phys. Rev. Lett.* **57**(23), 2983 (1986).
 - [161] M. L. Polianski and P. W. Brouwer, Scattering matrix ensemble for time-dependent transport through a chaotic quantum dot, *J. Phys. A: Math. Gen.* **36**, 3215–3236 (2003).
 - [162] M. L. Polianski and M. Büttiker, Mesoscopic Fluctuations of Nonlinear Conductance of Chaotic Quantum Dots, *Phys. Rev. Lett.* **96**, 156804 (2006).
 - [163] M. L. Polianski and M. Büttiker, Magnetic-field symmetries of mesoscopic non-linear conductance, *Physica E* **40**(1), 67–75 (2007).

- [164] M. L. Polianski and M. Büttiker, Rectification and nonlinear transport in chaotic dots and rings, *Phys. Rev. B* **76**, 205308 (2007).
- [165] P. Pradhan and N. Kumar, Localization of light in coherently amplifying random media, *Phys. Rev. B* **50**, 9644–9647 (1994).
- [166] A. Prêtre, H. Thomas and M. Büttiker, Dynamic admittance of mesoscopic conductors: Discrete-potential model, *Phys. Rev. B* **54**(11), 8130–8143 (1996).
- [167] X.-L. Qi and S.-C. Zhang, Topological insulators and superconductors, *Rev. Mod. Phys.* **83**, 1057–1110 (2011).
- [168] S. A. Ramakrishna and N. Kumar, Imaginary potential as a counter of delay time for wave reflection from a one-dimensional random potential, *Phys. Rev. B* **61**(5), 3163–3165 (2000).
- [169] S. A. Ramakrishna and N. Kumar, Distribution of the delay time and the dwell time for wave reflection from a long random potential, *Eur. Phys. J. B* **23**(4), 509–513 (2001).
- [170] K. Ramola and C. Texier, Fluctuations of random matrix products and 1D Dirac equation with random mass, *J. Stat. Phys.* **157**(3), 497–514 (2014).
- [171] Z. Ringel, Y. Imry and O. Entin-Wohlman, Delayed currents and interaction effects in mesoscopic capacitors, *Phys. Rev. B* **78**, 165304 (2008).
- [172] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, Topological insulators and superconductors: tenfold way and dimensional hierarchy, *New J. Phys.* **12**, 065010 (2010).
- [173] D. Sánchez and M. Büttiker, Magnetic-Field Asymmetry of Nonlinear Mesoscopic Transport, *Phys. Rev. Lett.* **93**, 106802 (2004).
- [174] D. V. Savin, private communication (2016).
- [175] D. V. Savin, Y. V. Fyodorov and H.-J. Sommers, Reducing nonideal to ideal coupling in random matrix description of chaotic scattering: Application to the time-delay problem, *Phys. Rev. E* **63**, 035202 (2001).
- [176] H. Schomerus, M. Marciani and C. W. J. Beenakker, Effect of Chiral Symmetry on Chaotic Scattering from Majorana Zero Modes, *Phys. Rev. Lett.* **114**, 166803 (2015).
- [177] H. Schomerus and M. Titov, Band-center anomaly of the conductance distribution in one-dimensional Anderson localization, *Phys. Rev. B* **67**, 100201 (2003).
- [178] P. Sebbah, O. Legrand and A. Z. Genack, Fluctuations in photon local delay time and their relation to phase spectra in random media, *Phys. Rev. E* **59**(2), 2406–2411 (1999).
- [179] U. Smilansky, Delay-time distribution in the scattering of time-narrow wave packets (I), *J. Phys. A: Math. Theor.* **50**, 215301 (2017).
- [180] F. T. Smith, Lifetime matrix in collision theory, *Phys. Rev.* **118**(1), 349–356 (1960).
- [181] H.-J. Sommers, D. V. Savin and V. V. Sokolov, Distribution of Proper Delay Times in Quantum Chaotic Scattering: A Crossover from Ideal to Weak Coupling, *Phys. Rev. Lett.* **87**, 094101 (2001).
- [182] S. Souma and A. Suzuki, Local density of states and scattering matrix in quasi-one-dimensional systems, *Phys. Rev. B* **65**, 115307 (2002).
- [183] B. Spivak and A. Zyuzin, Signature of the electron-electron interaction in the magnetic-field dependence of nonlinear I - V characteristics in mesoscopic systems, *Phys. Rev. Lett.* **93**, 226801 (2004).
- [184] M. Steiner, Y. Chen, M. Fabrizio and A. O. Gogolin, Statistical properties of localization-delocalization transition in one dimension, *Phys. Rev. B* **59**(23), 14848–14851 (1999).
- [185] F. H. Stillinger and D. R. Herrick, Bound states in the continuum, *Phys. Rev. A* **11**, 446–454 (1975).
- [186] T. Taniguchi, Charge current density from the scattering matrix, *Phys. Lett. A* **279**, 81–86 (2001).
- [187] T. Taniguchi and M. Büttiker, Friedel phases and phases of transmission amplitudes in quantum scattering systems, *Phys. Rev. B* **60**, 13814–13823 (1999).
- [188] C. Texier, *Quelques aspects du transport quantique dans les systèmes désordonnés de basse dimension*, Ph.D. thesis, Université Paris 6 (1999), http://lptms.u-psud.fr/christophe_texier/ or , <http://tel.archives-ouvertes.fr/tel-01088853>.
- [189] C. Texier, Scattering theory on graphs (2): the Friedel sum rule, *J. Phys. A: Math. Gen.* **35**, 3389–3407 (2002).
- [190] C. Texier, *Désordre, localisation et interaction – Transport quantique dans les réseaux métalliques* (Habilitation à Diriger des Recherches, Université Paris-Sud, 2010), <http://tel.archives-ouvertes.fr/tel-01091550>.
- [191] C. Texier, *Mécanique quantique*, Dunod, Paris, second edition (2015).
- [192] C. Texier and M. Büttiker (1999), unpublished.
- [193] C. Texier and M. Büttiker, Local Friedel sum rule in graphs, *Phys. Rev. B* **67**, 245410 (2003).
- [194] C. Texier and A. Comtet, Universality of the Wigner time delay distribution for one-dimensional random potentials, *Phys. Rev. Lett.* **82**(21), 4220–4223 (1999).
- [195] C. Texier and P. Degiovanni, Charge and current distribution in graphs, *J. Phys. A: Math. Gen.* **36**, 12425–12452 (2003).
- [196] C. Texier and C. Hagendorf, The effect of boundaries on the spectrum of a one-dimensional random mass Dirac Hamiltonian, *J. Phys. A: Math. Theor.* **43**, 025002 (2010).
- [197] C. Texier and S. N. Majumdar, Wigner time-delay distribution in chaotic cavities and freezing transition, *Phys. Rev. Lett.* **110**, 250602 (2013), Erratum: *ibid* **112**, 139902 (2014).
- [198] C. Texier and J. Mitscherling, Nonlinear conductance in weakly disordered mesoscopic wires: Interaction and magnetic field asymmetry, *Phys. Rev. B* **97**, 075306 (2018).
- [199] C. Texier and G. Montambaux, Scattering theory on graphs, *J. Phys. A: Math. Gen.* **34**, 10307–10326 (2001).
- [200] M. Titov, P. W. Brouwer, A. Furusaki and C. Mudry, Fokker-Planck equations and density of states in disordered quantum wires, *Phys. Rev. B* **63**, 235318 (2001).
- [201] M. Titov and Y. V. Fyodorov, Time-delay correlations and resonances in one-dimensional disordered systems, *Phys. Rev. B* **61**(4), R2444 (2000).
- [202] M. Titov and H. Schomerus, Anomalous Wave Function Statistics on a One-Dimensional Lattice with Power-Law Disorder, *Phys. Rev. Lett.* **91**, 176601 (2003).
- [203] T. Y. Tsang and T. A. Osborn, The spectral property of time delay, *Nucl. Phys. A* **247**, 43–50 (1975).
- [204] G. E. Uhlenbeck and E. Beth, The quantum theory of the non-ideal gas I. Deviations from the classical theory, *Physica* **3**(8), 729–745 (1936).

- [205] R. O. Vallejos, A. M. Ozorio de Almeida and C. H. Lewenkopf, Quantum time delay in chaotic scattering: a semiclassical approach, *J. Phys. A: Math. Gen.* **31**(21), 4885–4897 (1998).
- [206] B. A. van Tiggelen, P. Sebbah, M. Stoytchev and A. Z. Genack, Delay-time statistics for diffuse waves, *Phys. Rev. E* **59**, 7166–7172 (1999).
- [207] P. Vivo, private communication (2014).
- [208] J. von Neumann and E. Wigner, Über merkwürdige diskrete Eigenwerte, *Phys. Z.* **30**, 465 (1929).
- [209] P. Šeba, K. Życzkowski and J. Zakrzewski, Statistical properties of random scattering matrices, *Phys. Rev. E* **54**, 2438–2446 (1996).
- [210] E. P. Wigner, Lower limit for the energy derivative of the scattering phase shift, *Phys. Rev.* **98**(1), 145–147 (1955).
- [211] F. Xu and J. Wang, Statistics of Wigner delay time in Anderson disordered systems, *Phys. Rev. B* **84**, 024205 (2011).
- [212] M. Yor, *Exponential functionals of Brownian motion and related processes*, Springer (2000).
- [213] J. M. Ziman, *Principles of the theory of solids*, Cambridge University Press (1972).
- [214] M. R. Zirnbauer, Riemannian symmetric superspaces and their origin in random-matrix theory, *J. Math. Phys.* **37**(10), 4986–5018 (1996).
- [215] D. M. Zumbühl, C. M. Marcus, M. P. Hanson and A. C. Gossard, Asymmetry of Nonlinear Transport and Electron Interactions in Quantum Dots, *Phys. Rev. Lett.* **96**, 206802 (2006).