

Dynamics of Peccei-Quinn Breaking Field after Inflation and Axion Isocurvature Perturbations

Keisuke Harigaya^(a,b), Masahiro Ibe^(a,b),
Masahiro Kawasaki^(a,b) and Tsutomu T. Yanagida^(b)

^(a)*ICRR, University of Tokyo, Kashiwa, Chiba 277-8582, Japan*

^(b)*Kavli IPMU (WPI), UTIAS, University of Tokyo, Kashiwa, Chiba 277-8583, Japan*

The Peccei-Quinn mechanism suffers from the problem of the isocurvature perturbations. The isocurvature perturbations are suppressed if the Peccei-Quinn breaking scale is large during inflation. The oscillation of the Peccei-Quinn breaking field after inflation, however, leads to the formation of domain walls due to the parametric resonance effect. In this paper, we discuss the evolution of the Peccei-Quinn breaking field after inflation in detail, and propose a model where the parametric resonance is ineffective and hence domain walls are not formed. We also discuss consistency of our model with supersymmetric theory.

1 Introduction

The Peccei-Quinn (PQ) mechanism [1] is a compelling solution to the strong CP problem [2, 3, 4]. The mechanism not only solves the strong CP problem, but also predicts a Nambu-Goldstone boson associated with the spontaneous breaking of the PQ symmetry [5, 6], axion, which is a good candidate of dark matter [7, 8, 9].

The PQ mechanism, however, brings about cosmological problems. If the PQ symmetry is restored during inflation, the breaking of the PQ symmetry after inflation leads to formation of domain walls [10]. Unless the domain wall number is unity, the domain walls are stable and hence eventually dominate the energy density of the universe. If the PQ symmetry is broken during inflation, on the other hand, the quantum fluctuations of the axion induce the isocurvature perturbations of cold dark matter [11, 12, 13, 14, 15], which is strictly constrained by observations of the cosmic microwave background (CMB) [16]. This is called the isocurvature perturbation problem. To evade the constraint, the Hubble scale during inflation must be smaller than about 10^7 GeV (see Sec. 2). This strict bound is incompatible with many inflation models including chaotic inflation [17], which is the simplest inflation model without the initial condition problem and predicts a large Hubble scale during inflation, $H_{\text{inf}} \sim \text{several} \times 10^{13}$ GeV.

As was first pointed in [15] (see also [18, 19, 20, 21]), the isocurvature perturbations are suppressed if the PQ breaking scale is large during inflation. Due to the large PQ breaking scale, the fluctuations of the misalignment angle are suppressed and hence the axion density perturbations are also suppressed.¹ For this suppression mechanism to work, the large expectation value of the PQ breaking field should decrease to the present value after inflation. However, the relaxation of the PQ breaking scale may lead to the restoration of the PQ symmetry in the following way. After inflation, the PQ breaking field oscillates around the origin, which produces large fluctuations of the PQ breaking field through the parametric resonance effect [25]. The large fluctuations lead to the restoration of the PQ symmetry and hence the formation of domain walls [26, 27, 28, 29, 30]. To prevent the formation of domain walls, the PQ breaking scale during inflation cannot be arbitrary large. The recent study with lattice simulations [30] shows that the upper bound on the Hubble scale from the isocurvature perturbations is raised to at most 10^{12} GeV when the axion is the dominant component of dark matter.

In this paper, we discuss the evolution of the PQ breaking field after inflation in detail, and propose a model where the parametric resonance is ineffective and hence domain walls are not formed. Thus, the model allows for both axion dark matter and inflation models with large Hubble scale. We also discuss consistency of our model with supersymmetric theory.

This paper is organized as follows. In Sec. 2, we review the isocurvature perturbation problem of the PQ mechanism and suppression of isocurvature perturbations by a large PQ breaking scale during inflation. In Sec. 3, we discuss the evolution of the PQ breaking field and restoration of PQ symmetry after inflation. We propose a model of the PQ symmetry breaking without the restoration of the PQ symmetry. The final section is

¹ See Refs. [22, 23, 24] for other possible solutions to the isocurvature perturbation problem.

devoted to summary and discussion.

2 Isocurvature perturbation problem and a large PQ breaking scale

In this section, we review the problem of isocurvature perturbations in the PQ mechanism and suppression of their amplitude by a large scale of the PQ breaking during inflation.

2.1 QCD axion as a dark matter candidate

The axion a is the Nambu-Goldstone boson associated with the spontaneous breaking of the PQ symmetry [1, 5, 6]. It is assumed that the PQ symmetry has anomaly of QCD. Then the axion obtains its potential by non-perturbative QCD dynamics,

$$V(a) \sim m_\pi^2 f_\pi^2 (1 - \cos\theta_a), \quad \theta_a \equiv a/f_a, \quad (1)$$

where m_π , f_π and f_a are the mass of the pion, the pion decay constant, and the axion decay constant. Here, we have shifted the axion so that its vacuum expectation value (VEV) is zero.

In this paper, we assume that the PQ symmetry is spontaneously broken during inflation to avoid the domain wall problem. Then in the early universe, the axion in general has a non-zero field value a_i . After inflation, around a temperature of the QCD scale, the axion begins to oscillate and behaves as cold dark matter. The present density of the axion is given by [31]

$$\Omega_a h^2 = 0.2 \times \theta_i^2 \left(\frac{f_a}{10^{12} \text{ GeV}} \right)^{1.19}, \quad (2)$$

where $\theta_i \equiv a_i/f_a$ is the initial misalignment angle. For $f_a = 10^{11-12}$ GeV, the density of the axion oscillation is as large as that of dark matter.

2.2 Isocurvature perturbation

The initial axion field value a_i takes almost the same value in the whole observable universe. However, quantum effect during inflation inevitably induces fluctuations of the initial field value, $\delta a_i \sim H/(2\pi)$ [32, 33, 34, 35, 36]. Taking into account the fluctuations, the density parameter of the axion is modified as

$$\begin{aligned} \Omega_a h^2 &= 0.2 \times \theta_{i,\text{eff}}^2 \left(\frac{f_a}{10^{12} \text{ GeV}} \right)^{1.19}, \\ \theta_{i,\text{eff}}^2 &\equiv \theta_i^2 + \delta\theta_i^2 = \theta_i^2 + \left(\frac{H_{\text{inf}}}{2\pi f_a} \right)^2. \end{aligned} \quad (3)$$

The axion fluctuations are independent of those of the inflaton field in the flat time slice. Thus, the fluctuations of the axion result in un-correlated isocurvature perturbations of cold dark matter [11, 12, 13, 14, 15]. The power spectrum of the isocurvature perturbations is given by

$$\mathcal{P}_{S_c} = \left(\frac{\delta\Omega_a}{\Omega_c} \right)^2 = \frac{4}{\theta_{i,\text{eff}}^2} (\delta\theta_i)^2 \left(\frac{\Omega_a}{\Omega_c} \right)^2 = \frac{4}{\theta_{i,\text{eff}}^2} \left(\frac{H_{\text{inf}}}{2\pi f_a} \right)^2 \left(\frac{\Omega_a}{\Omega_c} \right)^2, \quad (4)$$

where Ω_c is the density parameter of cold dark matter.

Uncorrelated isocurvature perturbations of cold dark matter are constrained as $\alpha_c \equiv \mathcal{P}_{S_c}/(\mathcal{P}_\zeta + \mathcal{P}_{S_c}) < 0.033$, where $\mathcal{P}_\zeta \simeq 2 \times 10^{-9}$ is the power spectrum of the curvature perturbations [16]. In Figure 1, we show the bound on the axion decay constant f_a and the Hubble scale during inflation H_{inf} for $\theta_i = 1, 0.1$ and 0.01 . Shaded regions are excluded by the constraint from the isocurvature perturbations or too large cosmic density of the axion. We do not show the parameter region with $f_a < 10^9$ GeV because astrophysical constraints exclude that region. The Hubble scale during inflation must be smaller than about 10^7 GeV for $\theta_i = O(1)$, which severely constrains inflation models. For example, the simplest inflation model without the initial condition problem, namely chaotic inflation [17], predicts a large Hubble scale $H_{\text{inf}} \sim \text{several} \times 10^{13}$ GeV.




Figure 1: Bound on the axion decay constant f_a and the Hubble scale during inflation H_{inf} for $\theta_i = 1, 0.1$ and 0.01 . Shaded regions are excluded by the constraint on the isocurvature perturbations, $\alpha_c < 0.033$, or by too large cosmic density of the axion.

2.3 Large PQ breaking field value during inflation

The isocurvature perturbations can be suppressed if the PQ breaking scale is larger than the present one during inflation [15]. Suppose that the PQ breaking field P takes a large field value P_{inf} during inflation, which we take to be real by a PQ rotation. The large field value is explained by a negative Hubble induced mass of the PQ breaking field during inflation. Then the decay constant of the axion during inflation f_{inf} is given by

$$f_{\text{inf}} = \sqrt{2}P_{\text{inf}}/N_{\text{DW}}, \quad (5)$$

where N_{DW} is the domain wall number of the axion model. Then the fluctuation of the initial misalignment angle is written as

$$\delta\theta_i = \frac{H_{\text{inf}}}{2\pi f_{\text{inf}}} = \frac{H_{\text{inf}}}{2\pi\sqrt{2}P_{\text{inf}}} \times N_{\text{DW}}, \quad (6)$$

which leads to the power spectrum of the isocurvature perturbations as

$$\mathcal{P}_{S_c} = \frac{4}{\theta_{i,\text{eff}}^2} \left(\frac{N_{\text{DW}}H_{\text{inf}}}{2\pi\sqrt{2}P_{\text{inf}}} \right)^2 \left(\frac{\Omega_a}{\Omega_c} \right)^2. \quad (7)$$

In Figure 2, we show the bound on the axion decay constant f_a and the initial misalignment angle θ_i for $N_{\text{DW}} = 6$ (DFSZ model [37, 38]), $H_{\text{inf}} = 5 \times 10^{13}$ GeV (tensor-to-scalar ratio $r \simeq 0.04$) and $P_{\text{inf}} = M_{\text{Pl}}$ as well as $2M_{\text{Pl}}$. Even if the Hubble scale during inflation is such large, for $f_a \sim 10^{11}$ GeV and $\theta_i = O(1)$, the axion can be a sizable component of dark matter while the bound from the isocurvature perturbations is avoided. It should be remarked that the Hubble scale during inflation can be determined by measuring the tensor-to-scalar ratio r , whose present bound is $r \lesssim 0.1$ [16]. Future ground based experiments can detect $r \gtrsim 0.003$ [39], which corresponds to $H_{\text{inf}} \gtrsim 10^{13}$ GeV. Therefore, it is important to examine whether an axion model is consistent with high scale inflation with $H_{\text{inf}} \gtrsim 10^{13}$ GeV.

3 Evolution of PQ breaking field

In the previous section, we have shown that the bound from the isocurvature perturbations is evaded if the field value of the PQ breaking field is large during inflation. In this section, we discuss the evolution of the PQ breaking field after inflation in detail. We show that the oscillation of the PQ breaking field driven by a quartic potential leads to the formation of domain walls. Then we propose a model where domain walls are not formed. We also discuss the embedding of the model to supersymmetric theory.

3.1 General discussion

Let us discuss the dynamics of the PQ breaking field analytically, following Ref. [40]. We assume the following potential of the PQ breaking field P and the inflaton ϕ ,

$$V(\phi, P) = V_{\text{inf}}(\phi) + \lambda_{2n}^2 |P|^{2n} - \frac{cH}{3} V_{\text{inf}}(\phi) |P|^2, \quad (8)$$




Figure 2: Bound on the axion decay constant f_a and the initial misalignment angle θ_i for $N_{DW} = 6$, $H_{\text{inf}} = 5 \times 10^{13}$ GeV. The shaded region with a solid (dashed) line boundary shows the constraint from the isocurvature perturbation for $P_{\text{inf}} = (2)M_{\text{Pl}}$.

where $V_{\text{inf}}(\phi)$ is the potential of the inflaton, and λ_{2n} as well as $c_H (> 0)$ are constants.² Here, we neglect the mass term of the PQ breaking field, which is important only when the field value of the PQ breaking field becomes sufficiently small. With this potential, the PQ breaking field obtains a large field value during inflation,

$$P_{\text{inf}} = \left(\frac{c_H H_{\text{inf}}^2}{n \lambda_{2n}^2} \right)^{\frac{1}{2(n-1)}}, \quad (9)$$

where H_{inf} is the Hubble scale during inflation.

After inflation, the inflaton begins oscillation around the origin. During the oscillatory phase, the PQ breaking field may also begin oscillation around the origin. The equation of motion of the radial direction of P , $\chi = |P|/\sqrt{2}$, is

$$\ddot{\chi} + 3H\dot{\chi} + \frac{2n\lambda_{2n}^2}{2^n} \chi^{2n-1} - \frac{c_H}{2} H^2 \chi = 0, \quad (10)$$

where H is the Hubble scale. Here, we replace $V_{\text{inf}}(\phi)$ with its time averaged value, $3H^2/2$, which is a good approximation as long as the mass of the inflaton is much larger than the Hubble scale.

² Since the inflaton potential $V_{\text{inf}}(\phi)$ is neutral under any symmetries, it is natural that the interaction between the inflaton and the PQ breaking field in Eq. (8) is present. This is also the case even in supersymmetry theory.

It is convenient to use the number of e-folding $N \equiv \ln(R/R_i)$ as a time variable, where R is the scale factor and R_i is its initial value when the inflaton starts oscillation. In the oscillatory phase of the inflaton, namely in a matter dominant universe, the equation of motion of χ is then given by

$$\frac{d^2}{dN^2}\chi + \frac{3}{2}\frac{d}{dN}\chi + \frac{2n\lambda_{2n}^2}{2^n H_i^2}\chi^{2n-1} - \frac{c_H}{2}\chi = 0. \quad (11)$$

We further rewrite the equation of motion by $\psi \equiv \chi \exp(\frac{3N}{2(n-1)})$,

$$\frac{d^2}{dN^2}\psi + \frac{3(n-3)}{2(n-1)}\frac{d}{dN}\psi - \left(\frac{9(n-2)}{4(n-1)^2} + \frac{c_H}{2}\right)\psi + \frac{2n\lambda_{2n}^2}{2^n H_i^2}\psi^{2n-1} = 0, \quad (12)$$

where H_i is the Hubble scale when the inflaton starts oscillation. Finally, we normalize the field ψ so that the initial value of the normalized field is of $O(1)$,

$$\begin{aligned} \frac{d^2}{dN^2}\rho + \frac{3(n-3)}{2(n-1)}\frac{d}{dN}\rho - \left(\frac{9(n-2)}{4(n-1)^2} + \frac{c_H}{2}\right)\rho + \frac{c_H}{2}\rho^{2n-1} = 0, \\ \psi \equiv \left(\frac{2^n c_H H_i^2}{4n\lambda_{2n}^2}\right)^{\frac{1}{2n-2}} \rho. \end{aligned} \quad (13)$$

Note that all coefficients in Eq. (13) is N -independent.

The equation of motion of ρ is nothing but that of a particle with a potential

$$V(\rho) = -\frac{1}{2}\left(\frac{9(n-2)}{4(n-1)^2} + \frac{c_H}{2}\right)\rho^2 + \frac{c_H}{4n}\rho^{2n}, \quad (14)$$

whose minimum is at

$$\rho = \left(1 + \frac{9(n-2)}{2(n-1)^2 c_H}\right)^{\frac{1}{2n-2}} \equiv \rho_0. \quad (15)$$

The initial value of ρ , ρ_i , when the inflaton starts oscillation is expected be of $O(1)$. Its definite value, however, depends on the dynamics of the inflaton between the inflationary phase and the oscillatory phase.

For $n = 2$, the friction term has a wrong sign, and hence the field value of ρ grows and eventually starts oscillation around the origin. We will cross-check this conclusion numerically later. Due to the oscillation, the parametric resonance effect grows the fluctuations of the PQ breaking field exponentially in time [25]. The PQ field with large initial field value oscillates for a long time until it settles down to the minimum of the potential. Thus, the oscillation produces large fluctuations, which eventually leads to the restoration of the PQ symmetry and hence the formation of domain walls [26, 27, 28, 29, 30].

For $n = 3$, the friction term is absent and hence ρ oscillates continuously without attenuation. The potential of ρ is depicted in Figure 3. If ρ_i is within the range $|\rho_i| < \rho_1 \equiv [(27/8)c_H^{-1} + 3]^{1/4}$, ρ oscillates around ρ_0 . If $|\rho_i| > \rho_1$, ρ oscillates around the origin.

In both cases, the PQ breaking field oscillation might produce fluctuations.³ So let us consider the evolution of the fluctuations of the PQ breaking field around the zero mode for $n = 3$. The evolution is described by

$$\frac{d^2}{dN^2}\delta\rho_k + \frac{k^2}{H_i^2}e^N\delta\rho_k - \left(\frac{9}{16} + \frac{c_H}{2}\right)\delta\rho_k + \frac{5}{2}c_H\rho^4\delta\rho_k = 0, \quad (16)$$

where k is a comoving wave number of the fluctuation $\delta\rho_k$. The strength of the parametric resonance is expected to be estimated by the magnitude of the coefficient $5c_H\rho^4/2$, in analogy with the q parameter of the Mathieu's equation. In the present case, the magnitude is of $O(1)$. It is expected that resonance bands are not broad. Since effective wave numbers evolve in proportional to $\exp(N/2)$ [see, the 2nd term in Eq. (16)], given modes are in resonance bands at most during one oscillation. Thus the parametric resonance is considered to be ineffective. We have confirmed this by numerically solving the evolution equation.

For $n > 3$, the friction term has a correct sign. Thus, ρ quickly relaxes to the minimum ρ_0 . The PQ symmetry is not restored.




Figure 3: *The potential of ρ .*

3.2 More on Quartic potential

Let us discuss the case with $n = 2$, i.e. a quartic potential in detail, including the evolution of the inflaton from the inflationary phase to the oscillatory phase. To be definite, we assume the inflaton potential around the origin,

$$V_{\text{inf}}(\phi) = \frac{1}{2}m^2\phi^2, \quad (17)$$

³ See also Ref. [19] for discussion on the parametric resonance in a sextet potential, although the oscillation of the PQ breaking field is not taken into account.

where ϕ is the inflaton and m is its mass. Then the equations of motion of the inflaton and the PQ breaking field are given by

$$\ddot{\phi} + 3H\dot{\phi} + m^2\phi = 0, \quad (18)$$

$$\ddot{\chi} + 3H\dot{\chi} + \lambda_4^2\chi^3 - \frac{c_H}{3}m^2\phi^2\chi = 0. \quad (19)$$

Here, we neglect the back reaction of the evolution of the PQ breaking field on the dynamics of the inflaton.

We rewrite the equation of motion of χ to a more convenient form. For an initial condition of $\phi = \phi_0$, the field value of χ determined by Eq. (9) is

$$\chi_0 = \sqrt{\frac{c_H}{6} \frac{m\phi_0}{\lambda_4}}. \quad (20)$$

Then the equation of motion of $r \equiv \chi/\chi_0$ is

$$\ddot{r} + 3H\dot{r} + \frac{1}{6}c_H m^2 \phi_0^2 r^3 - \frac{c_H}{3} m^2 \phi^2 r = 0, \quad (21)$$

with the initial condition $r_0 = 1$. One can see that the dynamics of $r = \chi/\chi_0$ does not depend on λ_4 .

In Figure 4, we show the evolution of the PQ breaking field for $m = 10^{-5}M_{\text{Pl}}$, $c_H = 1$ with the unit system where the reduced Planck scale is unity. Here, we take the initial condition $\phi_0 = M_{\text{Pl}}$. It can be seen that the PQ breaking field starts oscillation around the origin, which is consistent with our analytical estimation.




Figure 4: The evolution of the PQ breaking field after inflation for the quartic potential. We also show the evolution of the inflaton.

As the PQ breaking field oscillates with the quartic potential, the oscillation produces the fluctuations of the PQ breaking field by the parametric resonance effect [25]. If the

oscillation lasts long, the fluctuations eventually restore the PQ symmetry and hence domain walls are formed [26, 27, 28, 29, 30]. In order to prevent the formation of domain walls, the field value of the PQ breaking field when it start oscillation, P_{osc} , must satisfy [30]

$$P_{\text{osc}} < 10^4 \langle P \rangle = 10^{16} \text{ GeV} \times \frac{\langle P \rangle}{10^{12} \text{ GeV}}, \quad (22)$$

where $\langle P \rangle (= N_{\text{DW}} f_a / \sqrt{2})$ is the present vacuum expectation value.

In Figure 5, we show the constraint on the axion decay constant f_a and the Hubble scale during inflation H_{inf} when the field value of the PQ breaking field during inflation P_{inf} is as large as $\min(10^4 \langle P \rangle, M_{\text{Pl}})$. Here, we require that $P_{\text{inf}} < M_{\text{Pl}}$ so that the PQ breaking field does not dominate the potential energy during inflation. For $\theta_i = 1$, the Hubble scale during inflation must be smaller than 10^{12} GeV. The constraint is more stringent ($H_{\text{inf}} < 2 \times 10^{11}$ GeV) if we require that axion is dark matter. For $\theta_i = 0.1$, a larger Hubble scale is allowed due to smaller abundance of the axion. For $\theta_i = 0.1$, we also show the upper bound on the Hubble scale with $P_{\text{inf}} = \min(10 \times 10^4 \langle P \rangle, M_{\text{Pl}})$ by a dashed line. We multiply the factor of 10 to take into account a larger Hubble scale when the CMB scale exits the horizon than that at the end of the inflation by a factor of about 10 in typical large field inflation models. (Thus, this loose upper bound should not be applied to small field models.) In this case the Hubble scale close to 10^{14} GeV is allowed although $H_{\text{inf}} < 10^{12}$ GeV is necessary for axion to account for dark matter. Therefore, even if we take small θ_i and adopt the loose bound, the Hubble scale during inflation should be less than 10^{12} GeV when the axion is a dominant component of dark matter.




Figure 5: The constraint on the axion decay constant f_a and the Hubble scale during inflation H_{inf} for the quartic potential. Here, we take $P_{\text{inf}} = \min(10^4 \langle P \rangle, M_{\text{Pl}})$. The dashed line in the right panel shows the upper bound on H_{inf} for $P_{\text{inf}} = \min(10 \times 10^4 \langle P \rangle, M_{\text{Pl}})$.

Let us discuss whether the beginning of the oscillation around the origin can be delayed

so that $P_{\text{osc}} \ll P_{\text{inf}}$. The reason why the PQ breaking field starts oscillation is that $\rho_i \neq \rho_0$ [see Sec. 3.1] due to the rapid change of the Hubble induced mass term of the PQ breaking field after inflation. To suppress the oscillation, we introduce a coupling between the PQ breaking field and the kinetic term of the inflaton,

$$\mathcal{L} = \frac{d}{2} \partial^\mu \phi \partial_\mu \phi \times |P|^2. \quad (23)$$

If $d \simeq c_H/3$, the Hubble induced mass term varies after inflation slowly and hence we may obtain $\rho_i \simeq \rho_0$. We further assume that the Hubble induced mass is large, which may help the PQ breaking field to follow the point ρ_0 adiabatically [41, 42]. In Figure 6, we show the evolution of the PQ breaking field for $m = 10^{-5} M_{\text{Pl}}$, $c_H = 16\pi^2$ and $d = c_H/3$. Still, the PQ breaking field starts oscillation around the origin when $\chi \sim 10^{-1} \chi_0$. Thus, the oscillation is inevitable for a quartic potential.⁴




Figure 6: The similar figure as Figure 4 but with a coupling of the PQ breaking field to the kinetic term of the inflaton.

3.3 Sextet and quartic potential

As we have discussed in Sec. 3.1, the growth of fluctuations is avoided if the oscillation of the PQ breaking field after inflation is driven by a sextet potential. We assume the following potential of the PQ breaking field P ,

$$V(P) = -m_P^2 |P|^2 + \lambda_4^2 |P|^4 + \lambda_6^2 |P|^6, \quad (24)$$

⁴ We find that a larger Hubble induced mass allows for smaller P_{osc} . However, in that case, the PQ breaking field value during inflation is required to be small so that the PQ breaking field does not dominate the potential energy during inflation.

where m_P^2 and, λ_4 and λ_6 are constants. We again assume the interaction with the inflaton in Eq. (8). Then the PQ breaking field obtains a large field value during inflation. We assume that λ_6 is sufficiently large so that the field value of the PQ breaking field during inflation is determined by a balance between the sextet term and the Hubble induced mass as

$$P_{\text{inf}} = \frac{c_H^{1/4} H_{\text{inf}}^{1/2}}{3^{1/4} \lambda_6^{1/2}} = M_{\text{Pl}} c_H^{1/4} \left(\frac{H_{\text{inf}}}{10^{14} \text{GeV}} \right)^{1/2} \left(\frac{2 \times 10^{-5} M_{\text{Pl}}^{-1}}{\lambda_6} \right)^{1/2}. \quad (25)$$

After inflation, the PQ breaking field starts oscillation driven by the sextet term. During the oscillation by the sextet term, the parametric resonance effect is ineffective and hence the PQ symmetry is not restored. The oscillation of the PQ breaking field is eventually driven by the quartic term when it dominates over the sextet term at $P = P_{\text{osc-4}}$,

$$P_{\text{osc-4}} = \frac{\lambda_4}{\lambda_6} = 10^{16} \text{ GeV} \times \frac{\lambda_4}{7 \times 10^{-8}} \frac{2 \times 10^{-5} M_{\text{Pl}}^{-1}}{\lambda_6}. \quad (26)$$

As long as $P_{\text{osc-4}} < 10^4 \langle P \rangle$, the PQ symmetry is not restored and hence domain walls are not formed. Therefore, this model can solve both isocurvature and domain wall problems for high scale inflation models.

Before closing this subsection, let us discuss the VEV of the PQ breaking field $\langle P \rangle$. If λ_4 is sufficiently small, the VEV is determined by a balance between the sextet and the quadratic terms. If not, the VEV is determined by a balance between the quartic and the quadratic terms;

$$\langle P \rangle = \begin{cases} \frac{m_P^{1/2}}{3^{1/4} \lambda_6^{1/2}} = 1 \times 10^{12} \text{GeV} \times \left(\frac{2 \times 10^{-5} M_{\text{Pl}}^{-1}}{\lambda_6} \right)^{1/2} \left(\frac{m_P}{20 \text{GeV}} \right)^{1/2} & (\lambda_4 < \sqrt{\lambda_6 m_P}), \\ \frac{1}{\sqrt{2} \lambda_4} m_P = 1 \times 10^{12} \text{GeV} \times \frac{7 \times 10^{-8}}{\lambda_4} \frac{m_P}{10^5 \text{GeV}} & (\lambda_4 > \sqrt{\lambda_6 m_P}). \end{cases} \quad (27)$$

In the former case, the mass of the PQ breaking field, m_P , is given by P_{inf} , $\langle P \rangle$ and H_{inf} as

$$m_P = c_H^{1/2} H_{\text{inf}} \frac{\langle P \rangle^2}{P_{\text{inf}}^2} = 10 \text{GeV} \times c_H^{1/2} \frac{H_{\text{inf}}}{10^{14} \text{GeV}} \left(\frac{\langle P \rangle}{10^{12} \text{GeV}} \right)^2 \left(\frac{M_{\text{Pl}}}{P_{\text{inf}}} \right)^2. \quad (28)$$

In the latter case, expressing λ_4 and λ_6 by $\langle P \rangle$ and P_{inf} respectively, and put the constraints $\lambda_4^2 > \lambda_6 m_P$ and $P_{\text{osc-4}} < 10^4 \langle P \rangle$, we obtain the bound on m_P ,⁵

$$\begin{aligned} 10 \text{ GeV} \times c_H^{1/2} \left(\frac{\langle P \rangle}{10^{12} \text{GeV}} \right)^2 \frac{H_{\text{inf}}}{10^{14} \text{GeV}} \times \left(\frac{M_{\text{Pl}}}{P_{\text{inf}}} \right)^2 &< m_P \\ &< 10^5 \text{ GeV} \times c_H^{1/2} \left(\frac{\langle P \rangle}{10^{12} \text{GeV}} \right)^2 \frac{H_{\text{inf}}}{10^{14} \text{GeV}} \times \left(\frac{M_{\text{Pl}}}{P_{\text{inf}}} \right)^2. \end{aligned} \quad (29)$$

⁵ If one adds a potential with a larger power than six instead of the sextet potential, the upper bound of m_P becomes severer.

3.4 Embedding in supersymmetric theory

Let us discuss the embedding of our model in supersymmetric theory. We introduce three chiral multiplets P , X and Y , whose PQ charges are 1, -2 and -3 , respectively. Then the superpotential of them is given by

$$W = \lambda_4 X P^2 + \lambda_6 Y P^3, \quad (30)$$

where λ_4 and λ_6 are constants. The F terms of X and Y yield the quartic and the sextet terms of P in Eq. (24), respectively. m_P^2 is provided by a soft supersymmetry breaking mass term of P . Assuming that the Hubble induced mass terms as well as soft supersymmetry breaking mass terms of X and Y are positive, they are fixed to their origin during and after inflation. Then, we obtain the potential in Eq. (24).

It should be noted that the allowed range of the mass of the PQ breaking field in Eqs. (28) and (29) is consistent with supersymmetry breaking and its mediation by Planck-suppressed interactions. With the mediation by Planck-suppressed interactions, the fermion superpartner of axion, axino, also has mass of order m_P . If it is known that the axino with mass less than $O(10^4)$ GeV, causes cosmological difficulties [43]. However, for $m_P \simeq 10^5$ GeV which is allowed for large λ_4 [Eq.(29)], neither the saxion nor the axino brings about cosmological problems.

Finally, we note that soft scalar masses of $\sim 10^5$ GeV fit in well with high scale supersymmetry breaking scenarios [44, 45, 46, 47, 48]. There, the observed Higgs mass of 125 GeV [49, 50] is explained by large quantum corrections from scalar tops with masses of 10^5 GeV [51, 52, 53]. Since anomaly mediation generates gaugino masses as large as 1 TeV [44, 54] (see also Refs. [55]), the supersymmetry breaking field does not have to be a singlet field. Then the Polonyi problem [56, 57] is absent. For the gravitino mass is as large as 10^5 GeV, the gravitino decays before the Big-Bang-Nucleosynthesis begins. Thus, our model of PQ symmetry breaking, combined with high scale supersymmetry breaking scenarios, is free from cosmological problems.

4 Summary and discussion

In this paper, we have discussed the solution to the isocurvature perturbation problem of the PQ mechanism by a large PQ breaking scale during inflation. We have investigated the evolution of the PQ breaking field after inflation and discussed whether the PQ symmetry restores after inflation, which leads to the formation of domain walls. We have proposed a model without the restoration of the PQ symmetry. Interestingly, the predicted mass of the PQ breaking field is not very far from the electroweak scale. Thus, our model is compatible with a supersymmetric theory where the mass of the PQ breaking field is given by supersymmetry breaking.

As we have discussed in Sec. 2, the field value of the PQ breaking field must be as large as the Planck scale, if the Hubble scale during inflation is of $O(10^{13})$ GeV. It is known that a field which obtains a Planck scale field value by a Hubble induced mass affects the prediction of large field inflation models [58]. With this effect, the chaotic

inflation model simply driven by a mass term of the inflaton is consistent with recent observations of the CMB [16].

Acknowledgements

This work is supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT), Japan, No. 24740151 and No. 25105011 (M. I.), No. 25400248 (M. K.) as well as No. 26104009 (T. T. Y.); Grant-in-Aid No. 26287039 (M. I. and T. T. Y.) from the Japan Society for the Promotion of Science (JSPS); and by the World Premier International Research Center Initiative (WPI), MEXT, Japan. K.H. is supported in part by a JSPS Research Fellowship for Young Scientists.

References

- [1] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977); R. D. Peccei and H. R. Quinn, Phys. Rev. D **16**, 1791 (1977).
- [2] G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976).
- [3] R. Jackiw and C. Rebbi, Phys. Rev. Lett. **37**, 172 (1976).
- [4] C. G. Callan, Jr., R. F. Dashen and D. J. Gross, Phys. Lett. B **63**, 334 (1976).
- [5] S. Weinberg, Phys. Rev. Lett. **40**, 223 (1978).
- [6] F. Wilczek, Phys. Rev. Lett. **40**, 279 (1978).
- [7] J. Preskill, M. B. Wise and F. Wilczek, Phys. Lett. B **120**, 127 (1983).
- [8] L. F. Abbott and P. Sikivie, Phys. Lett. B **120**, 133 (1983).
- [9] M. Dine and W. Fischler, Phys. Lett. B **120**, 137 (1983).
- [10] P. Sikivie, Phys. Rev. Lett. **48**, 1156 (1982).
- [11] A. D. Linde, JETP Lett. **40**, 1333 (1984) [Pisma Zh. Eksp. Teor. Fiz. **40**, 496 (1984)]; Phys. Lett. B **158**, 375 (1985).
- [12] D. Seckel and M. S. Turner, Phys. Rev. D **32**, 3178 (1985).
- [13] D. H. Lyth, Phys. Lett. B **236**, 408 (1990); Phys. Rev. D **45**, 3394 (1992).
- [14] M. S. Turner and F. Wilczek, Phys. Rev. Lett. **66**, 5 (1991).
- [15] A. D. Linde, Phys. Lett. B **259**, 38 (1991).
- [16] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.02114 [astro-ph.CO].
- [17] A. D. Linde, Phys. Lett. B **129**, 177 (1983).
- [18] K. Choi, K. S. Jeong and M. S. Seo, JHEP **1407**, 092 (2014) [arXiv:1404.3880 [hep-th]].
- [19] E. J. Chun, Phys. Lett. B **735**, 164 (2014) [arXiv:1404.4284 [hep-ph]].

- [20] M. Fairbairn, R. Hogan and D. J. E. Marsh, *Phys. Rev. D* **91**, no. 2, 023509 (2015) [arXiv:1410.1752 [hep-ph]].
- [21] K. Nakayama and M. Takimoto, arXiv:1505.02119 [hep-ph].
- [22] T. Higaki, K. S. Jeong and F. Takahashi, *Phys. Lett. B* **734**, 21 (2014) [arXiv:1403.4186 [hep-ph]].
- [23] N. Kitajima and F. Takahashi, *JCAP* **1501**, no. 01, 032 (2015) [arXiv:1411.2011 [hep-ph]].
- [24] M. Kawasaki, M. Yamada and T. T. Yanagida, arXiv:1506.05214 [hep-ph].
- [25] L. Kofman, A. D. Linde and A. A. Starobinsky, *Phys. Rev. Lett.* **73**, 3195 (1994) [hep-th/9405187]; L. Kofman, A. D. Linde and A. A. Starobinsky, *Phys. Rev. D* **56**, 3258 (1997) [hep-ph/9704452].
- [26] I. I. Tkachev, *Phys. Lett. B* **376**, 35 (1996) [hep-th/9510146].
- [27] S. Kasuya, M. Kawasaki and T. Yanagida, *Phys. Lett. B* **409**, 94 (1997) [hep-ph/9608405]; *Phys. Lett. B* **415**, 117 (1997) [hep-ph/9709202].
- [28] S. Kasuya and M. Kawasaki, *Phys. Rev. D* **56**, 7597 (1997) [hep-ph/9703354]; *Phys. Rev. D* **58**, 083516 (1998) [hep-ph/9804429]; *Phys. Rev. D* **61**, 083510 (2000) [hep-ph/9903324].
- [29] I. Tkachev, S. Khlebnikov, L. Kofman and A. D. Linde, *Phys. Lett. B* **440**, 262 (1998) [hep-ph/9805209].
- [30] M. Kawasaki, T. T. Yanagida and K. Yoshino, *JCAP* **1311**, 030 (2013) [arXiv:1305.5338 [hep-ph]].
- [31] K. J. Bae, J. H. Huh and J. E. Kim, *JCAP* **0809**, 005 (2008) [arXiv:0806.0497 [hep-ph]].
- [32] V. F. Mukhanov and G. V. Chibisov, *JETP Lett.* **33**, 532 (1981) [*Pisma Zh. Eksp. Teor. Fiz.* **33**, 549 (1981)];
- [33] S. W. Hawking, *Phys. Lett. B* **115**, 295 (1982).
- [34] A. A. Starobinsky, *Phys. Lett. B* **117**, 175 (1982).
- [35] A. H. Guth and S. Y. Pi, *Phys. Rev. Lett.* **49**, 1110 (1982).
- [36] J. M. Bardeen, P. J. Steinhardt and M. S. Turner, *Phys. Rev. D* **28**, 679 (1983).
- [37] M. Dine, W. Fischler and M. Srednicki, *Phys. Lett. B* **104**, 199 (1981).
- [38] A. R. Zhitnitsky, *Sov. J. Nucl. Phys.* **31**, 260 (1980) [*Yad. Fiz.* **31**, 497 (1980)].
- [39] P. Creminelli, D. L. Nacir, M. Simonovic, G. Trevisan and M. Zaldarriaga, arXiv:1502.01983 [astro-ph.CO].
- [40] M. Dine, L. Randall and S. D. Thomas, *Nucl. Phys. B* **458**, 291 (1996) [hep-ph/9507453].
- [41] A. D. Linde, *Phys. Rev. D* **53**, 4129 (1996) [hep-th/9601083].
- [42] K. Nakayama, F. Takahashi and T. T. Yanagida, *Phys. Rev. D* **84**, 123523 (2011) [arXiv:1109.2073 [hep-ph]].

- [43] M. Kawasaki, K. Nakayama and M. Senami, JCAP **0803**, 009 (2008) [arXiv:0711.3083 [hep-ph]].
- [44] G. F. Giudice, M. A. Luty, H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998) [hep-ph/9810442];
- [45] J. D. Wells, Phys. Rev. D **71**, 015013 (2005) [hep-ph/0411041].
- [46] M. Ibe, T. Moroi and T. T. Yanagida, Phys. Lett. B **644**, 355 (2007) [hep-ph/0610277]; M. Ibe and T. T. Yanagida, Phys. Lett. B **709**, 374 (2012) [arXiv:1112.2462 [hep-ph]];
- [47] L. J. Hall and Y. Nomura, JHEP **1201**, 082 (2012) [arXiv:1111.4519 [hep-ph]];
- [48] N. Arkani-Hamed, A. Gupta, D. E. Kaplan, N. Weiner and T. Zorawski, arXiv:1212.6971 [hep-ph].
- [49] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012) [arXiv:1207.7214 [hep-ex]]; Phys. Rev. D **90**, no. 5, 052004 (2014) [arXiv:1406.3827 [hep-ex]].
- [50] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012) [arXiv:1207.7235 [hep-ex]]; Phys. Rev. D **89**, no. 9, 092007 (2014) [arXiv:1312.5353 [hep-ex]]; V. Khachatryan *et al.* [CMS Collaboration], Eur. Phys. J. C **74**, no. 10, 3076 (2014) [arXiv:1407.0558 [hep-ex]].
- [51] Y. Okada, M. Yamaguchi, T. Yanagida, Prog. Theor. Phys. **85** (1991) 1-6.
- [52] J. R. Ellis, G. Ridolfi, F. Zwirner, Phys. Lett. **B257** (1991) 83-91.
- [53] H. E. Haber, R. Hempfling, Phys. Rev. Lett. **66** (1991) 1815-1818.
- [54] L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999) [hep-th/9810155];
- [55] M. Dine and D. MacIntire, Phys. Rev. D **46**, 2594 (1992) [hep-ph/9205227]; J. A. Bagger, T. Moroi and E. Poppitz, JHEP **0004**, 009 (2000) [hep-th/9911029]; F. D'Eramo, J. Thaler and Z. Thomas, JHEP **1309**, 125 (2013) [arXiv:1307.3251]; K. Harigaya and M. Ibe, Phys. Rev. D **90**, no. 8, 085028 (2014) [arXiv:1409.5029 [hep-th]].
- [56] G. D. Coughlan, W. Fischler, E. W. Kolb, S. Raby and G. G. Ross, Phys. Lett. B **131**, 59 (1983).
- [57] M. Ibe, Y. Shinbara and T. T. Yanagida, Phys. Lett. B **639**, 534 (2006) [hep-ph/0605252].
- [58] K. Harigaya, M. Ibe, M. Kawasaki and T. T. Yanagida, arXiv:1506.05250 [hep-ph].