

Quantum Zeno and anti-Zeno effects in an asymmetric nonlinear optical coupler

Kishore Thapliyal*, Anirban Pathak

Jaypee Institute of Information Technology, A-10, Sector-62, Noida, UP-201307, India

ABSTRACT

Quantum Zeno and anti-Zeno effects in an asymmetric nonlinear optical coupler are studied. The asymmetric nonlinear optical coupler is composed of a linear waveguide ($\chi^{(1)}$) and a nonlinear waveguide ($\chi^{(2)}$) interacting with each other through the evanescent waves. The nonlinear waveguide has quadratic nonlinearity and it operates under second harmonic generation. A completely quantum mechanical description is used to describe the system. The closed form analytic solutions of Heisenberg's equations of motion for the different field modes are obtained using Sen-Mandal perturbative approach. In the coupler, the linear waveguide acts as a probe on the system (nonlinear waveguide). The effect of the presence of the probe (linear waveguide) on the photon statistics of the second harmonic mode of the system is considered as quantum Zeno and anti-Zeno effects. Further, it is also shown that in the stimulated case, it is easy to switch between quantum Zeno and anti-Zeno effects just by controlling the phase of the second harmonic mode of the asymmetric coupler.

Keywords: Quantum Zeno effect, quantum anti-Zeno effect, optical coupler, waveguide.

1. INTRODUCTION

Zeno's paradoxes have been in discussion since fifth century. In 1977, Mishra and Sudarshan¹ introduced a quantum analogue of Zeno's paradox, which is later termed as quantum Zeno effect. Quantum Zeno effect in the original formulation refers to the inhibition of the temporal evolution of a system on continuous measurement,¹⁻³ while quantum anti-Zeno or inverse Zeno effect refers to the enhancement of the evolution instead of the inhibition⁴ (see^{5,6} for reviews on Zeno and anti-Zeno effect). In the last four decades, quantum Zeno effect has been studied in different physical systems, such as two coupled nonlinear optical processes,⁷ parametric down-conversion⁸ and cascaded parametric down-conversion with losses.⁹ Similarly, quantum anti-Zeno effect is also reported in various physical systems. Specifically, quantum anti-Zeno effect is observed in parametric down-conversion¹⁰ and in radioactive decay processes, where the measurement causes the system to disintegrate.¹¹ In Ref.,¹² both the effects (quantum Zeno and anti-Zeno) are reported in two-level systems. Further, a geometrical criterion for transition between the Zeno and anti-Zeno effects has also been discussed in Ref.¹³ Quantum Zeno effect using environment-induced decoherence theory has also been discussed in the past.¹⁴ Agarwal and Tewari proposed a scheme for an all optical realization of quantum Zeno effect using an arrangement of beam splitters.¹⁵ In addition to the above mentioned theoretical studies, quantum Zeno effect has also been experimentally realized in trapped beryllium ions¹⁶ and with the help of rotators and polarizers.¹⁷

Recently, the interest on quantum Zeno effect has increased by manifold as it has found its applications in counterfactual quantum computation,¹⁸ where computation is accomplished using the computer in superposition of running and not running states and later to infer the solution from it; counterfactual quantum communication, in which information is sent without sending the information encoded particles through the communication channel;¹⁹ and quantum Zeno tomography, which essentially uses interaction free measurement.²⁰ Interestingly, detection of an absorbing object without any interaction with light, using quantum Zeno effect with higher efficiency has already been demonstrated.²¹ These facts motivated the study of Zeno effect in macroscopic systems,²² too. Quantum Zeno and anti-Zeno effects in the nonlinear optical couplers have been studied in the recent past²³⁻²⁶ by considering that one of the mode in the nonlinear waveguide is coupled with the auxiliary mode in a linear waveguide, and the auxiliary linear mode acts as a probe (continuous observation) on the evolution of the system (nonlinear waveguide) and changes the photon statistics of the other modes of the nonlinear

*tkishore36@yahoo.com

waveguide, which are not coupled with the probe. This kind of continuous interaction with an external system is equivalent to the original inhibition or enhancement of evolution of the system on continuous measurement as measurement is strong coupling with a measuring device.^{27–29}

An important example of nonlinear optical coupler is an asymmetric nonlinear optical coupler, consisting of a nonlinear waveguide with $\chi^{(2)}$ nonlinearity operating under second harmonic generation coupled with a linear waveguide. This system is studied earlier and the nonclassical properties (such as squeezing, antibunching and entanglement) have been reported in this system for both codirectional and contradirectional propagation of the field in the linear waveguide (^{30–33} and references therein). We will consider here the codirectional propagation of the fields as in contradirectional propagation the solution was only valid at both the ends of the coupler, i.e., not valid for $0 < z < L$, where L is the interaction length of the coupler.^{32,33} It has already been established that the nonclassical effects can be observed in the asymmetric nonlinear coupler.^{30–33} The presence of quantum Zeno and anti-Zeno effects reported in this work further establishes the existence of nonclassicality in this asymmetric coupler.

To study the quantum Zeno and anti-Zeno effects closed form analytic expressions for different field operators are used here. These expressions were obtained earlier by using Sen-Mandal perturbative approach^{30,31} and a completely quantum mechanical description of the system. The solutions of Heisenberg's equations of motion used here are better than the conventional short-length solutions as these solutions are not restricted by length.³³ In what follows, we use the solutions reported in Refs.^{30,31} to establish the existence of quantum Zeno and anti-Zeno effects in the asymmetric nonlinear optical coupler.

The remaining part of this paper is organized as follows. In Section 2, we briefly describe the momentum operator that provides a completely quantum mechanical description of the system and also note the analytic expressions of the field operators required for the present study. In Section 3, we show the existence of quantum Zeno and anti-Zeno effects in the asymmetric nonlinear optical coupler and show the spatial evolution of the Zeno parameter. The variation of the Zeno parameter has also been studied with the phase of the coherent input in the second harmonic mode of the nonlinear waveguide, phase mismatch between fundamental and second harmonic modes in the nonlinear waveguide, and linear coupling between probe and the system. Finally, the paper is concluded in Section 4.

2. THE SYSTEM AND SOLUTION

The asymmetric nonlinear optical coupler, schematically shown in Fig. 1, is prepared by combining a linear waveguide with a nonlinear waveguide of $\chi^{(2)}$ nonlinearity. As $\chi^{(2)}$ medium can produce second harmonic generation, we may say that the codirectional asymmetric nonlinear optical coupler studied here for the investigation of possibility of observing quantum Zeno and anti-Zeno effects is operated under second harmonic generation. The momentum operator for this specific coupler in interaction picture is³⁰

$$G = -\hbar k a b_1^\dagger - \hbar \Gamma b_1^2 b_2^\dagger \exp(i\Delta k z) + \text{H.c.}, \quad (1)$$

where H.c. stands for the Hermitian conjugate, and $\Delta k = |2k_1 - k_2|$ is the phase mismatch between the fundamental and second harmonic beams, and k (Γ) being the linear (nonlinear) coupling constant is proportional to the linear (nonlinear) susceptibility $\chi^{(1)}$ ($\chi^{(2)}$). The value of $\chi^{(2)}$ is considerably smaller than $\chi^{(1)}$ (typically $\chi^{(2)}/\chi^{(1)} \simeq 10^{-6}$) which leads to $\Gamma \ll k$, unless an extremely strong pump is present in the nonlinear waveguide.

In Refs.^{30,31} closed form analytic expressions for the evolution of the field operators corresponding to the Hamiltonian (1) were obtained using Sen-Mandal perturbative approach valid up to the linear power of the nonlinear coupling coefficient Γ . The field operators reported there are:

$$\begin{aligned} a(z) &= f_1 a(0) + f_2 b_1(0) + f_3 b_1^\dagger(0) b_2(0) + f_4 a^\dagger(0) b_2(0), \\ b_1(z) &= g_1 a(0) + g_2 b_1(0) + g_3 b_1^\dagger(0) b_2(0) + g_4 a^\dagger(0) b_2(0), \\ b_2(z) &= h_1 b_2(0) + h_2 b_1^2(0) + h_3 b_1(0) a(0) + h_4 a^2(0), \end{aligned} \quad (2)$$




Figure 1. (Color online) Schematic diagram of an asymmetric nonlinear optical coupler of interaction length L in codirectional propagation prepared by combining a linear waveguide with a nonlinear (quadratic) waveguide operated by second harmonic generation.

where

$$\begin{aligned}
f_1 &= g_2 = \cos |k|z, \\
f_2 &= -g_1^* = -\frac{ik^*}{|k|} \sin |k|z, \\
f_3 &= \frac{2k^*\Gamma^*}{4|k|^2 - (\Delta k)^2} \left[G_- f_1 + \frac{f_2}{k^*} \left\{ \Delta k - \frac{2|k|^2}{\Delta k} G_- \right\} \right], \\
f_4 &= \frac{4k^{*2}\Gamma^*}{\Delta k[4|k|^2 - (\Delta k)^2]} G_- f_1 + \frac{2k^*\Gamma^*}{[4|k|^2 - (\Delta k)^2]} G_+ f_2, \\
g_3 &= \frac{2\Gamma^*k}{[4|k|^2 - (\Delta k)^2]} G_+ f_2 - \frac{2\Gamma^*(2|k|^2 - (\Delta k)^2)}{\Delta k[4|k|^2 - (\Delta k)^2]} f_1 G_-, \\
g_4 &= \frac{4\Gamma^*|k|^2}{\Delta k[4|k|^2 - (\Delta k)^2]} f_2 - \frac{2\Gamma^*(2|k|^2 - (\Delta k)^2)}{\Delta k[4|k|^2 - (\Delta k)^2]} (G_+ - 1) f_2 + \frac{2k^*\Gamma^*}{[4|k|^2 - (\Delta k)^2]} G_- f_1, \\
h_1 &= 1, \\
h_2 &= \frac{\Gamma G^*}{2\Delta k} - \frac{i\Gamma}{2[4|k|^2 - (\Delta k)^2]} \left\{ 2|k| (G_+^* - 1) \sin 2|k|z - i\Delta k [1 - (G_+^* - 1) \cos 2|k|z] \right\}, \\
h_3 &= \frac{-\Gamma|k|}{k^*[4|k|^2 - (\Delta k)^2]} \left\{ i\Delta k (G_+^* - 1) \sin 2|k|z + 2|k| [1 - (G_+^* - 1) \cos 2|k|z] \right\}, \\
h_4 &= -\frac{\Gamma|k|^2 G_-^*}{2k^{*2}\Delta k} - \frac{i\Gamma|k|^2}{2k^{*2}[4|k|^2 - (\Delta k)^2]} \left\{ 2|k| (G_+^* - 1) \sin 2|k|z - i\Delta k [1 - (G_+^* - 1) \cos 2|k|z] \right\},
\end{aligned} \tag{3}$$

where $G_{\pm} = [1 \pm \exp(-i\Delta kz)]$.

The number operator for the second harmonic field mode in the nonlinear waveguide, i.e., b_2 mode is given by

$$N_{b_2}(z) = b_2^\dagger(z) b_2(z) = b_2^\dagger(0) b_2(0) + \left[h_2 b_2^\dagger(0) b_1^2(0) + h_3 b_2^\dagger(0) b_1(0) a(0) + h_4 b_2^\dagger(0) a^2(0) + \text{H.c.} \right]. \tag{4}$$

The initial state being the multimode coherent state $|\alpha\rangle|\beta\rangle|\gamma\rangle$ with all three modes $|\alpha\rangle$, $|\beta\rangle$ and $|\gamma\rangle$ the eigenkets of annihilation operators a , b_1 and b_2 , respectively. For example, after the operation of the field operator $b_2(0)$ on such a multimode coherent state we would obtain

$$b_2(0)|\alpha\rangle|\beta\rangle|\gamma\rangle = \gamma|\alpha\rangle|\beta\rangle|\gamma\rangle, \tag{5}$$

where $|\alpha|^2$, $|\beta|^2$ and $|\gamma|^2$ are the initial number of photons in the field modes a , b_1 and b_2 , respectively. Further, the coupler discussed here can operate under two conditions: spontaneous and stimulated. In the spontaneous case, $|\alpha| \neq 0$, $|\beta| \neq 0$, and $|\gamma| = 0$, whereas in the stimulated case, $|\alpha| \neq 0$, $|\beta| \neq 0$, and $|\gamma| \neq 0$.

3. QUANTUM ZENO AND ANTI-ZENO EFFECTS

The number of photons in the second harmonic mode for the initial multimode coherent state (5) is given by

$$\langle N_{b_2}(z) \rangle = |\gamma|^2 + [h_2 \beta^2 \gamma^* + h_3 \alpha \beta \gamma^* + h_4 \alpha^2 \gamma^* + \text{c.c.}]. \tag{6}$$

In the absence of the probe mode, i.e., $k = 0$ and $\alpha = 0$, we have

$$\langle N_{b_2}(z) \rangle_{k=0} = |\gamma|^2 + [h_2' \beta^2 \gamma^* + \text{c.c.}], \tag{7}$$




Figure 2. (Color online) The spatial variation of the Zeno parameter (ΔN_Z) in the second harmonic (b_2) mode is shown with rescaled length (Γz) for the initial state $|\alpha\rangle|\beta\rangle|\gamma\rangle$ and $k = 0.1$, $\Gamma = 0.001$, $\Delta k = 10^{-4}$, $\alpha = 5$ and $\beta = 2$, where the smooth (blue) and dashed (red) lines correspond to $\gamma = 1$ (quantum Zeno effect) and $\gamma = -1$ (quantum anti-Zeno effect), respectively.

where $h'_2 = h_2(k=0) = \frac{\Gamma}{\Delta k} [1 - \exp(i\Delta k z)]$. The effect of the presence of the probe mode can be given as ($\Delta N_Z = \langle N_{b_2}(z) \rangle - \langle N_{b_2}(z) \rangle_{k=0}$), where ΔN_Z is the Zeno parameter. The non-zero value of the Zeno parameter implies that the presence of the probe affects the evolution of the photon statistics of the system, i.e., the positive (negative) value of the Zeno parameter means the photon generation is increased (decreased) due to the continuous measurement of the probe on the linear mode of the nonlinear waveguide. In the present case, using Eqs. (6) and (7), we may obtain the analytic expression for the Zeno parameter as

$$\Delta N_Z = [\{(h_2 - h'_2) \beta^2 + h_3 \alpha \beta + h_4 \alpha^2\} \gamma^* + \text{c.c.}] . \quad (8)$$

When the Zeno parameter becomes negative (positive), it implies the existence of quantum Zeno (anti-Zeno) effect. Here, the expression for the Zeno parameter is of the form $|\gamma| F(h_i, \phi)$, where we have considered $\gamma \equiv |\gamma| \exp(i\phi)$. So, the Zeno parameter becomes zero in the spontaneous case, and neither the quantum Zeno effect nor the anti-Zeno effect can be observed. In the stimulated case, the functional form of the Zeno parameter suggests that we can control the quantum Zeno and anti-Zeno effects in the asymmetric coupler just by controlling the phase of the input coherent beam of the b_2 mode, i.e., changing ϕ . As the phase change of π in γ changes the quantum Zeno effect into quantum anti-Zeno effect, and vice versa. For example, in Fig. 2, we show the spatial evolution of the Zeno parameter for $\phi = 0$ (smooth blue line) and $\phi = \pi$ (dashed red line). While the choice $\phi = 0$ illustrates the existence of quantum Zeno effect, $\phi = \pi$ is found to illustrate the existence of quantum anti-Zeno effect.

Further, the effect of change of the other parameters on the Zeno parameter can also be observed. Specifically, Fig. 3 shows that the transition between quantum Zeno and anti-Zeno effects can be observed with change in phase mismatch between fundamental and second harmonic modes in the nonlinear waveguide. Similarly, Fig. 4 illustrates the variation of quantum Zeno effect, as the Zeno parameter remains negative, with linear coupling between probe and the system. A similar behaviour can also be observed for quantum anti-Zeno effect. However, the variation of the Zeno parameter with change in nonlinear coupling constant is observed to be linear in nature and decreasing with nonlinear coupling constant (corresponding plot is not included in this paper).

4. CONCLUSION

Existence of quantum Zeno and anti-Zeno effects are reported in an asymmetric nonlinear optical coupler prepared by combing a linear and a nonlinear waveguide of $\chi^{(2)}$ nonlinearity (cf. Fig. 2). Further, it is also shown that in the stimulated case, it is easy to switch between quantum Zeno and anti-Zeno effects just by controlling the phase of the second harmonic mode in the asymmetric coupler discussed here. This flexibility to switch between the quantum Zeno and anti-Zeno effects was not observed in earlier studies on quantum Zeno and anti-Zeno effects in optical couplers.^{23, 24, 26} Further, the effects of change in linear coupling and phase mismatch on




Figure 3. (Color online) The spatial variation of the Zeno parameter (ΔN_Z) in the second harmonic (b_2) mode with rescaled length (Γz) and phase mismatch (Δk) between fundamental and second harmonic mode in the nonlinear waveguide for the initial state $|\alpha\rangle|\beta\rangle|\gamma\rangle$ and $k = 0.1$, $\Gamma = 0.001$, $\alpha = 5$ and $\beta = 2$, and $\gamma = 1$. Transition between quantum Zeno and anti-Zeno effects with change in phase mismatch can be observed.




Figure 4. (Color online) The spatial variation of the Zeno parameter (ΔN_Z) depicts quantum Zeno effect in the second harmonic (b_2) mode is shown with rescaled length (Γz) and linear coupling (k) between the waveguides for the initial state $|\alpha\rangle|\beta\rangle|\gamma\rangle$ and $\Gamma = 0.001$, $\Delta k = 10^{-4}$, $\alpha = 5$ and $\beta = 2$, and $\gamma = 1$.

the spatial variation of the Zeno parameter are also illustrated.

The approach adopted here and in Ref.³³ is quite general and same may be used for the similar studies on other nonlinear optical couplers. Further, the theoretical results reported here seem to be easily realizable in experiment as the coupler used here is commercially available and the photon number statistics required to study quantum Zeno and anti-Zeno effects can be obtained using high efficiency photon number resolving detectors.

ACKNOWLEDGMENTS

K. T. and A. P. thank Department of Science and Technology (DST), India for support provided through the DST project No. SR/S2/LOP-0012/2010. They also thank B. Sen, A. Venugopalan and J. Peřina for some useful discussions.

REFERENCES

- [1] Misra, B. and Sudarshan, E. C. G., "The Zeno's paradox in quantum theory," *Journal of Mathematical Physics* 18(4), 756-763 (1977).
- [2] Khalfin, L. A., "Zeno's Quantum Effect," *Usp. Fiz. Nauk* 160, 185-188 (1990).
- [3] Peres, A., "Zeno paradox in quantum theory," *American Journal of Physics* 48, 931 (1980).
- [4] Kaulakys, B. and Gontis, V., "Quantum anti-Zeno effect," *Phys. Rev. A* 56, 1131 (1997).
- [5] Venugopalan, A., "The quantum Zeno effect-watched pots in the quantum world," *Resonance* 12(4), 52-68 (2007).
- [6] Pascazio, S., "All You Ever Wanted to Know About the Quantum Zeno Effect in 70 Minutes," *Open Systems & Information Dynamics* 21, 1440007 (2014).
- [7] Chirkin, A. S. and Rodionov, A. V., "Quantum Zeno effect in two coupled nonlinear optical processes," *Journal of Russian Laser Research* 26, 83 (2005).
- [8] Luis, A. and Peřina, J., "Zeno Effect in Parametric Down-Conversion," *Phys. Rev. Lett.* 76, 4340 (1996).
- [9] Peřina, J., "Quantum Zeno effect in cascaded parametric down-conversion with losses," *Phys. Lett. A* 325, 16-20 (2004).
- [10] Luis, A. and Sánchez-Soto, L. L., "Anti-Zeno effect in parametric down-conversion," *Phys. Rev. A* 57, 781 (1998).
- [11] Kofman, A. G. and Kurizki, G., "Acceleration of quantum decay processes by frequent observations," *Nature* 405, 546-550 (2000).
- [12] Luis, A., "Zeno and anti-Zeno effects in two-level systems," *Phys. Rev. A* 67, 062113 (2003).
- [13] Facchi, P., Nakazato, H. and Pascazio, S., "From the Quantum Zeno to the Inverse Quantum Zeno Effect," *Phys. Rev. Lett.* 86, 2699 (2001).
- [14] Venugopalan, A. and Ghosh, R., "Decoherence and the Quantum Zeno Effect," *Phys. Lett. A* 204, 11 (1995).
- [15] Agarwal, G. S. and Tewari, S. P., "An all-optical realization of the quantum Zeno effect," *Phys. Lett. A* 185, 139-142 (1994).
- [16] Itano, W. M., Heinzen, D. J., Bollinger, J. J. and Wineland, D. J., "Quantum Zeno effect," *Phys. Rev. A* 41, 2295 (1990).
- [17] Kwiat, P., Weinfurter, H., Herzog, T., Zeilinger, A. and Kasevich, M., "Experimental Realization of Interaction-free Measurements," *Annals of the New York Academy of Sciences* 755, 383-393 (1995).
- [18] Hosten, O., Rakher, M. T., Barreiro, J. T., Peters, N. A. and Kwiat, P. G., "Counterfactual quantum computation through quantum interrogation," *Nature* 439, 949-952 (2006).
- [19] Salih, H., Li, Z.-H., Al-Amri, M. and Zubairy, M. S., "Protocol for Direct Counterfactual Quantum Communication," *Phys. Rev. Lett.* 110, 170502 (2013).
- [20] Facchi, P., Hradil, Z., Krenn, G., Pascazio, S. and Řeháček, J., "Quantum Zeno tomography," *Phys. Rev. A* 66, 012110 (2002).
- [21] Kwiat, P. G., White, A. G., Mitchell, J. R., Nairz, O., Weihs, G., Weinfurter, H. and Zeilinger, A., "High-Efficiency Quantum Interrogation Measurements via the Quantum Zeno Effect," *Phys. Rev. Lett.* 83, 4725 (1999).

- [22] Zezyulin, D. A., Konotop, V. V., Barontini, G. and Ott, H., “Macroscopic Zeno Effect and Stationary Flows in Nonlinear Waveguides with Localized Dissipation,” *Phys. Rev. Lett.* 109, 020405 (2012).
- [23] Thun, K., Peřina, J. and Křepelka, J., “Quantum Zeno effect in Raman scattering,” *Phys. Lett. A* 299(1), 19-30 (2002).
- [24] Řeháček, J., Peřina, J., Facchi, P., Pascazio, S. and Miřta Jr., L., “Quantum Zeno effect in a probed down-conversion process,” *Phys. Rev. A* 62, 013804 (2000); Řeháček, J., Peřina, J., Facchi, P., Pascazio, S. and Miřta Jr., L., “Quantum Zeno effect in a nonlinear coupler,” *Optics and Spectroscopy* 91, 501–507 (2001).
- [25] Facchi, P. and Pascazio, S., “Quantum Zeno and inverse quantum Zeno effects,” *Progress in Optics* 42, 147-218 (2001).
- [26] Miřta Jr., L., Herec, J., Jelínek, V., Řeháček, J. and Peřina, J., “Quantum dynamics and statistics of two coupled down-conversion processes,” *J. Opt. B: Quantum Semiclass. Opt.* 2, 726–734 (2000).
- [27] Schulman, L. S., “Continuous and pulsed observations in the quantum Zeno effect,” *Phys. Rev. A* 57, 1509 (1998).
- [28] Facchi, P. and Pascazio, S. “Quantum Zeno Subspaces,” *Phys. Rev. Lett.* 89, 080401 (2002).
- [29] Facchi, P. and Pascazio, S., “Three Different Manifestations of the Quantum Zeno Effect,” *Lecture Notes in Physics* 622, 141-156 (2003).
- [30] Mandal, S. and Peřina, J., “Approximate quantum statistical properties of a nonlinear optical coupler,” *Phys. Lett. A* 328(2), 144-156 (2004).
- [31] Thapliyal, K., Pathak, A., Sen, B. and Peřina, J., “Higher-order nonclassicalities in a codirectional nonlinear optical coupler: Quantum entanglement, squeezing, and antibunching,” *Phys. Rev. A* 90, 013808 (2014).
- [32] Thapliyal, K., Pathak, A., Sen, B. and Peřina, J., “Nonclassical properties of a contradirectional nonlinear optical coupler,” *Phys. Lett. A* 378(46), 3431-3440 (2014).
- [33] Peřina, J. and Peřina Jr., J., “Photon statistics of a contradirectional nonlinear coupler,” *Quantum and Semi-classical Optics: Journal of the European Optical Society Part B* 7(5), 849 (1995).