

Direct detection of singlet dark matter in classically scale-invariant standard model

Kazuhiro Endo^(a), Koji Ishiwata^(b)

^(a)*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

^(b)*Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192,
Japan*

Abstract

Classical scale invariance is one of the possible solutions to explain the origin of the electroweak scale. The simplest extension is the classically scale-invariant standard model augmented by a multiplet of gauge singlet real scalar. In the previous study it was shown that the properties of the Higgs potential deviate substantially, which can be observed in the International Linear Collider. On the other hand, since the multiplet does not acquire vacuum expectation value, the singlet components are stable and can be dark matter. In this letter we study the detectability of the real singlet scalar bosons in the experiment of the direct detection of dark matter. It is shown that a part of this model has already been excluded and the rest of the parameter space is within the reach of the future experiment.

1 Introduction

Higgs boson was discovered in 2012 at the CERN Large Hadron Collider [1,2]. Since then, its properties, such as spin, parity and couplings to the standard model fermions and gauge bosons, have been measured and it turned out that they are consistent with the standard model prediction. In spite of the success of the standard model up to now, however, it is commonly believed that the standard model is not the ultimate theory of particle physics. In fact there are lots of unsolved problems in the field of particle physics as well as cosmology.

One of them is the origin of the spontaneous symmetry breakdown of the electroweak gauge group. In the standard model, the electroweak symmetry is broken by Higgs field that has an ad hoc tachyonic mass term. One explanation for the tachyonic mass is supersymmetry. In supersymmetric extension of the standard model, the negative mass term is induced radiatively. On the other hand, radiative symmetry breaking is possible in non-supersymmetric theory, which is known as Coleman-Weinberg (CW) mechanism [3]. In the CW mechanism spontaneous symmetry breaking is induced at quantum level from classically scale-invariant scalar potential. Although it turned out that the CW mechanism with a Higgs does not work for the electroweak symmetry breaking, simple extensions of the Higgs sector are known to be phenomenologically viable (see, *e.g.*, [4–29]).

Recently Higgs properties were studied in a classically scale-invariant standard model augmented by an electroweak singlet scalars that form a multiplet of global $O(N)$ symmetry [30]. It was shown that the Higgs self-couplings deviate significantly from the standard model prediction. Such feature can be observed as a prominent signal of this model at the next-generation lepton collider experiment, such as the International Linear Collider (ILC) [31–33]. On the other hand, it was also shown that the singlet field does not get a vacuum expectation value (VEV). Then, other Higgs properties are unaffected since there is no mixing between the singlet and Higgs. Another important consequence is the stability of the singlet field due to unbroken $O(N)$ symmetry. If the reheating temperature of the universe is higher than the mass of the singlet, the singlet field is thermalized. Then non-vanishing thermal relic of the singlet remains, which can play a role of dark matter.

In this letter we study direct detection of the real singlet dark matter with $O(N)$ symmetry. It was pointed out in a similar framework where the thermal relic abundance of the singlet dark matter is too small to explain the present energy density of

dark matter by taking into account the 125 GeV Higgs [7, 10, 12]. This is due to an enhanced annihilation cross section caused by a large singlet-Higgs coupling. On the contrary, however, the large singlet-Higgs coupling may result in a large scattering cross section of the singlet with nucleon. According to the study of Ref. [30], the couplings of the singlet with the standard model particles are fixed for the successful electroweak symmetry breaking via the CW mechanism, which makes it possible to determine the relic abundance and the scattering cross section of the singlet with nucleon at a high precision. The scattering cross section of singlet scalar dark matter is also discussed in several literature mentioned above. We revise the calculation of the spin-independent cross section of singlet scalar particle by adopting the formalism given in Ref. [34] where next-to-leading order QCD effect is properly taken into account. It will be shown that part of the model has already been excluded by recent LUX result [35] and the future experiments will be able to probe almost the entire parameter space of the model.

Here is the organization of this letter. In Sec. 2 we briefly explain the model, including the prescription how to determine model parameters. Then the thermal relic and the scattering cross section of the singlet are calculated, and the detection of the singlet scalar is discussed in Sec. 3. Sec. 4 is dedicated to conclusion.

2 The Model

In the framework with classical scale invariance it is known that the standard model without the Higgs mass term has already been excluded. In order to construct phenomenologically viable model, therefore, it is necessary to extend the model, *e.g.*, by adding a new particle to the model. The simplest extension is to introduce a gauge singlet real scalar field. Such a singlet scalar can couple to the Higgs in general, then the singlet contributes to the CW potential. The effect strongly depends on the degree of freedom of the singlet field. To see the impact we introduce a fundamental representation of a global $O(N)$ symmetry, $S = (S_1, \dots, S_N)^T$. Consequently, the tree-level scalar potential which is allowed under the symmetry is

$$V = \lambda_H(H^\dagger H)^2 + \lambda_{HS}H^\dagger H S_i S_i + \frac{\lambda_S}{4}(S_i S_i)^2, \quad (2.1)$$

where H is the Higgs doublet field $H = (H^+, H^0)^T$, and summed over $i = 1, \dots, N$ for $N \geq 2$. Z_2 symmetry is assumed for $N = 1$ case, whereas it is also a subgroup

of $O(N)$ symmetry and always survives in the scale-invariant tree-level potential for $N \geq 2$.

The electroweak symmetry breaking is induced via the CW mechanism. To see this, the scalar fields can be taken without loss of generality as $H = (1/\sqrt{2})(0, \phi)^T$ and $S = (\varphi, 0, \dots, 0)^T$ where ϕ and φ are classical fields of the real scalars. Then the effective potential at one-loop level is given by

$$V_{\text{eff}}(\phi, \varphi) = V_{\text{tree}}(\phi, \varphi) + V_{1\text{-loop}}(\phi, \varphi), \quad (2.2)$$

with

$$V_{\text{tree}}(\phi, \varphi) = \frac{\lambda_H}{4}\phi^4 + \frac{\lambda_{HS}}{2}\phi^2\varphi^2 + \frac{\lambda_S}{4}\varphi^4, \quad (2.3)$$

$$V_{1\text{-loop}}(\phi, \varphi) = \frac{1}{4(4\pi)^2} \sum_i n_i M_i^4(\phi, \varphi) \left[\ln \frac{M_i^2(\phi, \varphi)}{\mu^2} - c_i \right], \quad (2.4)$$

in $\overline{\text{MS}}$ -scheme with renormalization scale μ . Index i denotes the fields which run in the loop diagrams. (n_i , M_i^2 and c_i are given in Appendix A.) The electroweak symmetry is spontaneously broken if $\partial V_{\text{eff}}/\partial\phi|_{\phi=\langle\phi\rangle} = 0$ with $\langle\phi\rangle \neq 0$, which implies $\lambda_H \sim \frac{N\lambda_{HS}^2}{16\pi^2} - \frac{3yt^4}{16\pi^2}$ as a necessary condition. Then λ_H should be regarded as the next-to-leading order in terms of the order counting of the dimensionless couplings. Consequently we rewrite the effective potential as

$$V_{\text{eff}} = V_{\text{LO}} + V_{\text{NLO}}, \quad (2.5)$$

with V_{LO} and V_{NLO} being regarded as leading order (LO) and next-to-leading order (NLO) of the scalar potential;

$$V_{\text{LO}} = \frac{\lambda_{HS}}{2}\phi^2\varphi^2 + \frac{\lambda_S}{4}\varphi^4, \quad (2.6)$$

$$\begin{aligned} V_{\text{NLO}} = & \frac{\lambda_H}{4}\phi^4 \\ & + \frac{F_{\text{+app}}^2(\phi, \varphi)}{64\pi^2} \left[\ln \left(\frac{F_{\text{+app}}(\phi, \varphi)}{\mu^2} \right) - \frac{3}{2} \right] + \frac{3}{64\pi^2} (\lambda_{HS}\varphi^2)^2 \left[\ln \left(\frac{\lambda_{HS}\varphi^2}{\mu^2} \right) - \frac{3}{2} \right] \\ & + \frac{N-1}{64\pi^2} (\lambda_{HS}\phi^2 + \lambda_S\varphi^2)^2 \left[\ln \left(\frac{\lambda_{HS}\phi^2 + \lambda_S\varphi^2}{\mu^2} \right) - \frac{3}{2} \right] \\ & - \frac{12}{64\pi^2} M_t^4(\phi) \left[\ln \left(\frac{M_t^2(\phi)}{\mu^2} \right) - \frac{3}{2} \right] \\ & + \frac{6}{64\pi^2} M_W^4(\phi) \left[\ln \left(\frac{M_W^2(\phi)}{\mu^2} \right) - \frac{5}{6} \right] + \frac{3}{64\pi^2} M_Z^4(\phi) \left[\ln \left(\frac{M_Z^2(\phi)}{\mu^2} \right) - \frac{5}{6} \right], \quad (2.7) \end{aligned}$$

where $F_{\text{+app}}$ is given in Appendix A. In Ref. [30] it is shown that the successful electroweak symmetry breaking with the Higgs mass $m_h \simeq 125$ GeV can be realized in a given number of N . Table 1 shows the results.^{#1} Roughly speaking, the Higgs mass is expected to be $m_h \sim (\sqrt{N}\lambda_{HS}/4\pi)v_H$ with $v_H = 246$ GeV, which is consistent with the numerical results in Table 1. With the proper order counting, the effective potential around the VEV is obtained by replacing the scalar fields as $\phi \rightarrow v_H + h$, $\varphi^2 \rightarrow s_i s_i$ and expanding by powers of h and $s_i s_i$;

$$V_{\text{eff}} = \text{const.} + \frac{1}{2}m_h^2 h^2 + \frac{1}{2}m_s^2 s_i s_i + \frac{\lambda_{hhh}}{3!} v_H h^3 + \frac{\lambda_{hhhh}}{4!} h^4 + \frac{\lambda_{hss}}{2} v_H h s_i s_i + \frac{\lambda_{hhss}}{4} h^2 s_i s_i + \frac{\lambda_{ssss}}{4!} (s_i s_i)^2 + \dots, \quad (2.8)$$

where m_s is the mass of singlet. We have taken $\mu = v_H$ and omitted irrelevant terms in our later discussion. The results for λ_{hhhh} , λ_{hhh} , λ_{hhss} , λ_{hss} , λ_{ssss} and m_s are summarized in Table 2 [30].^{#2} Basically λ_H and λ_{HS} are chosen to give rise to $v_H = 246$ GeV and $m_h = 125$ GeV, which determine the couplings (except for the singlet self-coupling) and the singlet mass. λ_S , on the other hand, has little impact on these results. Since the Higgs self-couplings λ_{hhhh} and λ_{hhh} significantly deviate from the SM prediction, the precise measurement of the Higgs self-couplings is a viable way to test this model.

Another important fact shown in Ref. [30] is that the singlet does not get VEV.^{#3} Without a VEV of the singlet, Higgs properties, such as Higgs production or decay rates, are unaffected. On the other hand, unbroken $O(N)$ symmetry forbids s_i to decay. Such stable particles can change the thermal history of the universe. If the reheating temperature in the early universe is higher than the singlet mass, the singlet particles are thermalized and their number densities freeze out eventually. Then the thermal relics can be components of dark matter. In this model the parameters which determine the interaction of s_i with the standard model particles (the Higgs field in our case) are completely fixed as discussed above. Therefore its nature is

^{#1}This is the results derived from the potential referred as (I) in Ref. [30].

^{#2}The couplings are obtained from the parameters shown in Table 1 which corresponds to case (I) in Ref. [30]. Since the couplings change by a few % in cases (II) or (III), we will use the couplings from case (I) in our later calculation.

^{#3} This fact is guaranteed to all orders in perturbative expansion. Strictly speaking, non-perturbative effect might break $O(N)$, which could allow non-zero VEV for the singlet. Possible (or known) non-perturbative effect is anomaly. In our model, however, $O(N)$ multiplet is scalar, thus it is anomaly free. Though one may concern another unknown non-perturbative effect, the situation is the same for the standard model, *i.e.* unbroken $U(1)_{\text{em}}$ symmetry.

N	1	4	12
μ	$v_H = 246 \text{ GeV}$		
y_t	0.919		
g	0.644		
g'	0.359		
λ_H	-0.11	-0.0045	0.075
λ_{HS}	4.8	2.4	1.4
λ_S	0.10	0.10	0.10

Table 1: The input parameters of the analysis in Ref. [30]. The values $\lambda_S = 0.10$ are benchmarks, but there are very few λ_S -dependences for λ_{hhh} , λ_{hhhh} , λ_{hss} .

highly predictable, and in fact we will see in the next section that the experiments of direct detection of dark matter provide the powerful tool to probe the model.

Before closing this section it is worth noting that $N = 1$ case is favored by two reasons [30]. In terms of Veltman's condition [36] the level of fine-tuning for the Higgs mass at the electroweak scale gets milder compared to larger N cases. Second, the fine-tuning is relaxed compared to the standard model and larger N case in a sense that the cutoff scale due to Landau pole is predicted to be around TeV. We will come back to this point later.

3 Detection of Singlet Scalars

As we discussed the singlet scalars are stable and they can play a role of dark matter. Since the singlet scalars interact with Higgs boson, they are thermalized in the early universe if the reheating temperature is higher than the singlet mass.^{#4} Then the relic abundance is determined by the conventional freeze-out scenario. What we need is the annihilation cross section of s_i . Relevant annihilation processes are $s_i s_i \rightarrow W^+ W^-$, ZZ , hh and $t\bar{t}$. The cross sections for the processes are given by

$$\sigma_{s_i s_i \rightarrow W^+ W^-} = \frac{\beta_f(s, m_W^2)}{4\pi s \beta_i} \left(\frac{\lambda_{hss} m_W^2}{s - m_h^2} \right)^2 \left[2 + \frac{1}{4} \left(\frac{s - 2m_W^2}{m_W^2} \right)^2 \right], \quad (3.1)$$

^{#4}To be specific, the reheating temperature should be larger than 300–600 GeV (see Table 2), which is a canonical case in the early universe.

N	1	4	12
λ_{hhh}	1.32	1.32	1.32
λ_{hhhh}	2.9	2.9	2.9
λ_{hss}	11.4	5.02	2.80
λ_{hhss}	14	5.6	3.0
λ_{ssss}	6.5	1.9	0.9
$m_s[\text{GeV}]$	556	378	285

Table 2: The results about the interactions among the Higgs boson and singlet scalar bosons derived in Ref. [30]. They are defined in Eq. (2.8). Some of them are also input parameters of the calculations in Section 3. These values are those in case (I), except for λ_{ssss} which is in (II) of Table 4 in Ref. [30]. In our notation of the SM Higgs potential the predicted self-couplings are $\lambda_{hhh}^{(\text{SM})} = \lambda_{hhhh}^{(\text{SM})} = 6\lambda_H = 3m_h^2/v_H^2 \simeq 0.78$.

$$\sigma_{s_i s_i \rightarrow ZZ} = \frac{\beta_f(s, m_Z^2)}{8\pi s \beta_i} \left(\frac{\lambda_{hss} m_Z^2}{s - m_h^2} \right)^2 \left[2 + \frac{1}{4} \left(\frac{s - 2m_Z^2}{m_Z^2} \right)^2 \right], \quad (3.2)$$

$$\begin{aligned} \sigma_{s_i s_i \rightarrow hh} = \frac{1}{16\pi s \beta_i} & \left[\beta_f(s, m_h^2) \tilde{\lambda}^2 + \frac{4\tilde{\lambda} \lambda_{hss}^2 v_H^2}{s \beta_i} \log \frac{t_+}{t_-} \right. \\ & \left. + \frac{2\lambda_{hss}^4 v_H^4}{s \beta_i} \left(\frac{s \beta_i \beta_f(s, m_h^2)}{t_+ t_-} + \frac{2}{2m_h^2 - s} \log \frac{t_+}{t_-} \right) \right], \quad (3.3) \end{aligned}$$

$$\sigma_{s_i s_i \rightarrow t\bar{t}} = \frac{3\beta_f^3(s, m_t^2)}{8\pi \beta_i} \left(\frac{\lambda_{hss} m_t}{s - m_h^2} \right)^2, \quad (3.4)$$

where m_W , m_Z and m_t are the masses of W , Z and t , respectively. s is the center-of-mass energy in the initial state, $\beta_i = (1 - 4m_s^2/s)^{1/2}$, $\beta_f(s, m^2) = (1 - 4m^2/s)^{1/2}$ and

$$\tilde{\lambda} = \lambda_{hhss} + \frac{\lambda_{hhh} \lambda_{hss} v_H^2}{s - m_h^2}, \quad t_{\pm} = m_h^2 - \frac{s}{2} [1 \mp \beta_i \beta_f(s, m_h^2)]. \quad (3.5)$$

Numerically it is found that the annihilation mode to $t\bar{t}$ is subdominant.^{#5} It is straightforward to compute the thermal-averaged cross section from above expression. We use the formula in Ref. [37] to get the relic abundance. (We have checked that the density parameter by solving the Boltzmann equation agrees with the approximated result within a few %.) The results are shown in Table 3. (We note

^{#5}This is expected since the amplitude of fermion pair final state is chirality suppressed, *i.e.*, the cross section is proportional to $m_t^2/m_{s_i}^4$ instead of $1/m_{s_i}^2$.

that when $N \geq 2$ all scalars s_1, \dots, s_N become dark matter.) It has figured out that the relic abundance Ω_{s_i} of s_i is much smaller than that of dark matter, which means that s_i cannot be the main component of dark matter. This is due to the large annihilation cross section enhanced by the large couplings (mainly λ_{hss}). On the other hand, however, s_i can be detected in the experiment of direct detection of dark matter, which we will discuss below.

For evaluation of the spin-independent cross section of s_i with nucleon, we adopt the formalism given in Ref. [34]. (See also Refs. [38–41] for earlier works.) In the present case only scalar-type operators are induced by Higgs-exchange diagram, then the effective Lagrangian for the scattering process is

$$\mathcal{L}_{\text{eff}} = \sum_{i=q,G} C_S^i \mathcal{O}_S^i, \quad (3.6)$$

where

$$\mathcal{O}_S^q = m_q s_i^2 \bar{q}q, \quad \mathcal{O}_S^G = \frac{\alpha_s}{\pi} s_i^2 G_{\mu\nu}^a G^{a\mu\nu}. \quad (3.7)$$

m_q is quark mass, $G_{\mu\nu}^a$ is the gluon field strength and α_s is the strong coupling constant. By integrating out the Higgs boson (and top quark), the Wilson coefficients at the electroweak scale $\mu_W \simeq m_Z$ at the next-to-leading order in α_s are given by

$$C_S^q(\mu_W) = \frac{\lambda_{hss}}{2m_h^2}, \quad (3.8)$$

$$C_S^G(\mu_W) = -\frac{\lambda_{hss}}{24m_h^2} \left[1 + \frac{11\alpha_s}{4\pi} \right]. \quad (3.9)$$

The amplitude is given by the hadronic matrix elements, *i.e.* $\langle N | m_q \bar{q}q | N \rangle$, $\langle N | \frac{\alpha_s}{\pi} s_i^2 G_{\mu\nu}^a G^{a\mu\nu} | N \rangle$ ($N = p, n$), which are obtained from lattice simulations [42, 43] and the QCD trace anomaly [44],^{#6}

$$\langle N | m_q \bar{q}q | N \rangle = m_N f_{T_q}^{(N)}, \quad (3.10)$$

$$\langle N | \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{a\mu\nu} | N \rangle = m_N \frac{4\alpha_s^2}{\pi \beta_s^{(N_f=3)}} \left[1 - (1 - \gamma_m) \sum_{q=u,d,s} f_{T_q}^{(N)} \right]. \quad (3.11)$$

Here m_N is nucleon mass, $f_{T_q}^{(N)}$ is mass fractions, *e.g.*, $f_{T_u}^{(p)} = 0.019(5)$, $f_{T_d}^{(p)} = 0.027(6)$ and $f_{T_s}^{(p)} = 0.009(22)$, which are evaluated in Ref. [41] based on Refs. [42, 43]. β_s and γ_m are the beta function of α_s and the anomalous dimension of quark mass defined

^{#6}It is given by $\Theta_\mu^\mu = \frac{\beta(\alpha_s)}{4\alpha_s} G_{\mu\nu}^a G^{a\mu\nu} + (1 - \gamma_m) \sum_q m_q \bar{q}q$, and $m_N = \langle N | \Theta_\mu^\mu | N \rangle$ to derive Eq. (3.11).

N	1	4	12
$\Omega_{s_i}/\Omega_{\text{DM}}$	2.01×10^{-4}	4.54×10^{-4}	8.07×10^{-4}
$\tilde{\sigma}_{\text{SI}}^{(p)} [10^{-46} \text{ cm}^2]$	6.77	25.6	74.5

Table 3: Fractions of the energy density of s_i in the observed dark matter density, and spin-independent cross sections of the scalars with proton multiplied by their fractions in the present dark matter density. The cross sections are computed with the effective couplings $f_S^{(p)}|_{\text{NLO}}$ shown in Table 4.

by $\beta_s = \mu \frac{d\alpha_s}{d\mu}$ and $\gamma_m m_q = \mu \frac{dm_q}{d\mu}$, respectively. In Eq. (3.11) the number of flavors $N_f = 3$ is taken since we only know the mass fractions for the light quarks, which means that the Wilson coefficients C_S^q and C_S^G should be evaluated at the hadronic scale $\mu_{\text{had}} \simeq 1 \text{ GeV}$. This can be done by the matching procedure at each quark threshold (bottom and charm quarks) and by solving the renormalization group equation for the Wilson coefficients. (For the details, such as the matching and renormalization group evolution at the next-to-leading order in α_s , see Ref. [34]). Finally the spin-independent cross section is given by

$$\sigma_{\text{SI}}^{(N)} = \frac{1}{\pi} \frac{m_N^2}{(m_s + m_N)^2} |f_S^{(N)}|^2, \quad (3.12)$$

with

$$f_S^{(N)} = \sum_{q=u,d,s} C_S^q(\mu_{\text{had}}) \langle N | m_q \bar{q} q | N \rangle + C_S^G(\mu_{\text{had}}) \langle N | \frac{\alpha_s}{\pi} G_{\mu\nu}^a G^{a\mu\nu} | N \rangle. \quad (3.13)$$

If renormalization group evolution is ignored as well as taking the leading order threshold matching, then the effective scattering amplitude is simply given by

$$\frac{f_S^{(N)}}{m_N} |_{\text{est}} \approx \frac{\lambda_{hss}}{2m_h^2} \left[\frac{2}{9} + \frac{7}{9} \sum_{q=u,d,s} f_{T_q}^{(N)} \right], \quad (3.14)$$

which is often used in the literature. To see the impact of the proper matching procedure to the effective coupling $f_S^{(N)}$, we show the numerical values in Table 4. The difference between the LO and the NLO results is about 4%, while the one between the rough estimation and the NLO is about 7%, which gives rise to about 14% deviation in the spin-independent cross section.

Now we are ready to see the experimental consequence of the model. Since the scalar fields are not the main component of the present DM, the “effective” cross

N	1	4	12
$f_S^{(p)} _{\text{est}} [10^{-5} \text{ GeV}^{-1}]$	9.07	3.99	2.23
$f_S^{(p)} _{\text{LO}} [10^{-5} \text{ GeV}^{-1}]$	10.2	4.49	2.50
$f_S^{(p)} _{\text{NLO}} [10^{-5} \text{ GeV}^{-1}]$	9.77	4.30	2.40

Table 4: Amplitude for s_i -proton scattering. $f_S^{(p)}|_{\text{est}}$ is given by Eq. (3.14) and $f_S^{(p)}|_{\text{LO}}$, $f_S^{(p)}|_{\text{NLO}}$ are the results obtained by appropriate matching at the leading order and the next-to-leading order, respectively.

section of the singlet with nucleon is obtained by multiplying the fraction of the total abundance of the scalars in the present DM density,

$$\tilde{\sigma}_{\text{SI}}^{(N)} = \sigma_{\text{SI}}^{(N)} \sum_{i=1, \dots, N} \frac{\Omega_{s_i}}{\Omega_{\text{DM}}}, \quad (3.15)$$

where $\Omega_{\text{DM}} = 0.264$ [45]. The results are given in Table 3 and Fig. 1. In Table 3 we have used the NLO result. Compared with the most stringent bound for the cross section [35], $N = 12$ case has already been excluded at 90% C.L. The others, *i.e.* $N = 1$ and 4 cases, are still viable, but $N = 4$ case is very close to the present bound. The cross section is far below the present bound for $N = 1$ case. However, it is much larger than the neutrino background in the direct detection experiments [46]. Therefore, our model (with any number of N) can be tested in ton-scale future experiments, such as LZ program [47]. Recall that $N = 1$ is favored in terms of Veltman's condition as well as the fine-tuning of the Higgs mass, which is discussed in the previous section. Thus the result shows that the most well-motivated case will be able to be examined in the future experiments.

4 Conclusion

In this letter we have studied direct detection of singlet scalar dark matter in a classically scale-invariant extension of the standard model. The model extends the Higgs sector to have an additional electroweak singlet scalars that form a multiplet of global $O(N)$ symmetry, and the electroweak symmetry is broken via Coleman-Weinberg mechanism. Recently the Higgs self-couplings as well as new couplings and the singlet mass were precisely computed in Ref. [30]. In the work it was shown the Higgs self-couplings deviate from the standard model prediction significantly,




Figure 1: Spin-independent cross sections of the singlet scalars compared with the LUX 90% C.L. bound [35]. Orange, green and purple points represent $N = 1, 4$ and 12 cases, respectively. Triangle and circle points represent rough estimation and the next-to-leading order calculation, respectively.

which can be observed at the next-generation collider experiments such as the ILC. Another important outcome of their analysis is unbroken $O(N)$. Consequently the singlet scalars are cosmologically stable and can play a role of dark matter. Since all couplings and mass parameters are fixed for given number of N [30], it is possible to precisely predict the nature of the singlet scalars. Therefore detection of the singlet scalars is complementary for the test of the model.

Assuming that the reheating temperature is above the singlet mass, we have computed the thermal relic abundance of the singlet scalars, and the scattering amplitude of the scalars with nucleon. For the precise determination of the scattering cross section we have used the formalism [34] which takes into account the next-to-leading order QCD effect in consistent way. We have focused on three benchmarks, *i.e.* $N = 1, 4$ and 12 (the singlet masses are predicted as 556, 378 and 285 GeV, respectively). Then it has been found that although the relic abundance is much

smaller than the present dark matter ($\Omega_{s_i}/\Omega_{\text{DM}} \sim \mathcal{O}(10^{-4})$), the scattering rate is enhanced due to the large Higgs-singlet coupling. To be concrete, $N = 12$ case has already been excluded by the LUX experiments, meanwhile $N = 4$ case is near the bound. In $N = 1$ case which is favored in terms of the fine-tuning regarding the Higgs mass, the effective spin-independent cross section ($\simeq 6.8 \times 10^{-46} \text{ cm}^2$) is far below the current bound. It is, however, much larger than the neutrino background. Thus it is concluded that the whole parameter space of this scenario is testable in the future ton-scale detector of dark matter direct detection.

Acknowledgement

The authors would like to express their special thanks to Yukinari Sumino for many helpful comments and careful reading of this paper. This work was supported in part by the German Science Foundation (DFG) within the Collaborative Research Center 676 ‘‘Particles, Strings and the Early Universe’’ (K.I.).

A Parameters in the effective potential

Here is the list of parameters of the one-loop potential in Eq. (2.2):

$$\begin{aligned}
n_W &= 6, & M_W^2 &= \frac{1}{4}g^2\phi^2, & c_W &= \frac{5}{6}; \\
n_Z &= 3, & M_Z^2 &= \frac{1}{4}(g^2 + g'^2)\phi^2, & c_Z &= \frac{5}{6}; \\
n_t &= -12, & M_t^2 &= \frac{1}{2}y_t^2\phi^2, & c_t &= \frac{3}{2}; \\
n_{\pm} &= 1, & M_{\pm}^2 &= F_{\pm}, & c_{\pm} &= \frac{3}{2}; \\
n_{H_{\text{NG}}} &= 3, & M_{H_{\text{NG}}}^2 &= \lambda_H\phi^2 + \lambda_{HS}\varphi^2, & c_{H_{\text{NG}}} &= \frac{3}{2}; \\
n_{S_{\text{NG}}} &= N - 1, & M_{S_{\text{NG}}}^2 &= \lambda_{HS}\phi^2 + \lambda_S\varphi^2, & c_{S_{\text{NG}}} &= \frac{3}{2},
\end{aligned} \tag{A.1}$$

where $i = W, Z, t$ show W, Z, t in the loop, respectively. $i = \pm$ indicates φ, ϕ , while $i = H_{\text{NG}}, S_{\text{NG}}$ stand for the degrees of freedom which are orthogonal to ϕ and φ , respectively. In the effective potential with the precise order counting (*i.e.*

Eq. (2.5)), we use

$$F_{\pm\text{app}}(\phi, \varphi) = \frac{\lambda_{HS}}{2}\phi^2 + \frac{\lambda_{HS} + 3\lambda_S}{2}\varphi^2 \pm \sqrt{\left[-\frac{\lambda_{HS}}{2}\phi^2 + \frac{\lambda_{HS} - 3\lambda_S}{2}\varphi^2\right]^2 + 4\lambda_{HS}^2\phi^2\varphi^2}, \quad (\text{A.2})$$

for F_{\pm} . The reason for dropping $F_{-\text{app}}$ in Eq. (2.5) is explained in Ref. [30] (see also Ref. [48]).

References

- [1] **ATLAS** Collaboration, G. Aad *et al.*, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC,” *Phys.Lett.* **B716** (2012) 1–29, [arXiv:1207.7214 \[hep-ex\]](#).
- [2] **CMS** Collaboration, S. Chatrchyan *et al.*, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC,” *Phys.Lett.* **B716** (2012) 30–61, [arXiv:1207.7235 \[hep-ex\]](#).
- [3] S. R. Coleman and E. J. Weinberg, “Radiative Corrections as the Origin of Spontaneous Symmetry Breaking,” *Phys.Rev.* **D7** (1973) 1888–1910.
- [4] R. Foot, A. Kobakhidze, and R. R. Volkas, “Electroweak Higgs as a pseudo-Goldstone boson of broken scale invariance,” *Phys.Lett.* **B655** (2007) 156–161, [arXiv:0704.1165 \[hep-ph\]](#).
- [5] J. R. Espinosa and M. Quiros, “Novel Effects in Electroweak Breaking from a Hidden Sector,” *Phys.Rev.* **D76** (2007) 076004, [arXiv:hep-ph/0701145 \[hep-ph\]](#).
- [6] L. Alexander-Nunneley and A. Pilaftsis, “The Minimal Scale Invariant Extension of the Standard Model,” *JHEP* **1009** (2010) 021, [arXiv:1006.5916 \[hep-ph\]](#).
- [7] D. Chway, T. H. Jung, H. D. Kim, and R. Dermisek, “Radiative Electroweak Symmetry Breaking Model Perturbative All the Way to the Planck Scale,” *Phys.Rev.Lett.* **113** no. 5, (2014) 051801, [arXiv:1308.0891 \[hep-ph\]](#).
- [8] I. Masina and M. Quiros, “On the Veltman Condition, the Hierarchy Problem and High-Scale Supersymmetry,” *Phys.Rev.* **D88** (2013) 093003, [arXiv:1308.1242 \[hep-ph\]](#).

- [9] O. Antipin, M. Mojaza, and F. Sannino, “Conformal Extensions of the Standard Model with Veltman Conditions,” *Phys.Rev.* **D89** (2014) 085015, [arXiv:1310.0957 \[hep-ph\]](#).
- [10] J. Guo and Z. Kang, “Higgs Naturalness and Dark Matter Stability by Scale Invariance,” [arXiv:1401.5609 \[hep-ph\]](#).
- [11] R. Hempfling, “The Next-to-minimal Coleman-Weinberg model,” *Phys.Lett.* **B379** (1996) 153–158, [arXiv:hep-ph/9604278 \[hep-ph\]](#).
- [12] C. Tamarit, “Higgs vacua with potential barriers,” *Phys.Rev.* **D90** (2014) 055024, [arXiv:1404.7673 \[hep-ph\]](#).
- [13] W.-F. Chang, J. N. Ng, and J. M. Wu, “Shadow Higgs from a scale-invariant hidden U(1)(s) model,” *Phys.Rev.* **D75** (2007) 115016, [arXiv:hep-ph/0701254 \[hep-ph\]](#).
- [14] K. A. Meissner and H. Nicolai, “Conformal Symmetry and the Standard Model,” *Phys.Lett.* **B648** (2007) 312–317, [arXiv:hep-th/0612165 \[hep-th\]](#).
- [15] R. Foot, A. Kobakhidze, K. L. McDonald, and R. R. Volkas, “A Solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory,” *Phys.Rev.* **D77** (2008) 035006, [arXiv:0709.2750 \[hep-ph\]](#).
- [16] S. Iso, N. Okada, and Y. Orikasa, “Classically conformal $B-L$ extended Standard Model,” *Phys.Lett.* **B676** (2009) 81–87, [arXiv:0902.4050 \[hep-ph\]](#).
- [17] R. Foot, A. Kobakhidze, and R. R. Volkas, “Stable mass hierarchies and dark matter from hidden sectors in the scale-invariant standard model,” *Phys. Rev.* **D82** (2010) 035005, [arXiv:1006.0131 \[hep-ph\]](#).
- [18] K. Ishiwata, “Dark Matter in Classically Scale-Invariant Two Singlets Standard Model,” *Phys.Lett.* **B710** (2012) 134–138, [arXiv:1112.2696 \[hep-ph\]](#).
- [19] S. Iso and Y. Orikasa, “TeV Scale B-L model with a flat Higgs potential at the Planck scale - in view of the hierarchy problem -,” *PTEP* **2013** (2013) 023B08, [arXiv:1210.2848 \[hep-ph\]](#).
- [20] C. Englert, J. Jaeckel, V. Khoze, and M. Spannowsky, “Emergence of the Electroweak Scale through the Higgs Portal,” *JHEP* **1304** (2013) 060, [arXiv:1301.4224 \[hep-ph\]](#).

- [21] T. Hambye and A. Strumia, “Dynamical generation of the weak and Dark Matter scale,” *Phys.Rev.* **D88** (2013) 055022, [arXiv:1306.2329 \[hep-ph\]](#).
- [22] C. D. Carone and R. Ramos, “Classical scale-invariance, the electroweak scale and vector dark matter,” *Phys.Rev.* **D88** (2013) 055020, [arXiv:1307.8428 \[hep-ph\]](#).
- [23] E. Gabrielli, M. Heikinheimo, K. Kannike, A. Racioppi, M. Raidal, *et al.*, “Towards Completing the Standard Model: Vacuum Stability, EWSB and Dark Matter,” *Phys.Rev.* **D89** no. 1, (2014) 015017, [arXiv:1309.6632 \[hep-ph\]](#).
- [24] S. Abel and A. Mariotti, “Novel Higgs Potentials from Gauge Mediation of Exact Scale Breaking,” *Phys.Rev.* **D89** no. 12, (2014) 125018, [arXiv:1312.5335 \[hep-ph\]](#).
- [25] S. Benic and B. Radovic, “Electroweak breaking and Dark Matter from the common scale,” *Phys.Lett.* **B732** (2014) 91–94, [arXiv:1401.8183 \[hep-ph\]](#).
- [26] V. V. Khoze, C. McCabe, and G. Ro, “Higgs vacuum stability from the dark matter portal,” *JHEP* **1408** (2014) 026, [arXiv:1403.4953 \[hep-ph\]](#).
- [27] A. Farzinnia and J. Ren, “Higgs Partner Searches and Dark Matter Phenomenology in a Classically Scale Invariant Higgs Boson Sector,” *Phys.Rev.* **D90** no. 1, (2014) 015019, [arXiv:1405.0498 \[hep-ph\]](#).
- [28] W. Altmannshofer, W. A. Bardeen, M. Bauer, M. Carena, and J. D. Lykken, “Light Dark Matter, Naturalness, and the Radiative Origin of the Electroweak Scale,” *JHEP* **1501** (2015) 032, [arXiv:1408.3429 \[hep-ph\]](#).
- [29] S. Benic and B. Radovic, “Majorana dark matter in a classically scale invariant model,” *JHEP* **1501** (2015) 143, [arXiv:1409.5776 \[hep-ph\]](#).
- [30] K. Endo and Y. Sumino, “A Scale-invariant Higgs Sector and Structure of the Vacuum,” *JHEP* **1505** (2015) 030, [arXiv:1503.02819 \[hep-ph\]](#).
- [31] H. Baer, T. Barklow, K. Fujii, Y. Gao, A. Hoang, *et al.*, “The International Linear Collider Technical Design Report - Volume 2: Physics,” [arXiv:1306.6352 \[hep-ph\]](#).
- [32] D. Asner, T. Barklow, C. Calancha, K. Fujii, N. Graf, *et al.*, “ILC Higgs White Paper,” [arXiv:1310.0763 \[hep-ph\]](#).
- [33] K. Fujii, C. Grojean, M. E. Peskin, T. Barklow, Y. Gao, *et al.*, “Physics Case for the International Linear Collider,” [arXiv:1506.05992 \[hep-ex\]](#).

- [34] J. Hisano, K. Ishiwata, and N. Nagata, “QCD Effects on Direct Detection of Wino Dark Matter,” *JHEP* **1506** (2015) 097, arXiv:1504.00915 [hep-ph].
- [35] **LUX** Collaboration, D. Akerib *et al.*, “First results from the LUX dark matter experiment at the Sanford Underground Research Facility,” *Phys.Rev.Lett.* **112** (2014) 091303, arXiv:1310.8214 [astro-ph.CO].
- [36] M. Veltman, “The Infrared - Ultraviolet Connection,” *Acta Phys.Polon.* **B12** (1981) 437.
- [37] E. W. Kolb and M. S. Turner, “The Early Universe,” *Front.Phys.* **69** (1990) 1–547.
- [38] J. Hisano, K. Ishiwata, and N. Nagata, “A complete calculation for direct detection of Wino dark matter,” *Phys.Lett.* **B690** (2010) 311–315, arXiv:1004.4090 [hep-ph].
- [39] J. Hisano, K. Ishiwata, and N. Nagata, “Gluon contribution to the dark matter direct detection,” *Phys.Rev.* **D82** (2010) 115007, arXiv:1007.2601 [hep-ph].
- [40] J. Hisano, K. Ishiwata, N. Nagata, and T. Takesako, “Direct Detection of Electroweak-Interacting Dark Matter,” *JHEP* **1107** (2011) 005, arXiv:1104.0228 [hep-ph].
- [41] J. Hisano, K. Ishiwata, and N. Nagata, “Direct Search of Dark Matter in High-Scale Supersymmetry,” *Phys.Rev.* **D87** (2013) 035020, arXiv:1210.5985 [hep-ph].
- [42] R. Young and A. Thomas, “Octet baryon masses and sigma terms from an SU(3) chiral extrapolation,” *Phys.Rev.* **D81** (2010) 014503, arXiv:0901.3310 [hep-lat].
- [43] **JLQCD** Collaboration, H. Ohki *et al.*, “Nucleon strange quark content from $N_f = 2 + 1$ lattice QCD with exact chiral symmetry,” *Phys.Rev.* **D87** (2013) 034509, arXiv:1208.4185 [hep-lat].
- [44] M. A. Shifman, A. Vainshtein, and V. I. Zakharov, “Remarks on Higgs Boson Interactions with Nucleons,” *Phys.Lett.* **B78** (1978) 443.
- [45] **Planck** Collaboration, P. Ade *et al.*, “Planck 2015 results. XIII. Cosmological parameters,” arXiv:1502.01589 [astro-ph.CO].
- [46] J. Billard, L. Strigari, and E. Figueroa-Feliciano, “Implication of neutrino backgrounds on the reach of next generation dark matter direct detection

experiments,” *Phys.Rev.* **D89** no. 2, (2014) 023524,
arXiv:1307.5458 [hep-ph].

[47] D. Malling, D. Akerib, H. Araujo, X. Bai, S. Bedikian, *et al.*, “After LUX: The LZ Program,” arXiv:1110.0103 [astro-ph.IM].

[48] S. P. Martin, “Taming the Goldstone contributions to the effective potential,” *Phys.Rev.* **D90** no. 1, (2014) 016013, arXiv:1406.2355 [hep-ph].