arXiv:1507.03419v1 [hep-ph] 13 Jul 2015

Neutrino Mixing from CP Symmetry

Peng Chen*, Chang-Yuan Yao†, Gui-Jun Ding‡

Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China

Abstract

The neutrino mass matrix has remnant CP symmetry expressed in terms of the lepton mixing matrix, and vice versa the remnant CP transformations allow us to reconstruct the mixing matrix. We study the scenario that all the four remnant CP transformations are preserved by the neutrino mass matrix. The most general parameterization of remnant CP transformations is presented. The lepton mixing matrix is completely fixed by the remnant CP, and its explicit form is derived. The necessary and sufficient condition for conserved Dirac CP violating phase is found. If the Klein four flavor symmetry generated by the postulated remnant CP transformations arises from a finite flavor symmetry group, the phenomenologically viable lepton flavor mixing would be the trimaximal pattern, both Dirac CP phase δ_{CP} and Majorana phase α_{31} are either 0 or π while another Majorana phase α_{21} is a rational multiple of π. These general results are confirmed to be true in the case that the finite flavor symmetry group is $\Delta(6n^2)$.

*E-mail: pche@mail.ustc.edu.cn
†E-mail: phyman@mail.ustc.edu.cn
‡E-mail: dinggj@ustc.edu.cn

1 Introduction

The lepton flavor mixing is described by the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix which contains three mixing angles θ_{12}, θ_{13} and θ_{23} and one Dirac CP phase δ_{CP} [1]. There are two additional CP violating Majorana phases α_{21} and α_{31} which do not have any effect in neutrino oscillations, if neutrinos are Majorana particles. Many neutrino oscillation experiments have been performed all over the world and the values of mixing angles have been determined to a quite good precision [2],

$$0.270 \leq \sin^2\theta_{12} \leq 0.344, \quad 0.0188 \leq \sin^2\theta_{13} \leq 0.0251, \quad 0.385 \leq \sin^2\theta_{23} \leq 0.644 \tag{1}$$

at 3σ confidence level. Other global fits give rise to the similar results [3,4]. On the other hand, the Dirac CP violating phase δ_{CP} is weakly constrained by the present neutrino oscillation data, and its 3σ interval is $[0, 2\pi]$ [2–4]. Therefore we don't know whether there is CP violation in the lepton sector at present. More recently, the T2K data show the hint that δ_{CP} is close to $-\pi/2$ (or $3\pi/2$) [5].

In the past years, much effort has been devoted to understanding the lepton mixing angles through the introduction of a finite discrete flavor symmetry G_f, which is subsequently broken down to a Klein subgroup $G_\nu = Z_2 \times Z_2$ in the neutrino sector and an abelian group G_l in the charged lepton sector [6] under the assumption of Majorana neutrinos. In this approach, the lepton flavor mixing is completely determined by the embedding of the residual symmetries G_ν and G_l into the original group G_f. A complete classification of the mixing patterns achievable in this way have been derived under the assumption that G_f is finite [7]. It turns out that the phenomenologically viable lepton mixing can only be of the trimaximal form and the Dirac phase δ_{CP} is trivial [7], and similar results are found even if neutrinos are Dirac particles [8].

If the signal of $\delta_{CP} \simeq -\pi/2$ from T2K is further confirmed or δ_{CP} is measured to take a nontrivial value by future long baseline neutrino oscillation experiments [9–11], the paradigm of discrete flavor symmetry would be disfavored, and we may have to resort to other new theoretical framework. Bottom-up analysis shows that the effective Majorana mass term of neutrino admits both Klein four flavor symmetry [12] and four different CP symmetry transformations [13, 14], and the flavor symmetry can be generated by performing two CP transformations in succession. Moreover, CP symmetry allows us to predict the values of Majorana CP phases which are not constrained by flavor symmetry at all. As a result, CP symmetry is a more general and more fundamental approach than flavor symmetry in explaining the lepton flavor mixing. Notice that only three of the four residual CP transformations of the neutrino mass term are independent. If two (or one) remnant CP transformations out of the original CP symmetry at high energy scale are preserved by the neutrino mass term in the charged lepton diagonal basis, the lepton mixing matrix would depend on one (or three) real parameters besides the parameters specifying the remnant CP transformations, and the explicit form of the mixing matrix has been presented in Ref. [13]. In this work, we shall consider the scenario that the maximal four CP transformations are preserved by the neutrino mass matrix. The most general parameterization of the CP transformations and the formula of constructing the PMNS matrix from the postulated CP transformations would be derived. Finally we would like to point that the phenomenological predictions and model building aspects of combining flavor symmetry with CP symmetry has been extensively discussed in the literatures [15–30] in recent years. The interplay between flavor symmetry and CP symmetry have been comprehensively studied [31–34].

The remaining part of this paper is organized as follows. In section 2, we present the most general parameterization of the full set of remnant CP transformations, and the explicit form of the PMNS matrix fixed by the remnant CP is derived. In section 3, we investigate the conditions for minimal or maximal CP violating phases δ_{CP}, α_{21} and α'_{31}. In section 4,

we discuss the scenario that the Klein four flavor symmetry induced by the remnant CP transformations originates from a finite flavor symmetry group. Then the viable lepton mixing matrix can only be of the trimaximal form, both δ_{CP} and α_{31} are conserved, and α_{21} is a rational angle. As a concrete example, we apply our formalism to the case that the Klein four flavor symmetry generated by remnant CP is a subgroup of the $\Delta(6n^2)$ flavor symmetry in section 5. Finally we present our conclusions in section 6.

2 Reconstructing PMNS matrix from residual CP transformations

We would assume in the present paper that neutrinos are Majorana particles. In the flavor basis, the Lagrangian describing the lepton masses reads

$$\mathcal{L}_{mass} = -\bar{l}_R m_l l_L + \frac{1}{2}\nu_L^T C^{-1} m_\nu \nu_L + h.c.\,. \tag{2}$$

Here C is the charge-conjugation matrix, l_L and l_R denote the vectors of the three left and right-handed charged lepton fields, ν_L refers to the three left-handed neutrino fields, and $m_l \equiv \mathrm{diag}\{m_e, m_\mu, m_\tau\}$. The neutrino mass matrix m_ν in the flavor basis, can be expressed via the mixing matrix U_{PMNS} as

$$m_\nu = U^*_{PMNS}\mathrm{diag}\,(m_1, m_2, m_3)\, U^\dagger_{PMNS}\,, \tag{3}$$

where m_1, m_2 and m_3 are the light neutrino mass eigenvalues. The generic neutrino mass term in Eq. (2) is invariant under both the CP transformations

$$\nu_L(x) \longmapsto i X_j \gamma^0 C \bar{\nu}_L^T(x_P), \qquad X_j = U_{PMNS}\, d_j\, U^T_{PMNS}, \quad j = 1, 2, 3, 4 \tag{4}$$

and the flavor transformations

$$\nu_L(x) \longmapsto G_i \nu_L(x), \qquad G_i = U_{PMNS}\, d_i\, U^\dagger_{PMNS}, \quad i = 1, 2, 3, 4\,, \tag{5}$$

where $x_P = (t, -\vec{x})$ and

$$\begin{aligned} d_1 &= \mathrm{diag}\,(1, -1, -1)\,, \qquad & d_2 &= \mathrm{diag}\,(-1, 1, -1)\,, \\ d_3 &= \mathrm{diag}\,(-1, -1, 1)\,, \qquad & d_4 &= \mathrm{diag}\,(1, 1, 1)\ . \end{aligned} \tag{6}$$

As has been shown in our previous work [13], only three of the four remnant CP transformations are independent. The reason is that any one of the residual CP transformations can be generated by the remaining three via

$$X_i = X_j X^*_m X_n, \qquad i \neq j \neq m \neq n\,. \tag{7}$$

The remnant flavor symmetry and remnant CP transformations are closely related with each other. The remnant flavor symmetry can be generated by performing two CP transformations as follows,

$$\begin{aligned} G_1 &= X_2 X_3^* = X_3 X_2^* = X_4 X_1^* = X_1 X_4^*, \\ G_2 &= X_1 X_3^* = X_3 X_1^* = X_4 X_2^* = X_2 X_4^*, \\ G_3 &= X_1 X_2^* = X_2 X_1^* = X_4 X_3^* = X_3 X_4^*, \\ G_4 &= X_1 X_1^* = X_2 X_2^* = X_3 X_3^* = X_4 X_4^* = 1\,. \end{aligned} \tag{8}$$

Furthermore, we can straightforwardly see that X_i and G_j fulfill the following relation

$$X_i G_j^* X_i^* = U_{PMNS}\, d_j\, U_{PMNS}^\dagger = G_j \quad \text{for} \quad i,j = 1,2,3,4\,. \tag{9}$$

This means that the residual flavor symmetry and residual CP symmetry should generally commute with each other in the neutrino sector. Given the experimentally measured mixing matrix U_{PMNS}, the CP transformation matrix X_i can be easily fixed by Eq. (4). Inversely, we shall show that U_{PMNS} can be deduced from any well-defined four CP transformations.

The scenario of one or two remnant CP transformations preserved by the neutrino mass matrix have been investigated in our previous work [13]. In the following, we shall consider the scenario that an original CP symmetry at high energy scale is broken down to four remnant CP transformations X_i ($i = 1,2,3,4$) in the neutrino sector by some scalar fields. Notice that if three CP transformations are preserved by the neutrino mass matrix, there are still four residual CP transformations since the fourth one can be generated via Eq. (7). Here we shall not consider how the remnant CP transformations X_i ($i = 1,2,3,4$) are dynamically achieved, since the lepton flavor mixing is completely fixed by the remnant CP and it is independent of the specific breaking mechanisms. X_i ($i = 1,2,3,4$) can be treated as remnant CP transformations only if the following consistency conditions are fulfilled,

$$\begin{aligned} &X_i = X_i^T, \quad X_i X_j^* = X_j X_i^* = X_m X_n^* = X_n X_m^*, \\ &\left(X_i X_j^*\right)^2 = 1, \quad X_i X_j^* \neq \pm 1, \quad \text{for} \quad i \neq j \neq m \neq n\,. \end{aligned} \tag{10}$$

In this case, a full Klein four flavor symmetry is generated with element of the form $X_i X_j^*$ ($i \neq j$). First of all, we present the parameterization of the whole set of CP transformations in the following. Using the freedom in redefining the phases of the charged lepton fields, one column of the PMNS matrix can always be set to be real and written as

$$v_1 = \begin{pmatrix} \cos\varphi \\ \sin\varphi\cos\phi \\ \sin\varphi\sin\phi \end{pmatrix}\,, \tag{11}$$

where both φ and ϕ are real parameters in the interval of $[0, 2\pi)$. As a result, one of the remnant flavor symmetry is given by

$$G_1 = 2 v_1 v_1^\dagger - 1 = 2 v_1 v_1^T - 1\,. \tag{12}$$

Obviously G_1 is a real matrix in our working basis, and then Eq. (9) implies that

$$[X_i, G_1] = 0, \quad \text{for} \quad i = 1,2,3,4, \tag{13}$$

Therefore v_1 should be a eigenvector of X_i as well. As the remnant CP transformations X_i are symmetric and unitary matrices, the eigenvalues of X_i must be of unit modulus and the eigenvectors of X_i can be chosen to be real. Following Ref. [13], we can conveniently parameterize the remnant CP transformation X_i in terms of its eigenvalues and eigenvectors. For instance, we can write X_1 as

$$X_1 = e^{i\kappa_1} v_1 v_1^T + e^{i\kappa_2} v_2 v_2^T + e^{i\kappa_3} v_3 v_3^T\,, \tag{14}$$

where κ_1, κ_2 and κ_3 are real, v_1, v_2 and v_3 are a set of most general real orthonormal vectors, and v_2 and v_3 are given by

$$v_2 = \begin{pmatrix} \sin\varphi\cos\rho \\ -\sin\phi\sin\rho - \cos\varphi\cos\phi\cos\rho \\ \cos\phi\sin\rho - \cos\varphi\sin\phi\cos\rho \end{pmatrix}\,, \quad v_3 = \begin{pmatrix} \sin\varphi\sin\rho \\ \sin\phi\cos\rho - \cos\varphi\cos\phi\sin\rho \\ -\cos\phi\cos\rho - \cos\varphi\sin\phi\sin\rho \end{pmatrix}\,. \tag{15}$$

From the expression of X_1 in Eq. (14) and the relation of Eq. (8), X_4 can be constructed directly

$$X_4 = G_1 X_1 = e^{i\kappa_1} v_1 v_1^T - e^{i\kappa_2} v_2 v_2^T - e^{i\kappa_3} v_3 v_3^T \,. \tag{16}$$

In a similar way, we can parameterize X_2 and X_3 as follows

$$\begin{aligned} X_2 &= e^{i\lambda_1} v_1 v_1^T + e^{i\lambda_2} w_2 w_2^T + e^{i\lambda_3} w_3 w_3^T \,, \\ X_3 &= e^{i\lambda_1} v_1 v_1^T - e^{i\lambda_2} w_2 w_2^T - e^{i\lambda_3} w_3 w_3^T \,, \end{aligned} \tag{17}$$

where v_1, w_2 and w_3 also form a set of real orthonormal vectors, and w_2 and w_3 take the form

$$w_2 = \cos\xi v_2 - \sin\xi v_3, \qquad w_3 = \sin\xi v_2 + \cos\xi v_3 \,, \tag{18}$$

The residual CP transformations X_1, X_2, X_3 and X_4 given by Eqs. (14, 16, 17) have to satisfy the consistency conditions in Eq. (10). We can straightforwardly obtain the expression for the product $X_i X_j^*$ as follows

$$\begin{aligned} X_1 X_1^* &= v_1 v_1^T + v_2 v_2^T + v_2 v_3^T = 1 \,, \\ X_1 X_2^* &= e^{i(\kappa_1-\lambda_1)} v_1 v_1^T + c_{22} v_2 v_2^T + c_{33} v_3 v_3^T + c_{23} v_2 v_3^T + c_{32} v_3 v_2^T \,, \\ X_1 X_3^* &= e^{i(\kappa_1-\lambda_1)} v_1 v_1^T - c_{22} v_2 v_2^T - c_{33} v_3 v_3^T - c_{23} v_2 v_3^T - c_{32} v_3 v_2^T \,, \\ X_1 X_4^* &= v_1 v_1^T - v_2 v_2^T - v_3 v_3^T \,, \\ X_2 X_1^* &= e^{i(\lambda_1-\kappa_1)} v_1 v_1^T + c_{22}^* v_2 v_2^T + c_{33}^* v_3 v_3^T + c_{32}^* v_2 v_3^T + c_{23}^* v_3 v_2^T \,, \\ X_2 X_2^* &= v_1 v_1^T + v_2 v_2^T + v_3 v_3^T = 1 \,, \\ X_2 X_3^* &= v_1 v_1^T - v_2 v_2^T - v_3 v_3^T \,, \\ X_2 X_4^* &= e^{i(\lambda_1-\kappa_1)} v_1 v_1^T - c_{22}^* v_2 v_2^T - c_{33}^* v_3 v_3^T - c_{32}^* v_2 v_3^T - c_{23}^* v_3 v_2^T \,, \\ X_3 X_1^* &= e^{i(\lambda_1-\kappa_1)} v_1 v_1^T - c_{22}^* v_2 v_2^T - c_{33}^* v_3 v_3^T - c_{32}^* v_2 v_3^T - c_{23}^* v_3 v_2^T \,, \\ X_3 X_2^* &= v_1 v_1^T - v_2 v_2^T - v_3 v_3^T \,, \\ X_3 X_3^* &= v_1 v_1^T + v_2 v_2^T + v_3 v_3^T = 1 \,, \\ X_3 X_4^* &= e^{i(\lambda_1-\kappa_1)} v_1 v_1^T + c_{22}^* v_2 v_2^T + c_{33}^* v_3 v_3^T + c_{32}^* v_2 v_3^T + c_{23}^* v_3 v_2^T \,, \\ X_4 X_1^* &= v_1 v_1^T - v_2 v_2^T - v_3 v_3^T \,, \\ X_4 X_2^* &= e^{i(\kappa_1-\lambda_1)} v_1 v_1^T - c_{22} v_2 v_2^T - c_{33} v_3 v_3^T - c_{23} v_2 v_3^T - c_{32} v_3 v_2^T \,, \\ X_4 X_3^* &= e^{i(\kappa_1-\lambda_1)} v_1 v_1^T + c_{22} v_2 v_2^T + c_{33} v_3 v_3^T + c_{23} v_2 v_3^T + c_{32} v_3 v_2^T \,, \\ X_4 X_4^* &= v_1 v_1^T + v_2 v_2^T + v_3 v_3^T = 1 \,, \end{aligned} \tag{19}$$

where we have defined

$$\begin{aligned} c_{22} &\equiv \left(e^{-i\lambda_2}\cos^2\xi + e^{-i\lambda_3}\sin^2\xi\right) e^{i\kappa_2} \,, \\ c_{33} &\equiv \left(e^{-i\lambda_2}\sin^2\xi + e^{-i\lambda_3}\cos^2\xi\right) e^{i\kappa_3} \,, \\ c_{23} &\equiv \left(e^{-i\lambda_3} - e^{-i\lambda_2}\right) e^{i\kappa_2}\cos\xi\sin\xi \,, \\ c_{32} &\equiv \left(e^{-i\lambda_3} - e^{-i\lambda_2}\right) e^{i\kappa_3}\cos\xi\sin\xi \,. \end{aligned} \tag{20}$$

The equations in the first line of Eq. (10) give rise to the following constraints

$$e^{i(\kappa_1-\lambda_1)} = e^{i(\lambda_1-\kappa_1)}, \quad c_{22}^* = c_{22}, \quad c_{33}^* = c_{33}, \quad c_{23}^* = c_{32} \,. \tag{21}$$

Furthermore, the requirement of $(X_i X_j^*)^2 = 1$ in the second line of Eq. (10) leads to

$$\begin{aligned} &e^{2i(\kappa_1-\lambda_1)} = 1, \quad c_{22}^2 + c_{23}c_{32} = 1, \quad c_{33}^2 + c_{23}c_{32} = 1, \\ &\left(c_{22} + c_{33}\right) c_{23} = 0, \qquad \left(c_{22} + c_{33}\right) c_{32} = 0 \,. \end{aligned} \tag{22}$$

The solutions to the above set of equations in Eqs. (21,22) are given by

$$e^{i\lambda_1}=\mp e^{i\kappa_1},\quad e^{i\lambda_2}=\mp\frac{e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|},\quad e^{i\lambda_3}=\pm\frac{e^{i\kappa_2}\sin^2\xi+e^{i\kappa_3}\cos^2\xi}{\left|e^{i\kappa_2}\sin^2\xi+e^{i\kappa_3}\cos^2\xi\right|}\,, \tag{23}$$

where the "$\mp$" signs in $e^{i\lambda_1}$ can be chosen independently. As a consequence, the parameters c_{22}, c_{23}, c_{32} and c_{33} are simplified into

$$c_{22}=\mp\frac{\cos 2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|}, \tag{24}$$

$$c_{23}=\pm\frac{\cos\left(\frac{\kappa_2-\kappa_3}{2}\right)e^{i\frac{\kappa_2-\kappa_3}{2}}\sin 2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|}, \tag{25}$$

$$c_{32}=\pm\frac{\cos\left(\frac{\kappa_2-\kappa_3}{2}\right)e^{-i\frac{\kappa_2-\kappa_3}{2}}\sin 2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|}, \tag{26}$$

$$c_{33}=\pm\frac{\cos 2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|}\,. \tag{27}$$

Since the above solutions give rise to the same set of residual CP transformations $X_{1,2,3,4}$ up to an irrelevant overall "-1" factor, without loss of generality, we could choose

$$e^{i\lambda_1}=-e^{i\kappa_1},\quad e^{i\lambda_2}=-\frac{e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi}{\left|e^{i\kappa_2}\cos^2\xi+e^{i\kappa_3}\sin^2\xi\right|},\quad e^{i\lambda_3}=\frac{e^{i\kappa_2}\sin^2\xi+e^{i\kappa_3}\cos^2\xi}{\left|e^{i\kappa_2}\sin^2\xi+e^{i\kappa_3}\cos^2\xi\right|}\,. \tag{28}$$

A remnant Klein four flavor symmetry $K_4\equiv\{1,G_1,G_2,G_3\}$ can be generated by performing two CP transformations, and the three nontrivial residual flavor symmetry transformations G_i for $i=1,2,3$ can be written into

$$\begin{aligned}
G_1&=X_1X_4^*=(v_1,v_2,v_3)\begin{pmatrix}1&0&0\\0&-1&0\\0&0&-1\end{pmatrix}\begin{pmatrix}v_1^T\\v_2^T\\v_3^T\end{pmatrix}\,,\\
G_2&=X_1X_3^*=(v_1,v_2,v_3)\begin{pmatrix}-1&0&0\\0&-c_{22}&-c_{23}\\0&-c_{32}&-c_{33}\end{pmatrix}\begin{pmatrix}v_1^T\\v_2^T\\v_3^T\end{pmatrix}\,,\\
G_3&=X_1X_2^*=(v_1,v_2,v_3)\begin{pmatrix}-1&0&0\\0&c_{22}&c_{23}\\0&c_{32}&c_{33}\end{pmatrix}\begin{pmatrix}v_1^T\\v_2^T\\v_3^T\end{pmatrix}\,.
\end{aligned} \tag{29}$$

We notice that the matrix

$$\begin{pmatrix}-1&0&0\\0&c_{22}&c_{23}\\0&c_{32}&c_{33}\end{pmatrix} \tag{30}$$

can be diagonalized by a 3×3 matrix $\Theta_{3\times 3}$ with

$$\Theta_{3\times 3}=\begin{pmatrix}1&0&0\\0&\cos\theta&\sin\theta e^{i\frac{\kappa_2-\kappa_3}{2}}\\0&-\sin\theta e^{-i\frac{\kappa_2-\kappa_3}{2}}&\cos\theta\end{pmatrix}\,, \tag{31}$$

where the rotation angle θ fulfills

$$\tan 2\theta = \cos\left(\frac{\kappa_2-\kappa_3}{2}\right)\tan 2\xi\,,$$
$$\cos 2\theta = \frac{\cos 2\xi}{\sqrt{\cos^2 2\xi + \cos^2\left(\frac{\kappa_2-\kappa_3}{2}\right)\sin^2 2\xi}}\,. \tag{32}$$

In other words, we have

$$\Theta^\dagger_{3\times 3}\begin{pmatrix} -1 & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & c_{32} & c_{33} \end{pmatrix}\Theta_{3\times 3} = \mathrm{diag}(-1,-1,1)\,. \tag{33}$$

We can introduce three column vectors $\mathscr{V}_1$, $\mathscr{V}_2$ and $\mathscr{V}_3$ with

$$(\mathscr{V}_1,\mathscr{V}_2,\mathscr{V}_3) = (v_1,v_2,v_3)\,\Theta_{3\times 3}\,. \tag{34}$$

Then the residual flavor symmetries G_1, G_2 and G_3 can be written as

$$G_1 = 2\mathscr{V}_1\mathscr{V}_1^\dagger - 1\,, \qquad G_2 = 2\mathscr{V}_2\mathscr{V}_2^\dagger - 1\,, \qquad G_3 = 2\mathscr{V}_3\mathscr{V}_3^\dagger - 1\,. \tag{35}$$

The three vectors $\mathscr{V}_1$, $\mathscr{V}_2$ and $\mathscr{V}_3$ are the unique eigenvectors of G_1, G_2 and G_3 respectively with eigenvalue $+1$, and they provide the three columns of the PMNS matrix up to permutations and phases. Therefore the PMNS matrix is determined to be

$$U_{PMNS} = (\mathscr{V}_1,\mathscr{V}_2,\mathscr{V}_3)\,\mathrm{diag}(e^{i\gamma_1},e^{i\gamma_2},e^{i\gamma_3})P = (v_1,v_2,v_3)\,\Theta_{3\times 3}\,\mathrm{diag}(e^{i\gamma_1},e^{i\gamma_2},e^{i\gamma_3})P\,, \tag{36}$$

where P is an arbitrary permutation matrix since the light neutrino masses are unconstrained in the present framework. The phases γ_1, γ_2 and γ_3 are further subject to the constraints from the postulated residual CP transformations as $U^\dagger_{PMNS}X_iU^*_{PMNS} = \mathrm{diag}(\pm 1,\pm 1,\pm 1)$ in Eq. (4), and they are fixed to be

$$\mathrm{diag}\left(e^{i\gamma_1},e^{i\gamma_2},e^{i\gamma_3}\right) = \mathrm{diag}\left(e^{i\kappa_1/2},e^{i\kappa_2/2},e^{i\kappa_3/2}\right)PKP^T\,, \tag{37}$$

where K is a diagonal phase matrix with entries ± 1 or $\pm i$, it render the light neutrino to be non-negative. As a result, the PMNS matrix is completely determined by the residual CP transformations to be

$$\begin{aligned} U_{PMNS} &= (v_1,v_2,v_3)\,\Theta_{3\times 3}\,\mathrm{diag}(e^{i\kappa_1/2},e^{i\kappa_2/2},e^{i\kappa_3/2})PK \\ &= \begin{pmatrix} \cos\varphi & \sin\varphi & 0 \\ \sin\varphi\cos\phi & -\cos\varphi\cos\phi & \sin\phi \\ \sin\varphi\sin\phi & -\cos\varphi\sin\phi & -\cos\phi \end{pmatrix}\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\rho & \sin\rho \\ 0 & -\sin\rho & \cos\rho \end{pmatrix}\begin{pmatrix} e^{i\kappa_1/2} & 0 & 0 \\ 0 & e^{i\kappa_2/2} & 0 \\ 0 & 0 & e^{i\kappa_3/2} \end{pmatrix} \\ &\quad\times\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & \sin\theta \\ 0 & -\sin\theta & \cos\theta \end{pmatrix}PK\,, \end{aligned} \tag{38}$$

where the contribution from the matrix K can only possibly shift the Majorana phases by π. The different arrangements of rows and columns can be related by parameter redefinition in the present framework. Without loss of generality, we choose the column permutation P in Eq. (38) to be

$$P = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \tag{39}$$

for illustration. The predictions for the lepton mixing parameters can be straightforwardly read out as follows:

$$
\begin{aligned}
&\sin^2\theta_{13}=\cos^2\varphi, \quad \sin^2\theta_{12}=\frac{1}{2}\left(1-\cos 2\rho\cos 2\theta+\cos\frac{\kappa^\prime_2-\kappa^\prime_3}{2}\sin 2\rho\sin 2\theta\right),\\
&\sin^2\theta_{23}=\cos^2\phi, \quad \tan\delta_{CP}=\frac{\sin(\frac{\kappa^\prime_2-\kappa^\prime_3}{2})}{\sin 2\rho\cot 2\theta+\cos(\frac{\kappa^\prime_2-\kappa^\prime_3}{2})\cos 2\rho},\\
&\tan\alpha_{21}=-\frac{2\sin(\kappa^\prime_2-\kappa^\prime_3)\sin^2 2\rho\cos 2\theta+2\sin(\frac{\kappa^\prime_2-\kappa^\prime_3}{2})\sin 4\rho\sin 2\theta}{(3\cos^2 2\rho-1)\sin^2 2\theta+\cos(\kappa^\prime_2-\kappa^\prime_3)\sin^2 2\rho(\cos^2 2\theta+1)+\cos(\frac{\kappa^\prime_2-\kappa^\prime_3}{2})\sin 4\rho\sin 4\theta},\\
&\tan\alpha^\prime_{31}=-\frac{2\cos^2\rho\cos^2\theta\sin\kappa^\prime_2+2\sin^2\rho\sin^2\theta\sin\kappa^\prime_3-\sin 2\rho\sin 2\theta\sin(\frac{\kappa^\prime_2+\kappa^\prime_3}{2})}{2\cos^2\rho\cos^2\theta\cos\kappa^\prime_2+2\sin^2\rho\sin^2\theta\cos\kappa^\prime_3-\sin 2\rho\sin 2\theta\cos(\frac{\kappa^\prime_2+\kappa^\prime_3}{2})},
\end{aligned}
\tag{40}
$$

where $\alpha^\prime_{31}=\alpha_{31}-2\delta_{CP}$, δ_{CP} is the Dirac CP violating phase in standard parameterization, α_{21} and α_{31} are the Majorana CP phases [1]. Since $e^{i\kappa_1}$ can be factorized out as an overall phase of the PMNS matrix in Eq. (38), all the mixing parameters only depend on the phase differences $\kappa^\prime_2\equiv\kappa_2-\kappa_1$ and $\kappa^\prime_3\equiv\kappa_3-\kappa_1$. Notice that θ_{12} and θ_{13} only relate to ϕ and φ respectively. Moreover, the well-known Jarlskog invariant [35] is

$$
J_{CP}=\frac{1}{4}\sin(\frac{\kappa^\prime_2-\kappa^\prime_3}{2})\cos\varphi\sin^2\varphi\sin 2\phi\sin 2\theta\,. \tag{41}
$$

Notice that only the overall sign of J_{CP} could possibly change for other permutation P distinct from that in Eq. (39).

3 Conditions for vanishing and maximal CP violation

From the analytical expressions of the mixing parameters shown in Eq. (40), we can straightforwardly derive the conditions for minimal or maximal CP phases. Firstly, we find that the Jarlskog invariant J_{CP} in Eq. (41) is vanishing such that the Dirac CP violating phase δ_{CP} would be conserved if and only if

$$
\xi=0,\ \frac{1}{2}\pi,\ \pi,\ \frac{3}{2}\pi, \quad \text{or} \quad \kappa_3=\kappa_2, \quad \text{or} \quad \kappa_3=\kappa_2\pm\pi\,. \tag{42}
$$

The resulting predictions for the remaining mixing parameters are collected in Table 1 for the column permutation P of Eq. (39). Note that one element of the PMNS matrix would be vanishing such that the value of δ_{CP} can not be fixed uniquely in case of $\phi=n\pi/2$ or $\varphi=n\pi/2$ where $n=0,1,2,3$. It is remarkable that δ_{CP} is always trivial with $\sin\delta_{CP}=0$ for any value of the permutation P once the conditions in Eq. (42) are fulfilled. In other words, the Dirac CP would be violated if the parameters ξ, κ_2 and κ_3 take values distinct from those in Eq. (42). In light of the weak evidence of $\delta_{CP}\sim-\pi/2$ from T2K experiment [5], we find that maximal Dirac CP phase $\delta_{CP}=\pm\pi/2$ would necessitate the following relation

$$
\cos^2\frac{\kappa_3-\kappa_2}{2}\tan 2\xi=-\tan 2\rho, \qquad \kappa_3-\kappa_2\neq n_1\pi, \quad \rho\neq\frac{n_2\pi}{2}, \quad n_{1,2}\in\mathbb{N}\,. \tag{43}
$$

If the permutation matrix P takes a value different from Eq. (39), the analytical expression of $\tan\delta_{CP}$ would differ from that in Eq. (40) so that the corresponding condition for maximal δ_{CP} would be distinct from Eq. (43). This implies that the condition for $\cos\delta_{CP}=0$ depends on the column arrangement P.

Minimal δ_{CP} with $\sin\delta_{CP}=0$			
Conditions	$\sin^2\theta_{12}$	$\tan\alpha_{21}$	$\tan\alpha_{31}$
$\xi = n\pi/2$	$\sin^2(\rho+\xi)$	$(-1)^{n-1}\tan(\kappa'_2-\kappa'_3)$	$\begin{cases} -\tan\kappa'_3, & \text{n is odd} \\ -\tan\kappa'_2, & \text{n is even} \end{cases}$
$\kappa_3 = \kappa_2 + n\pi$	$\begin{cases} \frac{1}{2}\left[1-\text{sign}(\cos 2\xi)\cos 2\rho\right], & \text{n is odd} \\ \sin^2(\rho+\xi), & \text{n is even} \end{cases}$	0	$-\tan\kappa'_2$

Minimal α_{21} with $\sin\alpha_{21}=0$			
Conditions	$\sin^2\theta_{12}$	$\tan\delta_{CP}$	$\tan\alpha_{31}$
$\kappa_3 = \kappa_2 + n\pi$	$\begin{cases} \frac{1}{2}\left[1-\text{sign}(\cos 2\xi)\cos 2\rho\right], & \text{n is odd} \\ \sin^2(\rho+\xi), & \text{n is even} \end{cases}$	0	$-\tan\kappa'_2$
$\rho = n\pi/2$	$\frac{1}{2}\left[1-\frac{(-1)^n\cos 2\xi}{\sqrt{\cos^2 2\xi+\cos^2\left(\frac{\kappa'_2-\kappa'_3}{2}\right)\sin^2 2\xi}}\right]$	$(-1)^n\tan\frac{\kappa'_2-\kappa'_3}{2}$	$\begin{cases} -\tan\kappa'_3, & \text{n is odd} \\ -\tan\kappa'_2, & \text{n is even} \end{cases}$
$\rho = -\xi + n\pi$	$\frac{1}{2}\left(1-\sqrt{\cos^2 2\xi+\cos^2\frac{\kappa'_2-\kappa'_3}{2}\sin^2 2\xi}\right)$	$-\cot\frac{\kappa'_2-\kappa'_3}{2}\sec 2\xi$	$-\frac{\cos^2\xi\sin\kappa'_2+\sin^2\xi\sin\kappa'_3}{\cos^2\xi\cos\kappa'_2+\sin^2\xi\cos\kappa'_3}$

Table 1: The conditions for trivial Dirac phase δ_{CP} and trivial Majorana phase α_{21} with $\sin\delta_{CP}=0$ and $\sin\alpha_{21}=0$ respectively, where n is an integer. The predictions for the mixing parameters are listed here. Note that we have $\sin^2\theta_{13}=\cos^2\varphi$ and $\sin^2\theta_{23}=\cos^2\phi$ for the PMNS matrix in Eq. (38) with the permutation matrix P given by Eq. (39).

In exactly a similar way, the necessary and sufficient conditions for trivial Majorana CP phase α_{21} are determined to be

$$\kappa_3 = \kappa_2 + n\pi, \quad \text{or} \quad \rho = n\pi/2, \quad \text{or} \quad \rho = -\xi + n\pi \quad \text{with} \quad n \in \mathbb{N}\,. \tag{44}$$

The corresponding predictions for the mixing parameters are listed in Table 1. The condition of maximal α_{21} is a bit complex as follows

$$a\cos^4\frac{\kappa_2-\kappa_3}{2} + b\cos^2\frac{\kappa_2-\kappa_3}{2} + c = 0\,, \tag{45}$$

where

$$\begin{aligned} a &= 2\sin^2 2\xi\sin^2 2\rho, \\ b &= 1+\sin 4\xi\sin 4\rho+\cos 4\xi\left(1-2\cos 4\rho\right), \\ c &= -2\cos^2 2\xi\sin^2 2\rho\,. \end{aligned} \tag{46}$$

Now we turn to another Majorana phase. The redefined Majorana phase α'_{31} would be conserved if

$$\tan\rho\tan\theta = \frac{\sin\frac{\kappa'_2}{2}}{\sin\frac{\kappa'_3}{2}} \quad \text{or} \quad \frac{\cos\frac{\kappa'_2}{2}}{\cos\frac{\kappa'_3}{2}}\,. \tag{47}$$

Maximal α'_{31} requires the following equations be fulfilled,

$$\tan\rho\tan\theta = \frac{\sin\left(\frac{\pi}{4}+\frac{\kappa'_2}{2}\right)}{\sin\left(\frac{\pi}{4}+\frac{\kappa'_3}{2}\right)}, \quad \text{or} \quad \frac{\cos\left(\frac{\pi}{4}+\frac{\kappa'_2}{2}\right)}{\cos\left(\frac{\pi}{4}+\frac{\kappa'_3}{2}\right)}\,. \tag{48}$$

Notice that the above conditions for minimal or maximal α_{21} and α'_{31} vary with the permutation matrix P.

4 Induced flavor symmetry arising from finite groups

Under the assumption of Majorana neutrinos, the residual flavor symmetry of the neutrino mass matrix should be a Klein group, i.e. $G_\nu = \{1, G_1, G_2, G_3\}$ with $G_i^2 = 1$ and $G_i G_j = G_k$ for $i \neq j \neq k$. If both remnant flavor symmetries G_ν in the neutrino sector and G_l in the charged lepton sector originate from a finite flavor symmetry group G_f, then G_f would be strongly constrained. In the basis where the neutrino mass matrix m_ν is diagonal, the residual flavor symmetry transformation $G_{1,2,3}$ are given by

$$G_1 = \mathrm{diag}(1,-1,-1)\,, \qquad G_2 = \mathrm{diag}(-1,1,-1)\,, \qquad G_3 = \mathrm{diag}(-1,-1,1)\,. \tag{49}$$

The three-dimensional representation matrix of any element g of G_f can only be of the following form [7]:

$$|g| = \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{2} \end{pmatrix}, \quad \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{5}-1}{4} & \frac{\sqrt{5}+1}{4} \\ \frac{\sqrt{5}+1}{4} & \frac{1}{2} & \frac{\sqrt{5}-1}{4} \\ \frac{\sqrt{5}-1}{4} & \frac{\sqrt{5}+1}{4} & \frac{1}{2} \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\theta & \sin\theta \\ 0 & \sin\theta & \cos\theta \end{pmatrix} \tag{50}$$

up to independent permutations of rows and columns, where θ/π is a rational number, and the abstract element g and its representation matrix have been denoted by the same notation. Note that $G_{1,2,3}$ belong to the last category with $\theta = 0$. In order to obtain a viable lepton mixing matrix in the experimentally preferred 3σ ranges, the residual flavor symmetry G_l in the charged lepton sector should be generated by a single matrix T with [7]

$$T = \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} e^{i\zeta_1} & 0 & 0 \\ 0 & e^{i\zeta_2} & 0 \\ 0 & 0 & -e^{i\zeta_2} \end{pmatrix}, \tag{51}$$

where both ζ_1/π and ζ_2/π are rational numbers. The charged lepton diagonalization matrix U_l is determined by the diagonalization of T, and consequently we have

$$U_l = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ -\sqrt{2}e^{-i\frac{\zeta_2'}{2}}\cos\frac{\zeta_2'}{6} & \sqrt{2}e^{-i\frac{\zeta_2'}{2}}\cos\left(\frac{\zeta_2'}{6}+\frac{\pi}{3}\right) & \sqrt{2}e^{-i\frac{\zeta_2'}{2}}\cos\left(\frac{\zeta_2'}{6}-\frac{\pi}{3}\right) \\ \sqrt{2}ie^{-i\frac{\zeta_2'}{2}}\sin\frac{\zeta_2'}{6} & -\sqrt{2}ie^{-i\frac{\zeta_2'}{2}}\sin\left(\frac{\zeta_2'}{6}+\frac{\pi}{3}\right) & -\sqrt{2}ie^{-i\frac{\zeta_2'}{2}}\sin\left(\frac{\zeta_2'}{6}-\frac{\pi}{3}\right) \end{pmatrix}, \tag{52}$$

with $U_l^\dagger T U_l = \mathrm{diag}(-e^{i(\zeta_1+2\zeta_2)/3}, e^{i(\pi+\zeta_1+2\zeta_2)/3}, e^{i(-\pi+\zeta_1+2\zeta_2)/3})$, where $\zeta_2' \equiv \zeta_2 - \zeta_1$ is defined for simplicity. In order to predict the values of the Majorana CP phases, we impose CP symmetry onto the theory. In the present m_ν diagonal basis, the residual CP transformations of the neutrino mass matrix can only be diagonal, i.e.

$$\begin{aligned} X_1 &= \mathrm{diag}(e^{i\alpha_1}, -e^{i\alpha_2}, -e^{i\alpha_3}), \qquad & X_2 &= \mathrm{diag}(-e^{i\alpha_1}, e^{i\alpha_2}, -e^{i\alpha_3}), \\ X_3 &= \mathrm{diag}(-e^{i\alpha_1}, -e^{i\alpha_2}, e^{i\alpha_3}), \qquad & X_4 &= \mathrm{diag}(e^{i\alpha_1}, e^{i\alpha_2}, e^{i\alpha_3})\,, \end{aligned} \tag{53}$$

where α_1, α_2 and α_3 are real parameters. The same set of residual CP transformations can be obtained by solving the equation $X_i G_j^* X_i^{-1} = G_j$ given in Eq. (9). Moreover, obviously the residual flavor symmetry $G_{1,2,3}$ can be generated by successively performing two CP transformations $X_i X_j^*$ as shown in Eq. (8). Now we shall proceed to study the constraints on the phases α_1, α_2 and α_3 provided that G_f is a finite group. If we firstly perform a CP transformation X_4, subsequently a flavor symmetry transformation T, and eventually an inverse CP transformation X_4^{-1}, the theory is still invariant. The total effect of this series of transformations should amount to a flavor symmetry transformation,

$$T' \equiv X_4 T^* X_4^{-1} \in G_f\,, \tag{54}$$

which is the so-called consistency condition [16, 33, 34]. Therefore $T'T$ is an element of G_f as well, and the absolute value of $T'T$ is

$$|T'T| = \begin{pmatrix} A & B & B \\ B & D & C \\ B & C & D \end{pmatrix} , \tag{55}$$

where

$$\begin{aligned} A &= |\sin\frac{\alpha_3-\alpha_2}{2}| \,, \\ B &= \frac{1}{\sqrt{2}}|\cos\frac{\alpha_3-\alpha_2}{2}| \,, \\ C &= \frac{1}{2\sqrt{2}}\sqrt{3-\cos(\alpha_3-\alpha_2)-2\cos(\alpha_2-\alpha_1+\zeta_2')+2\cos(\alpha_3-\alpha_1+\zeta_2')} \,, \\ D &= \frac{1}{2\sqrt{2}}\sqrt{3-\cos(\alpha_3-\alpha_2)-2\cos(\alpha_3-\alpha_1+\zeta_2')+2\cos(\alpha_2-\alpha_1+\zeta_2')} \,. \end{aligned} \tag{56}$$

If the flavor symmetry group generated by $G_{1,2,3}$, T and T' is finite, $|T'T|$ has to be of the forms in Eq. (50) up to row and column permutations. As a result, B can only be equal to 0, $\frac{1}{2}$ or $\frac{1}{\sqrt{2}}$. Then we can derive that

$$\alpha_3 = \alpha_2, \quad \text{or} \quad \alpha_3 = \alpha_2 \pm \pi \,. \tag{57}$$

Since both solutions $\alpha_3 = \alpha_2$ and $\alpha_3 = \alpha_2 \pm \pi$ give rise to the same set of residual CP transformations $X_{1,2,3,4}$ except an inessential overall "-1" factor, we shall choose $\alpha_3 = \alpha_2$ in the following without loss of generality. The eigenvalues of $T'^{\dagger}T$ are $e^{i(\alpha_1-\alpha_2+2\zeta_1)}$, $e^{2i\zeta_2}$ and $e^{i(\alpha_2-\alpha_1+2\zeta_2)}$. Once the flavor symmetry group G_f is finite, the order of $T'^{\dagger}T$ must be finite as well. As a consequence, $\alpha_2' \equiv \alpha_2 - \alpha_1$ a rational multiple of π. The neutrino mass matrix m_ν is constrained by the CP symmetry as

$$X_i^T m_\nu X_i = m_\nu^* \,. \tag{58}$$

Therefore m_ν is fixed to be

$$m_\nu = \text{diag}(m_1 e^{-i\alpha_1}, m_2 e^{-i\alpha_2}, m_3 e^{-i\alpha_2}) \,, \tag{59}$$

where m_1, m_2 and m_3 are real. The unitary diagonalization matrix U_ν, which fulfills $U_\nu^T m_\nu U_\nu = \text{diag}(|m_1|, |m_2|, |m_3|)$, is given by

$$U_\nu = \begin{pmatrix} e^{i\frac{\alpha_1}{2}} & 0 & 0 \\ 0 & e^{i\frac{\alpha_2}{2}} & 0 \\ 0 & 0 & e^{i\frac{\alpha_2}{2}} \end{pmatrix} K \,, \tag{60}$$

where K is a diagonal matrix with entries ± 1 or $\pm i$ which encode the CP parity of the neutrino states and it makes the light neutrino masses positive. Notice that U_ν is determined up to permutations and phases of its column vectors, since the order of the neutrino masses is undefined in this approach. Compared with the scenario with only flavor symmetry, the unitary matrix U_ν can be any diagonal phase matrix.

Combining the unitary transformations U_l in Eq. (52) and U_ν in Eq. (60), we can obtain the prediction for the lepton mixing matrix:

$$U_{PMNS} = \frac{1}{\sqrt{3}} \begin{pmatrix} -\sqrt{2}\cos\frac{\zeta_2'}{6} & 1 & \sqrt{2}\sin\frac{\zeta_2'}{6} \\ \sqrt{2}\cos\left(\frac{\zeta_2'}{6}+\frac{\pi}{3}\right) & 1 & -\sqrt{2}\sin\left(\frac{\zeta_2'}{6}+\frac{\pi}{3}\right) \\ \sqrt{2}\cos\left(\frac{\zeta_2'}{6}-\frac{\pi}{3}\right) & 1 & -\sqrt{2}\sin\left(\frac{\zeta_2'}{6}-\frac{\pi}{3}\right) \end{pmatrix} \begin{pmatrix} e^{i\frac{\zeta_2'+\alpha_2'}{2}} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -ie^{i\frac{\zeta_2'+\alpha_2'}{2}} \end{pmatrix} K \,. \tag{61}$$

Permuting the rows and columns of this mixing matrix while keeping the vector $(1,1,1)^T/\sqrt{3}$ in the second column, we find the resulting PMNS matrix can be obtained from Eq. (61) by redefinition of the parameters ζ_2' and α_2'. On the other hand, we can also turn to the charged lepton diagonal basis (i.e. the T diagonal basis) by performing the similarity transformation U_l of Eq. (52). The residual CP transformations X_i in Eq. (53) become $U_l^\dagger X_i U_l^*$ which are characterized by the parameter values,

$$\begin{aligned} &\varphi = \arccos\sqrt{\frac{1}{3}}, \quad \phi = \frac{\pi}{4}, \quad \rho = -\frac{\zeta_2'}{6}, \quad \xi = 0, \\ &\kappa_1 = \alpha_1, \quad \kappa_2 = \pi + \alpha_2 + \zeta_2', \quad \kappa_3 = \alpha_2 + \zeta_2' \,. \end{aligned} \tag{62}$$

Then the PMNS matrix can be easily obtained via the general formula of Eq. (38). We can further straightforwardly read out the lepton mixing parameters as follows

$$\begin{aligned} &\sin^2\theta_{13} = \frac{2}{3}\sin^2\frac{\zeta_2'}{6}, \quad \sin^2\theta_{12} = \frac{1}{2+\cos\frac{\zeta_2'}{3}}, \quad \sin^2\theta_{23} = \frac{1}{2}\left(1+\frac{\sqrt{3}\sin\frac{\zeta_2'}{3}}{2+\cos\frac{\zeta_2'}{3}}\right), \\ &\tan\delta_{CP} = 0, \quad \tan\alpha_{21} = -\tan\left(\zeta_2'+\alpha_2'\right), \quad \tan\alpha_{31} = 0\,. \end{aligned} \tag{63}$$

It is remarkable that both Dirac phase δ_{CP} and the Majorana phase α_{31} are trivial, and another Majorana phase $\alpha_{21} = -\zeta_2' - \alpha_2'$ or $\alpha_{21} = \pi - \zeta_2' - \alpha_2'$ is a rational angle. Note that these predictions for the CP phases are independent of the structure of G_f. These results are exactly found to be true in the context of $G_f = \Delta(6n^2)$ flavor symmetry with generalized CP, as will be discussed in the following section. We can see that the three mixing angles depend on only one parameter ζ_2' with period of 6π. Without loss of generality, the fundamental interval of the parameter ζ_2' can be taken to be $[-3\pi, 3\pi)$. Imposing the present experimentally favored 3σ experiment value [2], ζ_2'/π is a rational number satisfying $0.322 \leq \pm\zeta_2'/\pi \leq 0.373$. Moreover, the three mixing angles are strongly correlated with each other,

$$3\cos^2\theta_{13}\sin^2\theta_{12} = 1, \qquad \sin^2\theta_{23} = \frac{1}{2} \pm \frac{1}{2}\tan\theta_{13}\sqrt{2-\tan^2\theta_{13}}\,. \tag{64}$$

Given the 3σ range $0.0188 \leq \sin^2\theta_{13} \leq 0.0251$, we obtain $0.340 \leq \sin^2\theta_{12} \leq 0.342$, $0.387 \leq \sin^2\theta_{23} \leq 0.403$ or $0.597 \leq \sin^2\theta_{23} \leq 0.613$. Note that the atmospheric angle θ_{23} deviates from maximal mixing. These predictions for θ_{12} and θ_{23} can be tested by forthcoming neutrino oscillation experiments such as JUNO [36], LBNE [9] and LBNO [10]. Moreover, dedicated long baseline experiments LBNE [9], LBNO [10] and Hyper-Kamiokande [11] aim to measure the Dirac CP phase δ_{CP}, the present scenario will be ruled out if the signal of leptonic CP violation will be detected.

The neutrinoless double beta ($0\nu\beta\beta$) decay is the only feasible experiment which has the potential of establishing the Majorana nature of massive neutrinos. The dependence of the decay rate on the mixing parameters is specified by the effective Majorana mass,

$$|m_{ee}| = \left|m_1\cos^2\theta_{12}\cos^2\theta_{13} + m_2\sin^2\theta_{12}\cos^2\theta_{13}e^{i\alpha_{21}} + m_3\sin^2\theta_{13}e^{i(\alpha_{31}-2\delta_{CP})}\right| \,. \tag{65}$$

For the predicted mixing pattern in Eq. (61), we have

$$|m_{ee}| = \frac{1}{3}\left|2s_1m_1\cos^2\frac{\zeta_2'}{6} + m_2e^{-i(\zeta_2'+\alpha_2')} - 2s_3m_3\sin^2\frac{\zeta_2'}{6}\right| \,, \tag{66}$$

where $s_{1,3} = \pm 1$ arises from the ambiguity of the phase matrix K. The parameter α_2' freely varies in the region of $0 \leq \alpha_2' \leq 2\pi$ and ζ_2' is scattered in the viable ranges of $0.322\pi \leq \pm\zeta_2' \leq 0.373\pi$. The resulting predictions for the effective mass $|m_{ee}|$ are plotted in Fig. 1, where the 3σ uncertainties of the mass-squared splittings Δm^2_{21} and $\Delta m^2_{3\ell}$ with $\ell = 1$ for normal ordering and $\ell = 2$ for inverted ordering are included [2]. We see that all possible values of $|m_{ee}|$ allowed by experimental data at 3σ level can be achieved in case of inverted ordering mass spectrum.




Figure 1: The prediction for the $0\nu\beta\beta$ decay effective mass if the Klein flavor symmetry generated by the residual CP symmetry originates from a finite flavor symmetry group. In this case, the PMNS matrix is given in Eq. (61), both δ_{CP} and α_{31} are predicted to be trivial, and α_{21} is a rational angle. The blue dashed lines and the red dashed lines indicate the currently allowed 3σ regions for normal ordering and inverted ordering mass spectrum respectively [2]. The cyan and orange areas are theoretical predictions when α_2' freely varies in the interval $[0, 2\pi]$ and ζ_2' in the viable range $0.322\pi \leq \pm\zeta_2' \leq 0.373\pi$. Measurements of EXO-200 [37,38] in combination with KamLAND-ZEN [39] give rise to the upper bound of $|m_{ee}| < 0.120$ eV. The upper limit on the mass of the lightest neutrino is derived from the latest Planck result $m_1 + m_2 + m_3 < 0.230$ eV at 95% level [40].

5 Example with $G_f = \Delta(6n^2)$

In what follows, we shall substantiate the statement that the lepton mixing matrix must be the trimaximal pattern in Eq. (61) if the residual flavor symmetry generated by the residual CP transformations originates from a finite group G_f. As a example and a further check to our general results, we consider the case of $G_f = \Delta(6n^2)$. $\Delta(6n^2)$ is a series of non-abelian finite subgroup of $SU(3)$, and it can be generated by four generators a, b, c and d which fulfill [41]

$$\begin{gathered} a^3 = b^2 = (ab)^2 = 1, \\ c^n = d^n = 1, \quad cd = dc, \\ aca^{-1} = c^{-1}d^{-1}, \quad ada^{-1} = c, \quad bcb^{-1} = d^{-1}, \quad bdb^{-1} = c^{-1}\,. \end{gathered} \tag{67}$$

$\Delta(6n^2)$ group has $2n-2$ three dimensional irreducible representations denoted by $\mathbf{3}_{l,k}$, and representation matrices of the generators are given by

$$a = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad b = (-1)^l \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad c = \begin{pmatrix} \eta^k & 0 & 0 \\ 0 & \eta^{-k} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad d = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \eta^k & 0 \\ 0 & 0 & \eta^{-k} \end{pmatrix}, \tag{68}$$

where $\eta = e^{2\pi i/n}$, $l = 1, 2$ and $k = 1, 2, \ldots, n-1$. We see that the set of all matrices describing distinct triplet representations $\mathbf{3}_{l,k}$ is the same up to a possible overall "-1" factor. As a consequence, it is sufficient to focus on a single three-dimensional representation in the discussion of the mixing patterns. We shall assign the three generations of left-handed leptons to the three-dimensional representation $\mathbf{3}_{1,1}$ in the following. To simplify the notation, the abstract elements of $\Delta(6n^2)$ and their representation matrices have been denoted by the same symbol.

$\Delta(6n^2)$ as a flavor symmetry group has been studied comprehensively [17, 26, 27, 42]. It is found that CP symmetry can be consistently defined in the context of $\Delta(6n^2)$ family symmetry if n is not divisible by 3 or the fields transforming as two dimensional irreducible representation $\mathbf{2_2}$, $\mathbf{2_3}$ or $\mathbf{2_4}$ are not present in a concrete model [27]. In particular, the generalized CP transformations are of the same form as the flavor symmetry transformation in the working basis [27]. Phenomenologically viable lepton flavor mixing can be obtained if $\Delta(6n^2)$ is broken down to $G_l = Z_3^a \equiv \{1, a, a^2\}$ in the charged lepton sector and to Klein subgroup $G_\nu = K_4^{(c^{n/2}, abc^\gamma)} \equiv \left\{1, c^{n/2}, abc^\gamma, abc^{\gamma+n/2}\right\}$ in the neutrino sector [8,17,42], where $\gamma = 0, 1, \ldots, n-1$ and n should be even. In this work, we propose to start from CP symmetry rather than flavor symmetry. The remnant flavor symmetry $K_4^{(c^{n/2}, abc^\gamma)}$ can be generated if impose the following CP transformations

$$X_1 = c^s d^{2(s+\gamma)}, \quad X_2 = abc^{s+\gamma} d^{2(s+\gamma)}, \quad X_3 = abc^{s+\gamma+n/2} d^{2(s+\gamma)}, \quad X_4 = c^{s+n/2} d^{2(s+\gamma)}\,, \quad (69)$$

where $s = 0, 1, \ldots, n-1$. Note that the remnant symmetry $G_l = Z_3^a$ can be generated by the CP transformations $X_l = b,\ ab,\ a^2 b$. The generator a and the hermitian combination $m_l^\dagger m_l$ are diagonalized by the same unitary transformation U_l with

$$U_l = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \\ 1 & 1 & 1 \end{pmatrix}, \quad U_l^\dagger a U_l = \mathrm{diag}\left(1, \omega^2, \omega\right)\,, \quad (70)$$

where $\omega = e^{2\pi i/3}$. We now apply our method to this specific example. Firstly we perform a change of basis with the unitary matrix U_l, such that the charged lepton mass matrix would be diagonal. The postulated residual CP transformations in Eq. (69) transform into $U_l^\dagger X_i U_l^*$, they can be parameterized in the manner shown in section 2, and the parameter values are given by

$$\begin{aligned} &\varphi = \arccos\frac{1}{\sqrt{3}}, \quad \phi = \frac{\pi}{4}, \quad \rho = -\frac{\gamma\pi}{n}, \quad \xi = 0, \\ &\kappa_1 = -\frac{4(s+\gamma)\pi}{n}, \quad \kappa_2 = \frac{2(s+\gamma)\pi}{n}, \quad \kappa_3 = \pi + \frac{2(s+\gamma)\pi}{n}\,. \end{aligned} \quad (71)$$

Using our formula for the PMNS matrix in Eq. (38), we can construct the mixing matrix as

$$U_{PMNS} = \frac{1}{\sqrt{3}} \begin{pmatrix} -\sqrt{2}\cos\frac{\gamma\pi}{n} & 1 & \sqrt{2}\sin\frac{\gamma\pi}{n} \\ \sqrt{2}\cos\left(\frac{\gamma\pi}{n}+\frac{\pi}{3}\right) & 1 & -\sqrt{2}\sin\left(\frac{\gamma\pi}{n}+\frac{\pi}{3}\right) \\ \sqrt{2}\cos\left(\frac{\gamma\pi}{n}-\frac{\pi}{3}\right) & 1 & -\sqrt{2}\sin\left(\frac{\gamma\pi}{n}-\frac{\pi}{3}\right) \end{pmatrix} \begin{pmatrix} e^{\frac{3i\pi(s+\gamma)}{n}} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -ie^{\frac{3i\pi(s+\gamma)}{n}} \end{pmatrix} K\,, \quad (72)$$

which is of the same form as our general result of Eq. (61). As a consequence, the lepton mixing parameters are given by Eq. (63) with $\zeta_2' = \frac{6\gamma\pi}{n}$ and $\alpha_2' = \frac{6s\pi}{n}$. In particularly, both δ_{CP} and α_{31} are conserved, and $\alpha_{21} = -\frac{6(s+\gamma)\pi}{n}$ or $\alpha_{21} = \pi - \frac{6(s+\gamma)\pi}{n}$. The same results are obtained in Ref. [17]. We show the possible values of the Majorana phase α_{21} for each $\Delta(6n^2)$ group of even n in Fig. 2. For large n, we see that the predictions for α_{21} densely fill the whole range of $[0, 2\pi]$. For small values of n, the measured value of reactor angle θ_{13} can not be generated, and the corresponding values of α_{21} are plotted in blue color. The red points denote the possible values of α_{21} when the experimentally favored 3σ ranges of the mixing angles are taken into account.




Figure 2: The possible values of the Majorana phase α_{21} predicted by CP symmetry if the residual Klein flavor symmetry generated by the residual CP originates from the $\Delta(6n^2)$ group with even n. The red points denote the predictions for α_{21}, where the lepton mixing angles are required to lie in the experimentally favored 3σ intervals [2]. The blue points represent that the measured values of θ_{13} can not be accommodated.

6 Conclusions

CP symmetry is a more general framework than the flavor symmetry in constraining the lepton flavor mixing, since flavor symmetry can be generated by performing two CP transformations. Compared with flavor symmetry, all mixing parameters in particular the Majorana phases can be predicted by CP symmetry. In the charged lepton diagonal basis, a generic Majorana neutrino mass matrix has four remnant CP transformations which are determined by the experimentally measured PMNS matrix. Note that only three of the four remnant CP transformations are independent. Conversely the lepton flavor mixing PMNS matrix can be constructed from the postulated remnant CP symmetries. If only one CP transformation is preserved by the neutrino mass matrix, the PMNS matrix would be determined up to an arbitrary real orthogonal matrix. If two CP transformations are preserved in the neutrino sector, the PMNS matrix would depend on a single real free parameters besides the parameters characterizing the remnant CP transformations. The explicit form of the PMNS matrix for both one and two remnant CP has been derived in our previous work [13].

In the present work, we have considered the scenario that four CP transformations out of the original CP symmetry at high energy scale are conserved by the neutrino mass matrix. A remnant Klein four flavor symmetry would be generated in this case. We firstly present the most general parameterization of the four remnant CP transformations, and then the reconstruction formula for the PMNS matrix is derived. We see that the PMNS matrix including the Majorana phases are completely fixed by the postulated four remnant CP transformations. From the explicit form of the PMNS matrix, the necessary and sufficient condition for conserved Dirac CP violating phases is determined to be $\xi = 0,\ \pi/2,\ \pi,\ 3\pi/2$, $\kappa_3 = \kappa_2$ or $\kappa_3 = \kappa_2 \pm \pi$. In the same fashion, we find the conditions for maximal δ_{CP}, minimal Majorana phases and maximal Majorana phases for the column permutation P given by Eq. (39).

Furthermore, we discuss the situation that the Klein four flavor symmetry induced by the remnant CP originates from a finite flavor symmetry group. It turns out that the lepton flavor mixing would be strongly constrained. The phenomenologically viable PMNS matrix

can only take the trimaximal form, and it depends on two rational angles ζ'_2 and α'_2, as shown in Eq. (61). As a consequence, the three lepton mixing angles are correlated with each other. Given the measured values of the reactor angle θ_{13}, we have $\sin^2\theta_{12} \simeq 0.341$, $\sin^2\theta_{23} \simeq 0.395$ or $\sin^2\theta_{23} \simeq 0.605$. Regarding the CP violating phases, both δ_{CP} and α_{31} are determined to be 0 or π while another Majorana phase α_{21} can be any rational angle. These predictions can be tested by more precise neutrino oscillation experiments in near future. In addition, the corresponding predictions for the neutrinoless double beta decay are studied. The 3σ region of the effective mass $|m_{ee}|$ can be nearly reproduced in the case of inverted ordering neutrino mass spectrum. As a concrete example, we further consider the case that the induced Klein four flavor symmetry arises from the $\Delta(6n^2)$ flavor symmetry group. The PMNS matrix is really found to be the trimaximal pattern with $\zeta'_2 = 6\gamma\pi/n$ and $\alpha'_2 = 6s\pi/n$, where $\gamma, s = 0, 1, 2, \ldots, n-1$. The above general results are confirmed.

Acknowledgements

This work is supported by the National Natural Science Foundation of China under Grant Nos. 11275188 and 11179007.

References

[1] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014).

[2] M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, JHEP **1411**, 052 (2014) [arXiv:1409.5439 [hep-ph]].

[3] F. Capozzi, G. L. Fogli, E. Lisi, A. Marrone, D. Montanino and A. Palazzo, Phys. Rev. D **89**, 093018 (2014) [arXiv:1312.2878 [hep-ph]].

[4] D. V. Forero, M. Tortola and J. W. F. Valle, Phys. Rev. D **90**, no. 9, 093006 (2014) [arXiv:1405.7540 [hep-ph]].

[5] K. Abe *et al.* [T2K Collaboration], Phys. Rev. D **91**, no. 7, 072010 (2015) [arXiv:1502.01550 [hep-ex]].

[6] G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701 (2010) [arXiv:1002.0211 [hep-ph]]; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada and M. Tanimoto, Prog. Theor. Phys. Suppl. **183**, 1 (2010) [arXiv:1003.3552 [hep-th]]; S. F. King and C. Luhn, Rept. Prog. Phys. **76** (2013) 056201 [arXiv:1301.1340 [hep-ph]]; S. F. King, A. Merle, S. Morisi, Y. Shimizu and M. Tanimoto, New J. Phys. **16**, 045018 (2014) [arXiv:1402.4271 [hep-ph]].

[7] R. M. Fonseca and W. Grimus, JHEP **1409**, 033 (2014) [arXiv:1405.3678 [hep-ph]].

[8] C. Y. Yao and G. J. Ding, arXiv:1505.03798 [hep-ph].

[9] C. Adams *et al.* [LBNE Collaboration], arXiv:1307.7335 [hep-ex]; M. Bass *et al.* [LBNE Collaboration], arXiv:1311.0212 [hep-ex].

[10] S. K. Agarwalla *et al.* [LAGUNA-LBNO Collaboration], JHEP **1405**, 094 (2014) [arXiv:1312.6520 [hep-ph]]; L. Agostino, B. Andrieu, R. Asfandiyarov, D. Autiero, O. Bsida, F. Bay, R. Bayes and A. M. Blebea-Apostu *et al.*, arXiv:1409.4405 [physics.ins-det].

[11] K. Abe, T. Abe, H. Aihara, Y. Fukuda, Y. Hayato, K. Huang, A. K. Ichikawa and M. Ikeda *et al.*, arXiv:1109.3262 [hep-ex]; E. Kearns *et al.* [Hyper-Kamiokande Working Group Collaboration], arXiv:1309.0184 [hep-ex].

[12] C. S. Lam, Phys. Rev. D **87**, no. 1, 013001 (2013) [arXiv:1208.5527 [hep-ph]].

[13] P. Chen, C. C. Li and G. J. Ding, Phys. Rev. D **91**, no. 3, 033003 (2015) [arXiv:1412.8352 [hep-ph]].

[14] L. L. Everett, T. Garon and A. J. Stuart, JHEP **1504**, 069 (2015) [arXiv:1501.04336 [hep-ph]].

[15] R. N. Mohapatra and C. C. Nishi, Phys. Rev. D **86**, 073007 (2012) [arXiv:1208.2875 [hep-ph]].

[16] F. Feruglio, C. Hagedorn and R. Ziegler, JHEP **1307**, 027 (2013) [arXiv:1211.5560 [hep-ph]].

[17] S. F. King and T. Neder, Phys. Lett. B **736**, 308 (2014) [arXiv:1403.1758 [hep-ph]].

[18] G. J. Ding, S. F. King and A. J. Stuart, JHEP **1312**, 006 (2013) [arXiv:1307.4212 [hep-ph]].

[19] G. J. Ding, S. F. King, C. Luhn and A. J. Stuart, JHEP **1305**, 084 (2013) [arXiv:1303.6180 [hep-ph]].

[20] F. Feruglio, C. Hagedorn and R. Ziegler, Eur. Phys. J. C **74**, 2753 (2014) [arXiv:1303.7178 [hep-ph]].

[21] C. Luhn, Nucl. Phys. B **875**, 80 (2013) [arXiv:1306.2358 [hep-ph]].

[22] C. C. Li and G. J. Ding, Nucl. Phys. B **881**, 206 (2014) [arXiv:1312.4401 [hep-ph]].

[23] C. C. Li and G. J. Ding, arXiv:1408.0785 [hep-ph].

[24] G. J. Ding and Y. L. Zhou, Chin. Phys. C **39**, 021001 (2015) [arXiv:1312.5222 [hep-ph]]; G. J. Ding and Y. L. Zhou, JHEP **1406**, 023 (2014) [arXiv:1404.0592 [hep-ph]].

[25] G. J. Ding and S. F. King, Phys. Rev. D **89**, no. 9, 093020 (2014) [arXiv:1403.5846 [hep-ph]].

[26] C. Hagedorn, A. Meroni and E. Molinaro, Nucl. Phys. B **891**, 499 (2015) [arXiv:1408.7118 [hep-ph]].

[27] G. J. Ding, S. F. King and T. Neder, JHEP **1412**, 007 (2014) [arXiv:1409.8005 [hep-ph]].

[28] C. C. Li and G. J. Ding, JHEP **1505**, 100 (2015) [arXiv:1503.03711 [hep-ph]].

[29] A. Di Iura, C. Hagedorn and D. Meloni, arXiv:1503.04140 [hep-ph].

[30] P. Ballett, S. Pascoli and J. Turner, arXiv:1503.07543 [hep-ph].

[31] G. Ecker, W. Grimus and W. Konetschny, Nucl. Phys. B **191**, 465 (1981); G. Ecker, W. Grimus and H. Neufeld, Nucl. Phys. B **247**, 70 (1984); J. Phys. A **20**, L807 (1987); H. Neufeld, W. Grimus and G. Ecker, Int. J. Mod. Phys. A **3**, 603 (1988).

[32] W. Grimus and M. N. Rebelo, Phys. Rept. **281**, 239 (1997) [hep-ph/9506272].

[33] M. Holthausen, M. Lindner and M. A. Schmidt, JHEP **1304**, 122 (2013) [arXiv:1211.6953 [hep-ph]].

[34] M. C. Chen, M. Fallbacher, K. T. Mahanthappa, M. Ratz and A. Trautner, Nucl. Phys. B **883**, 267 (2014) [arXiv:1402.0507 [hep-ph]].

[35] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985).

[36] JUNO experiment, `http://english.ihep.cas.cn/rs/fs/juno0815/`.

[37] M. Auger *et al.* [EXO Collaboration], Phys. Rev. Lett. **109**, 032505 (2012) [arXiv:1205.5608 [hep-ex]].

[38] J. B. Albert *et al.* [EXO-200 Collaboration], Nature **510** (2014) 229-234 [arXiv:1402.6956 [nucl-ex]].

[39] A. Gando *et al.* [KamLAND-Zen Collaboration], Phys. Rev. Lett. **110** (2013) 062502 [arXiv:1211.3863 [hep-ex]].

[40] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571**, A16 (2014) [arXiv:1303.5076 [astro-ph.CO]].

[41] J. A. Escobar and C. Luhn, J. Math. Phys. **50**, 013524 (2009) [arXiv:0809.0639 [hep-th]].

[42] S. F. King, T. Neder and A. J. Stuart, Phys. Lett. B **726** (2013) 312 [arXiv:1305.3200 [hep-ph]].