

Some Nearly Quantum Theories

Howard Barnum
University of New Mexico
Albuquerque, USA
hnbarnum@ao1.com

Matthew A. Graydon
Perimeter Institute for Theoretical Physics
Waterloo, Canada
Institute for Quantum Computing, University of Waterloo
Waterloo, Canada
mgraydon@perimeterinstitute.ca

Alexander Wilce
Susquehanna University
Selinsgrove, USA
wilce@susqu.edu

We consider possible non-signaling composites of probabilistic models based on euclidean Jordan algebras. Subject to some reasonable constraints, we show that no such composite exists having the exceptional Jordan algebra as a direct summand. We then construct several dagger compact categories of such Jordan-algebraic models. One of these neatly unifies real, complex and quaternionic mixed-state quantum mechanics, with the exception of the quaternionic “bit”. Another is similar, except in that (i) it excludes the quaternionic bit, and (ii) the composite of two complex quantum systems comes with an extra classical bit. In both of these categories, states are morphisms from systems to the tensor unit, which helps give the categorical structure a clear operational interpretation. A no-go result shows that the first of these categories, at least, cannot be extended to include spin factors other than the (real, complex, and quaternionic) quantum bits, while preserving the representation of states as morphisms. The same is true for attempts to extend the second category to even-dimensional spin-factors. Interesting phenomena exhibited by some composites in these categories include failure of local tomography, supermultiplicativity of the maximal number of mutually distinguishable states, and mixed states whose marginals are pure.

1 Introduction

A series of recent papers [17, 12, 18, 16, 5] have shown that any of various packages of probabilistic or information-theoretic axioms force the state spaces of a finite-dimensional probabilistic theory to be those of formally real, or euclidean, Jordan algebras. Thus, euclidean Jordan algebras (hereafter, EJAs) form a natural class of probabilistic models. Moreover, it is one that keeps us in the general neighborhood of standard quantum mechanics, owing to the classification of simple EJAs as self-adjoint parts of real, complex and quaternionic matrix algebras (corresponding to real, complex and quaternionic quantum systems), the exceptional Jordan algebra of self-adjoint 3×3 matrices over the octonions, and one further class, the so-called *spin factors*. The latter are essentially “bits”: their state-spaces are balls of arbitrary dimension, with pairs of antipodal points representing maximal sets of sharply distinguishable states.¹

This raises the question of whether one can construct probabilistic *theories* (as opposed to a collection of models of individual systems) in which finite-dimensional complex quantum systems can be accommodated together with several — perhaps all — of the other basic types of EJAs listed above. Ideally, these would be symmetric monoidal categories; even better, we might hope to obtain compact closed, or still better, dagger-compact, categories of EJAs [1]. Also, one would like the resulting theory to embrace mixed states and CP mappings.

In this paper, we exhibit two dagger-compact categories of EJAs — called **URUE** and **URSE**, acronyms

¹Where this ball has dimension 2, 3 or 5, these are just the state spaces of real, complex and quaternionic quantum bits.

that will be explained below — that include all real, complex and quaternionic matrix algebras, with one exception: the quaternionic bit, or “qubit”, represented by $M_2(\mathbb{H})_{\text{sa}}$, cannot be added to **URUE** without destroying compact closure and the representation of states as morphisms. **URSE** includes a faithful copy of finite-dimensional complex quantum mechanics, while in **URUE**, composites of complex quantum systems come with an extra classical bit — that is, a $\{0, 1\}$ valued superselection rule.

There is scant hope of including more exotic Jordan algebras in a satisfactory categorical scheme. Even allowing for a very liberal definition of composite (our Definition 1 below), the exceptional Jordan algebra is ruled out altogether (Corollary 1), while non-quantum spin factors are ruled out if we want to regard states as morphisms — in particular, if we demand compact closure (see Example 1). Combined with the results of (any of) the papers cited above that derive a euclidean Jordan-algebraic structure from information-theoretic assumptions, these results provide a compelling motivation for a kind of unified quantum theory that accommodates real, complex and quaternionic quantum systems (possibly modulo the qubit) and permits the formation of composites of these.

A condition frequently invoked to rule out real and quaternionic QM is *local tomography*: the doctrine that the state of a composite of two systems should depend only on the joint probabilities it assigns to measurement outcomes on the component systems. Indeed, it can be shown [7] that standard complex QM with superselection rules is the only dagger-compact category of EJAs that includes the qubit. Accordingly, **URUE** and **URSE** are *not* locally tomographic. In our view, the very existence of these quite reasonable, well-behaved categories suggests that local tomography is not as well-motivated as is sometimes supposed.

A broadly similar proposal is advanced by Baez [3], who points out that one can view real and quaternionic quantum systems as pairs (\mathbf{H}, J) , where \mathbf{H} is a complex Hilbert space and J is an anti-unitary satisfying $J^2 = \mathbf{1}$ (the real case) or $J^2 = -\mathbf{1}$ (the quaternionic case). This yields a symmetric monoidal category in which objects are such pairs, morphisms $(\mathbf{H}_1, J_1) \rightarrow (\mathbf{H}_2, J_2)$ are linear mappings intertwining J_1 and J_2 , and $(\mathbf{H}_1, J_1) \otimes (\mathbf{H}_2, J_2) = (\mathbf{H}_1 \otimes \mathbf{H}_2, J_1 \otimes J_2)$. The precise connection between this approach and ours is still under study.

A forthcoming paper [4] will provide more details, including full proofs of the results presented here, as well as additional results obtained since our presentation of this paper at QPL XII.

2 Euclidean Jordan algebras

We begin with a concise review of some basic Jordan-algebraic background. References for this section are [2] and [8]. A *euclidean Jordan algebra* (hereafter: EJA) is a finite-dimensional commutative real algebra (A, \bullet) with a multiplicative unit element u , satisfying the *Jordan identity*

$$a^2 \bullet (a \bullet b) = a \bullet (a^2 \bullet b)$$

for all $a, b \in A$, and equipped with an inner product satisfying

$$\langle a \bullet b | c \rangle = \langle b | a \bullet c \rangle$$

for all $a, b, c \in A$. The basic example is the self-adjoint part M_{sa} of a real, complex or quaternionic matrix algebra M , with $a \bullet b = (ab + ba)/2$ and with $\langle a | b \rangle = \text{tr}(ab)$. Any Jordan subalgebra of an EJA is also an EJA. So, too, is the *spin factor* $V_n = \mathbb{R} \times \mathbb{R}^n$, with the obvious inner product and with

$$(t, x) \bullet (s, y) = (ts + \langle x | y \rangle, ty + sx) :$$

this can be embedded (universally) in $M_{2k}(\mathbb{C})_{\text{sa}}$ for $n = 2k$ and $M_{2k}(\mathbb{C})_{\text{sa}} \oplus M_{2k}(\mathbb{C})_{\text{sa}}$ for $n = 2k + 1$. Moreover, one can show that

$$V_2 \simeq M_2(\mathbb{R})_{\text{sa}}, V_3 \cong M_2(\mathbb{C})_{\text{sa}}, \text{ and } V_5 \simeq M_2(\mathbb{H})_{\text{sa}}.$$

Classification Direct sums of EJAs are also EJAs, so we can obtain more examples by forming direct sums of the EJAs of the types mentioned above. The *Jordan-von Neumann-Wigner Classification Theorem* (see [8] Chapter IV) provides a converse: every euclidean Jordan algebra is a direct sum of *simple* EJAs, each of which is isomorphic to a spin factor V_n , or to the self-adjoint part of a matrix algebra $M_n(\mathbb{K})$ where \mathbb{K} is one of the classical division rings \mathbb{R}, \mathbb{C} or \mathbb{H} , or, if $n = 3$, to the *Octonions*, \mathbb{O} . This last example, which is not embeddable into the self-adjoint part of a complex matrix algebra, is called the *exceptional Jordan algebra*, or the *Albert algebra*. A Jordan algebra that is embeddable in $M_n(\mathbb{C})_{\text{sa}}$ for some n , is said to be *special*. It follows from the classification theorem that any EJA decomposes as a direct sum $A_{\text{sp}} \oplus A_{\text{ex}}$ where A_{sp} is special and A_{ex} is a direct sum of copies of the exceptional Jordan algebra.

Projections and the Spectral Theorem A *projection* in an EJA A is an element $p \in A$ with $p^2 = p$. If p, q are projections with $p \bullet q = 0$, we say that p and q are orthogonal. In this case, $p + q$ is another projection. A projection not representable as a sum of other projections is said to be *minimal* or *primitive*. A *Jordan frame* is a set $E \subseteq A$ of pairwise orthogonal minimal projections that sum to the Jordan unit. The *Spectral Theorem* (cf. e.g. [8], Theorem III.1.1) for EJAs asserts that every element $a \in A$ can be expanded as a linear combination $a = \sum_{x \in E} t_x x$ where E is some Jordan frame.

One can show that all Jordan frames for a given Euclidean Jordan algebra A have the same number of elements. This number is called the *rank* of A . By the Classification Theorem, all simple Jordan algebras having rank 4 or higher are special.

Order Structure Any EJA A is at the same time an ordered real vector space, with positive cone $A_+ = \{a^2 | a \in A\}$; for $a, b \in A$, $a \leq b$ iff $b - a \in A_+$. This allows us to interpret A as a probabilistic model: an *effect* (measurement-outcome) in A is an element $a \in A_+$ with $a \leq u$. A *state* on A is a positive linear mapping $\alpha : A \rightarrow \mathbb{R}$ with $\alpha(u) = 1$. If a is an effect, we interpret $\alpha(a)$ as the probability that a will be observed (if tested) in the state α .

The cone A_+ is *self-dual* with respect to the given inner product on A : an element $a \in A$ belongs to A_+ iff $\langle a | b \rangle \geq 0$ for all $b \in A_+$. Every state α then corresponds to a unique $b \in A_+$ with $\alpha(a) = \langle a | b \rangle$.

Remark: Besides being self-dual, the cone A_+ is *homogeneous*: any element of the *interior* of A_+ can be obtained from any other by an order-automorphism of A , that is, a linear automorphism $\phi : A \rightarrow A$ with $\phi(A_+) = A_+$. The *Koecher-Vinberg Theorem* ([11, 15]; see [8] for a relatively accessible proof based on the one in [13]) identifies EJAs as precisely the finite-dimensional ordered linear spaces having homogeneous, self-dual positive cones. This fact underwrites the derivations in several of the papers cited above [17, 18, 16].²

Reversible and universally reversible EJAs A Jordan subalgebra of M_{sa} , where M is a complex $*$ -algebra, is *reversible* iff

$$a_1, \dots, a_k \in A \Rightarrow a_1 a_2 \cdots a_k + a_k \cdots a_2 a_1 \in A,$$

²A different characterization of EJAs, in terms of projections associated with faces of the state space, is invoked in [5].

where juxtaposition indicates multiplication in M . Note that with $k = 2$, this is just closure under the Jordan product on M_{sa} . An abstract EJA A is *reversible* iff it has a representation as a reversible Jordan subalgebra of some complex $*$ -algebra. A reversible EJA is *universally reversible* (UR) iff it has *only* reversible representations.

Universal reversibility will play a large role in what follows. Of the four basic types of special Euclidean Jordan algebra considered above, the only ones that are not UR are the spin factors V_k with $k \geq 4$. For $k = 4$ and $k > 5$, V_k is not even reversible; V_5 — equivalently, $M_2(\mathbb{H})_{\text{sa}}$ — has a reversible representation, but also non-reversible ones. Thus, if we adopt the shorthand

$$R_n = M_n(\mathbb{R})_{\text{sa}}, \quad C_n = M_n(\mathbb{C})_{\text{sa}}, \quad \text{and} \quad Q_n = M_n(\mathbb{H})_{\text{sa}},$$

we have R_n, C_n UR for all n , and Q_n UR for $n > 2$.

3 Composites of EJAs

A probabilistic theory must allow for some device for describing composite systems. Given EJAs A and B , understood as models for two physical systems, we'd like to construct an EJA AB that models the two systems considered together as a single entity. Is there any satisfactory way to do this? If so, how much latitude does one have?

A good candidate for such an EJA is provided by a construction due to Hanche-Olsen [9], which we now review. In the sequel, we show that, at least if no non-UR spin factors are involved, it does provide such a model.

The universal tensor product A *representation* of a Jordan algebra A is a Jordan homomorphism $\pi : A \rightarrow M_{\text{sa}}$, where \mathbf{M} is a complex $*$ -algebra. For any EJA A , there exists a (possibly trivial) $*$ -algebra $C^*(A)$ and a representation $\psi_A : A \rightarrow C^*(A)_{\text{sa}}$ with the universal property that any representation $\pi : A \rightarrow M_{\text{sa}}$, where \mathbf{M} is a C^* -algebra, decomposes uniquely as $\pi = \tilde{\pi} \circ \psi_A$, $\tilde{\pi} : C^*(A) \rightarrow \mathbf{M}$ a $*$ -homomorphism. Evidently, $(C^*(A), \psi_A)$ is unique up to a canonical $*$ -isomorphism. Since $\psi^{\text{op}} : A \rightarrow C^*(A)^{\text{op}}$ provides another solution to the same universal problem, there exists a canonical anti-automorphism Φ_A on $C^*(A)$, fixing every point of $\psi_A(A)$.

We refer to $(C^*(A), \psi_A)$ as the *universal representation* of A . A is exceptional iff $C^*(A) = \{0\}$. If A has no exceptional factors, then ψ_A is injective. In this case, we will routinely identify A with its image $\psi_A(A) \leq C^*(A)$.

In [9], Hanche-Olsen defines the *universal tensor product* of two special EJAs A and B to be the Jordan subalgebra of $C^*(A) \otimes C^*(B)$ generated by $A \otimes B$. This is denoted $A \tilde{\otimes} B$. It can be shown that

$$C^*(A \tilde{\otimes} B) = C^*(A) \tilde{\otimes} C^*(B) \quad \text{and} \quad \Phi_{A \tilde{\otimes} B} = \Phi_A \otimes \Phi_B.$$

Some further important facts about the universal tensor product are the following:

Proposition 1 Let A, B and C denote EJAs.

- (a) If $\phi : A \rightarrow C$, $\psi : B \rightarrow C$ are unital Jordan homomorphisms with operator-commuting ranges³, then there exists a unique Jordan homomorphism $A \tilde{\otimes} B \rightarrow C$ taking $a \otimes b$ to $\phi(a) \bullet \psi(b)$ for all $a \in A$, $b \in B$.

³Elements $x, y \in C$ operator commute iff $x \bullet (y \bullet z) = y \bullet (x \bullet z)$ for all $z \in C$.

- (b) $A \tilde{\otimes} B$ is UR unless one of the factors has a one-dimensional summand and the other has a representation onto a spin factor V_n with $n \geq 4$.
- (c) If A is UR, then $A \tilde{\otimes} M_n(\mathbb{C})_{\text{sa}} = (C^*(A) \otimes M_n(\mathbb{C}))_{\text{sa}}$.

These are Propositions 5.2, 5.3 and 5.4, respectively, in [9].

Note that part (b) implies that if A and B are irreducible and non-trivial, $A \tilde{\otimes} B$ will always be UR, hence, the fixed-point set of $\Phi_A \otimes \Phi_B$. Using this one can compute $A \tilde{\otimes} B$ for irreducible, universally reversible A and B [9]. Below, and for the balance of this paper, we use the shorthand $R_n := M_n(\mathbb{R})_{\text{sa}}$, $C_n = M_n(\mathbb{C})_{\text{sa}}$ and $Q_n = M_n(\mathbb{H})_{\text{sa}}$ (noting that Q_n is UR only for $n > 2$):

$\tilde{\otimes}$	R_m	C_m	Q_m
R_n	R_{nm}	C_{nm}	Q_{nm}
C_n	C_{nm}	$C_{nm} \oplus C_{nm}$	C_{2nm}
Q_n	Q_{nm}	C_{2nm}	R_{4nm}

Figure 2

For $Q_2 \tilde{\otimes} Q_2$, a bit more work is required, but one can show that $Q_2 \tilde{\otimes} Q_2$ is the direct sum of four copies of $R_{16} = M_{16}(\mathbb{R})_{\text{sa}}$ (stated in [9]; details will be provided in [4]).

General composites of EJAs The universal tensor product is an instance of the following (as it proves, only slightly) more general scheme. Recall that an order-automorphism of an EJA A is a linear bijection $\phi : A \rightarrow A$ taking A_+ onto itself. These form a Lie group, whose identity component we denote by $G(A)$.

Definition 1: A **composite** of EJAs A and B is a pair (AB, π) where AB is an EJA and $\pi : A \otimes B \rightarrow AB$ is a linear mapping such that

- (a) If $a \in A_+$ and $b \in B_+$, then $\pi(a \otimes b) \in (AB)_+$, with $\pi(u \otimes u)$ the Jordan unit of AB ;
- (b) for all states α on A , β on B , there exists a state γ on AB such that $\gamma(\pi(a \otimes b)) = \alpha(a)\beta(b)$;
- (c) for all automorphisms $\phi \in G(A)$ and $\psi \in G(B)$, there exists a preferred automorphism $\phi \otimes \psi \in G(AB)$ with $(\phi \otimes \psi)(\pi(a \otimes b)) = \pi(\phi(a) \otimes \psi(b))$. Moreover, we require that

$$(\phi_1 \otimes \psi_1) \circ (\phi_2 \otimes \psi_2) = (\phi_1 \circ \phi_2) \otimes (\psi_1 \circ \psi_2)$$

and

$$(\phi \otimes \psi)^\dagger = \phi^\dagger \otimes \psi^\dagger$$

for all $\phi, \phi_i \in G(A)$ and $\psi, \psi_i \in G(B)$.

It follows from (b) that π is injective (if $\pi(T) = 0$, then for any states α, β , there's a state γ of AB with $(\alpha \otimes \beta)(T) = \gamma(\pi(T)) = 0$; it follows that $T = 0$). Henceforth, we'll simply regard $A \otimes B$ as a subspace of AB .

Condition (c) calls for further comment. The dynamics of a physical system modeled by a Euclidean Jordan algebra A will naturally be represented by a one-parameter group $t \mapsto \phi_t$ of order automorphisms of A . As order-automorphisms in $G(A)$ are precisely the elements of such one-parameter groups, condition (c) is equivalent to the condition that, given dynamics $t \mapsto \phi_t$ and $t \mapsto \psi_t$ on A and B , respectively, there is a preferred dynamics on AB under which pure tensors $a \otimes b$ evolve according to $a \otimes b \mapsto \phi_t(a) \otimes \psi_t(b)$. In other words, there is a dynamics on AB under which A and B evolve independently.

Theorem 1: *If A and B are simple EJAs, then any composite AB is special, and an ideal in $A \widetilde{\otimes} B$.*

The basic idea of the proof is to show that if p_1, \dots, p_n is a Jordan frame in an irreducible summand of A , and q_1, \dots, q_m is a Jordan frame in an irreducible summand of B , then $\{p_i \otimes q_j | i = 1, \dots, n, j = 1, \dots, m\}$ is a pairwise orthogonal set of projections in AB , whence, the latter has rank at least four, and must therefore be special. For the details, we refer to [4].

Corollary 1: *If A is simple and B is exceptional, then no composite AB exists.*

In particular, if B is the exceptional factor, there exists no composite of B with itself.

Corollary 2: *If $A \widetilde{\otimes} B$ is simple, then $AB = A \widetilde{\otimes} B$ is the only possible composite of A and B .*

There are cases in which $A \widetilde{\otimes} B$ isn't simple, even where A and B are: namely, the cases in which A and B are both hermitian parts of complex matrix algebras. From table (2), we see that if $A = C_n$ and $B = C_m$, then $A \widetilde{\otimes} B = C_{nm} \oplus C_{nm}$. In this case, Proposition 1 gives us *two* choices for AB : either the entire direct sum above, or one of its isomorphic summands, i.e., the ‘‘obvious’’ composite $AB = C_{nm}$.

4 Embedded EJAs

Corollary 1 above justifies restricting our attention to special EJAs. These are precisely the finite dimensional JC-algebras, *i.e.* the ones isometrically isomorphic to norm closed Jordan subalgebras of self-adjoint linear operators on complex Hilbert spaces [9][10]. In fact, it will be helpful to consider *embedded* JC-algebras, that is, Jordan subalgebras of specified finite-dimensional C^* algebras.

Definition 2: An *embedded JC algebra*, or EJC, is a pair (A, \mathbf{M}_A) where A is a unital Jordan subalgebra of a unital finite-dimensional complex $*$ -algebra M_A (that is, a Jordan subalgebra such that the Jordan unit u_A coincides with the unit of M_A).

The notation \mathbf{M}_A is intended to emphasise that the embedding $A \mapsto \mathbf{M}_A$ is part of the structure of interest. Given A , there is always a canonical choice for \mathbf{M}_A , namely the universal enveloping $*$ -algebra $C^*(A)$ of A [9].

Definition 3: The *canonical product* of EJC's (A, \mathbf{M}_A) and (B, \mathbf{M}_B) is the pair $(A \odot B, \mathbf{M}_A \otimes \mathbf{M}_B)$ where $A \odot B$ is the Jordan subalgebra of $(\mathbf{M}_A \otimes \mathbf{M}_B)_{\text{sa}}$ generated by the subspace $A \otimes B$.

Note that, as a matter of definition, $\mathbf{M}_{A \odot B} = \mathbf{M}_A \otimes \mathbf{M}_B$. If $\mathbf{M}_A = C^*(A)$ and $\mathbf{M}_B = C^*(B)$, then $A \odot B$ is the Hanche-Olsen tensor product.

One would like to know that $A \odot B$ is in fact a composite of A and B in the sense of Definition 1. Using a result of Upmeyer [14], we can show that this is the case for *reversible* EJAs A and B . (that is, real, complex and quaternionic systems, and direct sums of these). Whether $A \odot B$ is a composite in the sense of Definition 1 when A or B is non-reversible spin factor remains an open question.

We can now form a category:

Definition 4: **EJC** is the category consisting of EJC's (A, \mathbf{M}_A) and completely positive (CP) maps $\phi : \mathbf{M}_A \rightarrow \mathbf{M}_B$ with $\phi(A) \subseteq B$. We refer to such maps as *Jordan preserving*.

Proposition 2: *The canonical product \odot is associative on **EJC**. More precisely, the associator mapping*

$$\alpha : \mathbf{M}_A \otimes (\mathbf{M}_B \otimes \mathbf{M}_C) \rightarrow (\mathbf{M}_A \otimes \mathbf{M}_B) \otimes \mathbf{M}_C$$

takes $A \odot (B \odot C)$ to $(A \odot B) \odot C$.

(Note that since the associator mapping is CP, this means that α is a morphism in **EJC**.) The proof is somewhat lengthy, so we refer the reader to the forthcoming paper [4].

Proposition 2 suggests that **EJC** might be symmetric monoidal under \odot . There is certainly a natural choice for the monoidal unit, namely $I = (\mathbb{R}, \mathbb{C})$. But the following example shows that tensor products of **EJC** morphisms are generally not morphisms:

Example 1: Let $(A, C^*(A))$ and $(B, C^*(B))$ be simple, universally embedded EJCs, and suppose that B is not UR (e.g., a spin factor V_n with $n > 3$). Let \widehat{B} be the set of fixed points of the canonical involution Φ_B . Then by Corollary 2, $A \odot B = A \widehat{\otimes} B$, the set of fixed points of $\Phi_A \otimes \Phi_B$. In particular, $u_A \otimes \widehat{B}$ is contained in $A \odot B$. Now let f be a state on $C^*(A)$: this is CP, and trivially Jordan-preserving, and so, a morphism in **EJC**. But

$$(f \otimes \text{id}_B)(u_A \otimes \widehat{B}) = f(u_A)\widehat{B} = \widehat{B},$$

which isn't contained in B . So $f \otimes \text{id}_B$ isn't Jordan-preserving.

5 Reversible and universally reversible EJCs

It seems that the category **EJC** is simply too large. We can try to obtain a better-behaved category by restricting the set of morphisms, or by restricting the set of objects, or both.

As a first pass, we might try this:

Definition 5: Let (A, \mathbf{M}_A) and (B, \mathbf{M}_B) be EJCs. A CP map $\phi : \mathbf{M}_A \rightarrow \mathbf{M}_B$ is *completely Jordan preserving* (CJP) iff $\phi \otimes 1_C$ takes $A \odot C$ to $B \odot C$ for all (C, \mathbf{M}_C) .

It is not hard to verify the following

Proposition 3: If $\phi : \mathbf{M}_A \rightarrow \mathbf{M}_B$ and $\psi : \mathbf{M}_C \rightarrow \mathbf{M}_D$ are CJP, then so is

$$\phi \otimes \psi : \mathbf{M}_{A \odot C} = \mathbf{M}_A \otimes \mathbf{M}_B \rightarrow \mathbf{M}_B \otimes \mathbf{M}_C = \mathbf{M}_{B \odot D}.$$

Thus, the category of EJC algebras and CJP mappings is symmetric monoidal.⁴

There are many examples of CJP mappings: Jordan homomorphisms are CJP maps. If $a \in A$, the mapping

$$U_a : A \rightarrow A$$

given by $U_a = 2L_{a^2} - L_a^2$, where $L_a(b) = ab$, is also CJP. On the other hand, by Example 1 above, **CJP**(A, I) is *empty* for universally embedded simple A !

So not all CP maps are CJP; for instance, states are never CJP. More seriously, we can't interpret states as morphisms in this category. The problem is the non-UR spin factors in **CJP**. If we remove these, things are much better.

Definition 6: Let \mathcal{C} be a subclass of embedded EJC algebras, closed under \odot and containing I . A linear mapping $\phi : \mathbf{M}_A \rightarrow \mathbf{M}_B$ is *CJP relative to \mathcal{C}* iff $\phi_A \otimes \text{id}_C$ is Jordan preserving for all C in \mathcal{C} . **CJP** $_{\mathcal{C}}$ is the

⁴Notice that scalars of this category are real numbers. It is sometimes suggested that quaternionic Hilbert spaces can't be accommodated in a symmetric monoidal category owing to the noncommutativity of \mathbb{H} , as the scalars in a symmetric monoidal category must always be commutative. As we are representing quaternionic quantum systems in terms of the associated real vector spaces of hermitian operators, this issue doesn't arise here.

category having objects elements of \mathcal{C} , mappings relatively CJP mappings.

Example 2: **URUE** is the class of universally reversible, universally embedded EJC algebras. **URSE** is the category of universally reversible, standardly embedded EJC algebras, and **RSE** is the category of reversible, standardly embedded EJC algebras. Equipped with relatively CJP mappings, both are symmetric monoidal categories.

Note that **RSE** consists of direct sums of real, complex and quaternionic quantum systems. **URSE** and **URUE** contain all real and complex quantum systems, and all quaternionic quantum systems *except* the “quabit”, i.e., the quaternionic bit $Q_2 := M_2(\mathbb{H})_{\text{sa}}$.

In both of the categories **URUE** and **URSE**, states are morphisms. However, Example 1 provides us with a no-go result for extensions of these categories that preserve the property that states are morphisms. For **URUE**, we cannot maintain this property while enlarging the class \mathcal{C} to include a spin factor other than the real, complex, and quaternionic bits, while for **URSE**, we cannot enlarge it to contain an even-dimensional one (it remains open whether we can include the additional odd-dimensional ones).

Not only do **URUE** and **URSE** include all states as morphisms, but we are going to see that they inherit compact closure from the category ***-ALG** of finite-dimensional, complex $*$ -algebras and CP maps, in which they are embedded. It’s worth taking a moment to review this compact structure. If \mathbf{M} is a finite-dimensional complex $*$ -algebra, let Tr denote the canonical trace on \mathbf{M} , regarded as acting on itself by left multiplication (so that $\text{Tr}(a) = \text{tr}(L_a)$, $L_a : \mathbf{M} \rightarrow \mathbf{M}$ being $L_a(b) = ab$ for all $b \in \mathbf{M}$). This induces an inner product on \mathbf{M} , given by $\langle a|b \rangle_{\mathbf{M}} = \text{Tr}(ab^*)^5$. Note that this inner product is self-dualizing, i.e., $a \in \mathbf{M}_+$ iff $\langle a|b \rangle \geq 0$ for all $b \in \mathbf{M}_+$. Now let $\overline{\mathbf{M}}$ be the conjugate algebra, writing \bar{a} for $a \in \mathbf{M}$ when regarded as belonging to $\overline{\mathbf{M}}$ (so that $\overline{\bar{a}} = a$ for scalars $a \in \mathbb{C}$ and $\overline{ab} = \bar{b}\bar{a}$ for $a, b \in \mathbf{M}$). Note that $\langle \bar{a}|\bar{b} \rangle = \langle b|a \rangle$. Now define

$$f_{\mathbf{M}} = \sum_{e \in E} e \otimes \bar{e} \in \mathbf{M} \otimes \overline{\mathbf{M}}$$

where E is any orthonormal basis for \mathbf{M} with respect to $\langle | \rangle_{\mathbf{M}}$. Then a computation shows that

$$\langle (a \otimes \bar{b}) f_{\mathbf{M}} | f_{\mathbf{M}} \rangle_{\mathbf{M} \otimes \overline{\mathbf{M}}} = \langle a|b \rangle_{\mathbf{M}}.$$

Since the left-hand side defines a positive linear functional on $\mathbf{M} \otimes \overline{\mathbf{M}}$, so does the right (remembering here that pure tensors generate $\mathbf{M} \otimes \overline{\mathbf{M}}$, as we’re working in finite dimensions). Call this functional $\eta_{\mathbf{M}}$. That is,

$$\eta_{\mathbf{M}} : \mathbf{M} \otimes \overline{\mathbf{M}} \rightarrow \mathbb{C} \text{ is given by } \eta_{\mathbf{M}}(a \otimes \bar{b}) = \langle a|b \rangle = \text{Tr}(ab^*)$$

and is, up to normalization, a state on $\mathbf{M} \otimes \overline{\mathbf{M}}$. A further computation now shows that

$$\langle a \otimes \bar{b} | f_{\mathbf{M}} \rangle_{\mathbf{M} \otimes \overline{\mathbf{M}}} = \eta(a \otimes \bar{b}).$$

It follows that $f_{\mathbf{M}}$ belongs to the positive cone of $\mathbf{M} \otimes \overline{\mathbf{M}}$, by self-duality of the latter. A final computation shows that, for any states α and $\bar{\alpha}$ on \mathbf{M} and $\overline{\mathbf{M}}$, respectively, and any $a \in \mathbf{M}, \bar{a} \in \overline{\mathbf{M}}$, we have

$$(\eta_{\mathbf{M}} \otimes \alpha)(a \otimes f_{\overline{\mathbf{M}}}) = \alpha(a) \text{ and } (\bar{\alpha} \otimes \eta_{\mathbf{M}})(f_{\overline{\mathbf{M}}} \otimes \bar{a}) = \bar{\alpha}(\bar{a}).$$

Thus, $\eta_{\mathbf{M}}$ and $f_{\overline{\mathbf{M}}}$ define a compact structure on ***-ALG**, for which the dual object of \mathbf{M} is given by $\overline{\mathbf{M}}$.

Definition 7: The *conjugate* of a EJC algebra (A, \mathbf{M}_A) is $(\bar{A}, \overline{\mathbf{M}}_A)$, where $\bar{A} = \{\bar{a} | a \in A\}$. We write η_A for $\eta_{\mathbf{M}_A}$ and f_A for $f_{\mathbf{M}_A}$.

⁵We are following the convention that complex inner products are conjugate linear in the *second* argument.

5.1 Universally-embedded, universally reversible EJAs

Now consider the category **URUE** of universally reversible, universally embedded EJAs A , i.e., pairs (A, \mathbf{M}_A) with A UR and $\mathbf{M}_A = C^*(A)$. Let Φ_A be the canonical involution on $C^*(A)$.

Lemma 1: *Let (A, \mathbf{M}_A) belong to **URUE**. Then*

- (a) $f_A \in A \odot \bar{A}$;
- (b) $\eta_A \circ (\Phi_A \otimes \Phi_{\bar{A}}) = \eta_A$.

Proof: (a) follows from the fact that Φ_A is unitary, so that if E is an orthonormal basis, then so is $\{\Phi_A(e) | e \in E\}$. Since f_A is independent of the choice of orthonormal basis, it follows that f is invariant under $\Phi_A \otimes \Phi_{\bar{A}}$, hence, an element of $A \otimes \bar{A}$. Now (b) follows from part (a) of the previous lemma. \square

Define $\gamma_A : C^*(\bar{A}) \rightarrow C^*(A)$ by $\gamma(\bar{a}) = \Phi_A(a^*)$. This is a $*$ -isomorphism, and intertwines Φ_A and $\Phi_{\bar{A}}$; hence, $\gamma_A \otimes \text{id}_B : C^*(\bar{A} \otimes B) \rightarrow C^*(A \otimes B)$ intertwines $\Phi_{\bar{A}} \otimes \Phi_B = \Phi_{\bar{A} \otimes B}$ and $\Phi_A \otimes \Phi_B = \Phi_{A \otimes B}$ — whence, takes $\bar{A} \otimes B$ to $A \otimes B$. In particular, γ_A is CJP relative to the class of UR, universally embedded EJC.

Lemma 2: *Let A be a universally embedded UR EJC. Then for all $\alpha \in \mathbf{CJP}(A, I)$, there exists $a \in A$ with $\alpha(b) = \langle b | a \rangle$ for all $b \in A$.*

Proof: Since $\alpha \in C^*(A)^*$, there is certainly some $a \in C^*(A)$ with $\alpha = |a\rangle$. We need to show that $a \in A$. Since α is CJP,

$$\gamma_A \otimes \alpha : C^*(\bar{A}) \otimes C^*(A) = C^*(\bar{A} \otimes A) \rightarrow C^*(A)$$

is Jordan-preserving. In particular, $(\alpha \otimes \gamma_A)(f_A) \in A$. But

$$\begin{aligned} (\alpha \otimes \gamma_A)(f_A) &= \sum_{e \in E} (\alpha \otimes \gamma_A)(e \otimes \bar{e}) \\ &= \sum_{e \in E} \langle e | a \rangle \Phi(e^*) \\ &= \Phi\left(\sum_{e \in E} \langle e | a \rangle e^*\right) \\ &= \Phi\left(\left(\sum_{e \in E} \langle a | e \rangle e\right)^*\right) = \Phi(a^*) = \gamma_A(\bar{a}). \end{aligned}$$

Hence, $\gamma_A(\bar{a}) \in A$, whence, $\bar{a} \in \bar{A}$, whence, $a \in A$. (Alternatively: $\Phi_A(a^*) \in A$ implies $a^* \in A$, whence, a is self-adjoint, whence, $a \in A$.) \square

It follows that η_A and f_A belong, as morphisms, to **URUE**. Hence, **URUE** inherits the compact structure from ***-ALG**, as promised.

The same holds for **URSE**. Specifically, we want to show that f_A belongs to $A \odot \bar{A}$ whenever A is a standardly embedded UR EJC.

Suppose that E is an orthonormal basis for \mathbf{M}_A : then so is $\{e^* | e \in E\}$; thus, since f_A is independent of the choice of basis, we have

$$f_A^* = \sum_{e \in E} e^* \otimes \bar{e}^* = \sum_{e^* \in E^*} e^* \otimes \bar{e}^* = f_A.$$

Thus, if $A \odot \hat{A}$ is the self-adjoint part of $\mathbf{M}_A \otimes \bar{\mathbf{M}}_A$, then $f_A \in A \otimes \bar{A}$. This covers the case where $A = C_n$. We also have, by the results above, that $f_A \in A \odot \bar{A}$ whenever the latter equals $A \otimes \bar{A}$. This covers $A = R_n$ and $A = Q_n$ for $n > 2$.

In fact, we can do a bit better. If \mathbf{M} and \mathbf{N} are finite-dimensional $*$ -algebras and $\phi : \mathbf{M} \rightarrow \mathbf{N}$ is a linear mapping, let ϕ^\dagger denote the adjoint of ϕ with respect to the natural trace inner products on \mathbf{M} and \mathbf{N} . It is not difficult to show that, for any \mathbf{M} in $*$ -ALG, $f_{\mathbf{M}}^\dagger = \eta_{\mathbf{M}}$ and vice versa; indeed, $*$ -ALG is dagger-compact.

Definition 8: Let (A, \mathbf{M}_A) and (B, \mathbf{M}_B) be EJC. A linear mapping $\phi : \mathbf{M}_A \rightarrow \mathbf{M}_B$ is \dagger -CJP iff both ϕ and ϕ^\dagger are CJP. If \mathcal{C} is a category of EJC and CJP mappings, we write \mathcal{C}^\dagger for the category having the same objects, but with morphisms restricted to \dagger -CJP mappings in \mathcal{C} .

If A belongs to **URUE** or **URSE**, then f_A and η_A are both CJP and, hence, are both \dagger -CJP with respect to the indicated category. Hence,

Theorem 2: *The categories \mathbf{URUE}^\dagger and \mathbf{URSE}^\dagger are dagger-compact.*

6 Conclusion

We have found two theories — the categories **URSE** and **URUE** of universally reversible euclidean Jordan algebras, standardly and universally embedded — that, in slightly different ways, unify finite-dimensional real, complex and (almost all of) quaternionic quantum mechanics. By virtue of being compact closed, both theories continue to enjoy many of the information-processing properties of standard complex QM, e.g., the existence of conclusive teleportation and entanglement-swapping protocols [1].

One property of standard QM that some of our composites do *not* share is the possibility of *local tomography*: a state ω on $A \odot B$ is not generally determined by the joint probability assignment $a, b \mapsto \omega(a \otimes b)$, where a and b are effects of A and B , respectively. Another way to put it is that $A \odot B$ is generally much larger than the vector-space tensor product $A \otimes B$. As local tomography is well known to separate complex QM from its real and quaternionic variants, this is hardly surprising. Perhaps more surprising is that some of our composites—those involving quaternionic factors—also exhibit supermultiplicativity of the maximal number of distinguishable states, and, relatedly, violate another property sometimes assumed as a basic desideratum for composites: that products of pure states are pure.

Neither theory includes the qubit, Q_2 . Example 1 shows that the qubit can't be added to **URUE** without a violation of compact closure. On the other hand, if f_{Q_2} belongs to the canonical composite $Q_2 \odot Q_2$, then the slightly larger category **RSE**, which consists of *all* finite-dimensional real, complex and quaternionic quantum systems, will be compact closed (indeed, dagger compact).⁶

The categories **URSE** and **URUE** contain interesting compact closed subcategories. In particular, real and quaternionic quantum systems (less the qubit), taken together, form a sub-theory, closed under composites. We *conjecture* that this is exactly what one gets by applying the CPM construction to Baez's (implicit) category of pairs (\mathbf{H}, J) , \mathbf{H} a finite-dimensional Hilbert space and J an anti-unitary with $J^2 = \pm 1$ — and, again, excluding the qubit. Should **RSE** prove to be compact closed, we could entertain the stronger conjecture that this is exactly what one obtains by applying CPM to Baez's category.

Complex quantum systems also form a monoidal subcategory of **URSE**, which we might call **CQM**: indeed, one that functions as an “ideal”, in that if $A \in \mathbf{URSE}$ and $B \in \mathbf{CQM}$, then $A \odot B \in \mathbf{CQM}$ as well. This is provocative, as it suggests that a universe initially consisting of many systems of all three types,

⁶Since this note was accepted for presentation at QPL XII we have settled this question in the affirmative. The details will be given in [4].

would eventually evolve into one in which complex systems greatly predominate.

The category **URUE** is somewhat mysterious. Like **URSE**, this encompasses real, complex and quaternionic quantum systems, excepting the qubit. In this theory, the composite of *complex* quantum systems comes with an extra classical bit — equivalently, a $\{0, 1\}$ -valued superselection rule. This functions to make the transpose operation — which is a Jordan automorphism of $\mathbf{M}_n(\mathbb{C})_{\text{sa}}$, but an antiautomorphism of $\mathbf{M}_n(\mathbb{C})$ — count as a morphism. The precise physical significance of this is a subject for further study.

Acknowledgements

We thank Cozmin Ududec for valuable discussions. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research & Innovation. MAG was supported by an NSERC Alexander Graham Bell Canada Graduate Scholarship. AW was supported by a grant from the FQXi foundation.

References

- [1] S. Abramsky & B. Coecke (2005): *Abstract Physical Traces. Theory and Applications of Categories* 14, pp. 111–124. Available at <http://arxiv.org/abs/0910.3144>.
- [2] E. M. Alfsen & F. W. Shultz (2003): *Geometry of state spaces of operator algebras*. Birkhäuser, doi:10.1007/978-1-4612-0019-2.
- [3] J. Baez (2012): *Division algebras and quantum theory. Foundations of Physics* 42, pp. 819–855, doi:10.1007/s10701-011-9566-z. Available at <http://arxiv.org/abs/1101.5690>.
- [4] H. Barnum, M. A. Graydon & A. Wilce (in preparation): *Composites and categories of Euclidean Jordan algebras*.
- [5] H. Barnum, M. Müller & C. Ududec (2014): *Higher-order interference and single-system postulates characterizing quantum theory. New Journal of Physics* 16, p. 123029, doi:10.1088/1367-2630/16/12/123029. Available at <http://arxiv.org/abs/1403.4147>.
- [6] H. Barnum & A. Wilce (2011): *Information processing in convex operational theories. Electronic Notes in Theoretical Computer Science* 270, pp. 3–15, doi:10.1016/j.entcs.2011.01.002. Available at <http://arxiv.org/abs/0908.2352>.
- [7] H. Barnum & A. Wilce (2014): *Local tomography and the Jordan structure of quantum theory. Foundations of Physics* 44, pp. 192–212, doi:10.1007/s10701-014-9777-1. Available at <http://arxiv.org/abs/1202.4513v2>.
- [8] J. Faraut & A. Koranyi (1994): *Analysis on Symmetric Cones*. Oxford.
- [9] H. Hanche-Olsen (1983): *On the structure and tensor products of JC-algebras. Can. J. Math.* 35, pp. 1059–1074, doi:10.4153/CJM-1983-059-8.
- [10] H. Hanche-Olsen & E. Størmer (1984): *Jordan Operator Algebras*. Pitman, Boston-London-Melbourne. Available at <http://www.math.ntnu.no/~hanche/joa/>.
- [11] M. Koecher (1958): *Die geodätischen von positivitätsbereichen. Math. Annalen* 135, pp. 192–202, doi:10.1007/BF01351796. See also: M. Koecher (1999): *The Minnesota Notes on Jordan Algebras and Their Applications*. Springer.
- [12] M. Müller & C. Ududec (2012): *The structure of reversible computation determines the self-duality of quantum theory. Phys. Rev. Lett.* 108, p. 130401, doi:10.1103/PhysRevLett.108.130401. Available at <http://arxiv.org/abs/1110.3516>.

- [13] I. Satake (1980): *Algebraic Structures of Symmetric Domains*. Publications of the Mathematical Society of Japan, No. 14, Iwanami Shoten and Princeton University Press.
- [14] H. Upmeyer (1980): *Derivations of Jordan C^* -algebras*. *Math. Scand.* 46, pp. 251–264.
- [15] E. Vinberg (1960): *Homogeneous cones*. *Dokl. Acad. Nauk. SSSR* 133, pp. 9–12. English translation: E. Vinberg (1961): *Soviet Math. Dokl.* 1, pp. 787-790.
- [16] A. Wilce (2012): *Conjugates, Filters and Quantum Mechanics*. Available at <http://arxiv.org/abs/1206.2897>.
- [17] A. Wilce (2012): *Four and a half axioms for quantum mechanics*. In Y. Ben-Menahem & M. Hemmo, editors: *Probability in Physics*, Springer, doi:10.1007/978-3-642-21329-8_17. Available at <http://arxiv.org/abs/0912.5530>.
- [18] A. Wilce (2012): *Symmetry and self-duality in categories of probabilistic models*. In B. Jacobs, P. Selinger & B. Spitters, editors: *Proceedings of QPL 2011, Series 95*, pp. 275–279, doi:10.4204/EPTCS.95. Available at <http://arxiv.org/abs/1210.0622>.